

# BRAIN COMMUNICATIONS

## The cognitive consequences of the COVID-19 epidemic: collateral damage?

Karen Ritchie,<sup>1,2</sup> Dennis Chan<sup>3</sup> and Tam Watermeyer<sup>2,4</sup>

See Lleó & Alcolea (<https://doi.org/10.1093/braincomms/fcaa072>) for a scientific commentary on this article.

Recovery from coronavirus disease 2019 (COVID-19) will be principally defined in terms of remission from respiratory symptoms; however, both clinical and animal studies have shown that coronaviruses may spread to the nervous system. A systematic search on previous viral epidemics revealed that while there has been relatively little research in this area, clinical studies have commonly reported neurological disorders and cognitive difficulties. Little is known with regard to their incidence, duration or underlying neural basis. The hippocampus appears to be particularly vulnerable to coronavirus infections, thus increasing the probability of post-infection memory impairment, and acceleration of neurodegenerative disorders such as Alzheimer's disease. Future knowledge of the impact of COVID-19, from epidemiological studies and clinical practice, will be needed to develop future screening and treatment programmes to minimize the long-term cognitive consequences of COVID-19.

1 INSERM, University of Montpellier, Neuropsychiatry: Epidemiological and Clinical Research, Montpellier, France

2 Centre for Dementia Prevention, University of Edinburgh, UK

3 Institute of Cognitive Neuroscience, University College London, London, UK

4 Department of Psychology, Faculty of Life Sciences, Northumbria University, Newcastle upon Tyne, UK

Correspondence to: Karen Ritchie, Inserm Unit1061: Neuropsychiatry, La Colombière Hospital

39 Avenue Charles Flahault, 34093 Montpellier Cedex 5, France

E-mail: karen.ritchie@inserm.fr

**Keywords:** COVID-19; neuropathology; cognition; hippocampus; coronavirus

**Abbreviations:** COVID-19 = coronavirus disease 19; HCoV = human coronaviruses; MERS-CoV = Middle East respiratory syndrome; SARS-CoV = severe acute respiratory syndrome

The majority of persons suffering from COVID-19 will recover; recovery being principally defined in terms of remission of respiratory tract symptoms. But is this the end of the story for these patients? There is increasing evidence that coronaviruses spread to extra-respiratory organs, notably the central nervous system (CNS) (Desforges *et al.*, 2014, 2019; Bohmwald *et al.*, 2018); however, little is currently known about the longer-term effects on the brain of coronavirus infection and its consequences in terms of cognitive functioning. The scarcity of research in this area precludes formal meta-analysis, yet a number of observations from a systematic literature

search suggest this to be potentially an important question for both clinical research and post-infection patient management. A literature search was undertaken from 2000 to ensure coverage of the principal recent coronavirus epidemics using Medline, SCOPUS and Google Scholar databases.

Neurological symptoms and sub-clinical cognitive dysfunction in the aftermath of COVID-19 infection are likely to result from multiple and interacting causes, notably direct damage by the virus to the cortex and adjacent subcortical structures, indirect effects due to non-CNS systemic impairment and psychological trauma.

Received April 16, 2020. Revised May 4, 2020. Accepted May 18, 2020. Advance Access publication May 28, 2020

© The Author(s) (2020). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

## Virus-induced CNS damage

Human coronaviruses (HCoV) are one of several virus groups which are considered to be potentially neurotrophic. It has been observed from previous epidemics that the respiratory coronaviruses may penetrate into the brain and cerebrospinal fluid, permeating the CNS in less than a week, and subsequently observable in cerebrospinal fluid (Bohmwald *et al.*, 2018). An autopsy series of severe acute respiratory syndrome (SARS-CoV) victims following the 2003 epidemic revealed SARS-CoV genome sequences throughout the cortex and hypothalamus (Gu *et al.*, 2005). In patients infected by Middle East respiratory syndrome (MERS-CoV), diffuse lesions were identified in several brain regions, including white matter and the subcortical areas of the frontal, temporal and parietal lobes (Arabi *et al.*, 2015). Two principal mechanisms of CNS invasion have been proposed:

- i. The blood–brain barrier, the first line of defense against viral infection, is composed in part of cerebral microvascular endothelium cells between which there are tight junctions controlling barrier permeability which appear to be compromised in the course of coronavirus infection, for example by inflammation (Koyuncu *et al.*, 2013; Miner and Diamond 2016).
- ii. The virus may directly infect neurons in the periphery or olfactory sensory neurones and thus use axonal transport to gain access to the CNS (Dahm *et al.*, 2016).

Coronaviruses produce a wide variety of acute CNS symptoms including headaches, epileptic seizures, cognitive dysfunction, motor difficulties and loss of consciousness, and may also contribute to respiratory difficulties through invasion of the brain stem and via a synapse-connected route to the medullary cardiorespiratory centre (Arabi *et al.*, 2015; Bohmwald *et al.*, 2018; Gandhi *et al.*, 2020; Li *et al.*, 2020). Although no clear aetiological pathway has been established between infection and human neurological diseases, the neuropathogenicity of HCoV is being increasingly recognized in humans, with several recent reports associating positive cases with multiple neurological disorders including encephalitis (Morfopoulou *et al.*, 2016), and Guillain–Barré syndrome (Sharma *et al.*, 2019). Other strains of coronavirus (e.g. 229E) have also been found in the brains of patients with Multiple Sclerosis (Arbour *et al.*, 2000). The presence and persistence of HCoV in human brains also appears to aggravate chronic neurological disorders such as Parkinson’s disease (Fazzini *et al.*, 1992). Already several clinical observations have been published regarding the neurological consequences of the current COVID-19 epidemic, including reports of loss of speech and comprehension, encephalopathy (Filatov *et al.*, 2020) and Guillain–Barre syndrome (Zhao *et al.*, 2020).

While HCoV infection appears to spread rapidly throughout the CNS, the temporal region appears to be a

consistent focus. Animal studies point more specifically to the vulnerability of the hippocampus with greater neuronal loss in CA1 and CA3 (Jacomy *et al.*, 2006) which would be predicted to have a detrimental effect on both learning and spatial orientation. The specific vulnerability of the hippocampus to respiratory virus infection has been previously observed in non-coronavirus infections. Studies of mice infected with influenza virus (Hosseini *et al.*, 2018) found changes in both hippocampal morphology and function, with short-term deterioration in hippocampus-dependent learning and reduced long-term potentiation associated with impairment in spatial memory. To date comprehensive neuropsychological assessment of patients, which would include tests of whole hippocampal function (such as tests of delayed recall or spatial memory) as well as novel tests probing hippocampal sub-regions such as tests of spatial memory which may aid detection of CA1 damage (Bartsch *et al.*, 2010) and a pattern completion task assessing CA3 function (Gold and Kesner 2005; Grande *et al.*, 2019), has not been carried out in relation to HCoV.

If hippocampal damage is indeed a consequence of HCoV infection then the question is raised as to whether this may lead to acceleration of hippocampal-related degeneration as occurs in Alzheimer’s disease and hasten disease onset in previously asymptomatic individuals. Animal studies have indicated that inflammation related to viral infection significantly worsens Alzheimer’s disease-related tau pathology and results in impairment of spatial memory (Sy *et al.*, 2011), now considered to be one of the first cognitive features of Alzheimer’s disease.

## Cognitive dysfunction due to non-CNS systemic impairment

Although numerous body organs are affected by coronaviruses, the respiratory system is the most severely compromised. A small study has recently estimated that 70% of critically ill patients admitted to intensive care with COVID-19 require mechanical ventilation (Arentz *et al.*, 2020), all of whom developed acute respiratory distress syndrome within 3 days. Previous neuropsychological studies of long-term outcomes for adults requiring ventilation for multiple causes observed impairments in attention, memory, verbal fluency, processing speed and executive functioning in 78% of patients 1 year after discharge and around half of patients up to 2 years (Hopkins *et al.*, 1999, 2005; Mikkelsen *et al.*, 2012). Adhikari *et al.* (2011) observed self-reported memory problems persisting up to 5 years after acute respiratory distress syndrome and impacting significantly on everyday functioning, notably taking medication and keeping medical appointments. While anxiety, depression and post-traumatic stress syndrome are also common in acute

respiratory distress syndrome patients, and may contribute to cognitive impairment (Adhikari *et al.*, 2011), there is some evidence to suggest that cognitive deficits occur independently of psychological problems, and are associated with severity of infection (Mikkelsen *et al.*, 2012).

Hypoxia, a common cause of neuropsychological changes observed in acute respiratory distress syndrome, has been associated with cerebral atrophy and ventricular enlargement (Hopkins *et al.*, 2006), with duration of hypoxia correlating with attention, verbal memory and executive functioning scores at discharge (Hopkins *et al.*, 2005). However, acute respiratory distress syndrome may also involve inflammatory responses (Han and Mallampalli, 2015) as well as anaemia and ischaemia, leading to cardiovascular and liver failure (Matthay and Zemans, 2011). Such a cascade of neurological and physiological events may further exacerbate neurological injury in acute stages to promote chronic cognitive dysfunction.

## Cognitive difficulties related to psychological distress

Cognitive difficulties are symptomatic features of all mental disorders. High rates of psychological symptoms, notably anxiety, depression, suicidal behaviour and post-traumatic stress syndrome have been reported in the general population following previous HCoV epidemics, irrespective of infectious status (Du *et al.*, 2003; Jeong *et al.*, 2016). A study of patients quarantined for suspected or confirmed MERS-CoV ( $n=40$ ), estimated that 70.8% of confirmed patients who survived the illness ( $n=24$ ) exhibited psychiatric symptoms, including hallucinations and psychosis, with 40% receiving a psychiatric diagnosis during their hospital admittance. Interestingly, none of the suspected but unconfirmed MERS-CoV patients exhibited any symptoms (Kim *et al.*, 2018), indicating a possible viral mechanism underlying psychiatric disturbance, a dose-response effect or a greater psychological impact from receiving a confirmed respiratory illness diagnosis. A study of 90 SARS-CoV cases with a 97% response rate similarly showed high levels of psychological distress with 59% diagnosed with psychiatric disorders and a continuing prevalence of 33% at 30 month follow-up. Severity of psychological symptoms was found to be related to severity of illness and functional impairment (Mak *et al.*, 2009; Wing and Leung, 2012).

Thus while higher rates of psychiatric symptoms might be expected in the general population following the epidemic due to exposure to traumatic life events (loss of income, fear, death of friends and relatives), nested within this group may be persons whose cognitive and psychological disorders are directly related to HCoV brain changes. The question might then be raised as to whether the latter group will respond to standard treatment for

example with anti-depressants, anxiolytics and cognitive therapies.

The small amount of information available from animal studies and previous respiratory epidemics suggests not only that HCoV may affect the brain, but that the consequent effect on cognitive functioning could potentially persist for a long period following recovery. For infected persons already suffering from CNS disorders the effects are likely to be even more debilitating. While new cases of well-characterized neurological disorders subsequent to the current epidemic may be relatively easy to identify, persisting sub-clinical disorders such as mild cognitive dysfunction, selective memory and speech impairments or exacerbation of pre-existing degenerative neuropathologies such as vascular dementia and Alzheimer's disease, presenting principally in general practice, may easily go undetected or be attributed to psychological reactions to the fear and social upheaval generated by the pandemic.

## Conclusion

In the face of increasing reports of CNS involvement in COVID-19 cases, the current epidemic is likely to be accompanied by a significant increase in the prevalence of longer-term cognitive dysfunction impacting on ability to return to everyday functioning. This is likely to be due not only to the behavioural consequences of incident neurological disorders directly related to the virus, but also secondary to damage to other body organs, psychiatric disorders and the worsening of pre-existing cognitive difficulties. The number of persons exposed to the virus likely to be affected and the protective factors operating in cases who do not experience cognitive changes is presently unknown. Studies of cognitive dysfunction related to previous epidemics are few and too small to make estimations at a population level.

Further research is needed to understand (i) the range of COVID-19 associated neurological disorders and their cognitive manifestations; (ii) the underlying associations between viral spread, associated proinflammatory changes and disease pathogenesis; (iii) the duration and extent of neurological and cognitive changes following resolution of the acute viral illness; (iv) the association between severity of the viral illness and subsequent cognitive dysfunction; (v) the effect of antiviral, psychological and other interventions (when available) on short- and long-term cognitive function.

The current coronavirus outbreak is unlikely to be the last (with SARS and MERS, COVID-19 this is the third coronavirus epidemic in 10 years). It is therefore imperative that the medical-scientific community look beyond the current acute crisis to the links between coronavirus infection and long-term neurological sequelae. While basic science research will deliver insights into potential mechanistic relationships between viral infections and neurological disease, better understanding of these associations

may inform treatment plans aimed both at treating the acute infection and limiting downstream cognitive decline. For instance, if future work showed that viral load and/or subsequent proinflammatory state correlated with long-term cognitive outcome then in future outbreaks treatment protocols could mitigate against the latter by adjusting the dose and duration of antiviral therapies or add second stage anti-inflammatory treatments.

This research could be taken forward in several ways. Existing longitudinal studies of neurological disorders could be expanded to include acquisition of data concerning COVID-19 exposure and antibody status into existing research protocols which already capture data on cognitive function, brain imaging and disease biomarkers. Data from large prospective cohorts such as UK Biobank could provide population-scale data with clearer information regarding prevalence and persistence of effects over extensive timelines. In parallel, preclinical studies aimed at uncovering the mechanistic association between coronavirus infection and neurological disease will be facilitated by the recent creation of mouse models of COVID-19, which will enable assessment of the interaction between viral pathology and neurodegeneration and will provide a resource for the development of new treatments (Wang, 2020). These will be complemented by the accumulation of information about disease onset and clinical progression, particularly from general practice, which will be crucially important for the implementation of future population-level screening and treatment programmes.

The scope and severity of the current COVID-19 pandemic are unparalleled in modern society. The downstream implications for neurological function may be equally grave. While the current focus is on acute disease management, in the near future attention will need to turn to the long-term consequences of COVID-19 infection and their mitigation.

## Competing interests

The authors report no competing interests.

## References

- Adhikari NKJ, Tansey CM, McAndrews MP, Matté A, Pinto R, Cheung AM, et al. Self-reported depressive symptoms and memory complaints in survivors five years after ARDS. *Chest* 2011; 140: 1484–93.
- Arabi YM, Harthi A, Hussein J, Bouchama A, Johani S, Hajeer AH, et al. Severe neurologic syndrome associated with Middle East respiratory syndrome corona virus (MERS-CoV). *Infection* 2015; 43: 495–501.
- Arbour N, Day R, Newcombe J, Talbot PJ. Neuroinvasion by human respiratory coronaviruses. *J Virol* 2000; 74: 8913–21.
- Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State. *JAMA* 2020; 323: 1612.
- Bartsch T, Schönfeld R, Müller FJ, Alfke K, Leprow B, Aldenhoff J, et al. Focal lesions of human hippocampal CA1 neurons in transient global amnesia impair place memory. *Science* 2010; 328: 1412–5.
- Bohmwald K, Galvez NMS, Rios M, Kalergis AM. Neurologic alterations due to respiratory virus infections. *Front Cell Neurosci* 2018; 12: 386.
- Dahm T, Rudolph H, Schwerk C, Schrotten H, Tenenbaum T. Neuroinvasion and inflammation in viral central nervous system infections. *Mediators Inflamm* 2016; 2016: 8562805.
- Desforges M, Le Coupanec A, Dubeau P, Bourgouin A, Lajoie L, Dubé M, et al. Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system? *Viruses* 2019; 12: 14.
- Desforges M, Le Coupanec A, Brison E, Meessen-Pinard M, Talbot PJ. Neuroinvasive and neurotropic human respiratory coronaviruses: potential neurovirulent agents in humans. *Adv Exp Med Biol* 2014; 807: 75–96.
- Du L, Zhao J, Shi Y, Xi Y, Zheng GG, Yi Y, et al. A report of 4 cases of severe acute respiratory syndrome patients with suicide tendency. *Acad J Second Mil Med Univ* 2003; 24: 636–7.
- Fazzini E, Fleming J, Fahn S. Cerebrospinal fluid antibodies to coronavirus in patients with Parkinson's disease. *Mov Disord* 1992; 7: 153–8.
- Filatov A, Sharma P, Hindi F. Neurological complications of coronavirus disease (COVID-19): encephalopathy. *Cureus* 2020; 70: 311–22.
- Gandhi S, Srivastava AK, Ray U, Tripathi PP. Is the collapse of the respiratory center in the brain responsible for respiratory breakdown in COVID-19 patients? *ACS Chem Neurosci* 2020; 11: 1379–81. [Epub ahead of print]
- Gold AE, Kesner RP. The role of the CA3 subregion of the dorsal hippocampus in spatial pattern completion in the rat. *Hippocampus* 2005; 15: 808–14.
- Grande X, Berron D, Horner AJ, Bisby JA, Düzel E, Burgess N. Holistic recollection via pattern completion involves hippocampal subfield CA3. *J Neurosci* 2019; 39: 8100–11.
- Gu J, Gong E, Zhang B, Zheng J, Gao Z, Zhong Y, et al. Multiple organ infection and the pathogenesis of SARS. *J Exp Med* 2005; 202: 415–24.
- Han S, Mallampalli RK. The acute respiratory distress syndrome: from mechanism to translation [published correction appears. *J Immunol* 2015; 194: 855–60.
- Hopkins RO, Weaver LK, Collingridge D, Parkinson RB, Chan KJ, Orme JF. Two-year cognitive, emotional and quality of life outcomes in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2005; 171: 340–7.
- Hopkins RO, Weaver LK, Pope D, Orme JF, Bigler ED, Larson-Lohr V. Neuropsychological sequelae and impaired health status in survivors of severe acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1999; 160: 50–6.
- Hopkins RO, Gale SD, Weaver LK. Brain atrophy and cognitive impairment in survivors of acute respiratory distress syndrome. *Brain Inj* 2006; 20: 263–71.
- Hosseini S, Wilk E, Michaelsen-Preusse K, Gerhauser I, Baumgärtner W, Geffers R, et al. Long-term neuroinflammation induced by influenza A virus infection and the impact on hippocampal neuron morphology and function. *J Neurosci* 2018; 38: 3060–80.
- Jacomy H, Fragoso G, Almazan G, Mushynski WE, Talbot PJ. Human coronavirus OC43 infection induces chronic encephalitis leading to disabilities in BALB/C mice. *Virology* 2006; 349: 335–46.
- Jeong H, Yim HW, Song YJ, Ki M, Min JA, Cho J, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health* 2016; 38: e2016048.
- Kim HC, Yoo SY, Lee BH, Lee SH, Shin HS. Psychiatric findings in suspected and confirmed middle east respiratory syndrome patients quarantined in hospital: a retrospective chart analysis. *Psychiatry Investig* 2018; 15: 355–60.

- Koyuncu OO, Hogue IB, Enquist LW. Virus infections in the nervous system. *Cell Host Microbe* 2013; 13: 379–93.
- Li YC, Bai WZ, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020; doi: 10.1002/jmv.25728.
- Mak IWC, Chu CM, Pan PC, Yiu MGC, Chan VL. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry* 2009; 31: 318–26.
- Matthay MA, Zemans RL. The acute respiratory distress syndrome: pathogenesis and treatment. *Annu Rev Pathol Mech Dis* 2011; 6: 147–63.
- Mikkelsen ME, Christie JD, Lanken PN, Biester RC, Thompson BT, Bellamy SL, et al. The adult respiratory distress syndrome cognitive outcomes study. *Am J Respir Crit Care Med* 2012; 185: 1307–15.
- Miner JJ, Diamond MS. Mechanisms of restriction of viral invasion at the blood-brain barrier. *Curr Opin Immunol* 2016; 38: 18–23.
- Morfopoulou S, Brown JR, Davies EG, Anderson G, Virasami A, Qasim W, et al. Human coronavirus OC43 associated with fatal encephalitis. *N Engl J Med* 2016; 375: 497–8.
- Sharma K, Tengsupakul S, Sanchez O, Phaltas R, Maertens P. Guillain-Barre syndrome with unilateral peripheral facial and bulbar palsy in a child: a case report. *SAGE Open Med Case Rep* 2019; 7: 2050313X19838750. doi: 10.1177/2050313X19838750.
- Sy M, Kitazawa M, Medeiros R, Whitman L, Cheng D, Lane TE, et al. Inflammation induced by infection potentiates tau pathological features in transgenic mice. *Am J Pathol* 2011; 178: 2811–22.
- Wang Q. HACE2 transgenic mouse model for coronavirus (Covid-19) research. The Jackson Laboratory. 2020. <https://www.jax.org/news-and-insights/2020/february/introducing-mouse-model-for-corona-virus> (24 February 2020, date last accessed).
- Wing YK, Leung CM. Mental health impact of severe acute respiratory syndrome: a prospective study. *Hong Kong Med J* 2012; 18: 24–7.
- Zhao H, Shen D, Zhou H, Liu J, Chen S. Guillain-Barré syndrome associated with SARS-CoV-2 infection: causality or coincidence? *Lancet Neurol* 2020; 19: 383–4.