



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

ventricular assessment, valve function, and pericardial effusion. In 12 patients, it was possible to acquire apical views and in 4 a subcostal view was feasible (Figure 1C).

The unconventional and innovative use of a TEE probe to perform transthoracic echocardiography with the patient in prone positioned and invasively ventilated intensive care unit patients, is feasible and can be of diagnostic quality in most cases. The technique described here can be an alternative or even a superior to conventional TTE in prone patients.

Apostolia Marvaki, PhD
Alexandros Papachristidis, MD*
Eleni Nakou, PhD
Eva Toth, MD
Kevin O’Gallagher, MBBS
Richard Fisher, MBBS
Ajay M. Shah, MD, FMedSci
Mark J. Monaghan, PhD

*Cardiology Department
King’s College Hospital NHS Foundation Trust
SE5 9RS London
United Kingdom
E-mail: alexandros.papachristidis@nhs.net

<https://doi.org/10.1016/j.jcmg.2020.06.005>

© 2020 by the American College of Cardiology Foundation. Published by Elsevier.

The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors’ institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the *JACC: Cardiovascular Imaging* author instructions page.

REFERENCES

1. Gibson PG, Qin L, Puah S. COVID-19 Acute Respiratory Distress Syndrome (ARDS): Clinical Features and Differences From Typical pre-COVID-19 ARDS. *Med J Aust* 2020;213:54–6.e1.
2. Giustiniano E, Padua E, Negri K, Bragato RM, Cecconi M. Echocardiography During Prone-Position Mechanical Ventilation in Patients With COVID-19: A Proposal for a New Approach. *J Am Soc Echocardiogr* 2020;33:905–6.
3. Papachristidis A, Galli E, Geleijnse ML, et al. Standardized delineation of endocardial boundaries in three-dimensional left ventricular echocardiograms. *J Am Soc Echocardiogr* 2017;30:1059–69.

Right Ventricular Function and Pulmonary Pressures as Independent Predictors of Survival in Patients With COVID-19 Pneumonia



Coronavirus disease-2019 (COVID-19) disease, caused by severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2), can lead to cardiac impairment, with increased troponin levels, left ventricular (LV) dysfunction and myocarditis (1,2). Information for right ventricular (RV) involvement and pulmonary pressures in these patients is limited, and this study

aimed to explore the possible association between these factors and mortality. This study was conducted from February 20, 2020, to April 2020 in 4 centers. The final follow-up date was April 20, 2020. All consecutive patients with test results positive for SARS-CoV-2, as well as laboratory and computed tomography-confirmed interstitial pneumonia were included. Clinical, laboratory, radiological, and ultrasonographic data were collected. Cardiac injury was defined by measuring blood concentrations of high-sensitivity troponin I (hs-TnI) using the 99th percentile upper reference limit of the electrochemiluminescence immunoassay method. RV end-diastolic chamber size was assessed by using basal-tract and mid-tract diameters in apical 4-chamber view. Tricuspid annular plane systolic excursion (TAPSE) was calculated as a bedside-feasible index of RV longitudinal systolic function by aligning an M-mode cursor parallel with the RV free wall as it met the tricuspid annulus.

Pulmonary artery systolic pressure (PASP) was calculated by adding the value of right atrial pressure to the systolic transtricuspid gradient. Mean pulmonary artery pressure (mPAP) was calculated as $0.6 \times \text{PASP} + 2$ mm Hg (3). Outcomes of patients with cardiac involvement were compared to those in patients without cardiac involvement. An ethics committee approved the study, and all individuals gave written informed consent.

A total of 115 patients with COVID-19 pneumonia were included in the analysis. The mean age was 64.6 years (range 20 to 88 years of age), and 45 (40%) were female. Twenty-six patients had cardiac injury, and those patients were older (mean age: 73.5 years of age [range 37 to 88] vs. 55.3 years of age [range 20 to 78], respectively; $p < 0.001$). They also more frequently had systemic hypertension (16 of 26 patients [61.5%] vs. 26 of 89 patients [29.2%], respectively; $p < 0.001$). They also had higher levels of C-reactive protein (mean: 98.8 [range 45.8 to 130.4] $\text{mg} \cdot \text{l}^{-1}$ vs. 38.9 [range 22.2 to 96.3] $\text{mg} \cdot \text{l}^{-1}$, respectively; $p < 0.001$); higher D-Dimer (mean: 4.8 [3.2 to 7.1] vs. 2.1 [0.5 to 3.4], respectively; $p < 0.001$); a higher proportion of multiple ground-glass opacities in computed tomography (CT) findings (18 of 26 patients [69.2%] vs. 30 of 89 patients [33.7%], respectively; $p < 0.001$); and multiple consolidations by lung ultrasonography (20 of 26 patients [76.6%] vs. 34 of 89 patients [38.2%], respectively; $p < 0.001$). Lower-limb compression ultrasonography (CUS) was positive (proximal deep vein thrombosis of the common femoral vein and/or popliteal vein) only in 3 patients with cardiac injury and in 3 other patients without cardiac injury. A single echocardiograph for each hospital was identified,

TABLE 1 Echocardiographic Structural and Systolic/Diastolic Functional Indices in COVID-19 Patients With and Without Cardiac Injury

| | Without Cardiac Injury (n = 89) | With Cardiac Injury (n = 26) | p Value |
|---------------------------------|------------------------------------|---------------------------------|---------|
| Septal wall thickness, mm | 10.7 ± 2.3 | 11.5 ± 2.3 | NS |
| LV posterior wall thickness, mm | 9.8 ± 1.2 | 10.8 ± 1.1 | NS |
| LV end-diastolic diameter, mm | 48.7 ± 4.8 | 49.9 ± 6.7 | NS |
| LV end-systolic diameter, mm | 33.4 ± 3.8 | 35.4 ± 5.2 | NS |
| Mitral peak A velocity, m/s | 0.72 ± 0.32 | 0.73 ± 0.3 | NS |
| Mitral peak E/A ratio | 0.91 ± 0.3 | 0.88 ± 0.3 | NS |
| LV ejection fraction % | 56.2 ± 4.4 | 54.4 ± 4.6 | NS |
| RV basal tract diameter, mm | 31.2 (2.6) | 35.8 (4.2) | <0.01 |
| RV mid-tract diameter, mm | 28.6 (3.7) | 32.4 (3.9) | <0.01 |
| Tricuspid peak E velocity, m/s | 0.15 ± 0.08 | 0.12 ± 0.06 | <0.01 |
| Tricuspid peak A velocity, m/s | 0.13 ± 0.08 | 0.16 ± 0.06 | <0.01 |
| Tricuspid peak E/A ratio | 1.1 ± 0.6 | 0.77 ± 0.5 | <0.01 |
| TRV, m/s | 2.9 ± 0.5 | 3.4 ± 0.6 | <0.001 |
| PASP, mm Hg | 36.1 ± 5.3 | 46.8 ± 4.8 | <0.001 |
| MPAP, mm Hg | 23.6 ± 2.7 | 29.6 ± 2.9 | <0.001 |
| TAPSE, mm | 20.3 ± 4.7 | 15.4 ± 3.2 | <0.001 |

Values are mean ± SD or n (%).

COVID-19 = coronavirus disease-2019; E/A = ratio of peak velocity blood flow from LV relaxation in early diastole [E wave] to peak velocity flow in late diastole caused by atrial contraction [A wave]; LV = left ventricle; MPAP = mean pulmonary artery pressure; NS = not significant; PASP = pulmonary artery systolic pressure; RV = right ventricle; TAPSE = tricuspid annular systolic plane excursion; TRV = tricuspid regurgitation velocity.

portably to facilitate sanitation, and the examination was performed by a single operator, reducing patient contact with the device. RV end-diastolic diameters and mPAP values were significantly increased in patients with cardiac injury, whereas the tricuspid inflow E/A ratio (the ratio of peak velocity blood flow from LV relaxation in early diastole [E wave] to peak velocity flow in late diastole caused by atrial contraction [A wave]) and TAPSE were reduced (Table 1). All patients were treated with oxygen, and the percentages of use of noninvasive ventilation and invasive mechanical ventilation were 16.5 % (19 patients) and 9.5 % (11 patients), respectively. A greater proportions of patients with cardiac injury required invasive or noninvasive mechanical ventilation (17 of 26 [65.4%] vs. 20 of 89 [22.4%], respectively; $p < 0.001$). The common complication in patients with cardiac injury was acute respiratory distress syndrome (14 of 26 [63.5%] vs. 15 of 89 [16.8%], respectively; $p < 0.001$). Patients with cardiac injury had higher mortality than those without cardiac injury (13 of 26 [50%] vs. 8 of 89 [7.8%], respectively; $p < 0.0001$). In a Cox regression model in COVID-19 patients, troponin levels (hazard ratio [HR]: 4.33 [95% confidence interval [CI]: 1.87 to 8.49]; $p < 0.001$), partial pressure of oxygen (P_{aO_2}) at admission (HR: 0.26 [95% CI: 0.15 to 0.74]; $p < 0.01$); mPAP (HR: 3.8 [95% CI: 1.72 to 6.39] $p < 0.001$), and RV TAPSE (HR: 0.5 [95% CI: 0.22 to 0.74] $p < 0.001$) emerged as the only independent predictors of in-hospital death.

Despite the high prevalence of normal CUS, it was found that increases in mPAP and RV dysfunction were common conditions among patients with COVID-19 pneumonia and cardiac injury, associated with higher risk of in-hospital mortality. Use of empiric therapeutic anticoagulation in certain COVID-19 patients who did not have deep vein thrombosis has been advocated, but this remains controversial because of the risk of major bleeding and lack of data about the true incidence of pulmonary embolism (4). However, assessment of RV function and of pulmonary pressures during the recovery of these patients may represent a key point in the prognostic stratification.

Antonello D'Andrea, MD, PhD*

Raffaella Scarafie, MD

Lucia Riegler, MD

Biagio Liccardo, MD

Fabio Crescibene, MD

Rosangela Cocchia, MD

Eduardo Bossone, MD

*Department of Cardiology

University of Campania Luigi Vanvitelli

Umberto I° Hospital - Nocera Inferiore (Salern)

Corso Vittorio Emanuele 121A

80121, Naples

Italy

E-mail: antonellodandrea@libero.it

<https://doi.org/10.1016/j.jcmg.2020.06.004>

© 2020 by the American College of Cardiology Foundation. Published by Elsevier.

The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the JACC: Cardiovascular Imaging author instructions page.

REFERENCES

- Guo T, Fan Y, Chen M, et al. Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol* 2020;5:811-8.
- Bonow RO, Fonarow GC, O'Gara PT, Yancy CW. Association of coronavirus disease 2019 (COVID-19) with myocardial injury and mortality. *JAMA Cardiol* 2020;5:751-3.
- Bossone E, D'Andrea A, D'Alto M, et al. Echocardiography in pulmonary arterial hypertension: from diagnosis to prognosis. *J Am Soc Echocardiogr* 2013;26:1-14.
- Obi AT, Tignanelli CJ, Jacobs BN, et al. Empirical systemic anticoagulation is associated with decreased venous thromboembolism in critically ill influenza A H1N1 acute respiratory distress syndrome patients. *J Vasc Surg Venous Lymphat Disord* 2019;7:317-24.

Coronary Artery Calcification and Complications in Patients With COVID-19



In the coronavirus disease-2019 (COVID-19) pandemic, early risk stratification of severe forms is essential. Patients with COVID-19 and cardiovascular