



Modeling and forecasting trend of COVID-19 epidemic in Iran until May 13, 2020

Ali Ahmadi*¹, Yasin Fadaei², Majid Shirani³, Fereydoon Rahmani⁴

Received: 18 Mar 2020

Published: 31 Mar 2020

Abstract

Background: COVID-19 is a new disease and precise data are not available about this illness in Iran and in the world. Thus, this study aimed to determine the epidemic trend and prediction of COVID-19 in Iran.

Methods: This was a secondary data analysis and modeling study. The daily reports of definitive COVID-19 patients released by Iran Ministry of Health and Medical Education were used in this study. Epidemic projection models of Gompertz, von Bertalanffy, and least squared error (LSE) with percentage error were used to predict the number of hospitalization cases from April 3, 2020 until May 13, 2020.

Results: The prediction of the number of patients on April 3, 2020 by von Bertalanffy, Gompertz, and LSE, with 95% confidence interval (CI), were estimated at 44 200 (39 208-53 809), 47 500 (38 907-52 640), and 48 000 (40 000-57 560), respectively. The number of deceased COVID-19 patients was also estimated to be 3100 (2633-3717) individuals by the von Bertalanffy model, 3700 (2900-4310) by Gompertz's model, and 3850 (3200-4580) by LSE. Making predictions about the flat epidemic curve and number of patients based on Gompertz model, will project 67 000 (61 500-87 000) cases. Based on Gompertz and von models, 7900 (6200- 9300) and 4620 (3930-5550) deaths will occur from May 13 to June 1, 2020, respectively, and then the curve will flatten.

Conclusion: In this study, estimations were made based on severely ill patients who were in need of hospitalization. If enforcement and public behavior interventions continue with current trends, the COVID-19 epidemic will be flat from May 13 until July, 2020 in Iran.

Keywords: COVID-19, Coronavirus, Prediction, Modeling, Epidemiology

Conflicts of Interest: None declared

Funding: This study was financially supported by the Deputy of Research and Technology of Shahrekord University of Medical Sciences, Shahrekord, Iran (Grant no. 254).

*This work has been published under CC BY-NC-SA 1.0 license.

Copyright© Iran University of Medical Sciences

Cite this article as: Ahmadi A, Fadaei Y, Shirani M, Rahmani F. Modeling and forecasting trend of COVID-19 epidemic in Iran until May 13, 2020. *Med J Islam Repub Iran.* 2020 (31 Mar);34:27. <https://doi.org/10.34171/mjiri.34.27>

Introduction

Coronaviruses are a large family of viruses that have been identified since 1965; to date, 7 species of them have been reported to affect humans. These viruses have 3 genotypes of alpha, beta, and gamma. The natural reservoirs of these diseases are mammals and birds, and thus they are considered as zoonotic diseases (1, 2). Severe acute respiratory syndrome (SARS) is caused by a species of coronavirus that infects humans, bats, and certain other mammals, which has led to epidemics in 2002 and 2003 (2-4); Middle

East respiratory syndrome (MERS) caused an epidemic in Saudi Arabia in 2012 (5); and more recently their newest variant Coronavirus Disease-2019 (COVID-19) has led to the recent pandemic in China, Iran, Italy, and across the world (6-8). COVID-19 causes a direct and indirect transmission of respiratory diseases with a wide range of symptoms, including cold symptoms, respiratory/fever symptoms, cough, shortness of breath, kidney failure, and even death (9). The complete clinical manifestation of COVID-

Corresponding author: Dr Ali Ahmadi, ahmadi@skums.ac.ir

¹ Modeling in Health Research Center and Department of Epidemiology and Biostatistics, School of Public Health, Shahrekord University of Medical Sciences, Shahrekord, Iran

² Modeling in Health Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

³ Department of Urology, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁴ Department of Infectious Disease, School of Medicine, Shahrekord University of Medical Sciences, Shahrekord, Iran

↑What is “already known” in this topic:

COVID-19 is a new pandemic disease and precise data on its epidemic spread are not available in Iran and in the world.

→What this article adds:

This study found that if interventions continue at the same rate, the spread of the Covid-19 epidemic would stop during May 13 and early July, and the curve will be flattened.

COVID-19 is not clear yet and its transmission risk is not completely understood (10). However, the virus is believed to be transmitted mostly via contact, droplets, aspirates, feces. Generally, everyone is prone to this viral disease. The mean incubation period of COVID-19 was 5.2 days (4.1–7 days) and the basic reproductive number (R_0) was reported 2.2 (95% CI: 1.4 to 3.9) (11). In another study, the mean incubation period ranged from 0–24 days, with the mean of 6.4 days. The R_0 of COVID-19 at the early phase, regardless of different prediction models, was higher than SARS and MERS, and the majority of patients (80.9%) were considered asymptomatic or mild pneumonia (12). The case fatality ratio was 2% (12), 2.3% (8), 3.46% (13), and elderly men with underlying diseases were at a higher risk of death (13). As of March 29, 2020, COVID-19 pandemic was declared by the World Health Organization (WHO) in more than 100 countries (most prevalent in the United States, Italy, China, Spain, Germany, Iran, and France) (14). In Iran, the first case of COVID-19 was February 19, 2020 in Qom, and we used the reported data until March 29, 2020 in Iran. Until March 29, 2020 (The date when this manuscript was being conducted), according to the daily reports, 38 309 cases of Covid-19 and 2640 related deaths were reported in Iran (15, 16). As of February 29, 2020, all schools and universities and as of March 7, 2020 almost all public places and shrines have been closed. On March 2, 2020, a team of WHO experts landed in Tehran, Iran, to support the ongoing response to the COVID-19 outbreak in the country (17). Currently, people are referring to health centers and hospitals, and the public is almost alarmed by the epidemic of panic and inaccurate reporting in cyberspace. Important questions in people's mind are as follow: How many people have COVID-19 in Iran? What is the status of COVID-19 epidemic curve in Iran? When will the epidemic will go and how it ends? We cannot answer these questions with certainty, but they will be investigated in terms of pathogenic agents (coronavirus), host conditions, behavior (human), and environmental factors of coronavirus transmission, daily reports of definitive COVID-19 patients released by Iran Ministry of Health and Medical Education, and the use of modeling given the assumptions and the percentage of error. Although the models are different, multiple and changeable in nature and do not insist on the correctness of the forecasts, the decision-making conditions for health policymakers and authorities are more transparent and helpful (18). This study aimed to model and determine the epidemic trend and predict the number of patients hospitalized due to COVID-19 in Iran using mathematical and statistical modeling.

Methods

This was a secondary data analysis and mathematical modeling study based on a research proposal approved by Shahrekord University of Medical Sciences (Code of Ethics Committee on Biological Research: IR.SKUMS.REC 1398.254) (19).

For the statistical analysis of definitive COVID-19 patients in Iran, daily reports of the Ministry of Health and Medical Education were used (15). The definitive diagnosis of COVID-19 was made using virus isolates from patients'

biological samples in hospitals. When real time polymerase chain reaction (PCR) test turned positive for COVID-19 in patients with respiratory symptoms and confirmed by the reference laboratory in School of Public Health, Tehran University Medical of Sciences and Pasteur Institute of Iran (20), they were used for analysis. Patient population growth, epidemic curves, and recovered, and deceased individuals were used to form a conceptual framework of an epidemic and predict the COVID-19 epidemic trend. Quasi classical infectious disease (Susceptible→Exposed→Infected→Removed: SEIR) model was used (21). Different scenarios were designed and implemented for modeling and forecasting. First, based on a search for reliable sources of disease trends and epidemic curves across the world, the curve of Iran was drawn (10, 18, 22). Focused and scientific group discussion sessions were held with experts on epidemiology, biostatistics, and mathematics, infectious diseases specialists, and health care managers on the topic. Different scenarios were discussed and agreement was reached on the application of final scenarios. To predict the growth of this epidemic, different models were used. In the first scenario, the most optimistic estimation and control of the epidemic was during an incubation period (von Bertalanffy model, the most ideal model). In this scenario, traced contacts are isolated immediately on symptom onset (and not before) and isolation prevents disease transmission. In the second scenario, an intermediate and fit-to-data model (Gompertz) was used. In the third scenario, the use of the growth rate is greater than the first and second models, and it is the opposite of the first scenario (LSE). To select the scenarios, fit the data with the models and growth rate of the cases were used. The Gompertz growth, von Bertalanffy growth equation, and curve fitting by LSE method with cubic polynomial for Epidemic forecasts were run in MATLAB software. Models are presented as the following differential equations:

Gompertz Differential Equation:

$$\frac{d}{dt}P(t) = aP \ln(1/bP)$$

Von Bertalanffy's differential growth equation:

$$\frac{d}{dt}P(t) = aP ((bP)^c - 1)$$

Cubic polynomial polynomials:

$$P(t) = at^3 + bt^2 + ct + e$$

Where p represents the number of individuals in each population, a , b , c , and e represent unknown parameters and t time. d/dt is a derivative of time (23,24).

The assumptions of the model are as follow:

- Models are based on official reported data and recruitment testing from hospitalized cases.
- Any manipulation and misinformation will affect the model.
- The method for finding patients is fixed.

The unknown parameters were estimated by running the `fminsearch`, a MATLAB function, which is a least squares algorithm. The parameters were estimated based on the official reported data of infected, cured, and dead cases. The estimated values of the parameters for different scenarios are reported in Table 1. Also, MATLAB software was used to fit data and solve the equations. MATLAB codes are presented in Appendix. Moreover, the percentage of the root

Table 1. Estimated Parameters (values of the inputs used) for modeling

Model	Parameter	Infected cases			Recovered cases			Death cases		
		a	B	c	a	B	c	a	B	c
Gompertz		0.1	2.9939e+04	-	0.1	1.8107e+04	-	0.0660	2.6710e+04	-
VonBertalanffy		0.8702	2.7109e-05	0.2109	0.5564	7.5133e-05	0.3708	0.7321	2.2840e-05	0.1239

mean square error (RMSE) was used to validate the models and 95% confidence interval (CI) was utilized to calculate the coefCI MATLAB function.

The basic reproduction number R_0 was calculated using the following formula (25):

$$R_0 = 1 + r * T_c$$

where, T_c and r are the mean generation interval of the infected and the growth rate, respectively, $T_c = 7.5$, and $r = 0.1$. The growth rate of Gompertz model is $r = 0.1$; thus, the number of R_0 is 1.75. All estimations and detections of COVID-19 were made based on the current conditions of laboratory sampling from critically ill and hospitalized patients (tip of the iceberg spread of the disease). In this modeling, asymptomatic patients and those with moderate symptoms from whom no samples were taken were excluded. Also, the data of patients diagnosed based on CT scans were not included in this study. The forecast dates were selected based on the end of New Year (April 3, 2020)

holidays in Iran and the onset of epidemic curve flattening (May 13, 2020).

Results

Frequency of daily statistics of COVID-19, including definite new cases, number of deaths, and recovered cases in Iran are shown in Table 2. The trend of this epidemic spread in Iran (daily linear and cumulative trend) is illustrated in Figure 1. According to data released on COVID-19 in Iran as of March 29, 2020, the following forecasts for April 3 and May 13, 2020 were reported (Figs. 2, 3, and 4).

According to the Gompertz model, in the most optimistic perspective, the maximum number of infected people until April 3, 2020 is 47 500, with 95% confidence interval (CI: 38907- 52640) (Fig. 2 A1). The percentage of the root mean square error (RMSE) for Gompertz model is 12%. Based on von Bertalanffy's growth model (the most ideal model with high isolation of patients and others intervention such

Table 2. The frequency of COVID-19 new cases, cumulative cases, and deceased cases in Iran

Date	New cases	Cumulative number cases	New deceased	Cumulative deceased	Cumulative recovered cases
19-Feb-20	2	2	2	2	0
20-Feb-20	3	5	0	2	0
21-Feb-20	13	18	2	4	0
22-Feb-20	10	28	1	5	0
23-Feb-20	15	43	3	8	0
24-Feb-20	18	61	4	12	0
25-Feb-20	34	95	4	16	0
26-Feb-20	44	139	3	19	0
27-Feb-20	106	245	7	26	0
28-Feb-20	143	388	8	34	73
29-Feb-20	205	593	9	43	123
1-Mar-20	385	978	11	54	175
2-Mar-20	523	1501	12	66	291
3-Mar-20	835	2336	11	77	435
4-Mar-20	586	2922	15	92	552
5-Mar-20	591	3513	15	107	739
6-Mar-20	1234	4747	17	124	913
7-Mar-20	1076	5823	21	145	1669
8-Mar-20	743	6566	49	194	2134
9-Mar-20	595	7161	43	237	2394
10-Mar-20	881	8042	54	291	2731
11-Mar-20	958	9000	63	354	2959
12-Mar-20	1075	10075	75	429	3276
13-Mar-20	1289	11364	85	514	3529
14-Mar-20	1365	12729	97	611	4339
15-Mar-20	1209	13938	113	724	4790
16-Mar-20	1053	14991	129	853	4996
17-Mar-20	1178	16169	135	988	5389
18-Mar-20	1192	17361	147	1135	5710
19-Mar-20	1046	18407	149	1284	5979
20-Mar-20	1237	19644	149	1433	6745
21-Mar-20	966	20610	123	1556	7635
22-Mar-20	1028	21638	129	1685	7913
23-Mar-20	1411	23049	127	1812	8376
24-Mar-20	1762	24811	122	1934	8913
25-Mar-20	2206	27017	143	2077	9625
26-Mar-20	2389	29406	157	2234	10457
27-Mar-20	2926	32322	144	2378	11133
28-Mar-20	3076	35408	139	2517	11679
29-Mar-20	2901	38309	123	2640	12391




Fig. 1. The trend of new and cumulative cases of COVID-19 in Iran




Fig 2. Prediction of COVID-19 by Gompertz and von Bertalanffy models in Iran, April 3, 2020

as China experience), the maximum number of COVID-19 patients is 44 200, with 95% confidence interval (CI:




Fig. 3. Prediction of the flat of epidemic curve of COVID-19 in Iran, May 13, 2020

39208-53809), (Fig. 2 A2) and 17% RMSE. According to the method of the least squared error, this value was estimated to be 48 000 (CI: 40000 - 57560), with 19% RMSE (Fig. 4 H).

According to Gompertz and von models on prediction of recovered patients, and the models estimated the number of recovered individuals to be 15 300, with 95% confidence interval (CI: 8600-23000) and 3% RMSE, and 15 950, with 95% confidence interval (CI: 13000-24000), and 3.7% RMSE until April 3, 2020, respectively (Fig. 2 B1, B2).

Moreover, according to Figure 4 G, the maximum population of recovered individuals was estimated to be 15 900, with 95% confidence interval (CI: 13500-19000), according to the method of least squared error.

To predict and estimate the number of deaths due to COVID-19, 2 growth models (Fig. 2 C1, C2) and the least

squared error (Fig. 4 I) were used to maximize von's growth in the most optimistic manner. According to von, Gompertz, and least squared error methods, the number of deaths, with 95% confidence interval, will be 3100, (CI: 2633-3717) with 17% RMSE, 3700 (CI: 2900-4310) with 12% RMSE, and 3850 (CI: 3200-4580) on April 3, 2020, respectively. To make predictions for May 13, 2020, when the epidemic curve will be flat, 3 models of growth with 95% confidence interval were used. Based on Gompertz model, the epidemic curve will be flattened on May 13, 2020 with about 67 000 patients (CI: 61500- 87000), with 10% RMSE (Fig. 3 D). Based on von and Gompertz models, the number of cured cases will be 22 000 (CI: (12500 - 32000), with 3.1% MSRE, and 26 000 (CI: 21700-39500), with 3.7% MSRE, respectively (Fig. 3 E2, E1). To predict the number of deaths due to COVID-19 when the epidemic




Fig. 4. Prediction of COVID-19 by least squared error (LSE) cubic polynomial model in Iran, April 3, 2020

curve is flat based on Gompertz and von models, there will be 7900 deaths (CI: 6200-9300), with 10% MSRE, and 4620 deaths (CI: 3930-5550), with 9.1% MSRE, until May 13 and June 1, 2020, respectively. (Fig. 3 F1, F2).

Discussion

In this study, the trend of COVID-19 epidemic prediction and estimation of the number of patients, R_0 , deaths, and recovered individuals were performed and reported based on mathematical and statistical models. Although this prediction may be associated with random errors, it was made with assumptions about the past trends of the COVID-19 epidemic in Iran as well as the behavior of the people and government interventions (sampling of severe cases and hospitalization). Implementation of government interventions such as social distancing, isolation of patients, and follow-up of those around them based on the epidemic management protocol in Iran is of high importance. Thus, observance of these interventions by the public and the government has an impact on modeling predictions. Moreover, according to a valid scientific report, delay in the onset of symptoms until the isolation of patients plays an important role in controlling the epidemic (18). To control the majority of outbreaks, for R_0 of 2.5, more than 70% of contacts had to be traced, and for R_0 of 3.5, more than 90% of contacts had to be traced. The delay between symptom onset and isolation had the largest role in determining whether an outbreak was controllable when R_0 was 1.5. For R_0 values of 2.5 or 3.5, if there were 40 initial cases, contact tracing and isolation were only potentially feasible when less than

1% of transmission occurred before symptom onset (18). Therefore, efforts should be made to control this epidemic with greater vigor and urgency and to conduct a daily risk assessment. In the current epidemiological situation (26) in the world and Iran, fear control and avoidance of rumors are very important for COVID-19 prevention and control.

There are 3 important and debatable points of view about this epidemic in Iran: First, to avoid tension in the society; second, to properly interpret the COVID-19 case fatality ratio (CFR) in Iran and calculate CFR tactfully; and third, to recommend personal hygiene, including hand washing and avoiding contact with suspected patients, social distancing, discovering unknown cases of infection and early detection, tracing direct contact and isolation of patients, all of which have been emphasized by the health care officials to overcome this disease. Moreover, in interpreting this index, since the denominator of the fraction is only positive cases in hospital beds and the numerator is the number of patients who died of COVID-19. This index should also be calculated until the end of the epidemic period, and if it is until the end of the epidemic and their outcome (death/recovery) is determined, this indicator will approach the real number. The estimated case fatality ratio among medically attended patients was reported to be approximately 2% (12) and the true ratio may not be known for some time (27). The underreporting estimation is very sensitive to the baseline CFR, meaning that small errors lead to large errors in the estimate for underreporting (28).

Underreporting of COVID-19 patients (1.38% / cCFR) and modification of disease mortality had previously been

estimated by the Center for Mathematical Modeling of Infectious Diseases, London School of Health and Tropical (28). If a country has a higher adjusted CFR (eg, 20.02%), it means that only a fraction of cases have been reported (in this case, $1.38 / 20.02 = 6.89\%$ cases reported approximately). This formula can accurately estimate the statistics of all patients with COVID-19 (from asymptomatic and mild to severe cases and death).

One article reported that up to 70% of the supply chain could be cut off and the epidemic could be controlled if contact and isolation, quarantine and isolation were appropriately accomplished (18). We think the top priorities in Iran are now circular and comprehensive efforts to conduct epidemiological studies and identify all aspects of the disease (source of disease, reservoir, pathways, infectivity, incubation period, incidence and prevalence, pathogenicity, immunogenicity, herd immunity, causes, epidemic and pandemic pattern, primary and secondary attack rates, response time, time needed for isolation and quarantine, treatment regimens, vaccines and other prevention methods, disease surveillance, and statistical reporting) and evidence-based interventions and epidemic control. The experience of China and South Korea should be used to control this epidemic disease in Southeast Asian countries and in Iran, as South Korea and China were successful in controlling the disease. Cultural conditions are also effective. Given the prediction and modeling of the number of Coronavirus cases in Iran and because the virus is going to circulate in the country for at least a few weeks, we will have an ascending trend in the coming weeks.

We recommend using the WHO guideline to properly manage patients (29). Considering China's experience and the fact that it took about 70 days for the epidemic curve to flatten in China, and based on a search of scientific texts, this study provides the following recommendations:

(1) Up-to-date and accurate data on definitions related to suspected, probable, and definitive people with Coronavirus should be collected at all provincial levels in the health care system.

(2) Percentage of completion and accuracy of assessments and data should be monitored precisely and the epidemic curve should be drawn based on district, province, rural, and urban divisions and be provided to provincial and academic headquarters in an updated dashboard format.

(3) Data should be carefully recorded and analyzed regarding the pathology, time of onset of symptoms, natural course of the disease, and the outcome of the disease to determine effective strategies to prevent and determine the necessity of intervention to control the spread of the disease at different levels. At all levels of the health system (governmental and nongovernmental), medical and diagnostic interventions and their outcomes should be recorded for all patients. These records should be based on the date of onset of symptoms, the date of referral, the date of diagnosis, method of diagnosis, the date of intervention and the outcome (eg, death, recovery, discharge). Such records can widely be used to compare and evaluate the cost-effectiveness of various diagnostic and therapeutic methods. Health system staff should be trained to appropriately and accu-

rately record data, especially by means of web-based networks. This will certainly improve the quality of data recording. All epidemiological indicators that determine the epidemic pattern, including baseline R zero, attack rate, incubation period, index case, primary cases, secondary cases, and GIS mapping should be determined in provinces, cities, and nationwide, and epidemic trends should be monitored.

Access to the results of the analysis and data should be provided for researchers and experts on the basis of specific protocols available for this purpose in the world and Iran, and a thorough critique and creative theories and ideas should be elicited from all university training and research groups. The models used to predict the end of the epidemic and control it should be evaluated as well. The results of our study are inconsistent with Zhuang brief report (30), in which the data were collected from the World Health Organization (WHO). However, this report may not be accurate, as the WHO has not reported it.

Limitations

Given the urgency of the need for valid and transparent models to informed interventions and policies, some further considerations like the no consideration or account of systematic cases, testing coverage and time delay to the test results availability, seasonality, and comorbidities have not been included in this study. However, it may be feasible to consider them in revisions of the models or future studies. Moreover, the progression of disease epidemic across space-time has not been seen in Iran, as we used the parameters of Chinese models in modeling estimates in Iran to calculated R_0 . In this study, no attempt was made to detect and report cases and deaths in Iran. Although this could have been performed, given that the study has already been designed, conducted, and being reported, and given the exceptional mortality and morbidity situation, it was not feasible to go back and report the cases and deaths due to COVID-19. However, it can be done for the future similar studies. There is no fixed page in Ministry of Health and Medical Education web site for reference of daily reports and they are scattered on various pages of this site. If screening is done in the community along with biological sampling to diagnose COVID-19, the number of cases will certainly be higher and the results of modeling will change.

Conclusion

The actual trend of detecting COVID-19 cases in Iran, which has been based on people's health behaviors and government interventions, has been increasing. In this study, estimates were based on current trends, social distancing, sampling of severe cases, hospitalization, and tip of iceberg spread disease, and thus asymptomatic, mild, and moderate cases could not be calculated. We used the reports of positive COVID-19 cases in hospitals; thus, the prediction model in this study can be used for patients hospitalized due to COVID-19. Complete reliance on any type of model will lead to systematic and random error, unless modeling provides a prediction with precise and clear assumptions and inputs and outputs. To predict the flattening of the epidemic

curve, 3 models of growth with percentage of the root mean square error (RMSE) were used. Based on RMSE, Gompertz growth model was valid and predicted that the epidemic curve will be around May 13, 2020 with about 67 000 hospitalized patients and 7900 deaths (RMSE= 10%), respectively. This study suggests that government interventions and people's behaviors determine the persistence of the epidemic, and thus they should be addressed with greater responsibility, accountability, rigor, and quality.

Acknowledgments

We thank the Modeling in Health Research Center, School of Health, Deputy of Research and Technology, Shahrekord University of Medical Sciences. Furthermore, we truly appreciate the Ministry of Health and Medical Education, especially the team of daily information reports of COVID-19 in Iran and reviewers of Medical Journal of the Islamic Republic of Iran.

Also, the authors would like to extend their thanks to Dr. Hadi Raeisi and Dr. Victoria Atabaki Kianpour.

Funding

This study was financially supported by the Deputy of Research and Technology of Shahrekord University of Medical Sciences, Shahrekord, Iran (Grant no. 254).

Conflict of Interests

The authors declare that they have no competing interests.

References

- Anderson RM, Fraser C, Ghani AC, Donnelly CA, Riley S, Ferguson NM, et al. Epidemiology, transmission dynamics and control of SARS: the 2002-2003 epidemic. *Philos Trans R Soc Lond B, Biol Sci* 2004;359(1447):1091-105.
- Peeri NC, Shrestha N, Rahman MS, Zaki R, Tan Z, Bibi S, et al. The SARS, MERS and novel coronavirus (COVID-19) epidemics, the newest and biggest global health threats: what lessons have we learned? *Int J Epidemiol*. 2020 impress. DOI: 10.1093/ije/dyaa033
- Hsieh YH, Lee JY, Chang HL. SARS epidemiology modeling. *Emerging Infect Dis*. 2004;10(6):1165-7
- Zhou G, Yan G. Severe acute respiratory syndrome epidemic in Asia. *Emerging Infect Dis*. 2003;9(12):1608-10.
- Mobaraki K, Ahmadzadeh J. Current epidemiological status of Middle East respiratory syndrome coronavirus in the World From 1.1.2017 To 17.1.2018: A Cross-Sectional Study. *BMC Infect dis*. 2019;19(1):351.
- Zu ZY, Jiang MD, Xu PP, Chen W, Ni QQ, Lu GM, et al. Coronavirus Disease 2019 (COVID-19): A Perspective from China. *Radiology*. 2020:200490.
- CDC. 2019 Novel coronavirus, Wuhan, China2020. [cited 29 March 2020]. Available from: <https://www.cdc.gov/coronavirus/2019-nCoV/summary.html>.
- Porcheddu R, Serra C, Kelvin D, Kelvin N, Rubino S. Similarity in Case Fatality Rates of COVID-19/SARS-COV-2 in Italy and China. *J Infect Dev Ctries*. 2020;14:125-128.
- Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020;S1473-3099(20):30232-2.
- Adhikari SP, Meng S, Wu YJ, Mao YP, Ye RX, Wang QZ, et al. Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: a scoping review. *Infect Dis Poverty*. 2020;9(1):29
- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020. 10.1056/NEJMoa2001316
- Lipsitch M, Swerdlow DL, Finelli L. Defining the Epidemiology of Covid-19 - Studies Needed. *N Engl J Med*. 2020 Mar 26;382(13):1194-1196.
- Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *J Med Virol*. 2020, in press. doi:10.1002/jmv.25748
- Worldometer. COVID-19 CORONAVIRUS OUTBREAK. [cited: March29, 2020]. Available from: <https://www.worldometers.info/coronavirus/#countries>.
- National Committee on COVID-19 Epidemiology, Daily Situation Report on Coronavirus disease (COVID-19) in Iran; March 22, 2020. *Arch Acad Emerg Med*. 2020;8(1):e32.
- Arab-Mazar Z, Sah R, Rabaan AA, Dhama K, Rodriguez-Morales AJ. Mapping the incidence of the COVID-19 hotspot in Iran - Implications for Travellers. *Travel Med Infect Dis*. 2020;101630.
- World Health Organization. Countries, Islamic republic of Iran. WHO team arrives in Tehran to support the COVID-19 response [cited 2020 March 27]. Available from: <http://www.emro.who.int/irn/iran-news/who-team-arrives-in-tehran-to-support-the-covid-19-response.html>
- Hellewell J, Abbott S, Gimma A, Bosse NI, Jarvis CI, Russell TW, et al. Feasibility of controlling COVID-19 outbreaks by isolation of cases and contacts. *Lancet Global Health*. 2020;8(4):e488-e496.
- Research Ethic Committee. The epidemiological survey for COVID-19 in Chaharmahal and Bakhtiari & Prediction its with modelling in Iran. [cited: 2020 March 27]. Available from: <http://ethics.research.ac.ir/ProposalCertificateEn.php?id=124057&Print=true&NoPrintHeader=true&NoPrintFooter=true&NoPrintPageBorder=true&LetterPrint=true>.
- the National Committee on COVID-19 Epidemiology in Ministry of Health and Medical Education. [cited: March29, 2020]. Available from: <http://corona.behdasht.gov.ir>
- Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*. 2020;395(10225):689-697
- Novel Coronavirus (COVID-19) Cases, GitHub repository. Provided by Johns Hopkins, [cited: March29, 2020]. Available from: <https://github.com/CSSEGISandData/COVID-19>.
- Brauer F, van den Driessche P, Wu J. *Mathematical Epidemiology*, Springer-Verlag Berlin Heidelberg, 2008.
- Brauer F. *Mathematical epidemiology: Past, present, and future*. *Infect Dis Model*. 2017;2(2):113-127.
- Wang H, Wang Z, Dong Y, Chang R, Xu C, Yu X, et al. Phase-adjusted estimation of the number of Coronavirus Disease 2019 cases in Wuhan, China. *Cell Discov*. 2020 Feb 24;6:10
- Wu Z, McGoogan JM. Characteristics of and Important Lessons from the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020 Feb 24. doi: 10.1001/jama.2020.2648.
- Lipsitch M, Swerdlow DL, Finelli L. Defining the Epidemiology of Covid-19 - Studies Needed. *N Engl J Med*. 2020 Mar 26;382(13):1194-1196.
- The Centre for the Mathematical Modelling of Infectious Diseases at the London School of Hygiene & Tropical Medicine. Using a delay-adjusted case fatality ratio to estimate under-reporting. [cited: March 25, 2020]. Available from: https://cmmid.github.io/topics/covid19/severity/global_cfr_estimates.html
- World Health Organization. Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance. [cited: 28 January 2020]. Available from: <https://apps.who.int/iris/handle/10665/330893>
- Zhuang Z, Zhao S, Lin Q, Cao P, Lou Y, Yang L, et al. Preliminary estimation of the novel coronavirus disease (COVID-19) cases in Iran: A modelling analysis based on overseas cases and air travel data. *Int J Infect Dis*. 2020;20:30138-7.

Appendix. Matlab codes

```

clc
clear all
close all
%Infected Data:
data = [0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39
;2 5 18 28 43 61 95 139 245 388 593 978 1501 2336 2922 3513 4747 5823 6566 7161 8042 9000 10078 11364 12729 13938...
14991 16169 17361 18407 19644 20610 21638 23049 24811 27017 29406 32322 35408 38303 ];
%Cured Data:
dataI=[0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39
;2 2 7 18 29 35 48 53 62 73 123 175 291 435 522 739 913 1669 2134 2394 2731 2959 3276 3529 4339 4790 4996 5389 5710...
5979 6745 7635 7913 8376 8913 9625 10457 11133 11679 12391 ];
%Death Data:
dataD=[0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39;
2 2 4 6 8 12 15 19 26 34 43 54 66 77 92 107 124 145 194 237 291 354 429 514 611 724 853 988 1135 1284 1433 1556 1685...
1811 1934 2077 2234 2378 2517 2640];
% Parameter Estimation:
[x1,fval1] = fminsearch(@vonBertalanffydist,[0.2,1,2],[],data);
[x2,fval2] = fminsearch(@logisticdist,[.18,1],[],data);
[x3,fval3] = fminsearch(@gompertzdist,[1,2],[],data);
% Solve equations
tvec = [0:0.1:data(1,end)+59]; % go one time step beyond the data time
P2=logisticsoln(tvec,data(2,:),x2(1),x2(2))
P3=gompertzsoln(tvec,data(2,:),x3(1),x3(2))
P1=vonBertalanffysoln(tvec,data(2,:),x1(1),x1(2),x1(3));
%Confidence Intervals
alpha=0.05;
coefCI(P3,alpha);
% Plot curves:
plot(tvec,P3,'b',data(1,:),data(2,:),'ro','LineWidth',2.3);
legend('Gompertz Growth','Real Data',2);
dateaxis('x',6,[num2str(18),'-Feb']);
xlabel('Days');ylabel('Number of Infected Cases');
h=title(['The cumulative growth models of COVID-19 in Iran, SKUMS']);
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
function x = gompertzsoln(t,x0,a,b);
x = b.*(x0./b).^exp(-a*t) ;
function d = gompertzdist(x,data);
% vector x = [a,b]
% data should have the form: data = [t1 t2 ... tn; d1 d2 ... dn], so 2 rows, n cols
a = x(1);
b = x(2);
x0 = data(2,1); %assume (t1,d1) = (0,x0)
xt = gompertzsoln(data(1,2:end),x0,a,b);
d = norm(xt - data(2,2:end))^2; % sum of squares
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
function d = vonBertalanffydist(x,data);
% vector x = [a,b]
% data should have the form: data = [t1 t2 ... tn; d1 d2 ... dn], so 2 rows, n cols
a = x(1);
b = x(2);
c=x(3);
x0 = data(2,1); %assume (t1,d1) = (0,x0)
xt = vonBertalanffysoln(data(1,2:end),x0,a,b,c);
d = norm(xt - data(2,2:end))^2; % sum of squares
function x = vonBertalanffysoln(t,x0,a,b,c);
x = 1./(b.*(x0.^c).*(1-exp(-a.*c*t)) + (exp(-a*c*t))/b.^c).^.(1./c));
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%
%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%%

```