

Isoform-specific roles of AMPK catalytic α subunits in Alzheimer's disease

Fanpeng Zhao,¹ Chunyu Wang,^{2,3} and Xiongwei Zhu¹

¹Department of Pathology, School of Medicine, Case Western Reserve University, Cleveland, Ohio, USA. ²Department of Neurology and ³Department of Medical Genetics, Second Xiangya Hospital, Central South University, Changsha, People's Republic of China.

AMPK is a heterotrimeric complex that serves as a major sensor of energy status in eukaryotic cells. Accumulating evidence depicts a complex role of dysregulated AMPK signaling in Alzheimer's disease (AD). In this issue of the *JCI*, Zimmermann et al. report on their investigation of AD-specific differential expression of AMPK α 1 and AMPK α 2 isoforms of the catalytic subunit and demonstrate that genetic reduction of AMPK α 1, but not AMPK α 2, rescued cognitive decline in AD mouse models. These findings reveal an isoform-specific role of AMPK α in the pathogenesis of AD, which likely provides a more precise target for future therapeutic development.

AMPK activation and outcome are likely context dependent

AMPK is an evolutionally conserved Ser/Thr kinase that serves a crucial physiological function as a cellular energy sensor (1). It is a heterotrimeric complex composed of a catalytic α subunit and two regulatory β and γ subunits at 1:1:1 ratio. The β subunit acts as a scaffold, interacting with the α subunit containing the kinase domain and the γ subunit containing four nucleotide-binding sites that allow AMPK to sense the status of cellular energy state. Two isoforms of α and β subunits and three isoforms of the γ subunit have been identified, giving rise to 12 distinct AMPK complexes (1). Despite functional redundancy, these complexes may differ in their tissue- and cell-specific distribution, regulation, subcellular localization, and biochemical properties and function (2). Cellular ATP concentration is kept at a constant level to ensure adequate ATP supply, which is essential to the survival of mammalian cells. Energy stress causes changes in AMP/ATP and ADP/ATP ratios. AMP binding to the γ subunit allosterically acti-

vates the complex and promotes the phosphorylation of Thr172 at the kinase domain of α subunit by upstream kinase of LKB1, which is a step required for catalytic activation of the complex (1). A rise in intracellular Ca^{2+} also activates AMPK through CAMKK2-mediated phosphorylation of Thr172. AMPK activation is instrumental in restoration of energy balance, acting by turning off energy-consuming processes, such as protein and lipid synthesis, and turning on energy-generating processes such as glucose metabolism, mitochondrial biogenesis, and autophagy, through a myriad of effectors (1).

As one of the high energy-consuming organs, the brain is vulnerable to disturbance in energy metabolism. A large body of evidence has demonstrated reduced energy metabolism as an early and consistent feature in Alzheimer's disease (AD) (3). Insulin resistance, abnormal glucose transport, mitochondrial dysfunction, dysregulated cholesterol metabolism, and calcium homeostasis are also prominent features in the AD brain (3, 4). These studies implicate a potential role of AMPK sig-

naling in AD. Indeed, recent studies found putative pathogenic somatic mutations in the AD brain enriched in the AMPK pathway genes (5). Robust AMPK activation, as evidenced by increased Thr172 phosphorylation, was found in tangle- and pre-tangle-bearing neurons in AD brain (6). However, accumulating evidence appears to depict a complex role of dysregulated AMPK signaling in AD: AMPK directly phosphorylates tau protein, but it is also able to inhibit tau phosphorylation/aggregation through GSK3 β inhibition or SIRT1 activation-mediated deacetylation of tau (7). AMPK activation could increase A β generation through ER stress (8) or transcriptional upregulation of BACE1 (9); however, it can also reduce A β levels by reducing lipid raft localization of APP protein through modulation of sphingomyelin levels or by enhancing A β clearance. AMPK activation by metformin alleviated mitochondrial, pathological, and cognitive deficits in AD models (10, 11), but AMPK inhibitor compound C treatment corrected toxic effects of A β on synaptic function (12, 13). While these studies underscore the critical roles of AMPK in the development of many aspects of AD, the controversy highlights the notion that AMPK activation and outcome are likely context dependent. It is possible that different isoforms and/or distinct AMPK complexes may contribute to the complexity (14).

AMPK isoform differences in Alzheimer's disease

Neurons show widespread expression of both AMPK α 1 and AMPK α 2 isoforms of the catalytic subunit. The AMPK α 2 isoform is the predominant subunit in the brain, with constantly high neuronal expression, especially in the cortex and hippocampus, while AMPK α 1 isoform expression peaks in embryos and declines during development (15, 16). Although AMPK is activated in AD, it is not known which AMPK α isoforms contribute to the

► **Related Article:** p. 3511

Conflict of interest: The authors have declared that no conflict of interest exists.

Copyright: © 2020, American Society for Clinical Investigation.

Reference information: *J Clin Invest.* 2020;130(7):3403–3405. <https://doi.org/10.1172/JCI137908>.

disease. In this issue of the *JCI*, Zimmermann et al. (17) aimed to fill this important gap in our knowledge. The researchers found that expression of the AMPK α 1 isoform was consistently elevated in the hippocampus from both sporadic and familial AD patients and in the Tg19959 APP transgenic mouse model. In contrast, AMPK α 2 expression was decreased in the hippocampus from sporadic AD patients. These changes in AMPK α expression appeared to be AD specific, since AMPK α 1/2 levels were unaffected in either Lewy body dementia (LBD) or frontotemporal dementia (FTD). Importantly, reduced expression of AMPK α 1, but not AMPK α 2, in hippocampal and cortical neurons effectively rescued synaptic deficits and memory decline in 2 different amyloid- β -bearing AD mouse models. However, it is somewhat surprising that no effects on amyloid pathology were found. Nevertheless, the reduction of AMPK α 1, but not AMPK α 2, corrected AD-associated hyperphosphorylation of eukaryotic elongation factor 2 (eEF2) and restored de novo protein synthesis (17). This change in protein synthesis is consistent with the rescuing effects on synaptic function and cognition, since nascent protein synthesis is essential for producing the synaptic modification needed for long-term memory storage (18). Given that prior studies from this same group demonstrated a critical role of aberrant eEF2 signaling in causing cognitive and pathological deficits in AD (19), these data suggest that upregulation of AMPK α 1 in AD caused cognitive declines at least partly through eEF2-mediated chronic repression of protein synthesis. Therefore, this study provided a first set of evidence identifying an isoform-specific pathogenic role of AMPK α 1 in AD, which paves the road for a more precise target development in the fight against AD (17).

Conclusions and future directions

While finding and characterizing an isoform-specific role of AMPK α in AD is a leap forward (17), future efforts to address some additional questions are certainly needed before we move ahead with translational studies: (a) reduced AMPK α 1 rescued cognitive decline in APP/PS1 mice, yet metformin protected against cognitive and pathological deficits in this same

mouse model, presumably through AMPK activation (10). How can we reconcile these contradictory findings? Does metformin selectively act on specific AMPK complexes other than those containing AMPK α 1? Or, perhaps, did metformin mobilize additional pathways in concert with AMPK activation that are needed for protection? (b) It is unclear whether the expression pattern of AMPK α isoforms is the same in rodents and in humans, affecting whether this study (17) is applicable to human disease. This isoform expression ambiguity is not trivial, since prior studies demonstrated species-specific roles of different AMPK subunits. For example, AMPK γ 2 is highly expressed in the human heart, but not in the mouse heart (2). AMPK γ 2 mutations caused human Wolff-Parkinson-White syndrome, affecting the heart, but knockin mice with these mutations failed to recapitulate the heart phenotype (2). (c) The two APP transgenic mouse models (Tg19959 and APP/PS1) used in this study (17) are models with relatively rapidly progressive amyloid pathology and cognitive dysfunction. Slower progressive disease mouse models may need to be used to confirm the finding. It is worth noting that AMPK activation appears to be detrimental in a quickly developing mouse model of advanced Huntington disease (20), while it is protective in progressive animal models of early phase Huntington disease (21). Detailed characterization of AMPK signaling in different AD stages will also be of value.

Unlike reduced AMPK α 1 expression, reduced AMPK α 2 expression had no beneficial effect in AD mouse models. Does the result suggest AMPK α 2 is not involved in AD? Probably not. It should be noted that the protein level of AMPK α 2 was dramatically decreased in the hippocampus of sporadic AD patients. It is unlikely this decrease was a compensation for AMPK α 1 elevation, since AMPK α 2 levels were maintained when AMPK α 1 levels were changed, as occurred in the brain from either familial AD patients or AMPK α 1 haploinsufficient mice. More importantly, reduced AMPK α 2 expression by itself impaired long-term synaptic plasticity and caused memory deficits, highlighting the importance of AMPK α 2 as the predominant isoform in the brain. These data hint that AMPK α 2 reduction may

contribute to AD. Indeed, reduced AMPK α 2 expression exacerbated eEF2 hyperphosphorylation and protein synthesis inhibition in the AD mouse model. Therefore, future studies to restore AMPK α 2 signaling in AD models are warranted to clarify the role of AMPK α 2 in AD.

Given that changes in energy metabolism are implicated in many neurodegenerative diseases, it is puzzling that changes of AMPK α were only found in AD and not in other neurodegenerative diseases, such as FTD and DLB. This AD-specific effect may prompt one to speculate that different patterns of brain energy crisis accompany different types of diseases. However, prior studies demonstrated AMPK activation, measured by Thr172 phosphorylation, in cerebral neurons of multiple tauopathies, including Pick's disease and progressive supranuclear palsy (6). Since Zimmermann et al. (17) did not determine AMPK activation/phosphorylation in the human brain, there is a possibility that phosphorylation, rather than expressional change, may activate AMPK in FTD or DLB samples. Nevertheless, the isoform-specific dysregulation of AMPK in postmortem AD brain tissue raised the important question of how AMPK α 1/2 is differentially regulated in AD. While A β could induce AMPK α phosphorylation via Ca²⁺-mediated CAMKK2 activation (12), not much is known about how AMPK α expression may be modulated. In this regard, it is of interest to mention that AMPK α expression could be induced in activated astrocytes in the brain (15).

Zimmermann et al. (17) explored the mechanisms that underlie the protective effects of reduced AMPK α 1 in APP Tg mice by focusing on protein synthesis. Given the distinct effects of AMPK α 1 and AMPK α 2 in the AD mouse, the authors performed mass spectrometry analysis and revealed that expression of ten proteins was uniquely restored by AMPK α 1, but not by AMPK α 2, which is worth further investigation. However, AMPK is a versatile signaling protein with multiple effectors that regulate various important cellular functions (7), many of which are altered in AD. Therefore, it remains to be determined what other downstream pathways, such as mitochondrial biogenesis (22), autophagy, cholesterol dyshomeostasis (23), etc., are selectively rescued by AMPK α 1 reduction.

Acknowledgments

This work was supported in part by the NIH (AG049479 and AG056363) and the Alzheimer's Association (AARG-16-443584).

Address correspondence to: Xiongwei Zhu, Department of Pathology, Case Western Reserve University, 2103 Cornell Road, Cleveland, Ohio 44106, USA. Phone: 216.368.5903; Email: xiongwei.zhu@case.edu. Or to: Chunyu Wang, Department of Neurology, Second Xiangya Hospital, Central South University, 139 People Road, Changsha, Hunan, China. Phone: 86.15073144560; Email: wangchunyu@csu.edu.cn.

- Lin SC, Hardie DG. AMPK: sensing glucose as well as cellular energy status. *Cell Metab.* 2018;27(2):299–313.
- Carling D. AMPK signalling in health and disease. *Curr Opin Cell Biol.* 2017;45:31–37.
- Wang W, Zhao F, Ma X, Perry G, Zhu X. Mitochondria dysfunction in the pathogenesis of Alzheimer's disease: recent advances. *Mol Neurodegener.* In press.
- Jiang S, Shao C, Tang F, Wang W, Zhu X. Dynamin-like protein 1 cleavage by calpain in Alzheimer's disease. *Aging Cell.* 2019;18(3):e12912.
- Park JS, et al. Brain somatic mutations observed in Alzheimer's disease associated with aging and dysregulation of tau phosphorylation. *Nat Commun.* 2019;10(1):3090.
- Vingtdeux V, Davies P, Dickson DW, Marambaud P. AMPK is abnormally activated in tangle- and pre-tangle-bearing neurons in Alzheimer's disease and other tauopathies. *Acta Neuropathol.* 2011;121(3):337–349.
- Salminen A, Kaamiranta K, Haapasalo A, Soininen H, Hiltunen M. AMP-activated protein kinase: a potential player in Alzheimer's disease. *J Neurochem.* 2011;118(4):460–474.
- Yoon SO, et al. JNK3 perpetuates metabolic stress induced by A β peptides. *Neuron.* 2012;75(5):824–837.
- Chen Y, et al. Antidiabetic drug metformin (GlucophageR) increases biogenesis of Alzheimer's amyloid peptides via up-regulating BACE1 transcription. *Proc Natl Acad Sci U S A.* 2009;106(10):3907–3912.
- Ou Z, et al. Metformin treatment prevents amyloid plaque deposition and memory impairment in APP/PS1 mice. *Brain Behav Immun.* 2018;69:351–363.
- Chiang MC, Cheng YC, Chen SJ, Yen CH, Huang RN. Metformin activation of AMPK-dependent pathways is neuroprotective in human neural stem cells against Amyloid-beta-induced mitochondrial dysfunction. *Exp Cell Res.* 2016;347(2):322–331.
- Mairet-Coello G, Courchet J, Pieraut S, Courchet V, Maximov A, Polleux F. The CAMKK2-AMPK kinase pathway mediates the synaptotoxic effects of A β oligomers through Tau phosphorylation. *Neuron.* 2013;78(1):94–108.
- Ma T, et al. Inhibition of AMP-activated protein kinase signaling alleviates impairments in hippocampal synaptic plasticity induced by amyloid β . *J Neurosci.* 2014;34(36):12230–12238.
- Rajamohan F, et al. Probing the enzyme kinetics, allosteric modulation and activation of α 1- and α 2-subunit-containing AMP-activated protein kinase (AMPK) heterotrimeric complexes by pharmacological and physiological activators. *Biochem J.* 2016;473(5):581–592.
- Turnley AM, Stapleton D, Mann RJ, Witters LA, Kemp BE, Bartlett PF. Cellular distribution and developmental expression of AMP-activated protein kinase isoforms in mouse central nervous system. *J Neurochem.* 1999;72(4):1707–1716.
- Culmsee C, Monnig J, Kemp BE, Mattson MP. AMP-activated protein kinase is highly expressed in neurons in the developing rat brain and promotes neuronal survival following glucose deprivation. *J Mol Neurosci.* 2001;17(1):45–58.
- Zimmermann HR, et al. Brain-specific repression of AMPK α 1 alleviates pathophysiology in Alzheimer's model mice. *J Clin Invest.* 2020;130(7):3511–3527.
- Hernandez PJ, Abel T. The role of protein synthesis in memory consolidation: progress amid decades of debate. *Neurobiol Learn Mem.* 2008;89(3):293–311.
- Beckelman BC, et al. Genetic reduction of eEF2 kinase alleviates pathophysiology in Alzheimer's disease model mice. *J Clin Invest.* 2019;129(2):820–833.
- Ju TC, et al. Nuclear translocation of AMPK-alpha1 potentiates striatal neurodegeneration in Huntington's disease. *J Cell Biol.* 2011;194(2):209–227.
- Vázquez-Manrique RP, et al. AMPK activation protects from neuronal dysfunction and vulnerability across nematode, cellular and mouse models of Huntington's disease. *Hum Mol Genet.* 2016;25(6):1043–1058.
- Sheng B, et al. Impaired mitochondrial biogenesis contributes to mitochondrial dysfunction in Alzheimer's disease. *J Neurochem.* 2012;120(3):419–429.
- Williams T, Borchelt DR, Chakrabarty P. Therapeutic approaches targeting apolipoprotein E function in Alzheimer's disease. *Mol Neurodegener.* 2020;15(1):8.