

VOLUME 295 (2020) PAGES 2270-2284

DOI 10.1074/jbc.AAC120.014344

Correction: DAPLE protein inhibits nucleotide exchange on G α s and G α q via the same motif that activates G α i.

Arthur Marivin, Marcin Maziarz, Jingyi Zhao, Vincent DiGiacomo, Isabel Olmos Calvo, Emily A. Mann, Jason Ear, Juan B. Blanco-Canosa, Elliott M. Ross, Pradipta Ghosh, and Mikel Garcia-Marcos

MgCl₂ was omitted as a reagent in two buffers described under “*In vitro* protein-binding assays with GST-fused proteins” section (under “Experimental procedures”). “Beads were washed twice with PBS and resuspended in 300 μ l of binding buffer (50 mM Tris-HCl (pH 7.4), 100 mM NaCl, 0.4% (vol:vol) NP-40, 5 mM EDTA, 2 mM DTT)” should read as “Beads were washed twice with PBS and resuspended in 300 μ l of binding buffer (50 mM Tris-HCl (pH 7.4), 100 mM NaCl, 10 mM MgCl₂, 0.4% (vol:vol) NP-40, 5 mM EDTA, 2 mM DTT).” “Beads were washed four times with 1 ml of wash buffer (4.3 mM Na₂HPO₄, 1.4 mM KH₂PO₄, pH 7.4, 137 mM NaCl, 2.7 mM KCl, 0.1% (vol:vol) Tween-20, 5 mM EDTA, 1 mM DTT)” should read as “Beads were washed four times with 1 ml of wash buffer (4.3 mM Na₂HPO₄, 1.4 mM KH₂PO₄, pH 7.4, 137 mM NaCl, 2.7 mM KCl, 0.1% (vol:vol) Tween-20, 10 mM MgCl₂, 5 mM EDTA, 1 mM DTT).”