




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short communication

Ultrasound findings in patients with COVID-19 pneumonia in early and late stages: Two case-reports

Laurent Zieleskiewicz^{a,b}, Gary Duclos^a, Ophélie Dransart-Rayé^c, Nicolas Nowobilski^c, Belaid Bouhemad^{c,d,*}

^a Department of Anaesthesia and Intensive Care Medicine, University Hospital of Marseille, Aix-Marseille University, Marseille, France

^b C2VN Inra Inserm, Faculty of medicine, Aix-Marseille University, Marseille, France

^c Department of Anaesthesiology and Intensive Care, University Hospital of Dijon, Dijon, France

^d Lipness Team, Inserm Research Centre LNC-UMR1231 and LabEx LipSTIC, Université Bourgogne Franche-Comté, Dijon, France



ARTICLE INFO

Article history:

Received 30 April 2020

Accepted 1st May 2020

Available online 3 July 2020

Morphology assessment of COVID-19 pneumonia may be crucial for its diagnosis and management. Chest computed tomography (CT) scan has a high diagnosis accuracy and may be associated with patient clinical condition [1]. However, during a pandemic the safe use of CT is questionable. Given its characteristics, CT scan cannot be considered as a screening tool (too wide target population) or a monitoring tool (intra-hospital transfer, radiation, availability).

Lung ultrasound (LUS) has been validated for the diagnosis and the management of acute respiratory failure, pneumonia [2], even during viral pandemic [3]. LUS has a high diagnosis accuracy for interstitial syndrome and alveolar consolidations, which are the most common radiological signs of COVID-19 pneumonia [4]. Moreover, preliminary reports are in favour of a predominant involvement of the pulmonary periphery, which could reinforce LUS interest in the actual pandemic.

Here two cases of confirmed COVID-19 patients are described at early and late stages of the disease. We compared the CT scan and ultrasound features performed at the same time.

In a 70 year-old woman, at admission, CT scan showed the typical pattern of COVID-19 with peripheral ground-glass opacities that were bilateral and multilobar (Fig. 1A and B). Ground-glass opacities appear as coalescent B-line areas and are associated with irregularities of pleural line (white arrows, Fig. 1A, B and C). This pattern alternates with normal area. Nodular or mass-like ground-

glass opacities can also reach the periphery and be depicted as small consolidations in subpleural areas (red arrows, Fig. 1A, B, D and E) or consolidation (yellow arrows, Fig. 1A, B and E).

A 65-year old male, after 25 days of mechanical ventilation for acute respiratory distress syndrome, developed on the CT scan fibrosis (Fig. 2A, B) consisting on subpleural fibrosis, honeycombing, traction bronchiectasis with anterior distribution and thickened interlobular septal [5]. These abnormalities using LUS were reported as irregularities of pleural line associated with coalescent B-lines (red arrows Fig. 2A, C and D) or multifocal subpleural consolidation (red arrows, Fig. 2C and D). Thickened interlobular septal seen as regularly spaced B-lines (yellow arrows Fig. 2E) combined with thickened pleural line allows the visualisation of secondary pulmonary lobules (white arrow; Fig. 2E). Thickened interlobular septal and small subpleural consolidations evidenced the major fissure (red arrows Fig. 2B and E).

These two suggestive cases underline the potential of LUS for COVID-19 pneumonia assessment at different stages of the disease. Even if all LUS signs are not specific, presence of these various patterns in a clinical context is very suggestive of viral pneumonia. When the clinical context is evocative, LUS may therefore help physicians to identify COVID-19 patients. After a proper validation of LUS against chest CT scan, one can believe that LUS could serve as a diagnostic and a triage tool detecting at risk patients, thus allowing a better allocation of ICU resources as previously described in another context [6]. Finally, it seems important to underline that LUS should also be associated with multi-organ point-of-care ultrasound. In particular, COVID-19 infection is associated with myocarditis and an elevated incidence of thromboembolic events. Multi-organ point-of-care ultrasound could therefore be of high interest for the screening of these complications at the bedside with a minimum exposition of healthcare providers.

* Corresponding author at: Department of Anaesthesiology and Intensive Care, Unity of Cardiovascular Anaesthesiology and Intensive Care, CHU de Dijon, BP 77908, 21709 Dijon Cedex, France.

E-mail address: belaid_bouhemad@hotmail.com (B. Bouhemad).




Fig. 1. CT scan and lung ultrasound in a 70-year old woman, at admission revealing ground-glass opacities.




Fig. 2. CT scan and lung ultrasound in a 65-year old male, after 25 days of mechanical ventilation revealing fibrosis.

Human and animal rights

Not applicable.

Informed consent and patient details

Not applicable.

Disclosure of interest

LZ received fees for teaching ultrasound to GE healthcare costumers. The other authors declare that they have no competing interest.

Funding

This work did not receive any grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions

All authors attest that they meet the current International Committee of Medical Journal Editors (ICMJE) criteria for Authorship.

Prior abstract publication/presentation

No.

Ethics approval and consent to participate

Written consents from the patients were waived, because of entirely anonymised images from which the individual cannot be identified.

Acknowledgements

None.

References

- [1] Zhao W, Zhong Z, Xie X, Yu Q, Liu J. Relation between chest CT findings and clinical conditions of Coronavirus Disease (COVID-19) pneumonia: a multicenter study. *AJR Am J Roentgenol* 2020;214(5):1072–7.
- [2] Mojoli F, Bouhemad B, Mongodi S, Lichtenstein D. Lung ultrasound for critically ill patients. *Am J Respir Crit Care Med* 2019;199(6):701–14.
- [3] Testa A, Soldati G, Copetti R, Giannuzzi R, Portale G, Gentiloni-Silveri N. Early recognition of the 2009 pandemic influenza A (H1N1) pneumonia by chest ultrasound. *Crit Care* 2012;16(1):R30.
- [4] Wu J, Wu X, Zeng W, Guo D, Fang Z, Chen L, et al. Findings in patients with Coronavirus Disease 2019 and its relationship with clinical features. *Invest Radiol* 2020;55(5):257–61.
- [5] Desai SR, Wells AU, Rubens MB, Evans TW, Hansell DM. Acute respiratory distress syndrome: CT abnormalities at long-term follow-up. *Radiology* 1999;210(1):29–35.
- [6] Markarian T, Zieleskiewicz L, Perrin G, Claret PG, Loundou A, Michelet P, et al. A lung ultrasound score for early triage of elderly patients with acute dyspnea. *CJEM* 2019;21(3):399–405.