CORRECTION

Correction to: Predictive validity of a novel non-invasive estimation of effective shunt fraction in critically ill patients

Emma M. Chang[1†], Andrew Bretherick[1,2†], Gordon B. Drummond[1] and J. Kenneth Baillie[1,3*]

Correction to: ICMx 7, 49 (2019)
https://doi.org/10.1186/s40635-019-0262-1

Following publication of the original article [1], the authors reported an error in Eq. 2 of the article; the article is missing brackets.

Please find the correct version (with the brackets added) of the equation in this correction.

The authors apologize for any inconvenience caused.

$$\frac{Q_S}{Q_T} = \frac{C_{c'}O_2 - C_aO_2}{C_{c'}O_2 - \left(C_aO_2 - \frac{VO_2}{Q}\right)}$$

Author details
¹Anaesthesia, Critical Care and Pain Medicine, Royal Infirmary of Edinburgh, Edinburgh EH16 4SA, UK. ²MRC Institute of Genetics and Molecular Medicine, The University of Edinburgh, Edinburgh EH4 2XU, UK. ³The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush, Edinburgh EH25 9RG, UK.

Published online: 10 July 2020

Reference
1. Chang EM, Bretherick A, Drummond GB et al (2019) Predictive validity of a novel non-invasive estimation of effective shunt fraction in critically ill patients. ICMx 7:49 https://doi.org/10.1186/s40635-019-0262-1