



In-Depth Longitudinal Study of *Listeria monocytogenes* ST9 Isolates from the Meat Processing Industry: Resolving Diversity and Transmission Patterns Using Whole-Genome Sequencing

Annette Fagerlund,^a Solveig Langsrud,^a Trond Mørretø^a

^aNofima—Norwegian Institute of Food, Fisheries and Aquaculture Research, Ås, Norway

ABSTRACT *Listeria monocytogenes* is a pathogen mostly associated with the consumption of ready-to-eat foods and can cause severe disease and death. It can be introduced into food chains from raw materials, but often the contamination source is the food production environment, where certain clones can persist for years. In the meat chain, ST9 is one of the most commonly encountered *L. monocytogenes* sequence types, and for effective source tracking, the divergence and spread of ST9 must be understood. In this study, whole-genome sequencing (WGS) was used to characterize and track 252 *L. monocytogenes* ST9 isolates collected from four Norwegian meat processing plants between 2009 and 2017. The isolates formed distinct clusters relative to genomes found in public databases, and all but three isolates clustered into two major clonal populations. Different contamination patterns were revealed, e.g., evidence of contamination of two factories with a clone that diverged from its ancestor in the late 1990s through a common source of raw materials; breach of hygienic barriers within a factory, leading to repeated detection of two clones in the high-risk zone during a 4- to 6-year period; entry through the purchase and installation of second-hand equipment harboring a previously established clonal population; and spreading and diversification of two clones from two reservoirs within the same production room over a 9-year period. The present work provides data on the diversity of ST9, which is crucial for epidemiological investigations and highlights how WGS can be used for source tracking within food processing factories.

IMPORTANCE *Listeria monocytogenes* is a deadly foodborne pathogen that is widespread in the environment, and certain types can be established in food factories. The sequence type ST9 dominates in meat processing environments, and this work was undertaken to obtain data needed for the tracking of this subtype. By using whole-genome sequencing (WGS), we revealed the presence of cross-contamination routes between meat factories as well as within a single factory, including the spread from different reservoirs within the same room. It was also possible to estimate the time frame of persistence in the factory, as well as when and how new clones had entered. The present work contributes valuable information about the diversity of ST9 and exemplifies the potential power of WGS in food safety management, allowing the determination of relationships between strains both in an international context and locally between and within factories.

KEYWORDS CC9, *Listeria monocytogenes*, ST9, WGS, food processing environment, food safety, meat industry, meat processing, whole-genome sequencing

L*isteria monocytogenes* is a pathogenic bacterium causing the severe foodborne disease listeriosis, which is mostly linked to the consumption of ready-to-eat (RTE) food like fresh produce, soft cheeses, lightly processed fish products, and meat prod-

Citation Fagerlund A, Langsrud S, Mørretø T. 2020. In-depth longitudinal study of *Listeria monocytogenes* ST9 isolates from the meat processing industry: resolving diversity and transmission patterns using whole-genome sequencing. *Appl Environ Microbiol* 86:e00579-20. <https://doi.org/10.1128/AEM.00579-20>.

Editor Christopher A. Elkins, Centers for Disease Control and Prevention

Copyright © 2020 Fagerlund et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Annette Fagerlund, annette.fagerlund@nofima.no.

Received 9 March 2020

Accepted 6 May 2020

Accepted manuscript posted online 15 May 2020

Published 2 July 2020

ucts, such as cooked sausages and cold cuts. In 2017 and 2018, the largest foodborne outbreak of listeriosis reported so far, with 1,060 confirmed cases and 216 deaths, was reported in South Africa and linked to the consumption of an RTE meat product (1). Several other listeriosis outbreaks also have been linked to RTE meat products (2, 3). The main route of contamination of *L. monocytogenes* to RTE food products is through cross-contamination from the processing environment (4, 5). Many studies have reported that clones of *L. monocytogenes* may establish and persist for years in processing environments, including in factories in the meat industry (5–15).

Isolates of *L. monocytogenes* can be subdivided into genetic groups, including different sequence types (STs) from multilocus sequence typing (MLST), in which several related STs comprise a central clonal complex (CC). CC9, which includes ST9, is a frequently detected clone in meat products and in meat industry environments; it is common in many countries and was ranked the fourth most common clone worldwide (16–22). For instance, CC9 was the second most prevalent clone isolated from foods in a study comprising 3,333 food isolates collected in France, and the majority of the CC9 isolates (66.4%) were from meat products (23). ST9 has also been shown to be the most frequently isolated (33%) *L. monocytogenes* type in Spanish meat processing plants, detected in 14 of 18 sampled plants (10). Several other studies also report ST9 contaminations in meat industry environments (12–15, 24, 25). For example, in a previous study (26), we subtyped a total of 680 *L. monocytogenes* isolates from Norwegian meat and salmon processing plants and found that 70% of the isolates from the meat industry (obtained mainly from two extensively sampled factories) belonged to ST9.

To better understand contamination routes within food industry environments and during foodborne outbreaks of *L. monocytogenes*, there is a need for more information about the genetic variation between environmental isolates over time, both within single facilities and between different processing plants. Whole-genome sequencing (WGS) analysis, employing either genomic MLST or single-nucleotide polymorphism (SNP) analyses, has become the gold standard for typing of bacteria. However, the majority of publicly available WGS data from *L. monocytogenes* environmental isolates are from inspections by authorities, often performed as part of outbreak investigations, resulting in a sampling bias for data deposited in public databases (27). Furthermore, a study examining the genetic distances between *L. monocytogenes* collected by the FDA during inspections in the United States during 1999 to 2017 covered WGS data for 5,321 isolates from 846 facilities (28); however, for 93% of the facilities included in the study, fewer than 20 isolates were collected (isolates per facility, median of 2 and range of 1 to 98). This suggests that the number of isolates collected by authorities from each facility is usually limited. In most studies where WGS-based strategies have been employed to investigate *L. monocytogenes* clones from food processing environments, only isolates from a single processing facility were considered, and only limited numbers of isolates belonging to the same cluster or strain were analyzed (12, 15, 24, 29–33). Thus, the description of contamination patterns is, in most cases, primarily based on looking at the occurrence of certain sequence types in different parts of the factory. For example, in a study that sampled *L. monocytogenes* from a meat establishment with slaughter, deboning, and meat processing units for 1 year, eight *L. monocytogenes* isolates were isolated and subjected to WGS. Five ST9 isolates, originating from both the slaughtering line and the meat processing units, formed a cluster whose maximum pairwise SNP distance was five, leading to the conclusion that the strains from the processing units originated from the slaughter line (24). Although this study used WGS to provide insight into contamination patterns, sampling was performed on a limited number of occasions (4 days during 1 year), resulting in a small number of obtained isolates, likely limiting insight into the genetic diversity of strains in the processing plant.

As an empirical practice used in outbreak investigations, it has been suggested that for *L. monocytogenes*, a threshold of fewer than 20 SNPs indicates an epidemiological link between isolates (28, 34). However, in contrast to that observed in typical point-

TABLE 1 Overview of *L. monocytogenes* ST9 isolates subjected to WGS analysis

Plant	No. of isolates for each plant and yr ^a									Total	
	2009	2010	2011	2012	2013	2014	2015	2016	2017		
M1	2	2	5	39	13	12				34	107
M4				35	18	74	10			1	138
M5					4						4
M7							3				3

^aIsolates from plant M7 and isolates from 2017 are from the present study, and the remaining isolates were from Mørretrø et al. (26).

source outbreaks, outbreaks caused by a population of persistent strains, which have had time to evolve in their environment, may have genetic diversity greater than 20 SNPs (35, 36). Additional evidence, such as epidemiological data and traceback evidence, as well as the phylogenetic tree topology, is, in any case, required to establish the contamination source during outbreaks (27, 37), as more than one facility may be home to the same clone of *L. monocytogenes*. This may occur, for example, in cases where suppliers of contaminated raw materials have more than one customer. A few WGS-based studies have considered the diversity of *L. monocytogenes* isolates across multiple food-associated facilities. For example, in a longitudinal study by Stasiewicz et al. (38), 175 isolates of *L. monocytogenes* sampled over 2 years in 30 retail delis were subjected to WGS. The results showed the presence of both persistent isolates within a deli and strains repeatedly introduced from external sources. The finding of almost identical isolates in different delis also indicated a common source for the introduction of strains. In a study by Morganti et al. (25), 108 *L. monocytogenes* isolates were obtained from 38 Parma ham production facilities, and isolates with pulsed-field gel electrophoresis (PFGE) profiles found in more than one facility ($n = 69$) were subjected to WGS analysis. Seven different STs were identified, of which the most prevalent was ST9 ($n = 24$), and three tight clusters, differing in only 1 or 2 core SNPs, were detected in more than one facility, suggesting interlinks between the establishments. However, in general, studies investigating the diversity of persistent clones found in more than one facility are lacking. To avoid misleading conclusions regarding contamination sources, increased knowledge about the diversity of closely related *L. monocytogenes* isolates in practical cases is needed.

In the present study, 252 isolates of *L. monocytogenes* ST9 from four Norwegian meat processing plants were subjected to WGS analysis to get further insight into the diversity, transmission patterns, and entry routes. The goals of the present study were to (i) apply whole-genome MLST (wgMLST), core genome MLST (cgMLST), and SNP analyses to assess the genetic diversity of the ST9 strains; (ii) determine whether certain genetic clusters of ST9 were factory specific, to what extent they were present in the same factory during extended periods of time, and how they related to isolates from other sources; and (iii) explore whether WGS analyses can be used to differentiate between contamination scenarios: hygienic breach between zones, dissemination of persistent listeria from a specific niche, or wide dissemination of a specific variant within a zone.

RESULTS

WGS of Norwegian ST9 isolates from meat processing industry. A total of 252 *L. monocytogenes* ST9 isolates from four Norwegian meat processing plants were subjected to whole-genome sequencing (WGS) analysis. The four processing plants were not related; none belonged to the same company, and they were located in different geographic areas of Norway. The isolates were mainly from samples taken from floors, drains, or food processing equipment (26). The majority ($n = 245$) were collected from two plants, M1 and M4, over a period of 9 and 6 years, respectively (Table 1; see also Table S1 in the supplemental material). Isolates from 2017 were from the plants' regular environmental hygiene monitoring programs. ST9 isolates had also been found in two less intensively sampled facilities (M5 and M7), and these were included for comparison.




FIG 1 Minimum spanning tree based on cgMLST allelic profiles, showing the relationship between the 252 isolates from the Norwegian meat processing industry, colored according to the factory where they were isolated, and 202 CC9 reference genomes obtained from public databases (white nodes). The area of each circle is proportional to the number of isolates represented, and the number of allelic differences between isolates is indicated on the edges connecting two nodes. Labels Clade A and Clade B refer to the two main clusters of isolates from the Norwegian meat processing industry described in the text. Selected reference strains are indicated with red arrows.

In silico MLST analysis confirmed that all isolates belonged to ST9, except one isolate (MF6279), which was technically not ST9, as it lacked the *bgIA* gene.

The Norwegian clones formed distinct clusters relative to public reference strains. To examine the relationship between the ST9 isolates from Norwegian meat processing plants and those from other sources, a representative subset of CC9 genomes found in public databases ($n = 202$) were selected for comparative analysis. A core genome MLST (cgMLST) analysis was performed using the subset of 1,748 cgMLST loci from the scheme described by Moura et al. (19). In this analysis, 88 polymorphic alleles were detected within the Norwegian data set, while a maximum of 160 allelic differences were identified between any pair from the 454 CC9 isolates included in the analysis. With the exception of three outliers, the Norwegian isolates clustered into two major genetic lineages, here referred to as Clade A and Clade B (Fig. 1). Isolates in these two clusters were separated by between 25 and 33 (median, 31) cgMLST alleles. Since the majority of the Norwegian isolates ($n = 246$) were represented by only four cgMLST profiles, three of which comprised isolates from two different meat processing plants, cgMLST typing was not sufficiently discriminative for the character-

ization of the diversity within the Norwegian data set. However, the analysis was sufficiently sensitive for differentiating between the Norwegian clones and those in the publicly available reference data set (Fig. 1). The reference strains most closely related to the Norwegian isolates were *L. monocytogenes* AT3E and NRRL B-57115, which were each separated by 3 or 4 alleles from isolates within clade B. AT3E was isolated from a food product in Finland in 1995 (39), while NRRL B-57115 (alias BCW_4316 [40]) was isolated from Italian roast beef in the United States. For the Norwegian clade A strains, the seven most closely related reference strains had 7 to 10 pairwise allelic differences compared to the closest Norwegian isolates. The two outlier strains MF6338 and MF6339, from plant M7, were separated by 1 and 2 cgMLST alleles from *L. monocytogenes* BCW_4274 (strain origin not indicated [40]). In conclusion, the Norwegian isolates in clades A and B were more related to each other than to any external/foreign strain from the cross-section of CC9 genomes from public databases.

The ST9 isolates belong to two major clones. To differentiate between the Norwegian isolates, whole-genome MLST (wgMLST) and single-nucleotide polymorphism (SNP) analyses were performed. The wgMLST analysis detected a total of 3,334 loci, of which 746 were polymorphic, and between any pair of isolates a maximum of 179 (median, 95) allelic differences were found. As also observed from the cgMLST analysis, all but three of the isolates clustered into two major genetic lineages, clade A and clade B (Fig. 2A and B). Clade A includes isolates from all four investigated meat processing plants ($n = 112$), while only representatives from plants M1 and M4 were found in clade B ($n = 137$). Any pair of isolates within clade A differed by up to 59 (median, 26) wgMLST alleles, while the pairs of isolates in clade B showed a maximum of 129 (median, 34) allelic differences. The number of allelic differences separating isolates from clade A and clade B ranged from 73 to 165 (median, 119). With the exception of a subcluster of clade A isolates from plant M4, isolated during an intensively sampled 3-month period in 2014 (four groups comprising a total of 16 isolates), and two clade B isolates from 2015, all isolates could be distinguished using the wgMLST analysis.

The SNP analysis was performed by mapping raw reads from all 252 Norwegian isolates to the genome of the ST9 isolate *L. monocytogenes* SLCC 2479, for which a complete genome sequence is available (41). In addition, SNP analyses were performed separately for the isolates within clades A and B, using representatives from within each clade as reference genomes, MF4562 from clade A and MF4697 from clade B (42). Due to high sequence variability within prophage regions and plasmids, these regions were excluded from the SNP analyses. In the analysis performed for all genomes using SLCC 2479 as the reference (Fig. 2C), a total of 554 high-quality SNP positions were identified, of which 516 SNPs were present in all 252 isolates. The remaining 38 variant positions had missing data in up to 35% of the isolates, including a stretch of 23 consecutive SNP sites covering an ~70-kb region, which included the missing *bgIA* locus in MF6279. In comparison, of the 746 identified polymorphic wgMLST loci, only 494 were present in all isolates, reflecting the inclusion of accessory genes such as those found within prophages in the wgMLST scheme. In this SNP analysis, any pair of isolates differed by up to 123 (median, 58) variants while in the wgMLST analysis, they differed by up to 179 (median, 95) alleles, and a greater number of isolates remained undistinguishable in the SNP analysis than in wgMLST (Fig. 2). The number of SNP differences separating any pair of isolates from clade A and clade B ranged from 51 to 93 (median, 74), with 100% bootstrap support for the separation of clade A, clade B, and the three outlier strains.

The SNP analyses performed using representatives within clades A and B as reference genomes (Fig. S1) were more sensitive than the analysis performed using the external reference SLCC 2479 (Fig. 2C). A total of 121 SNPs were detected when reads from the clade A isolates were mapped to the MF4562 genome, compared to 106 SNPs detected using the SLCC 2479 reference. Any pair of isolates within clade A differed by up to 26 (median, 13) and 22 (median, 11) SNPs in the two analyses. Within clade B, the total numbers of detected SNPs were 331 and 287, and any pair of isolates differed by 77 (median, 27) and 64 (median, 24) SNPs, using MF4697 and SLCC 2479 as references,




FIG 2 Phylogenetic trees for ST9 isolates from Norwegian meat processing industry. Shown are minimum spanning trees based on wgMLST (A) and cgMLST (B) analyses and a maximum parsimony tree based on an alignment of the 554 SNP positions detected using SLCC 2479 as the reference genome, showing the tree with the highest resampling support (C). Red arrows point to outliers. The area of each circle is proportional to the number of isolates represented, and the factory of origin for each isolate is indicated by the color. In panels A and B, branch labels indicate the numbers of target genes with differing alleles. In panel C, branch labels represent bootstrap resampling support values, showing only values of >50%. Branch lengths were scaled using logarithmic (A) and square root (B and C) scaling. Labels next to selected clusters refer to the subclusters described in the text. Selected isolate names are indicated in blue.

respectively. SNP counts reported below for strains within clade A or B are based on the analyses performed using MF4562 and MF4697 as reference genomes.

All three genomic analyses (wgMLST, cgMLST, and SNP) indicated that a subcluster within clade B comprising 19 isolates from plant M4, referred to as subcluster B1 (Fig. 2), could be considered a third major clade within this data set. This subgroup was separated from the remaining genomes in clade B with 100% bootstrap support (Fig. 2C and Fig. S1), between 45 and 77 (median, 61) SNPs, between 59 and 129 (median, 86) wgMLST alleles, and with 8 cgMLST alleles. Moura et al. (19) observed that isolates with no documented epidemiological link generally differed by more than ten cgMLST alleles. We conclude that a total of 249/252 (99%) of the *L. monocytogenes* ST9 isolates collected over a 9-year period (2009 to 2017) in four Norwegian meat processing plants belonged to one of two clonal lineages, referred to here as clades A and B.

MLVA typing was not in accordance with the WGS phylogeny. In previous work (26), the ST9 isolates collected between 2009 and 2015 were typed using the multiple-locus variable-number tandem repeat analysis (MLVA) scheme from Lindstedt et al. (43), also used by the Norwegian Institute of Public Health until March 2018. The majority of isolates belonged to three major MLVA profiles (26) (Table S1). The MLVA typing was, however, not fully concordant with the population structure determined using genomic analyses and was not sufficiently discriminatory to differentiate between isolates originating from different facilities (Fig. S2). This was not unexpected, since markers used in MLVA evolve by the addition or removal of repeat units. As a result, MLVA data comprise a substantial degree of phylogenetic noise (44).

Spatiotemporal spread of ST9 clones within Norwegian meat supply chains. When the year of isolation was projected onto the wgMLST and SNP-based phylogenies (Fig. S3), no specific clustering of isolates according to isolation year was seen, except for subcluster A-M4, for which all isolates were obtained during 2014. To further investigate the dissemination and temporal history of the Norwegian ST9 isolates, a Bayesian tip-dated phylogenetic analysis (BEAST [45]) was conducted using the 554 differentiating SNPs, the date of isolation for each isolate, and assuming a strict molecular clock (Fig. 3 and Table 2 and Fig. S4). The analysis estimated that the most recent common ancestor (MRCA) of the set of 252 analyzed Norwegian ST9 isolates existed around the year 1935 (95% highest posterior density [HPD], 1907 to 1955; Table 2), while the division between the two main lineages (clades A and B) appeared to have occurred around 1955 (Fig. 3). The time of the MRCA was estimated to be 1998 for clade A (95% HPD, 1992 to 2004) and 1981 for clade B (95% HPD, 1970 to 1991). Since both clades contain isolates from multiple meat processing plants, cross-contamination between facilities, or contamination from a common primary source of *L. monocytogenes*, appears to have occurred in relatively recent years.

Groups of isolates from all four factories were closely related. The phylogenies based on the SNP analyses (Fig. 2C and 3 and Fig. S1 and S4) widely concur with the clustering of isolates obtained using wgMLST analysis (Fig. 2A), with incongruences in tree topology typically linked to branching points for outlier strains and smaller groups of isolates in clades with lower bootstrap and posterior probability support values. For instance, within clade A (comprising isolates from all four factories), the phylogenetic reconstructions based on the SNP matrices strongly support a monophyletic origin for a group of 55 isolates from plant M4 (named subcluster A-M4), presenting 99% bootstrap support (Fig. 2C and Fig. S1A), while the wgMLST analysis suggests that the factory M5 isolates belong to the same cluster as A-M4 (Fig. 2A). Support values for the branch delineating a second clade A subcluster, containing 50 isolates from plant M1 (named subcluster A-M1), as well as for the branches for the clade A isolates from factories M5 and M7, were low and/or showed conflicting results (Fig. 2C and Fig. S1A and S4). The pairwise distances between the factory M7 isolate (MF6343) and the isolates from factory M5, subclusters A-M1 and subcluster A-M4, ranged from 7 to 18 SNPs and 9 to 20 wgMLST loci. Therefore, although isolates within clade A largely




FIG 3 Time-scaled phylogenetic reconstruction, showing a maximum clade credibility tree produced from the concatenated SNP alignment obtained with SLCC 2479 as the reference genome, using the Bayesian Markov chain Monte Carlo method available in BEAST (45). The inset shows the whole tree, while the main figure shows the section containing the tips, enlarged for clarity. Figure S4 shows the same tree, labeled with the posterior probability values for each split and node labels containing strain names. Isolates within cluster B2 were isolated from raw zone sampling points, except for the three nodes labeled with C (cooked zone). All other isolates were from cooked zones, except the five nodes labeled with circles and an R (raw zone). The tips of the branches correspond to the sampling date, while tip colors indicate factory of origin for each strain, colored as described for Fig. 1 and Fig. 2. The scale is in years.

clustered according to factory of origin, the small number of genetic differences between the groups indicated close genetic relationships between isolates from all four factories. Furthermore, the precise evolutionary relationship between these groups remains ambiguous. The BEAST analysis suggested that their time of divergence was

TABLE 2 Time of MRCA for the isolates in each clade estimated using BEAST analysis

Summary statistic	Mean time of MRCA ^a	95% HPD interval ^b	Mean SE	SD	ESS ^c
Age(root)	1934	1907.3, 1955.4	0.32	12.55	1,539
Age(cladeA)	1998	1991.8, 2004.0	0.06	3.22	3,212
Age(A-M1)	2007	2004.7, 2008.6	0.02	1.03	3,194
Age(A-M4)	2011	2009.2, 2012.5	0.01	0.89	3,956
Age(cladeB)	1981	1970.2, 1991.4	0.12	5.55	1,983
Age(B1)	2008	2004.1, 2011.1	0.03	1.88	4,796
Age(B2/B3)	1994	1987.0, 2000.6	0.07	3.56	2,294
Age(B2)	1999	1993.7, 2004.5	0.05	2.83	2,708
Age(B3)	2004	1999.9, 2007.2	0.04	1.93	2,173

^aMRCA, most recent common ancestor.

^bHPD, highest posterior density interval for the divergence time.

^cESS, effective sample size.

approximately 10 to 15 years prior to the collection date for the most recently sampled isolates (Fig. 3).

Clade B harbors clusters comprising isolates from both M1 and M4. In clade B, both the wgMLST and SNP analyses show separation into three clearly delineated subclusters, named subclusters B1 ($n = 19$), B2 ($n = 54$), and B3 ($n = 64$) (Fig. 2 and Fig. S1B). Bootstrap support for the delineation of subclusters B1 and B3 from the remaining isolates in the tree was 100%, while support for the branch leading to subcluster B2 was lower (66% to 67%) (Fig. 2C and Fig. S1B), indicating that the grouping of cluster B2 was not as strongly supported. The cgMLST analysis was not able to differentiate between subclusters B2 and B3 (Fig. 2B). Subcluster B1 was estimated to have split from subclusters B2 and B3 around 1981, while the time of divergence of the B2 and B3 lineages was around 1994 (Table 2 lists 95% HPD intervals for the estimated times of MRCAs). Subcluster B1, with an MRCA estimated to have existed in 2008, exclusively contained isolates from only one factory (M4). Subclusters B2 and B3, however, both contained isolates from factories M1 and M4.

The subcluster B2 isolates from factories M1 ($n = 3$) and M4 ($n = 51$) were significantly associated with raw side production rooms ($P = 1.04 \times 10^{-37}$ by two-sided Fisher's exact test). Only three subcluster B2 isolates were from high-risk zones in which cooked meat was handled (in factory M4; see below), and subcluster B2 contained 51 of the 63 total isolates obtained from raw meat zones in the present study. The pairwise distances between the factory M1 and M4 isolates within subcluster B2 ranged from 8 to 24 SNPs and 12 to 23 wgMLST loci (median of 13 SNPs and 17 wgMLST loci), and the MRCA was estimated to have existed in 1999. The association of this subcluster with raw meat zones in both M1 and M4, along with the relatively close genetic relationship between isolates from both factories, indicates that for this clone the common link between the facilities is a contamination source related to raw materials.

Subcluster B3 contained 54 isolates from factory M1, most of them from the high-risk zone where cooked products were processed, plus a well-supported monophyletic cluster harboring ten factory M4 isolates (94% to 97% bootstrap support) (Fig. 2C and Fig. S1B). The pairwise distances between these factory M1 and M4 isolates ranged from 9 to 33 SNPs and 14 to 43 wgMLST loci (median of 21 SNPs and 26 wgMLST loci), and the MRCA was estimated to have existed in 2004. Thus, also within this subcluster, pairs of isolates from two different facilities had sufficiently close genetic relationships for a mistaken association of isolates to occur during a potential outbreak investigation, as a threshold genetic distance of <20 SNPs is commonly used to formulate a hypothesis that two isolates originate from the same source (28). The evidence of close association between isolates from more than one facility, within almost all branches of the phylogeny of *L. monocytogenes* ST9 from Norwegian meat processing facilities, further points to a widely interconnected meat industry sector in Norway.

Related isolates in multiple areas of factory M4 suggests breach of hygienic barriers. Factory M4 produces RTE meat products, like cold cuts, sausages, and liver

pâté, as well as raw meat products, such as loin steaks and minced meat. The building structure houses two separate raw meat departments, one for sausage production and one for cutting and marinating raw beef. These departments are separated by hygienic barriers from the high-risk areas, where heat-treated products are processed and handled (Fig. 4A).

The high-risk zone and the raw sausage room harbored isolates belonging to subcluster B1 ($n = 19$, 2012 to 2015) and B3 ($n = 10$, 2012 to 2014) over a period of 3 to 4 years. None of these isolates appear to cluster in accordance with sampling area or the year of isolation (Fig. 3 and 4A and Fig. S3). The MRCA of the subcluster B1 isolates (for which isolates had a median pairwise distance of 3 SNPs [range, 0 to 14] and 5 wgMLST loci [range, 0 to 40]) was estimated to have existed around 2008 (Table 2), suggesting that this clone had persisted in the facility from around 2008 to 2015. The delineation of the M4 isolates within subcluster B3 (median pairwise distance, 8 SNPs [range, 0 to 14] and 9 wgMLST loci [range, 2 to 16]) had fairly good bootstrap support (94% to 97%; Fig. 2C and Fig. S1B), and the cluster had an estimated MRCA from about 2010 (Fig. 3), suggesting persistence for this clone from around 2010 to 2014. The analysis indicated that the breach of the hygienic barriers separating the raw sausage and high-risk zones had occurred on more than one occasion.

All the isolates from the second raw meat department in M4 ($n = 48$, 2012 to 2017) belonged to subcluster B2. Isolates from this subcluster were also found in three relatively recent samples (2014 and 2017) from the high-risk areas in the factory and formed a polyphyletic group within subcluster B2 (nodes labeled C in Fig. 3). The median pairwise distance between subcluster B2 isolates from M4 was 5 SNPs (range, 0 to 16) and 7 wgMLST loci (range, 1 to 18). These observations suggest that the three recent isolates originated from the raw beef cutting department, representing additional breaches of hygienic barriers in the factory.

In sum, the WGS data suggested that the clade B ST9 isolates found in the high-risk zone were related to those found in both adjacent departments handling raw meats. In 2016, as a response to these results, the factory decided to improve the hygienic barriers between departments to reduce contamination between the areas handling raw meat and the areas used for the processing of cooked products. Subsequently, only a single ST9 (subcluster B2) isolate was identified in the high-risk zone of factory M4 between January and October 2017 through the routine environmental monitoring program. The raw meat departments were not sampled after the reinforcement of hygienic barriers, but overall, the data suggest that the origin of the clade B ST9 isolates in factory M4 was in the raw meat departments.

Contamination event related to installation of a second-hand slicer line. The majority of the clade B isolates from factory M4 discussed above were from floors and drains. However, in 2014, factory M4 bought a used slicer line and installed it in the department where heat-treated products were processed. Within a week after starting production on the line, *L. monocytogenes* was detected in products (different types of sliced deli meats), product residues, and on processing equipment associated with the slicer line. The plant recalled products from the market after the first *L. monocytogenes*-positive samples. Despite several rounds with thorough cleaning and disinfection, including dismantling of the line, *L. monocytogenes* ST9 was repeatedly detected over the coming weeks. WGS analysis of the 55 ST9 isolates obtained in association with the introduction of the slicer line showed that they formed a separate and dense cluster within clade A, referred to as subcluster A-M4 (Fig. 2 and 4A). All isolates were closely related; a total of 41 SNP sites were identified, the median pairwise distance was 5 SNPs (range, 0 to 15) and 5 wgMLST loci (range, 0 to 12), and the MRCA of the A-M4 cluster was estimated to have existed in 2011 (Table 2). Isolates belonging to this subtype had not previously been detected in the factory.

After 8 weeks, the slicer line was discarded and completely removed from the processing plant, and the last subcluster A-M4 isolate was found 4 weeks after the slicer line was removed. Only three isolates obtained from the area of the factory where the




FIG 4 Minimum spanning trees based on wgMLST alleles, colored according to the location within factory M4 (A) and factory M1 (B) at which each *L. monocytogenes* isolate was obtained. Insets show sketches of the factory layouts, with arrows indicating the direction of movement of products and equipment through processing.

line was installed, during the time it was present, were not part of subcluster A-M4 (they were typed as other STs). Thus, the WGS analysis confirmed that a persistent clone was introduced into the factory through the used slicer line. Similar cases of transfer of *L. monocytogenes* between factories have been documented in previous reports (8, 46). Most probably, the slicer line was contaminated with a population of closely related *L. monocytogenes* isolates when purchased, as the level of genetic variation between isolates was greater than what would be expected to develop within the short period the line was in use at plant M4. Unfortunately, the cluster could not be traced back to the factory where the slicer line originated, as this factory did not monitor *L. monocytogenes* in the processing environment, because they produced low-risk products (fermented meats) in which *L. monocytogenes* isolates are unlikely to grow.

Two ST9 clones have diversified within the high-risk zone of factory M1. Factory M1 produces various types of cooked sausages and pork liver pâté. The raw meat department is located next to a large high-risk production room used for processing and packaging of heat-treated products. The two departments are separated physically by large double-entry ovens as well as walls/doors and hygienic zone barriers (Fig. 4B). Sausages and other products enter the ovens from the raw side. After heat treatment, the oven doors on the cooked side are opened, and the products are taken into the high-risk production room. Racks used for carrying products inside the ovens are pushed back to the low-risk side after use. The majority of the isolates obtained from factory M1 ($n = 107$) were from two regions in the high-risk production room: an area for removing casings from sausages ($n = 28$; isolates were mainly from floors and drains) and an area containing conveyor-based packaging lines ($n = 67$; isolates were from floors, drains, and processing equipment/machines). The regions are referred to as area I and area II, respectively, in Fig. 4B. Although the areas had separate drains and equipment, flow of water, personnel, and trolleys could occur across the two regions, as they were not physically separated.

All isolates from the high-risk production room belonged to either subcluster A-M1 or subcluster B3. Subcluster A-M1 isolates (detected from 2009 to 2017) had a median distance between any pair of isolates of 4 SNPs (range, 0 to 12) and 11 wgMLST loci (range, 1 to 50), and the estimated time of existence of the MRCA was around 2007 (Table 2). For the subcluster B3 isolates from factory M1 (isolated from 2011 to 2017), the median pairwise distance was 11 SNPs (range, 0 to 27) and 16 wgMLST loci (range, 2 to 32), and the MRCA was estimated to have existed around 2004 (Table 2). Although isolates from both sampling areas I and II were found in both subclusters (as expected, since there were no barriers between these areas), isolates within subcluster B3 were more strongly associated with area I and the A-M1 isolates with area II ($P = 0.004$ by two-sided Fisher's exact test). There was, however, no clear pattern with respect to diversity between specific locations (Fig. 4B) and no apparent clustering of isolates according to isolation year (Fig. 3 and Fig. S3). The WGS results strongly indicated that the high-risk production room in factory M1 was home to two different ST9 lineages, both of which had diversified within the factory for a period of around ten years and were found in two distinct reservoirs (located in areas I and II) from which isolates were spread to other areas of the high-risk zone.

Only a few positive samples from plant M1 were obtained from the raw meat department ($n = 8$), as it was not part of the plant's monitoring program for *L. monocytogenes*. Sampling was performed mainly in locations close to the high-risk zone at sites mentioned by factory personnel to have suboptimal hygienic zoning between the two departments. The ST9 isolates from the raw side were from three subclusters: A-M1 ($n = 3$), B3 ($n = 2$), and B2 ($n = 3$). Thus, five isolates belonged to the two ST9 subclusters found in the high-risk zone, indicating the occurrence of contamination between the two zones. Subcluster B2 isolates, in contrast, were not found in the more frequently sampled high-risk production room, suggesting that the direction of transmission of ST9 *L. monocytogenes* was more likely to be from the cooked to the raw side rather than *vice versa*.

DISCUSSION

WGS was employed in the current study to investigate the diversity and transmission patterns of ST9 strains widely disseminated within the Norwegian meat processing industry and provides new insights into plant-to-plant and within-plant variation of *L. monocytogenes*. Several reports have shown that WGS may be used as a surveillance tool for tracking *L. monocytogenes* in food processing environments, and that high-resolution subtyping of bacterial isolates is necessary to unravel contamination routes within processing plants, as well as to track sources of foodborne outbreaks (24, 33, 34, 37, 47). The use of high-resolution WGS-based typing strategies is particularly appropriate for the investigation of widely distributed clones. However, many different bioinformatic methods can be used to analyze WGS data, and the various pipelines may not always yield concordant results (29, 48). In the current study, cgMLST did not yield sufficient resolution to differentiate between isolates originating from different processing plants, although it could separate the Norwegian clones from other publicly available CC9 reference genomes (Fig. 1). Furthermore, highly comparable results were obtained with the wgMLST- and SNP-based approaches (Fig. 2A and C; see also Fig. S1 in the supplemental material), largely grouping isolates according to their factory of origin. The wgMLST approach detected higher numbers of differentiating loci than the SNP analyses, even when SNP analysis was performed using reference genomes found within the same clusters as the examined isolates. Discrepancies between the two analyses were likely due to the exclusion of variable elements such as prophage and plasmid regions in the SNP analysis, while the employed wgMLST scheme includes accessory genes. The dependence on selection (or availability) of suitable reference genomes in SNP-based analyses may complicate interpretations of genetic distances between isolates. However, while the calculation of confidence support values for topologies of phylogenetic trees is straightforward for trees built using SNP analyses, it is less so for minimum spanning trees based on genomic MLST data (49). Confidence values should be taken into account when interpreting WGS analyses (27, 29).

The obtained results highlight the importance of considering evidence such as epidemiological data, traceback evidence, and phylogenetic tree topology, in addition to SNP counts, when seeking to establish potential sources of bacterial contaminations or outbreaks. The closest pairwise differences between any two isolates originating from two different facilities among the four examined in the current study ranged from 7 to 11 SNPs and from 9 to 11 wgMLST loci. These numbers are well below the threshold of 20 SNPs commonly employed during outbreak investigations as an indication that two isolates have originated in the same facility (28, 34). Our study also demonstrates the advantage of having a broad overview of the genotypes of strains circulating in food chains to prevent the misinterpretation of data during the investigation of contamination sources or in the event of potential outbreak scenarios. A more representative overview of circulating bacterial populations may both reveal the presence of closely related isolates present in multiple facilities and allow the detection of closely related but genetically distinct subpopulations through the building of more detailed phylogenies. A WGS-based analysis of a less deeply sampled data set than the ones currently considered (within clade A or B) could have resulted in the conclusion that isolates originated from the same source when in reality they belong to distinct subpopulations. The risk of misinterpretations would be even greater in the event that a potential outbreak was suspected of originating from a persistent clone, subject to diversification events within a factory, because in such cases the use of even more relaxed genetic distance thresholds has been recommended (35).

One of the questions posed in the current study was whether WGS could be used to differentiate between different contamination scenarios, including whether the presence of ST9 clones in an area of a factory was likely a result of persistence, multiple reintroduction events from an external source, or a combination of both. In the present work, the isolates formed five distinct subclusters, and for four of the clusters, closely related isolates were detected over the course of several (four to nine) years in the same

meat processing facility. This typically indicates that the persistence of strains is the most common scenario. An alternative explanation is that isolates are more or less continuously introduced from an outside source in which the strain is persistent.

One clear indication that raw meat is a common source of the introduction of *L. monocytogenes* into more than one factory was the observation that subcluster B2 isolates, mainly found in the raw meat department used for cutting and marinating raw beef in factory M4, also were found on the raw side, but not in the high-risk zone, of factory M1. *L. monocytogenes* strains, including ST9 strains, have frequently been found to persist in meat slaughterhouses (5, 6, 15, 50–56). Therefore, a common primary source of a certain *L. monocytogenes* strain present in more than one meat processing plant potentially could be one that persists in a slaughterhouse supplying raw meat to multiple processing plants. The raw meat distribution chain in Norway is complex, where the same slaughterhouses deliver raw meat to many meat processing plants. The largest meat-producing company owns several slaughterhouses and sells raw meat to most meat processing plants. They also have a role as a market regulator and freeze meat for long-term storage during periods of high production. For beef, about 20% to 30% of the raw meat on the Norwegian market is imported, from lamb/mutton at 2% to 10% and <3% for pork and poultry, with some variation in import between years based on market demands (57). The M1 plant has its own slaughter department but also acquires raw meat from other slaughterhouses and sells raw meat to other processing plants, while M4 and M5 do not have slaughtering departments. Factory M7 has its own slaughter department for poultry but occasionally also buys raw poultry meat from others and routinely sells to other processing plants (>10 customers). Thus, it is evident that different factories may receive raw meat from the same supplier, which may explain why closely related isolates are found in multiple processing plants. The examination of *L. monocytogenes* isolates found upstream in the meat chain is needed to potentially identify the primary sources of the examined strains.

The primary origin of ST9 entering the food chain is not known. While *L. monocytogenes* is frequently found in animals, on farms, and in the natural environment, the frequency of the different genetic groups of *L. monocytogenes* appears to differ significantly in these environments compared to that observed in meat products and the meat processing industry. *L. monocytogenes* ST9 is rarely found in pig or ruminant farming environments, animals, or wild natural surroundings (22, 58–60). This sequence type has, however, frequently been found to dominate in raw pork meat (21, 52, 61) and is also commonly found to persist in slaughterhouses (14, 15, 24, 62) as well as in downstream compartments in the meat processing chain (10, 22, 26). In contrast, several studies report an absence or low abundance of ST9 in other food types and processing environments, such as fish/salmon, dairy, and fresh produce (22, 26, 31, 47, 63). These observations suggest that ST9 isolates are highly adapted to survival in meat production environments. We previously showed that an ST9 isolate included in the present study (MF4562, from subcluster A-M1) outcompeted isolates of other genotypes when grown in a meat-based growth medium (brain heart infusion [BHI]) in the presence of *Listeria innocua* (64). It also has been hypothesized that *L. monocytogenes* strains harboring *comK* prophages were adapted to meat environments (65), but information about the STs of examined strains was not provided. The genome data acquired in the current study will provide a basis for future studies aiming to determine potential underlying mechanisms responsible for the tight association of ST9 strains with the meat industry.

In conclusion, the current study confirms that isolates with few genetic differences were found across multiple meat processing plants and emphasizes the importance of having an overview of clones found in a given food chain to correctly interpret WGS analyses and prevent false-positive associations between isolates. The use of epidemiological data and traceback evidence is also fundamental for an accurate assessment of the origin of a given isolate. WGS analyses were shown to successfully distinguish between different contamination scenarios both between and within factories and can be used to estimate the time frame of persistence. Such information may be employed

to significantly strengthen food safety management efforts within processing plants in the food industry.

MATERIALS AND METHODS

DNA isolation and whole-genome sequencing. Bacteria were grown on BHI agar overnight at 37°C before a loopful of cells were suspended in 500 μ l 2 \times Tris-EDTA (TE) buffer with 1.2% Triton X-100. Cells were lysed using lysing matrix B and a FastPrep instrument (both MP Biomedicals), and genomic DNA was isolated using the DNeasy blood and tissue kit (Qiagen). Libraries were prepared using the Nextera XT DNA sample preparation kit (Illumina) and sequenced either on a MiSeq platform with 300-bp paired-end (PE) reads or on a HiSeq platform with 150-bp PE reads (see Table S1 in the supplemental material). Sequencing on the HiSeq platform, and one of the MiSeq runs, were performed at the Norwegian Sequencing Centre (<http://www.sequencing.uio.no>); otherwise, sequencing was performed in-house. Raw reads were filtered on q15 and trimmed of adaptors using fastq-mcf from the ea-utils package (66).

Genome assembly. For 6 of the 252 isolates (MF4545, MF4562, MF4624, MF4697, MF4626, and MF6172), complete genome sequence assemblies were previously prepared using both Illumina and Nanopore sequencing data (42). For the remaining 246 isolates, *de novo* genome assemblies were generated using SPAdes v.3.10 (67) with the careful option and kmer sizes of 21, 33, 55, 77, 99, and 127. Contigs with sizes of <500 bp and kmer coverage of <5 were removed from the assemblies. The genome assemblies were evaluated using QUAST v2.2 (68).

Genomic MLST analyses. The wgMLST analysis was performed using a whole-genome schema containing 4,797 coding loci from the *L. monocytogenes* pangenome and the assembly-based BLAST approach, implemented in BioNumerics 7.6 (69). The cgMLST analysis was performed using the scheme described by Moura et al. (19), and results were either extracted from the wgMLST analysis (for isolates sequenced in the current study) or (for reference genomes) obtained using the KMA alignment method and the cgMLSTfinder program, which maps the filtered fastq reads to a database of the cgMLST alleles (70, 71). Minimum spanning trees were constructed using BioNumerics 7.6 based on the categorical differences in the allelic cgMLST or wgMLST profiles for each isolate. Loci with no allele calls were not considered in the pairwise comparison between two genomes. The number of allelic differences between isolates was read from genetic distance matrices computed from the absolute number of categorical differences between genomes.

Selection of public reference genomes for comparison. Reference genomes included in the cgMLST analysis of a cross-section of CC9 strains were selected and identified using the following sources. (i) Typing data were downloaded from the BIGSdb-*Lm* database (<http://bigsdb.pasteur.fr/listeria/listeria.html>) for the 19 *L. monocytogenes* sublineage 9 (SL9) isolates with associated cgMLST profiles publicly available on 19 March 2019. (ii) The *L. monocytogenes* AT3E genome (39) was selected for inclusion in the analysis, since it was previously identified as being related to isolates examined in the current study (42). As raw reads were not publicly available for AT3E, the cgMLST profile was obtained using the BioNumerics assembly-based wgMLST BLAST approach. For the remaining *L. monocytogenes* CC9 reference strains, publicly available raw fastq data were downloaded from the Sequence Read Archive (SRA) database (<https://www.ncbi.nlm.nih.gov/sra>). (iii) Of the 1,696 isolates included in the study by Moura et al. (19), 78 belonged to CC9, and of these, fastq read data were available for 55 genomes. cgMLST data for one strain (FSL R2-561) were also present in the BIGSdb data set; thus, 54 genomes were included. (iv) Among the 1,231 *L. monocytogenes* genome assemblies present in NCBI GenBank as of 6 December 2016, 87 were determined by *in silico* MLST analysis (72, 73) to belong to CC9, and of these, fastq read data were available for 27 isolates, of which nine were also present in the study by Moura et al. (19); thus, 18 genomes were obtained from this data set. (v) The LiSEQ project (74) included 110 *L. monocytogenes* CC9 strains from European countries that were included in the current analysis. In total, 202 CC9 reference genomes were selected for comparative analysis using cgMLST.

SNP analysis. The reference genomes used were *L. monocytogenes* SLCC 2479 (ST9; GenBank accession number [NC_018589](https://www.ncbi.nlm.nih.gov/nuccore/NC_018589) [41]) and the chromosomes of *L. monocytogenes* MF4562 (clade A; GenBank accession number [CP025442](https://www.ncbi.nlm.nih.gov/nuccore/CP025442) [42]) and *L. monocytogenes* MF4697 (clade B, GenBank accession number [CP025438](https://www.ncbi.nlm.nih.gov/nuccore/CP025438) [42]). Each reference genome had two prophage regions, which were not conserved between strains and were excluded from the analysis. For each of the 252 ST9 isolates, Illumina reads filtered using fastq-mcf (described above) were mapped against the references using Bowtie2 v.2.2.6 or v.2.3.5.1 (75). Variants were called using FreeBayes v1.2.0-17 (76) with the ploidy option set to 1. The called variants were filtered using VCFtools (77) to remove indels, SNPs with quality values of <30, SNPs with missing data in >35% of genomes, and genotypes with <10 reads. SNPs clustered closer than 1,000 bp in the same individual isolate were excluded. The SNPs in each genome were extracted from the filtered FreeBayes outputs and concatenated into fasta pseudosequences containing only SNP sites (78), which then were used to generate multiple-sequence alignments. Maximum parsimony trees were built from the concatenated SNP alignments using BioNumerics 7.6 with default parameter values and 100 \times bootstrap resampling. In each analysis, the tree with the highest resampling support was computed. The SNP distance matrices summarizing the number of sites that differ between each pair of sequences were built from the filtered FreeBayes outputs using the `dist.dna()` function from the R package `ape` v.5.3 (79), using pairwise deletion of sites with missing data.

Tip-dated phylogenetic analysis. Timed phylogeny was reconstructed from the concatenated SNP alignment obtained with the SLCC 2479 reference genome, using the Bayesian Markov chain Monte Carlo (MCMC) method available in BEAST v1.10.4 (45). The sampling date for each isolate was entered as the tip date. For 21 isolates collected between 2009 and 2012, only the sampling year was known, and for

these isolates, the sampling date was set to July 1 and the uncertainty value associated with the date was set to 1. For two isolates, only the sampling month and year were known, and for these, the 1st of the month was used. The most recently sampled isolate was sampled on 18 November 2017, which was set as the current date. To estimate the time of the most recent common ancestor (MRCA) for subclades, taxon sets were defined as comprising the isolates belonging to the clades and subclusters described in the study (clades A and B and subclusters A-M1, A-M4, B1, B2, B3, and B2/B3) (Table 2). Settings used were the HKY nucleotide substitution model, a strict molecular clock, and a coalescent constant size tree prior with a random starting tree. The analysis was run with a chain length of 10^9 , a sampling frequency of 10^5 , and a 5% burn-in. The effective sample size (ESS) values and trace files were monitored to ensure sufficient mixing (ESS of $>1,000$ for all statistics). The MCMC output was summarized as a maximum clade credibility tree, i.e., the tree with the highest product of the posterior probabilities of all its nodes.

Data availability. The raw reads have been deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under BioProject accession no. [PRJNA419519](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA419519). Accession numbers for individual genomes have been submitted to NCBI's Sequence Read Archive and BioSample databases (accession numbers [SRR11261968](https://www.ncbi.nlm.nih.gov/submit/sra/SRR11261968) to [SRR11262219](https://www.ncbi.nlm.nih.gov/submit/sra/SRR11262219) and accession numbers [SAMN08056483](https://www.ncbi.nlm.nih.gov/submit/sra/SAMN08056483) to [SAMN08056488](https://www.ncbi.nlm.nih.gov/submit/sra/SAMN08056488) and [SAMN14314680](https://www.ncbi.nlm.nih.gov/submit/sra/SAMN14314680) to [SAMN14314925](https://www.ncbi.nlm.nih.gov/submit/sra/SAMN14314925); see Table S1 for details).

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

SUPPLEMENTAL FILE 1, PDF file, 1.9 MB.

ACKNOWLEDGMENTS

We thank Merete Rusås Jensen, Tove Maugesten, and Anette Wold Åsli for excellent technical assistance.

This work was funded by the Norwegian Research Funding for Agriculture and Food Industry, grant numbers 207765 and 262306.

REFERENCES

- Smith AM, Tau NP, Smouse SL, Allam M, Ismail A, Ramalwa NR, Disenyeng B, Ngomane M, Thomas J. 2019. Outbreak of *Listeria monocytogenes* in South Africa, 2017-2018: laboratory activities and experiences associated with whole-genome sequencing analysis of isolates. *Foodborne Pathog Dis* 16:524–530. <https://doi.org/10.1089/fpd.2018.2586>.
- Gilmour MW, Graham M, Van Domselaar G, Tyler S, Kent H, Trout-Yakel KM, Larios O, Allen V, Lee B, Nadon C. 2010. High-throughput genome sequencing of two *Listeria monocytogenes* clinical isolates during a large foodborne outbreak. *BMC Genomics* 11:120. <https://doi.org/10.1186/1471-2164-11-120>.
- Olsen SJ, Patrick M, Hunter SB, Reddy V, Kornstein L, MacKenzie WR, Lane K, Bidol S, Stoltman GA, Frye DM, Lee I, Hurd S, Jones TF, LaPorte TN, Dewitt W, Graves L, Wiedmann M, Schoonmaker-Bopp DJ, Huang AJ, Vincent C, Bugenhagen A, Corby J, Carloni ER, Holcomb ME, Woron RF, Zansky SM, Dowdle G, Smith F, Ahrabi-Fard S, Ong AR, Tucker N, Hynes NA, Mead P. 2005. Multistate outbreak of *Listeria monocytogenes* infection linked to delicatessen turkey meat. *Clin Infect Dis* 40:962–967. <https://doi.org/10.1086/428575>.
- Mørtrø T, Langsrud S. 2004. *Listeria monocytogenes*: biofilm formation and persistence in food processing environments. *Biofilms* 1:107–121. <https://doi.org/10.1017/S1479050504001322>.
- Ferreira V, Wiedmann M, Teixeira P, Stasiewicz MJ. 2014. *Listeria monocytogenes* persistence in food-associated environments: epidemiology, strain characteristics, and implications for public health. *J Food Prot* 77:150–170. <https://doi.org/10.4315/0362-028X.JFP-13-150>.
- Nesbakken T, Kapperud G, Caugant D. 1996. Pathways of *Listeria monocytogenes* contamination in the meat processing industry. *Int J Food Microbiol* 31:161–171. [https://doi.org/10.1016/0168-1605\(96\)00978-6](https://doi.org/10.1016/0168-1605(96)00978-6).
- López V, Villatoro D, Ortiz S, López P, Navas J, Dávila JC, Martínez-Suárez JV. 2008. Molecular tracking of *Listeria monocytogenes* in an Iberian pig abattoir and processing plant. *Meat Sci* 78:130–134. <https://doi.org/10.1016/j.meatsci.2007.05.002>.
- Lundén JM, Autio TJ, Korkeala HJ. 2002. Transfer of persistent *Listeria monocytogenes* contamination between food-processing plants associated with a dicing machine. *J Food Prot* 65:1129–1133. <https://doi.org/10.4315/0362-028X-65.7.1129>.
- Ferreira V, Barbosa J, Stasiewicz M, Vongkamjan K, Switt AM, Hogg T, Gibbs P, Teixeira P, Wiedmann M. 2011. Diverse geno- and phenotypes of persistent *Listeria monocytogenes* isolates from fermented meat sausage production facilities in Portugal. *Appl Environ Microbiol* 77:2701–2715. <https://doi.org/10.1128/AEM.02553-10>.
- Martin B, Perich A, Gómez D, Yangüela J, Rodríguez A, Garriga M, Aymerich T. 2014. Diversity and distribution of *Listeria monocytogenes* in meat processing plants. *Food Microbiol* 44:119–127. <https://doi.org/10.1016/j.fm.2014.05.014>.
- Bolocan AS, Oniciuc EA, Alvarez-Ordóñez A, Wagner M, Rychli K, Jordan K, Nicolau AI. 2015. Putative cross-contamination routes of *Listeria monocytogenes* in a meat processing facility in Romania. *J Food Prot* 78:1664–1674. <https://doi.org/10.4315/0362-028X.JFP-14-539>.
- Stoller A, Stevens MJA, Stephan R, Guldemann C. 2019. Characteristics of *Listeria monocytogenes* strains persisting in a meat processing facility over a 4-year period. *Pathogens* 8:32. <https://doi.org/10.3390/pathogens8010032>.
- Véghová A, Minarovičová J, Koreňová J, Drahovská H, Kacliková E. 2017. Prevalence and tracing of persistent *Listeria monocytogenes* strains in meat processing facility production chain. *J Food Saf* 37:e12315. <https://doi.org/10.1111/jfs.12315>.
- Melero B, Manso B, Stessl B, Hernández M, Wagner M, Rovira J, Rodríguez-Lázaro D. 2019. Distribution and persistence of *Listeria monocytogenes* in a heavily contaminated poultry processing facility. *J Food Prot* 82:1524–1531. <https://doi.org/10.4315/0362-028X.JFP-19-087>.
- Cherifi T, Carrillo C, Lambert D, Miniaï I, Quessy S, Larivière-Gauthier G, Blais B, Fravalto P. 2018. Genomic characterization of *Listeria monocytogenes* isolates reveals that their persistence in a pig slaughterhouse is linked to the presence of benzalkonium chloride resistance genes. *BMC Microbiol* 18:220. <https://doi.org/10.1186/s12866-018-1363-9>.
- Chenal-Francois V, Lopez J, Cantinelli T, Caro V, Tran C, Leclercq A, Lecuit M, Brisse S. 2011. Worldwide distribution of major clones of *Listeria monocytogenes*. *Emerg Infect Dis* 17:1110–1112. <https://doi.org/10.3201/eid1706.101778>.
- Parisi A, Latorre L, Normanno G, Miccolupo A, Fracalvieri R, Lorusso V, Santagada G. 2010. Amplified fragment length polymorphism and multi-locus sequence typing for high-resolution genotyping of *Listeria monocytogenes* from foods and the environment. *Food Microbiol* 27:101–108. <https://doi.org/10.1016/j.fm.2009.09.001>.
- Ragon M, Wirth T, Hollandt F, Lavenir R, Lecuit M, Le Monnier A, Brisse S. 2008. A new perspective on *Listeria monocytogenes* evolution. *PLoS Pathog* 4:e1000146. <https://doi.org/10.1371/journal.ppat.1000146>.
- Moura A, Criscuolo A, Pouseele H, Maury MM, Leclercq A, Tarr C, Björkman JT, Dallman T, Reimer A, Enouf V, Larssonneur E, Carleton H, Bracq-

- Dieye H, Katz LS, Jones L, Touchon M, Toudjman M, Walker M, Stroika S, Cantinelli T, Chenal-Francisque V, Kucerova Z, Rocha EPC, Nadon C, Grant K, Nielsen EM, Pot B, Gerner-Smidt P, Lecuit M, Brisse S. 2016. Whole genome-based population biology and epidemiological surveillance of *Listeria monocytogenes*. *Nat Microbiol* 2:16185. <https://doi.org/10.1038/nmicrobiol.2016.185>.
20. Wang Y, Zhao AL, Zhu RF, Lan RT, Jin D, Cui ZG, Wang YL, Li ZC, Wang YT, Xu JG, Ye CY. 2012. Genetic diversity and molecular typing of *Listeria monocytogenes* in China. *BMC Microbiol* 12:119. <https://doi.org/10.1186/1471-2180-12-119>.
21. Chen M, Cheng J, Zhang J, Chen Y, Zeng H, Xue L, Lei T, Pang R, Wu S, Wu H, Zhang S, Wei X, Zhang Y, Ding Y, Wu Q. 2019. Isolation, potential virulence, and population diversity of *Listeria monocytogenes* from meat and meat products in China. *Front Microbiol* 10:946. <https://doi.org/10.3389/fmicb.2019.00946>.
22. Félix B, Feurer C, Maillat E, Guillier L, Boscher E, Kerouanton A, Denis M, Roussel S. 2018. Population genetic structure of *Listeria monocytogenes* strains isolated from the pig and pork production chain in France. *Front Microbiol* 9:684. <https://doi.org/10.3389/fmicb.2018.00684>.
23. Maury MM, Bracq-Dieye H, Huang L, Vales G, Lavina M, Thouvenot P, Disson O, Leclercq A, Brisse S, Lecuit M. 2019. Hypervirulent *Listeria monocytogenes* clones' adaptation to mammalian gut accounts for their association with dairy products. *Nat Commun* 10:2488. <https://doi.org/10.1038/s41467-019-10380-0>.
24. Nastasijevic I, Milanov D, Velebit B, Djordjevic V, Swift C, Painset A, Lakicevic B. 2017. Tracking of *Listeria monocytogenes* in meat establishment using whole genome sequencing as a food safety management tool: a proof of concept. *Int J Food Microbiol* 257:157–164. <https://doi.org/10.1016/j.jfoodmicro.2017.06.015>.
25. Morganti M, Scaltriti E, Cozzolino P, Bolzoni L, Casadei G, Pierantoni M, Foni E, Pongolini S. 2016. Processing-dependent and clonal contamination patterns of *Listeria monocytogenes* in the cured ham food chain revealed by genetic analysis. *Appl Environ Microbiol* 82:822–831. <https://doi.org/10.1128/AEM.03103-15>.
26. Mørseth T, Schirmer BCT, Heir E, Fagerlund A, Hjemli P, Langsrud S. 2017. Tolerance to quaternary ammonium compound disinfectants may enhance growth of *Listeria monocytogenes* in the food industry. *Int J Food Microbiol* 241:215–224. <https://doi.org/10.1016/j.jfoodmicro.2016.10.025>.
27. Pightling AW, Pettengill JB, Luo Y, Baugher JD, Rand H, Strain E. 2018. Interpreting whole-genome sequence analyses of foodborne bacteria for regulatory applications and outbreak investigations. *Front Microbiol* 9:1482. <https://doi.org/10.3389/fmicb.2018.01482>.
28. Wang Y, Pettengill JB, Pightling A, Timme R, Allard M, Strain E, Rand H. 2018. Genetic diversity of *Salmonella* and *Listeria* isolates from food facilities. *J Food Prot* 81:2082–2089. <https://doi.org/10.4315/0362-028X.JFP-18-093>.
29. Jagadeesan B, Baert L, Wiedmann M, Orsi RH. 2019. Comparative analysis of tools and approaches for source tracking *Listeria monocytogenes* in a food facility using whole-genome sequence data. *Front Microbiol* 10:947. <https://doi.org/10.3389/fmicb.2019.00947>.
30. Palma F, Pasquali F, Lucchi A, De Cesare A, Manfreda G. 2017. Whole genome sequencing for typing and characterisation of *Listeria monocytogenes* isolated in a rabbit meat processing plant. *Ital J Food Saf* 6:6879. <https://doi.org/10.4081/ijfs.2017.6879>.
31. Smith A, Hearn J, Taylor C, Wheelhouse N, Kaczmarek M, Moorhouse E, Singleton I. 2019. *Listeria monocytogenes* isolates from ready to eat plant produce are diverse and have virulence potential. *Int J Food Microbiol* 299:23–32. <https://doi.org/10.1016/j.jfoodmicro.2019.03.013>.
32. Gelbíčová T, Florianová M, Tomáščíková Z, Pospíšilová L, Koláčková I, Karpíšková R. 2019. Prediction of persistence of *Listeria monocytogenes* ST451 in a rabbit meat processing plant in the Czech Republic. *J Food Prot* 82:1350–1356. <https://doi.org/10.4315/0362-028X.JFP-19-030>.
33. Hurley D, Luque-Sastre L, Parker CT, Huynh S, Eshwar AK, Nguyen SV, Andrews N, Moura A, Fox EM, Jordan K, Lehner A, Stephan R, Fanning S. 2019. Whole-genome sequencing-based characterization of 100 *Listeria monocytogenes* isolates collected from food processing environments over a four-year period. *mSphere* 4:e00252-19. <https://doi.org/10.1128/mSphere.00252-19>.
34. Allard MW, Strain E, Rand H, Melka D, Correll WA, Hintz L, Stevens E, Timme R, Lomonaco S, Chen Y, Musser SM, Brown EW. 2019. Whole genome sequencing uses for foodborne contamination and compliance: discovery of an emerging contamination event in an ice cream facility using whole genome sequencing. *Infect Genet Evol* 73:214–220. <https://doi.org/10.1016/j.meegid.2019.04.026>.
35. Gerner-Smidt P, Besser J, Concepción-Acevedo J, Folster JP, Huffman J, Joseph LA, Kucerova Z, Nichols MC, Schwensohn CA, Tolar B. 2019. Whole genome sequencing: bridging one-health surveillance of food-borne diseases. *Front Public Health* 7:172. <https://doi.org/10.3389/fpubh.2019.00172>.
36. Chen Y, Luo Y, Curry P, Timme R, Melka D, Doyle M, Parish M, Hammack TS, Allard MW, Brown EW, Strain EA. 2017. Assessing the genome level diversity of *Listeria monocytogenes* from contaminated ice cream and environmental samples linked to a listeriosis outbreak in the United States. *PLoS One* 12:e0171389. <https://doi.org/10.1371/journal.pone.0171389>.
37. Jagadeesan B, Gerner-Smidt P, Allard MW, Leuillet S, Winkler A, Xiao Y, Chaffron S, Van Der Vossen J, Tang S, Katase M, McClure P, Kimura B, Ching Chai L, Chapman J, Grant K. 2019. The use of next generation sequencing for improving food safety: translation into practice. *Food Microbiol* 79:96–115. <https://doi.org/10.1016/j.fm.2018.11.005>.
38. Stasiewicz MJ, Oliver HF, Wiedmann M, den Bakker HC. 2015. Whole-genome sequencing allows for improved identification of persistent *Listeria monocytogenes* in food-associated environments. *Appl Environ Microbiol* 81:6024–6037. <https://doi.org/10.1128/AEM.01049-15>.
39. Pöntinen A, Aalto-Araneda M, Lindström M, Korkeala H. 2017. Heat resistance mediated by pLM58 plasmid-borne ClpL in *Listeria monocytogenes*. *mSphere* 2:e00364-17. <https://doi.org/10.1128/mSphere.00364-17>.
40. Chen P, Kong N, Huang B, Thao K, Ng W, Storey DB, Arabyan N, Foutouhi A, Foutouhi S, Weimer BC. 2017. 100K pathogen genome project: 306 *Listeria* draft genome sequences for food safety and public health. *Genome Announc* 5:e00967-16. <https://doi.org/10.1128/genomeA.00967-16>.
41. Kuenne C, Billion A, Mraheil MA, Strittmatter A, Daniel R, Goesmann A, Barbuddhe S, Hain T, Chakraborty T. 2013. Reassessment of the *Listeria monocytogenes* pan-genome reveals dynamic integration hotspots and mobile genetic elements as major components of the accessory genome. *BMC Genomics* 14:47. <https://doi.org/10.1186/1471-2164-14-47>.
42. Fagerlund A, Langsrud S, Moen B, Heir E, Mørseth T. 2018. Complete genome sequences of six *Listeria monocytogenes* sequence type 9 isolates from meat processing plants in Norway. *Genome Announc* 6:e00016-18. <https://doi.org/10.1128/genomeA.00016-18>.
43. Lindstedt BA, Tham W, Danielsson-Tham ML, Vardund T, Helmersson S, Kapperud G. 2008. Multiple-locus variable-number tandem-repeats analysis of *Listeria monocytogenes* using multicolour capillary electrophoresis and comparison with pulsed-field gel electrophoresis typing. *J Microbiol Methods* 72:141–148. <https://doi.org/10.1016/j.mimet.2007.11.012>.
44. Chenal-Francisque V, Diancourt L, Cantinelli T, Passet V, Tran-Hykes C, Bracq-Dieye H, Leclercq A, Pourcel C, Lecuit M, Brisse S. 2013. Optimized multilocus variable-number tandem-repeat analysis assay and its complementarity with pulsed-field gel electrophoresis and multilocus sequence typing for *Listeria monocytogenes* clone identification and surveillance. *J Clin Microbiol* 51:1868–1880. <https://doi.org/10.1128/JCM.00606-13>.
45. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. 2018. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evol* 4:vey016. <https://doi.org/10.1093/ve/vey016>.
46. Fagerlund A, Langsrud S, Schirmer BCT, Mørseth T, Heir E. 2016. Genome analysis of *Listeria monocytogenes* sequence type 8 strains persisting in salmon and poultry processing environments and comparison with related strains. *PLoS One* 11:e0151117. <https://doi.org/10.1371/journal.pone.0151117>.
47. Harrand AS, Jagadeesan B, Baert L, Wiedmann M, Orsi RH. 2020. Evolution of *Listeria monocytogenes* in a food processing plant involves limited single-nucleotide substitutions but considerable diversification by gain and loss of prophages. *Appl Environ Microbiol* 86:e02493-19. <https://doi.org/10.1128/AEM.02493-19>.
48. Henri C, Leekitcharoenphon P, Carleton HA, Radomski N, Kaas RS, Mariet JF, Felten A, Aarestrup FM, Gerner Smidt P, Roussel S, Guillier L, Mistou MY, Hendriksen RS. 2017. An assessment of different genomic approaches for inferring phylogeny of *Listeria monocytogenes*. *Front Microbiol* 8:2351. <https://doi.org/10.3389/fmicb.2017.02351>.
49. Teixeira AS, Monteiro PT, Carriço JA, Ramirez M, Francisco AP. 2015. Not seeing the forest for the trees: size of the minimum spanning trees (MSTs) forest and branch significance in MST-based phylogenetic analysis. *PLoS One* 10:e0119315. <https://doi.org/10.1371/journal.pone.0119315>.
50. Thevenot D, Dernburg A, Vernozy-Rozand C. 2006. An updated review of *Listeria monocytogenes* in the pork meat industry and its products. *J Appl Microbiol* 101:7–17. <https://doi.org/10.1111/j.1365-2672.2006.02962.x>.
51. Rothrock MJ, Davis ML, Locatelli A, Bodie A, McIntosh TG, Donaldson JR,

- Ricke SC. 2017. *Listeria* occurrence in poultry flocks: detection and potential implications. *Front Vet Sci* 4:125. <https://doi.org/10.3389/fvets.2017.00125>.
52. Luo LJ, Zhang ZD, Wang H, Wang PF, Lan RT, Deng JP, Miao YM, Wang Y, Wang Y, Xu JG, Zhang L, Sun SS, Liu X, Zhou Y, Chen X, Li Q, Ye CY. 2017. A 12-month longitudinal study of *Listeria monocytogenes* contamination and persistence in pork retail markets in China. *Food Control* 76:66–73. <https://doi.org/10.1016/j.foodcont.2016.12.037>.
 53. Autio T, Säteri T, Fredriksson-Ahomaa M, Rahkio M, Lundén J, Korkeala H. 2000. *Listeria monocytogenes* contamination pattern in pig slaughterhouses. *J Food Prot* 63:1438–1442. <https://doi.org/10.4315/0362-028x-63.10.1438>.
 54. Santorum P, García R, López V, Martínez-Suárez JV. 2012. Review. Dairy farm management and production practices associated with the presence of *Listeria monocytogenes* in raw milk and beef. *Span J Agric Res* 10:360–371. <https://doi.org/10.5424/sjar/2012102-314-11>.
 55. Hellström S, Laukkanen R, Siekkinen KM, Ranta J, Majjala R, Korkeala H. 2010. *Listeria monocytogenes* contamination in pork can originate from farms. *J Food Prot* 73:641–648. <https://doi.org/10.4315/0362-028x-73.4.641>.
 56. Camargo AC, Dias MR, Cossi MVC, Lanna F, Cavicchioli VQ, Vallim DC, Pinto PSD, Hofer E, Nero LA. 2015. Serotypes and pulsotypes diversity of *Listeria monocytogenes* in a beef-processing environment. *Foodborne Pathog Dis* 12:323–326. <https://doi.org/10.1089/fpd.2014.1875>.
 57. Kjos AK, Nafstad O, Odden H, Ruud TA, Saltnes T, Ytterdahl M. 2019. Kjøttets tilstand 2019. Status i norsk kjøtt- og eggproduksjon. *Animalia AS*, Oslo, Norway. <https://www.animalia.no/contentassets/3dce35cde68a47b091097fa8c6ec2dd5/kjottets-tilstand-2019.pdf>.
 58. Steckler AJ, Cardenas-Alvarez MX, Townsend Ramsett MK, Dyer N, Bergholz TM. 2018. Genetic characterization of *Listeria monocytogenes* from ruminant listeriosis from different geographical regions in the U.S. *Vet Microbiol* 215:93–97. <https://doi.org/10.1016/j.vetmic.2017.12.021>.
 59. Papić B, Pate M, Félix B, Kušar D. 2019. Genetic diversity of *Listeria monocytogenes* strains in ruminant abortion and rhombencephalitis cases in comparison with the natural environment. *BMC Microbiol* 19:299. <https://doi.org/10.1186/s12866-019-1676-3>.
 60. Gismervik K, Aspholm M, Rørvik LM, Bruheim T, Andersen A, Skaar I. 2015. Invading slugs (*Arion vulgaris*) can be vectors for *Listeria monocytogenes*. *J Appl Microbiol* 118:809–816. <https://doi.org/10.1111/jam.12750>.
 61. Ma A, Wang Y, Wang Y, Li D, Xu H, Yuan X, Liu K, Ye C. 2015. Molecular epidemiological characteristics of *Listeria monocytogenes* isolated from raw meat samples in some regions of Beijing, China. *Chinese J Zoonoses* 31:403–407.
 62. De Cesare A, Parisi A, Mioni R, Comin D, Lucchi A, Manfreda G. 2017. *Listeria monocytogenes* circulating in rabbit meat products and slaughterhouses in Italy: prevalence data and comparison among typing results. *Foodborne Pathog Dis* 14:167–176. <https://doi.org/10.1089/fpd.2016.2211>.
 63. Palma F, Brauge T, Radomski N, Mallet L, Felten A, Mistou MY, Brisabois A, Guillier L, Midelet-Bourdin G. 2020. Dynamics of mobile genetic elements of *Listeria monocytogenes* persisting in ready-to-eat seafood processing plants in France. *BMC Genomics* 21:130. <https://doi.org/10.1186/s12864-020-6544-x>.
 64. Heir E, Mørtrø T, Simensen A, Langsrud S. 2018. *Listeria monocytogenes* strains show large variations in competitive growth in mixed culture biofilms and suspensions with bacteria from food processing environments. *Int J Food Microbiol* 275:46–55. <https://doi.org/10.1016/j.ijfoodmicro.2018.03.026>.
 65. Verghese B, Lok M, Wen J, Alessandria V, Chen Y, Kathariou S, Knabel S. 2011. *comK* prophage junction fragments as markers for *Listeria monocytogenes* genotypes unique to individual meat and poultry processing plants and a model for rapid niche-specific adaptation, biofilm formation, and persistence. *Appl Environ Microbiol* 77:3279–3292. <https://doi.org/10.1128/AEM.00546-11>.
 66. Aronesty E. 2011. ea-utils: command-line tools for processing biological sequencing data. <https://github.com/ExpressionAnalysis/ea-utils>. Accessed 3 April 2014.
 67. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Son P, Pribelski AD, Pyshtkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
 68. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
 69. Applied Maths. 2015. *Listeria monocytogenes* whole genome sequence typing. <http://www.applied-maths.com/news/listeria-monocytogenes-whole-genome-sequence-typing>. Accessed 2 December 2019.
 70. Clausen P, Aarestrup FM, Lund O. 2018. Rapid and precise alignment of raw reads against redundant databases with KMA. *BMC Bioinformatics* 19:307. <https://doi.org/10.1186/s12859-018-2336-6>.
 71. Lund O. 2014. cgMLSTfinder. <https://bitbucket.org/genomicpidemiology/cgmlstfinder>. Accessed 6 February 2019.
 72. Seemann T. 2014. MLST. <https://github.com/tseemann/mlst>. Accessed 19 April 2017.
 73. Jolley KA, Maiden MC. 2010. BIGSdb: scalable analysis of bacterial genome variation at the population level. *BMC Bioinformatics* 11:595. <https://doi.org/10.1186/1471-2105-11-595>.
 74. Painset A, Björkman JT, Kiil K, Guillier L, Mariet JF, Félix B, Amar C, Rotariu O, Roussel S, Perez-Reche F, Brisse S, Moura A, Lecuit M, Forbes K, Strachan N, Grant K, Møller-Nielsen E, Dallman TJ. 2019. LiSEQ—whole-genome sequencing of a cross-sectional survey of *Listeria monocytogenes* in ready-to-eat foods and human clinical cases in Europe. *Microb Genom* 5:000257. <https://doi.org/10.1099/mgen.0.000257>.
 75. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>.
 76. Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv 1207.3907 [q-bio.GN]* <https://arxiv.org/abs/1207.3907>.
 77. Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, Genomes Project Analysis Group. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>.
 78. Bergey CM. 2012. vcf-tab-to-fastq. <http://code.google.com/p/vcf-tab-to-fastq>. Accessed 12 November 2015.
 79. Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35:526–528. <https://doi.org/10.1093/bioinformatics/bty633>.