

Evolutionary history of a Scottish harbour seal population

Natacha Nikolic^{1,2}, Paul Thompson³, Mark de Bruyn⁴, Matthias Macé⁵ and Claude Chevalet²

¹ ARBRE (Reunion Island Biodiversity Research Agency), Saint-Leu, La Réunion

² Génétique Physiologie et Systèmes d'Élevage - UMR1388, INRAE, Castanet Tolosan, France

³ Lighthouse Field Station, Sciences School of Biological Sciences, University of Aberdeen, Cromarty, United Kingdom

⁴ School of Life and Environmental Sciences, University of Sydney, Sydney, Australia

⁵ Laboratoire d'Anthropologie Moléculaire et d'Imagerie de Synthèse - UMR 5288, CNRS, Toulouse, France

ABSTRACT

Efforts to conserve marine mammals are often constrained by uncertainty over their population history. Here, we examine the evolutionary history of a harbour seal (*Phoca vitulina*) population in the Moray Firth, northeast Scotland using genetic tools and microsatellite markers to explore population change. Previous fine-scale analysis of UK harbour seal populations revealed three clusters in the UK, with a northeastern cluster that included our Moray Firth study population. Our analysis revealed that the Moray Firth cluster is an independent genetic group, with similar levels of genetic diversity across each of the localities sampled. These samples were used to assess historic abundance and demographic events in the Moray Firth population. Estimates of current genetic diversity and effective population size were low, but the results indicated that this population has remained at broadly similar levels following the population bottleneck that occurred after post-glacial recolonization of the area.

Subjects Evolutionary Studies, Genetics, Population Biology

Keywords Evolution, Genetic, Seal

Submitted 15 January 2020

Accepted 19 April 2020

Published 10 July 2020

Corresponding author
Natacha Nikolic,
natachanikolic@hotmail.com

Academic editor
Elizabeth King

Additional Information and
Declarations can be found on
page 20

DOI 10.7717/peerj.9167

© Copyright
2020 Nikolic et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

INTRODUCTION

Efforts to conserve marine animals are frequently constrained by uncertainty over historic baselines and the factors driving changes in abundance (Lotze & Worm, 2009). A variety of techniques have been developed to address this issue, including archaeological investigations (Rick & Lockwood, 2013), studies based upon historical records or traditional knowledge (McClenachan, Ferretti & Baum, 2012), and molecular analyses of changes in genetic diversity (Roman & Palumbi, 2003;ref-101).

Harbour seals (*Phoca vitulina*) are widely distributed around the North Atlantic and North Pacific coasts, but studies over recent decades have identified wide variations in the status of these populations. Abundance in many parts of Europe was severely reduced by successive outbreaks of Phocine Distemper Virus (PDV), but mortality in some populations was low (Heide-Jørgensen, Härkönen & Aberg, 1992; Härkönen et al., 2006). Similarly, longer-term trends in harbour seal abundance show steady increases in some parts of their range (Jeffries et al., 2003; Aarts et al., 2019), whereas other regions have

experienced unexplained declines ([Boveng et al., 2003](#); [Lonergan et al., 2007](#); [Hanson et al., 2015](#); [Thompson et al., 2019](#)).

These changes have resulted in new conservation measures in many areas, the most significant of which in European waters is the EU Habitats Directive ([Baxter, 2001](#)). This requires EU Member States to develop a NATURA 2000 network of Special Areas of Conservation (SACs) for a range of key species, including harbour seals ([Thompson et al., 2019](#)). In UK waters, SACs have been established within eleven management units that reflect the current understanding of genetic structure ([Olsen et al., 2017](#)), but there is uncertainty over the causes underlying different regional trends in abundance and historic baselines ([Matthiopoulos et al., 2014](#); [Thompson et al., 2019](#)). Here, we focus on the evolutionary history of a population that uses one of these SACs, in the Moray Firth, NE Scotland. In this area, individual-based studies since 2006 have provided detailed estimates of contemporary vital rates ([Cordes & Thompson, 2014](#); [Matthiopoulos et al., 2014](#)) during a period where there has been no clear trend in abundance ([Thompson et al., 2019](#)). The most recent counts indicate that the population has declined by around 40% since the mid 1990's ([Thompson et al., 2019](#)), but direct information on the historic abundance and evolution over multiple generations is lacking given the lack of comparable survey data prior to this. An alternative approach is to assess historic changes in the genetic-based indicator, effective population size (N_e). The effective size reflects the abundance and evolutionary history of the population, and can inform conservation efforts because it affects the degree to which a population can respond to selection ([Berthier et al., 2002](#)).

Analysis of neutral molecular markers such as microsatellites can be used to calculate N_e and provide an evolutionary perspective to these conservation and management issues ([King et al., 2001](#)). Genetic studies conducted on harbour seals have generally used less than 15 microsatellite markers for “Atlantic” harbour seals (12 in [Olsen et al. \(2017\)](#); 15 in [Anderse et al. \(2011\)](#); 7 in [Goodman \(1997\)](#); [Goodman \(1998\)](#)) and less than 20 for “Pacific” harbour seals (5 in [Burg et al. \(1999\)](#), 8 in [Curtis, Stewart & Karl \(2011\)](#), 7 in [Dishman \(2011\)](#), 20 in [Hayes et al. \(2006\)](#) and 6 in [Herreman et al., 2009](#)). We used the markers developed specifically for harbour seal microsatellites ([Allen et al., 1995](#); [Coltman, Bowen & Wright, 1996](#); [Goodman, 1997](#)) and a mix of pinniped microsatellite markers ([Allen et al., 1995](#); [Buchanan et al., 1998](#); [Davis et al., 2002](#); [Gelatt et al., 2001](#)), as other harbour seal microsatellite papers (ex. [Olsen et al., 2017](#)). We therefore tested 30 markers, which allowed us to utilise 17 polymorphic microsatellite markers (25 markers minus 8 monomorphic markers) for genetic analysis of harbour seals. Previous study of genetic structuring of UK harbour seals suggests that there are two main initial groups consisting of localities in the northern UK, and the southern UK and mainland Europe ([Olsen et al., 2017](#)). However, [Olsen et al. \(2017\)](#) further divided these groups into geographically distinct genetic clusters, including the North-East England and Eastern Scotland cluster (cf. Fig. 2 of [Olsen et al., 2017](#)) that includes our Moray Firth study sites (cf. Fig. 1 of [Hanson et al., 2015](#)). Here, we extended the number of microsatellite markers available for harbour seals in the Moray Firth to estimate historic changes in effective population size in this region.

Using this extended set of microsatellite markers, the objectives of the present study were to address the following questions: (i) what does current genetic diversity tell us about




Figure 1 Mean geographic location of the three areas sampled (Dornoch –13 individuals, Cromarty –12 individuals and Inverness –68 individuals) from Moray Firth in north-east Scotland.

Full-size [DOI: 10.7717/peerj.9167/fig-1](https://doi.org/10.7717/peerj.9167/fig-1)

historic changes of harbour seal effective population sizes; and (ii) how might genetic data contribute to understanding the decline of seals in the north-east of Scotland?

MATERIAL & METHODS

Study area

The Moray Firth study population (Fig. 1) contained an estimated 1,653 harbour seals in 1993 (Thompson *et al.*, 1997a) when the samples used in this study were collected. Seals come ashore at inter-tidal sites throughout the year, with most pups born in three sub-areas spaced approximately 50 km apart (Dornoch Firth, Cromarty Firth, and Beaully Firth). One of these sub-areas, in the Dornoch Firth, has been designated as a SAC to protect harbour seal populations under the EU Habitats Directive (Cordes *et al.*, 2011; Thompson *et al.*, 2019).

Samples and DNA extraction

Archived blood samples were collected from 93 harbour seals that had been captured, sampled and released as part of earlier ecological studies (Thompson *et al.*, 1997b; Hall *et al.*, 2019). Samples were collected between 1992 and 1995 (see Thompson *et al.*, 1997b for details of capture and sampling methods), and constituted 47 females and 46 males (48 juveniles, 37 adults, 8 sub-adults, data are available at <https://doi.org/10.15454/AOZ7JI>); representing approximately 6% of the population. All capture and handling methods were

carried out in accordance with the approved guidelines and conducted under licences from the UK Home Office. Blood samples were collected under Home Office licence issued to the University of Aberdeen under the Animal (Scientific Procedures) Act 1986 (PPL number 60/01351). These samples were preserved at -20°C and hydrated with PBS (Phosphate Buffered Saline) before DNA extraction. Genomic DNA was extracted by QIAamp DNA Blood Mini Kit (QIAGEN) (see the [Text S1](#) for more details).

Marker selection and genotyping

Because the number of available harbour seal microsatellite markers was low, we considered all potential microsatellites for this species and additionally from other pinnipeds—*Halichoerus grypus*, *Hydrurga leptonyx* (two species belonging to the family Phocidae) and *Odobenus rosmarus rosmarus* (Odobenidae). A total of 30 markers were tested [Appendix S1A](#). Previously some primers were designed using Primer 3 ([Rozen & Skaletsky, 2000](#)) because only the amplified sequences were provided by the authors (ie the primers H12, HL20, HL16, HL15) [Appendix S1B](#). The primers PVC63 were redefined using Primer 3 to optimize amplification. We retained 25 loci that amplified reliably (meaning that on 12 individuals tested a minimum of 10 were successfully amplified and visualized on agarose gel) for harbour seals [Appendix S1B](#), with a minimum of twelve repeat motifs and a GC percentage of approximately 50%.

All 93 harbour seal samples were successfully genotyped with these 25 microsatellite markers using fluorescent labelled primers and multiplex PCR pools ([Appendices S1A, S1B](#)). PCR amplification was carried out using an Applied Biosystems 2720 Thermal Cycler with 10 μl reaction volume containing ~ 50 ng DNA, 1.5 mM MgCl_2 , 1x Promega buffer, 200 μM dNTPs, 0.5 μM each primer, 0.5 U Taq DNA polymerase (Promega). An initial denaturation step at 94°C for 5 min was followed by 42–45 cycles of 30 s at 94°C , 30 s at $T_m^{\circ}\text{C}$ (annealing temperature), 30 s at 72°C , followed by a final elongation step of 30 min at 72°C . The annealing temperatures were optimised for each locus [Appendix S1B](#). The PCR products (2 μl) were added to a mixture of deionised formamide and the internal size standard GENESCAN-400HD Rox (Applied Biosystems) (8 μl), then denatured for 5 min at 95°C . For this mixture (formamide and internal size standard), we prepared a final volume of 1000 μl with 982.5 μl of formamide and 17.5 μl GENESCAN-400HD Rox. Individual electropherograms were obtained using an ABI 3730 multi-capillary sequencer. PCR products were visualised by GeneMapper v4.0 software (Applied Biosystems) (see the data and profile examples per pool at <https://doi.org/10.15454/AOZ7JI>). Across the panel of 25 markers, 8 were monomorphic [Appendix S1B](#) in both male and female individuals. The genetic analysis was performed on the overall panel (25 markers) and on the polymorphic markers (17 markers).

Analysis of diversity and genetic differentiation

To ensure that the number of loci was sufficient, we calculated the probability of individual identity (PI; the probability that two individuals in a population have identical genotypes) for each locus and their combinations with the program GENALEX ([Peakall & Smouse, 2006](#); [Peakall & Smouse, 2012](#)). SPOTG ([Hoban, Gaggiotti & Bertorelle, 2013](#)) was used to

estimate the power of individual assignment, using 1,000 runs, the number of sampled individuals (93), the mean number of alleles (4), and the F_{ST} values found in this study under a normal allele frequencies model.

Genetic diversity was measured as the mean number of alleles per locus (A) for 25 markers. Excluding the monomorphic markers we measured observed heterozygosity (H_o), expected heterozygosity (H_e), and Nei 's (1978) unbiased heterozygosity ($H_{n.b}$) using GENETIX 4.05.2 (Belkhir et al., 1998). Estimates of homozygote and heterozygote excess that differed significantly from zero ($p < 0.05$) were calculated from the standard error rates with confidence intervals in Pedant (Johnson & Haydon, 2007). Deviations from Hardy-Weinberg Equilibrium (HWE) were assessed using polymorphic loci, and exact tests with p -values and their standard errors were computed using ARLEQUIN version 3.1 (Excoffier, Laval & Schneider, 2005; Excoffier & Lischer, 2010) with permutations (1,000,000 chains and 100,000 steps). Polymorphism information content (PIC) was generated in Cervus (Kalinowski, Taper & Marshall, 2007). Probability of parentage exclusion (PE1, single parent (Jamieson & Taylor, 1997); PE2, a second parent given a first parent assigned (Jamieson, 1994); PE3, a pair of parents (Jamieson & Taylor, 1997)) was estimated per locus using INEst (Chybicki & Burczyk, 2009). The potential occurrence of null alleles and scoring errors due to stuttering or large allele dropout in the data set was assessed using the software MICRO-CHECKER (Oosterhout et al., 2004), and the significance of null allele frequency (F_{null}) was estimated with INEst using the individual inbreeding model with 100,000 iterations (estimates significantly different from zero, $p < 0.05$). The inbreeding coefficient F_{IS} was estimated from polymorphic loci with 10,000 bootstraps by GENETIX software. F_{IS} measures the decrease in heterozygosity due to inbreeding, assortative mating, or selection. Genotyping error rate per allele, E1 referring to allelic dropout rate and E2 to the false allele rate, and the 95% confidence interval (CI) with 10,000 permutations, were evaluated with maximum likelihood from 10 individuals' random replicates by marker based on H_e computed in Pedant. The number of repeated genotypes (Nrep) and the percentage (%) of the total number of individuals genotyped for each loci were also estimated.

Our findings were consistent with the previous study identifying Moray Firth sites as part of one genetic cluster (Olsen et al., 2017). Thus, we also investigated whether external migrations (e.g., gene flow from another cluster) were possible because grouping several different populations could skew our analyses of evolutionary history. Genetic differentiation was visualized by Factorial Correspondence Analysis (FCA) in GENETIX using the polymorphic loci ($n = 17$). Pairwise Wright's F -statistics F_{ST} (Weir & Cockerham, 1984) and their levels of significance were assessed based on 10,000 permutations using ARLEQUIN 3.1 (Excoffier, Laval & Schneider, 2005), considering geographical location per individual. Number of migrants (N_m) was estimated through the frequency of private alleles with GENEPOP (Raymond & Rousset, 1995; Rousset, 2008) online (<http://genepop.curtin.edu.au>). We estimated only the level of migration rates (m) (superior or inferior to 0.1, and not the values of gene flow due to uncertainty with a weak genetic structure) using the Bayesian multi-locus genotyping approach implemented in BAYESASS 3.0.4 (Wilson & Rannala, 2003) with different random number seed values

(S) and sampling frequency (n), 10,000,000 iterations, and a burn-in period of 1,000,000. To detect immigration into the Moray Firth, we used *Rannala & Mountain's (1997)* estimate and ran Monte-Carlo simulations with 10,000 repetitions implemented in GENECLASS2 software (*Piry et al., 2004*), considering geographical location per individual.

Finally, we used a population assignment test to check for sex-biased dispersal using the software GENALEX. This method produces an Assignment Index correction (*AIc*) for each sex following the method of *Mossman & Waser (1999)*. Negative *AIc* values characterize individuals with a higher probability of being migrants (high dispersion) and positive values characterize individuals with a lower probability of being migrants. Mean *AIc* values were compared for each sex with a non-parametric Mann–Whitney U-test using R version 3.0.2 software. Moreover, hierarchical Analysis of Molecular Variance (AMOVA) analyses were performed, with 9,999 permutations, for each sex on their own, and for the combined male and female data set to determine whether genetic variation was similar for males and females.

Clustering analysis

STRUCTURE 2.3 (*Pritchard, Stephens & Donnelly, 2000*) was used to identify population structure and individual admixture coefficients. Five independent runs were performed on STRUCTURE for each assumed number of population(s) $K = 1-6$ under an admixture model. All runs were executed with 50,000 burn-in periods and 200,000 MCMC (Markov chain Monte Carlo) repetitions, using the three regions defined earlier as prior information. STRUCTURE HARVESTER 0.6.94 (*Earl Dent & VonHoldt Bridgett, 2012*) was used to visualise results and assess *K*, the number of genetic populations that best fit the data, based on Maximum Likelihood (*Evanno & Goudet, 2005*). Structure Selector (<http://lmme.qdio.ac.cn/StructureSelector/>) was also used to process the *Puechmaile (2016)* approach on uneven sampling and to also estimate the best *K*. Plots for optimal *K* was performed with CLUMPAK (*Kopelman et al., 2015*).

In addition, a discriminant analysis of principal components (DAPC, *Jombart, Devillard & Balloux, 2010*) was performed using R package ADEGENET (*Jombart, 2008; Jombart & Ahmed, 2011*). DAPC is a multivariate analysis that integrates principal component analysis (PCA) with discriminant analysis to summarize genetic differentiation between groups (*Jombart, 2008*). Sampling location was used as prior. While STRUCTURE forms genetic clusters of individuals by minimizing departure from Hardy–Weinberg and linkage disequilibria, DAPC maximizes genetic separation among groups and minimizes variation within groups (*Jombart, Devillard & Balloux, 2010*) which may constitute a more accurate approach for species exhibiting potentially high gene flow (*Bailleul et al., 2018*).

Isolation by distance was tested using a Mantel test between genetic (Euclidean Edwards' distance) and geographic distances with 10,000 resamplings between individuals and regions using ADEGENET, ADE4 (*Thioulouse et al., 1997*) and GRDEVICES (R Development Core Team and contributors worldwide) R packages.

Effective population sizes and evolutionary history

The average mutation rate on the set of loci (μ), ancestral time (*Tf*), and present and ancestral effective size (*No* and *Na*) were estimated for the seals using MSVAR (*Beaumont,*

1999), running 80,000 MCMC chains and 20,000 iterations between chains. This method is suited to microsatellite data that are assumed to be evolving by a stepwise mutation model (SMM), sampled from a population that has varied in size. When compared with other classic coalescence models (TM3 and DIYABC) aimed at estimating the present and ancestral effective sizes, MSVAR was most efficient for small populations (Nikolic et al., 2009). Because we used this model only to get preliminary results for the seal mutation rate, and computation times are very long (here more than 3 days), this analysis was run only with the polymorphic panel (17 markers). MSVAR estimates the effective population size for the present and ancestral effective sizes but does not infer fluctuation of effective size between them. We applied Gelman & Rubin, 1992's (1992) test to monitor convergence of MCMC output.

To check for historic population declines in the seals, we used BOTTLENECK software (Piry, Luikart & Cornuet, 1999) with 10,000 iterations and we applied three tests—the sign test, the standardized differences test (Cornuet & Luikart, 1997), and the Wilcoxon sign-rank test (Luikart et al., 1997) to analyze the presence of heterozygote excess resulting from perturbation of allele frequencies. We applied the Wilcoxon signed-rank test specifically as it does not require a large number of polymorphic loci which are scarce in a population with low variability (Han et al., 2010). Inferences from heterozygosity excess or deficiency tests are heavily influenced by the mutational model (Busch, Waser & DeWoody, 2007). Here, we used the Stepwise Mutation model as it is thought to be a more appropriate model for use with microsatellites (Nikolic & Chevalet, 2014b).

Bayesian methods using coalescence theory and MCMC sampling to estimate posterior distributions of demographic parameters and history, seem more robust to certain violations of mutation model assumptions (Girod et al., 2011) and bottleneck duration (Peery et al., 2012). Hence, to estimate historic fluctuations of effective population size from microsatellite markers, we used the algorithm and method VarEff (Chevalet & Nikolic, 2010; Nikolic & Chevalet, 2014b) programming in R package. The model assumes a stepwise mutation model for microsatellites and makes use of an approximate likelihood of data based on theoretical results. It is then implemented in a MCMC framework which simulates past demography by sampling step functions. The model is freely available as an R package (<https://forge-dga.jouy.inra.fr/projects/package-vareff-variation-of-effective-population-size/>). Results were based on MCMC chains including 10,000 dememorization steps, a total length of 1,000,000 and the extraction of 10,000 uncorrelated states as suggested by previous analysis (Nikolic & Chevalet, 2014b). Results are provided in normalized scales $\hat{\theta} = 4N\mu$ for population size ($4 \times$ effective size \times mutation rate) and $T = g\mu$ for time (number of generations \times mutation rate), or in the natural scale, effective size N_e and generation number g provided the mutation rate μ is known. For harbour seals, we set $\mu = 0.00015$ and used the generation time of 8.75 years based upon the value used in Swart, Reijnders & Van Delden (1996).

VarEff offers several functions to characterize the posterior distribution of effective size at several times in the past: arithmetic and harmonic means, median and mode of the distribution, detailed distribution at specified times, with the quantiles and standard deviation. In addition, the results of VarEff allow the posterior distribution of the Time

to the Most Recent Common Ancestor (TMRCA) of two random alleles to be recovered, which provides complementary information on the occurrence and time of past bottlenecks, since peaks in this distribution indicate times when coalescence events likely occurred, and intervals between peaks may indicate periods when bottlenecks occurred. Depending on the intensity and shape of the peak and the value of effective size (obtained with the previous functions), information on the bottleneck event(s) can be recovered. Furthermore, because migration can mimic the effects of bottlenecks (Nikolic & Chevalet, 2014b), we repeated the same analysis discarding the immigrants that were detected (2 individuals).

RESULTS

Genetic polymorphism and panel of markers

Of the 25 microsatellite loci, eight loci were monomorphic for both sexes (*PVC63*, *PVC74*, *SGPV3*, *PVC26*, *PVC29*, *GS1*, *HL-16*, and *OrrFCB23*) and one marker *HL-20* was monomorphic in males (Appendix S1B). These markers were hence excluded from the analyses resulting in a final panel of 17 polymorphic markers (Table 1). Our final genotyping data contained only 3.2% missing data for 93 individuals and 17 polymorphic markers.

The probability of identity (PI) values ranged from 0.023 to 0.927 and the probability of exclusion (E1, E2, E3) from 0.001 to 0.921 (Table 1). A total of 9 markers had a probability of identity in the higher range at 0.5 (Table 1). The analysis based on the cumulated probability of individual identity (PI), an indication of the statistical power of marker loci, revealed that 10 to 17 polymorphic markers were sufficient to carry out the population genetics analysis of harbour seals in the Moray Firth (Fig. 2). SPOTG estimated that 93 individuals and 17 microsatellite markers could detect power of individual assignment >57% and connectivity >87%. However, the theoretical study by Nikolic & Chevalet (2014b) shows that the number of markers to assess accurate effective population size must be higher than the basic diversity analyses. It was therefore important to have more than 10 markers for this study to reduce the variance of the estimators around the effective size values.

The total number of alleles for the remaining 17 loci ranged between 2 and 14 (Table 1), while the mean number of alleles was 3.7 (Table 1). The global mean of observed heterozygosity was 0.384 for the 17 polymorphic microsatellites. Estimates of homozygote and heterozygote excess were not significant except *OrrFCB24* with a heterozygote excess (Table 1).

Genotyping error rates and associated 95% confidence intervals were very close to zero for all loci except *HL20* and *OrrFCB1* (Table 1). Most loci were close to HW equilibrium ($P > 0.05$, Bonferroni correction applied) (Table 1) except *SGPV11*, *SGPV17*, *PVC78* and *OrrFCB24* (Table 1). The locus *SGPV17* has been suspected to be X-linked in pinniped species (Coltman, Bowen & Wright, 1996; Gemmell et al., 1997; Pastor et al., 2004), although this remains equivocal (Herreman, Blundell & Ben-David, 2008). Such a linkage would imply that males are homozygous for this marker. In the present study, we identified 3 alleles (153, 155 and 161) at *SGPV17* with similar allelic frequencies in males and females. Heterozygote frequencies were 14/46 in males and 24/47 in females suggesting no X-linkage

Table 1 Summary statistics of the 17 microsatellite markers selected for Harbour seals (*Phoca vitulina*). Sample sizes per locus (S). Number of alleles (A). Expected (He), unbiased Nei's⁹⁵ expected (H.n.b) and observed (HO) heterozygosity. Hardy–Weinberg equilibrium (HWE) *p*-values (P) with the standard error in parentheses. Polymorphism information content (PIC). Probability of identity (PI). Probability of parentage exclusion (PE1, single parent; PE2, a second parent given a first parent assigned; PE3, a pair of parents). Null allele frequency (Fnull). Number of repeated genotypes (Nrep and percentage (%)) of the total number of individuals genotyped for each loci). Genotyping error rate per allele, E1 referring to allelic dropout rate and E2 to the false allele rate, and the 95% confidence interval (CI). Significant values are highlighted in bold ($P < 0.05$) for heterozygote excess.

Locus	S	A	He	H.n.b	Ho	P	PIC	PI	PE1	PE2	PE3	Fnull	Nrep(%)	Genotyping error rate	
														E1 (CI 95%)	E2 (CI 95%)
SGPv9	85	3	0.277	0.279	0.329	0.254 (0.00038)	0.242	0.558	0.038	0.123	0.205	0.023	10 (12)	0.00 (0.00–0.41)	0.00 (0.00–0.07)
SGPV11	92	3	0.275	0.276	0.174	0.002 (0.00004)	0.240	0.561	0.038	0.122	0.203	0.107	10 (11)	0.00 (–0.00–0.78)	0.00 (0.00–0.07)
SGPV17	92	3	0.409	0.411	0.413	0.019 (0.00013)	0.345	0.413	0.084	0.184	0.292	0.039	10 (11)	0.00 (–0.00–0.20)	0.00 (–0.00–0.07)
SGPV10	90	2	0.231	0.232	0.267	0.352 (0.00047)	0.204	0.618	0.027	0.102	0.174	0.026	10 (11)	0.00 (0.00–0.40)	0.00 (–0.00–0.07)
SGPV16	74	14	0.886	0.892	0.919	0.659 (0.00033)	0.876	0.023	0.628	0.772	0.921	0.013	10 (14)	0.00 (0.00–0.08)	0.00 (–0.00–0.07)
PVC19	93	2	0.350	0.352	0.344	1.000 (0.00000)	0.289	0.484	0.061	0.144	0.229	0.039	10 (11)	0.00 (–0.00–0.19)	0.00 (0.00–0.07)
PVC30	82	4	0.493	0.496	0.476	0.767 (0.00037)	0.388	0.362	0.122	0.204	0.310	0.039	10 (12)	0.00 (–0.00–0.21)	0.00 (0.00–0.07)
PVC78	93	4	0.243	0.245	0.204	0.003 (0.00005)	0.219	0.597	0.030	0.113	0.193	0.058	10 (11)	0.00 (–0.00–0.19)	0.00 (0.00–0.07)
GS7	93	2	0.157	0.158	0.172	1.000 (0.00000)	0.145	0.723	0.012	0.072	0.129	0.033	10 (11)	0.00 (–0.00–0.71)	0.00 (0.00–0.07)
GS2	91	3	0.173	0.174	0.165	0.557 (0.00049)	0.163	0.694	0.015	0.086	0.156	0.047	10 (11)	0.00 (–0.00–0.25)	0.00 (0.00–0.07)
GS3	93	4	0.560	0.563	0.570	0.579 (0.00048)	0.491	0.262	0.158	0.294	0.442	0.022	10 (11)	0.00 (–0.00–0.16)	0.00 (0.00–0.07)
H12	93	5	0.614	0.617	0.613	0.901 (0.00025)	0.550	0.213	0.194	0.345	0.504	0.029	10 (11)	0.00 (–0.00–0.09)	0.00 (0.00–0.07)
HL20	90	2	0.043	0.044	0.044	1.000 (0.00000)	0.043	0.916	0.001	0.021	0.041	0.050	9 (10)	0.70 (–0.03–1.81)	0.00 (–0.00–0.14)
HL15	93	3	0.242	0.243	0.237	0.707 (0.00043)	0.215	0.601	0.029	0.109	0.186	0.042	10 (11)	0.00 (–0.00–0.76)	0.00 (0.00–0.07)
OrrFCB2	90	4	0.547	0.550	0.633	0.070 (0.00022)	0.489	0.263	0.151	0.296	0.449	0.016	10 (11)	0.00 (–0.00–0.11)	0.00 (–0.00–0.07)
OrrFCB1	79	2	0.013	0.013	0.013	1.000 (0.00000)	0.012	0.927	0.001	0.019	0.037	0.055	10 (13)	0.70 (–0.03–1.81)	0.00 (0.00–0.12)
OrrFCB24	90	3	0.510	0.513	0.956	0.000 (0.00000)	0.391	0.359	0.130	0.202	0.303	0.009	9 (10)	0.00 (0.00–0.08)	0.00 (–0.00–0.08)
Mean	89	3.7	0.354	0.356	0.384	0.522 (0.000)	0.312								




Figure 2 Probability of Identity for each Locus (PI) and for Increasing Combinations (PIsibs) with the 17 polymorphic microsatellite markers genotyped on harbour seals in the Moray Firth.

Full-size [DOI: 10.7717/peerj.9167/fig-2](https://doi.org/10.7717/peerj.9167/fig-2)

for this locus. Analysis of sex-biased dispersal using the marker *SGPV17* showed there were no significant differences in the mean AIC with the polymorphic panel (Mann–Whitney U -test, p -value = 0.78). This result indicates no sex-biased dispersal. Thus, we concluded that we have no reason to exclude the marker *SGPV17* in the polymorphic panel on the basis of linkage.

MICROCHECKER estimated that only *SGPV11* showed evidence for a null allele but the test by permutations with INest revealed that the result was not significant (Table 1). Null allele analyses were not significant for all loci (Table 1). Hence, there was no evidence for scoring error due to stuttering and no evidence of large allele dropout. The harbour seal population appears to be in Hardy Weinberg equilibrium with the global test ($P = 0.519$).

With respect to the distribution of repeat numbers Appendix S1C, harbour seals in the Moray Firth exhibited quite large distances between microsatellite alleles (up to 37 repeat numbers) and a large range of missing values (between 10 and 24, and between 26 and 36) that may be the signature of bottleneck events.

Genetic differentiation and clustering analysis

Analysis of sex-biased dispersal (Mossman & Waser, 1999) showed no significant differences in the mean AIC with the polymorphic panel (Mann–Whitney U -test, p -value = 0.69), indicating no sex-biased dispersal. Furthermore, when males and females were considered separately in AMOVA analyses, no genetic differentiation was found between the sexes for the panel of polymorphic markers.

FIS values by permutations (sampling with replacement of individuals with all loci) were negative -0.081 (with CI narrow and negative, -0.129 and -0.044) and by Jackknife result on locus -0.079 (variance = 0.006). Negative FIS values indicate that there were more heterozygotes than expected and individuals in the population may be less related than expected under a model of random mating. Global frequency distribution of observed and permuted FST values were not significant, as they lie well inside the distribution of FST for the null hypothesis. FST were <0.05 between the three geographic areas with only the value between Dornoch and Inverness (0.014) being significant according to permutation

tests [Appendix S1D](#). Corrected average pairwise differences were low and not significant [Appendix S1D](#). These values indicate an absence of significant differentiation. Clustering analyses (STRUCTURE and DAPC) supported the existence of one main cluster with no spatial differentiation and an absence of structuring between sampling localities [Appendix S1E](#) and sex. Moreover, the isolation by distance (Mantel test) was clearly not significant (p -value = 0.12).

Concerning migration, FCA on individuals identified 4 axes with eigenvalues of 0.15 (*axis 1*), 0.089 (*axis 2*), 0.078 (*axis 3*), and 0.072 (*axis 4*). The three dimensional FCA [Appendix S1F](#) revealed two migrants as immigrants (one female from Cromarty and one male from Inverness), i.e., a rate of 2.15%. The mean Nm value using private alleles, after correction for size, and considering the individual geographic localisation was $Nm = 4.52$. According to [Wright \(1969\)](#), $Nm < 1$ indicates strong genetic differentiation, and Nm much larger than 1 means that panmixia can be assumed for the localities of the population. Based on [Hastings \(1993\)](#), [Faubet, Waples & Gaggiotti \(2007\)](#) suggest that $m < 0.1$ is needed to ensure demographic independence of populations. In this study, m were higher than 0.1 between sampling localisations in the Moray Firth.

Thus, the genetic differentiation and clustering analysis indicate that the harbour seal populations in the Moray Firth behave as a single genetic group. When considering the Moray Firth as a single genetic group, mean frequency of private alleles was equal to 0.03 which is consistent with the two immigrants detected by FCA. The Bayesian method of [Rannala & Mountain \(1997\)](#) did not detect immigrants into the Moray Firth. However, based on frequencies of [Paetkau et al. \(1995\)](#) and 10,000 resamplings, 5 of the 93 individuals were detected as immigrants (4 males and one female: 5%). Nei's standard distance (1972) with 10,000 resamplings with the algorithm of [Paetkau et al. \(2004\)](#) showed 4 immigrant individuals (2 females and 2 males), equivalent to around 4%. Hence, the global immigration rate in the Moray Firth is in the range of 2 to 4%.

Evolutionary trends

Most markers shared a similar range of mutation rates of around 0.0001–0.0002 with MSVAR estimates. VarEff provided global estimates of $\hat{\theta} = 4 \hat{N}u$, roughly corresponding to present ($\hat{\theta}_o = 0.77$), ancestral ($\hat{\theta}_a = 38.90$), and intermediate times ($\hat{\theta}_I = 8.56$) that help fix priors for effective size.

The global estimates in the present effective sizes of VarEff provided results in the same order of magnitude as those from MSVAR for seals particularly in terms of median but with lower standard deviations (Sd) ([Table 2](#)).

According to Gelman and Rubin's test ([Gelman & Rubin, 1992](#)), convergence in the MSVAR estimations of the effective size occurred after 20,000 iterations; and run 4 was the best supported [Appendix S1G](#). The results from MSVAR suggested that the effective size of the Moray Firth harbour seal population during our study period (1990–1995) was 821–1,669 (N_o) (sd 2,979), and that the ancestral effective size was much higher around 128,000–340,000 (N_a) (sd 846,836) ([Table 2](#)). Values for mutation rates were around 0.0001–0.0002 (μ) (sd 0.00026), and the ancestral coalescent time was estimated to occur 37,000–96,000 years ago (T_f) (sd 219,518) ([Table 2](#)).

Table 2 Mean (arithmetic), median and standard deviation of present (N_o) and ancestral (N_a) effective population size, the mutation rate (μ), and ancestral time in years (T_f) of Harbour seals in Moray Firth (left part, MSVAR analysis). Mean (arithmetic and harmonic), median and standard deviation (Sd) of present (N_o) and the estimated size 5,000 generations ago (N_{5000}), and the global present (\hat{N}_o) and ancestral estimate (\hat{N}_a) (right part, VarEff analysis).

	MSVAR				VarEff			
	N_o	N_a	μ	T_f	N_o	N_{5000}	\hat{N}_o	\hat{N}_a
Mean	1,669	342,874	0.00018	96,416	988 and 649	14,824		
Median	821	128,186	0.00010	37,235	714	11,298	1,287	64,826
Sd	2,979	846,836	0.00026	219,518	2,089	230,55		

VarEff harbour seal effective size priors were set between \hat{N}_o and \hat{N}_i from the global current ($\hat{\theta}_o$) and intermediate ($\hat{\theta}_i$) theta ($4\hat{N}u$) estimates: the prior mean was set to $\hat{N} = 5,000$ with a large variance (equal to 3 as suggested by [Nikolic & Chevalet \(2014a\)](#) for the logarithms of N). The estimate of present effective size (mean harmonic (N_o) and \hat{N}_o) was around 700 and 1,300 and the ancestral effective size (\hat{N}_a) 65,000 (Table 2). We also analysed females and males separately and obtained similar effective sizes which is consistent with no sex-biased dispersal. The arithmetic mean, harmonic mean, mode, and median of the posterior distributions for effective population sizes of harbour seals revealed a trend pattern that looks similar with a recent decrease from a higher effective size (Fig. 3A). A decrease in effective size of around 2.5% was suggested during the last 25 generations. The effective size at generation 0 (N_o) was 649 for the harmonic mean, 988 for the arithmetic mean, 714 for the median and 729 for the mode. This represents a 388–2,181 95% confidence interval. The difference observed between estimators of past N_e (mean, median and mode) (Fig. 3A) is likely to be due to the long tail of the posterior probability distribution ([Nikolic & Chevalet, 2014a](#)).

According to global estimates ($\hat{\theta}_o$, $\hat{\theta}_i$, $\hat{\theta}_a$) and effective size distribution (Fig. 3) from VarEff, harbour seals in the Moray Firth came from a large ancestral population. The estimated effective sizes in the past suggest that the extant population derived from an ancestral population of around 10,000–15,000 individuals approximately 5,000–30,000 generations ago (Fig. 3). This seems to be preceded by a lower ancient size (around 2,000–3,000 individuals) some 30,000–100,000 generations ago (Fig. 3C); however this should be considered with caution, given the flat posterior distribution in ancient times. Concerning the recent and historical effective size, several decreases of the effective size were observed. Firstly, a long period of decline over the last 2,000 generations (17,500 years) was suggested (Figs. 3A, 3B), when population size decreased from tens of thousands to less than 1,000 (Figs. 3A, 4A) and followed by a further decline prior to 600 generations ago (Fig. 4), estimated at around 800 generations ago (Fig. 5). Finally, a recent accelerated decline during the last 100 generations was followed by a period when population size remained approximately constant (N_e between ≈ 665 and 685) (Fig. 4).

The VarEff results describing the global evolution of the population are consistent with the tests of heterozygosity excess (sign, standardized and Wilcoxon tests) assuming SMM as proposed in BOTTLENECK Appendix S1H. The latter analysis suggested a




Figure 3 Effective size (N_e) of harbour seals in the north-east UK (from Moray Firth genetic group) as a function of past generation time (G) using 17 microsatellites (VarEff analysis). (A) Arithmetic (red) and harmonic (green) mean, mode (blue), and median (black) from sampling time (0) to 30,000 generations ago. (B) Posterior densities at the past generation time (G): 2,000 (black), 3,000 (blue), 4,000 (red), 5,000 (green), 6,000 (grey), 7,000 (purple), 8,000 (orange), 9,000 (pink), and 10,000 (red) generations ago. (C) Posterior densities at the past generation time (G) 10,000 (black), 20,000 (blue), 30,000 (red), 40,000 (blue), 50,000 (grey), 60,000 (purple), 70,000 (orange), 80,000 (green), 90,000 (brown), and 100,000 (red) generations ago.

Full-size DOI: 10.7717/peerj.9167/fig-3




Figure 4 Harbour seal's effective size (harmonic mean of the posterior distribution, VarEff analysis) within Moray Firth (using 17 microsatellites) in generation time (from 0 to 1,000 generations ago) (A) and in calendar years (from 1995 (sampling date) to 500 AD). The arrows represent the main trends: reduction (red) and increase (blue). (B) shows the latest tendency enlarged, the last red arrow.

Full-size DOI: 10.7717/peerj.9167/fig-4




Figure 5 Posterior distribution of the Time to Most Recent Common Ancestor allele (TMRCA, VarEff analysis). Suggested coalescent events are given as generation numbers and as years for harbour seals in the Moray Firth (Scotland). Each peak, in the posterior distribution, represents a potential bottleneck.

Full-size DOI: [10.7717/peerj.9167/fig-5](https://doi.org/10.7717/peerj.9167/fig-5)

recent bottleneck in the north-eastern UK population. Bayesian methods make use of coalescence theory and MCMC sampling to characterize demographic history and can have a higher probability of detecting bottlenecks than heterozygosity excess tests (contrasts heterozygosity expected under Hardy–Weinberg equilibrium with heterozygosity expected under mutation-drift equilibrium calculated from observed number of alleles) (*Cornuet & Luikart, 1997; Giroud et al., 2011; Peery et al., 2012*). Concerning the coalescent results from VarEff, Fig. 4A represents the harmonic mean of the last 1,000 generations (A), which was characterized by a decrease followed by a slight increase before a further decrease in the last 100 generations. Variation over the last 1,500 years is also shown in Fig. 4B, based upon the assumed generation time of 8.75 years.

According to Fig. 5, which shows the distributions of TMRCA, the harbour seal population underwent drastic bottlenecks. The more recent times of the bottlenecks were estimated in generation time (G) and in years, assuming a generation time of 8.75 years. Figure 5 also reveals that huge coalescence events occurred between 7,000–10,000 years (around the first peak in Fig. 5). These coalescence events occurred after a period of decline of the population size (the drastic bottleneck seen around 17,000 years ago, the time when the density is at a minimum between the two peaks). Before that period, coalescence events occurred at times given by the second peak of the distribution, which corresponds to the large ancestral population size, as illustrated in Figs. 3 and 4. Harbour seal showed declines over the last 1,000 years, with a steeper decline over the last few centuries. Furthermore, the analysis run without the two detected immigrants provided exactly the same results.

DISCUSSION

Genetic diversity

Results from the PI and SPOTG simulations suggest that the number of individuals and polymorphic markers are sufficient to provide high assignment discrimination and to detect evolutionary and ecological processes. The low genetic diversity in Moray Firth harbour seals, with an overall average heterozygosity of 38% and 3.7 alleles per microsatellite locus, was lower than in many pinnipeds (summarised in [Curtis, Stewart & Karl, 2011](#)). However, this value falls within the range recorded for other North Atlantic harbour seal populations, which were estimated between 24% ([Coltman, Bowen & Wright, 1996](#)), 38% ([Olsen et al., 2017](#)) and 50% ([Goodman, 1998](#)). In contrast, higher levels of heterozygosity (>65%) have been recorded in the North Pacific ([Burg et al., 1999](#)). There is concern that small populations with low genetic diversity may become susceptible to environmental, demographic, and genetic stochasticity, increasing their risks for extinction ([Franklin, 1980](#); [Newman & Pilson, 1997](#); [Brook et al., 2002](#)). However, low genetic diversity appears to be common amongst many marine mammals ([Hoelzel, Goldsworthy & Fleischer, 2002](#)), and the conservation implications of contemporary patterns are difficult to assess without some understanding of the relative role of natural and anthropogenic influences on historic population sizes. For example, low levels of genetic diversity may simply exist in some populations because of the nature of their social structure ([Whitehead, 1998](#)), or due to founder effects that occurred following natural post-glacial habitat change ([Palsbøll, Heide-Jørgensen & Dietz, 1997](#)). It is likely to be of greater concern to managers where low diversity has resulted from more recent over-exploitation ([Hoelzel et al., 1993](#)) or bycatch ([Pichler & Baker, 2000](#)).

Genetic structure

Our data suggest that the Moray Firth acts as one genetic group, in accordance with the results of [Olsen et al., 2017](#). Migration per generation above 5–10% can strongly influence the estimate of effective size ([Waples & England, 2011](#)) and it was therefore important to exclude this potential bias by evaluating gene flow. Current gene flow of harbour seals into the Moray Firth was low (immigration 2–4%), likely because the adults are strongly philopatric ([Thompson & Hall, 1993](#); [Härkönen & Harding, 2001](#)). [Goodman's \(1998\)](#) study of European harbour seal population differentiation suggests a critical range over which animals are philopatric of around 485 km. In that study, samples collected in the Moray Firth (Scottish east coast) were further than 500 km from neighbouring genetic populations (Scottish west coast, [Olsen et al., 2017](#)). In a Neighbor-joining phenogram presented in the [Goodman](#) study ([1998](#)), harbour seals from the Scottish east coast were distinguishable from the Scottish west coast with only 7 microsatellite loci, even though the number of seals from the Scottish west coast was very low (18 individuals). In a recent genetic clustering analysis with 12 microsatellite loci ([Olsen et al., 2017](#)), north-west and north-east UK were considered two geographically distinct genetic clusters. The genetic homogeneity between the Moray Firth and other north-eastern localities (Orkney and Shetland; [Olsen et al., 2017](#)), and associated low immigration, suggest that we can develop a robust reconstruction of evolutionary history for the north-eastern Scottish population.

Effective population size (N_e)

Inferring N_e from an estimate of θ can be obtained assuming a mutation rate. We estimated the mutation rate at around 0.0001–0.0002, similar to that found in ringed seals (*Phoca hispida*) (Palo *et al.*, 2003), allowing us to estimate effective population size. Using the complete panel of polymorphic markers with and without immigrants identified by the assignment method, the effective size estimated by the coalescent tool (VarEff) at generation 0 (N_0) was around 500–1,000 (CI [388–2181]). The advantage of VarEff is that it can estimate the recent historic abundance from theta ($\hat{\theta}_0$) (average of the effective size on the last evolutionary stage) and the current abundance from N_0 (effective size at generation time 0). N_e is a theoretical measure of an idealized population size that would be expected to experience the same rate of genetic diversity loss due to genetic drift (Wright, 1931). N_e is hence defined as the size of an ‘ideal’ population with the observed rate of genetic drift (Wright, 1931; Wright, 1969) and thus does not represent census population size (Palsbøll *et al.*, 2013). In ecology, N_e is usually thought of as the number of breeding individuals that successfully transmit their genes to the next generation (Frankham, 1995) and thus should equal the ‘genetic’ effective population size (Palsbøll *et al.*, 2013). In practice, the exact relationship is rarely known and most studies apply a generic ratio representing a range of estimates (Palsbøll *et al.*, 2013). Small N_e with no or limited gene flow among populations tend to accelerate stochastic loss of genetic diversity and can increase population risk as it leads to inbreeding depression and reduced fertility, and increases the potential fixation of deleterious alleles (Fagan & Holmes, 2006; Gilpin & Soulé, 1986; Palstra & Ruzzante, 2008; Lonsinger, Adams & Waits, 2018). Franklin (1980) suggested a minimum $N_e \geq 50$ may be required to avoid short-term inbreeding depression, and an $N_e \geq 500$ may be necessary to maintain long-term adaptive potential. However this rule (50/500) is open to criticism as the estimates do not take into account the force of selection. According to Lande (1995), wild populations could not bear the same consanguinity as farm populations (purged by humans); wild populations would fall more quickly into inbreeding depression and therefore the 50/500 rule would be underestimated. Lande (1995) recommended an N_e of 5,000 for long-term viability. Other analyses (Allendorf & Ryman, 2002) suggested an N_e of 1,000 to prevent accumulation of harmful mutations. All these recommended values must be taken with caution, as they ignore uncertainty arising from environmental and demographic factors, but they encourage conservation of small populations (Grove, 2003).

N_e is typically smaller than census population size (N) and this ratio may help for assessing the genetic health of a population and for predicting short-term and long-term risk (Palstra & Ruzzante, 2008). This ratio varies from 10^{-5} in many marine invertebrate species to nearly 1.0 in some terrestrial vertebrates (Frankham, 1995; Hedrick, 2005), but comparative data for other pinnipeds are difficult to interpret due to the high level of uncertainty in census population size (Curtis, Stewart & Karl, 2011). Surveys during the sampling period resulted in a direct estimate (N , census population size) of approximately 1,650 individuals (1993) for Moray Firth locality (Thompson *et al.*, 1997a), 773 individuals in 1992 and 575 in 1994 for Firth of Tay and Eden Estuary (Hanson *et al.*, 2015), 9,000 and 6,000 individuals in 1997 for north coast and Orkney, then Shetland respectively (Fig. 4 from Duck & Morris, 2014). These different values suggest that the ratio N_0/N can be

very different depending on sampling location considered. The N_0 estimate in this study represents one genetic cluster, the north-east UK, and including all N would be unwise until we verify that N_0 from other localities are in the same general estimate. Although N is a relatively straightforward entity, it can be difficult to obtain (Palsbøll *et al.*, 2013). Hence, in order to infer N from N_e , a rigorous estimate of the ratio is warranted (Palsbøll *et al.*, 2013).

Evolutionary trends

Over 80% of UK harbour seals are found in Scotland and estimating the historic effective size of harbour seal in the Moray Firth showed that seals have recently experienced a decline, as also revealed by direct counts in parts of both the north-east Atlantic (Thompson *et al.*, 2019; Lonergan *et al.*, 2007; Hanson *et al.*, 2015) and the north-west Atlantic (Bowen *et al.*, 2003). Concerning the ancestral harbour seal population size (MRCA) in the Moray Firth, both models suggested that effective population size had once been extremely large; in the region of 65,000–130,000 derived from ancestral theta for VarEff and the median for MSVAR. One advantage of VarEff is the ability to infer demographic history from a single temporal sample, rather than depending on two or more samples that span multiple generations. Detecting past bottlenecks using the method relies on observing pairs of alleles that have coalesced during the bottleneck event (Nikolic & Chevalet, 2014a). The VarEff method enables various mutation models to be considered, and some trials determined that the risk of false bottleneck detection due to an inappropriate mutation model would be unlikely. Results also indicated that the Moray Firth harbour seal population has undergone a drastic bottleneck, which is concordant with the analysis of detection by the heterozygote deficiency tests. Figures 3A and 3B show that a recent strong bottleneck occurred approximately 2,000 generations ago (17,500 years), when population size decreased from more than 10,000 to less than 1,000 in about 1,000 generations. The model suggested an ancestral population size of about 15,000 in ancient times, 30,000 generations or 262,000 years ago (the ultimate peak in the TMRCA distribution, Fig. 5). However, other evolutionary schemes could lead to a similar distribution of TMRCA, such as recurrent immigration from a large population into the Moray Firth. Thus, there are several possible interpretations for this pattern. The population may derive from ancestral fragmentation followed by permanent introductions from a metapopulation made of similar colonies, or it may be the result of a sharp decline some time in the past. Several arguments support the bottleneck hypothesis: the gaps in the distribution of repeat numbers Appendix S1C, the rather low immigration rate, and the high probability that coalescent events occurred recently. A more detailed answer will require archaeological samples (ancient DNA) or samples from a number of distant populations.

During the period corresponding to the drastic bottleneck detected, the current geographical range of this population would have been uninhabitable as it was covered by the last British Ice Sheet (Bradwell *et al.*, 2008), and this ancestral population would likely have re-distributed to the margins of the ice sheet. Whilst the size of large terrestrial mammal populations in Europe is likely to have been reduced during this period (Marshall *et al.*, 1982), harbour seals breed on ice in certain areas, and these colder conditions could

have led to population expansion. For example, analysis of mtDNA variation in Antarctic ice-breeding seals indicated that effective population size expanded suddenly during times of intensified glaciations (Curtis, Stewart & Karl, 2009), while southern elephant seals, which require access to open beaches, show the opposite trend (De Bruyn et al., 2009). The strong bottleneck detected at around 17,500 years ago appears to have occurred in the period following the retreat of ice from the current study area, as glacial moraines in the Inverness Firth indicate that the ice margin occurred here around 15,000 years ago (Merritt, Auton & Firth, 1995). Subsequently, the harbour seals may have recolonized this area once the ice retreated, as was the case for ringed seals which colonized the Baltic Sea basin soon after deglaciation (11,500 years) (Palo et al., 2001).

The analysis of TMRCA revealed a peak of coalescence events around 800 generations (7,000 years) ago in a period when mean effective size was approximately 5,000, and may correspond to a transient sharp decrease of effective size. This coincides with the warmest stages in the post-glacial period in the mid-Holocene 8,000-7,000 years ago (Andersson, 1902; Andersson, 1909; Seppä, Bjune AE & Veski, 2009). Current global distribution patterns indicate that harbour seals could have been restricted to more northerly waters at this time. Archaeological records also suggest that populations of harbour seals in the eastern Baltic were founded around 8,000 years ago (Härkönen & Johannesson, 2005), which may also reflect a northward shift in distribution during this period. Here, we see that this period may correspond to a transient decrease of the harbour seal population, long after it was derived from a larger ancestral population. Several dramatic events therefore appear to have influenced the structure and effective size of this population. This complexity inevitably brings uncertainty, particularly as the model cannot fully account for migration events.

Assuming immigration from a large external population, the distribution of coalescence time TMRCA depends on the proportion m of immigrants per generation and on the ratio ε of the considered population size (N_e) to the size of the larger external population (Nikolic & Chevalet, 2014a). Using VarEff provides the distribution of TMRCA which is converted to past effective sizes in an isolated population. In the case of small $4 * N_e * m$, such as found from the harbour seal analysis, the estimated population size remains in the order of magnitude of the true N_e while increasing with m (Nikolic & Chevalet, 2014a). However, for larger immigration rates from a larger external population, a false current bottleneck could be predicted in a small population. Applying the equation (8) of Nikolic & Chevalet (2014a), with hypothetical immigration (2–4%) from larger populations of harbour seals around the UK (Lonergan et al., 2007, which brings ε about 0.05 to 0.10) and specifically around our North-eastern Scottish population, we estimated the time of a potentially false bottleneck at 100–200 generations ago. This period matches with the last event detected in Fig. 4. Thus, immigration from an external population could potentially lead to a false current bottleneck (100–200 generations ago), and we have to be cautious in the final decline in effective size observed (Fig. 4). We encourage studies on seal migration in this area by the monitoring of individuals between the geographically distinct genetic clusters in UK and neighbouring waters.

Following the same analysis, the older bottlenecks detected at 800 and 2,000 generations ago (Figs. 3 and 5) should not be an effect of immigration. Either way, it is clear that

these past events led to a decrease in population size resulting from natural environmental changes. This study suggests that, more recently, the harbour seal population has declined over the last century at a rate of 2–6% in terms of effective size. This downward trend could also result from natural changes in environmental conditions, or from human exploitation that is known to have occurred at many European and Arctic sites over the last millennia (*Härkönen & Johannesson, 2005; Murray, 2008*).

Conservation implications

Previous population genetic studies of UK harbour seals have focused on using molecular tools to support conservation management by identifying appropriate spatial management units (*Olsen et al., 2017*). Our study illustrates how genetic tools based on coalescent theory can also provide insights into evolutionary history and temporal changes in population size within those management units. Critically, they contribute knowledge about the state of a population by estimating changes in effective size. Estimates of the current effective population size of this harbour seal population are small compared to theoretical estimates of the minimum viable population size (*Reed et al., 2003*). Our analyses indicate that the harbour seal population in the north-east UK has remained at a broadly similar level following the bottleneck that occurred after post-glacial recolonization of the area but have slowly declined more recently. Subsequent demographic studies indicate that the Moray Firth population has declined further, by approximately 40%, since the early 1990's when our DNA samples were collected (*Thompson et al., 2019*). Thus, whilst contemporary estimates of N_e are expected to remain well above the critical threshold of $N_e = 50$ suggested to avoid short-term effects of inbreeding (*Franklin, 1980*) they will now be closer to the $N_e = 500$ potential threshold for maintenance of long-term adaptive potential (*Franklin, 1980*). Molecular evidence of connectivity within the much larger European metapopulation exists (*Olsen et al., 2017*), but further work is required to better understand the extent to which contemporary movements between management units may offset these risks. Contrasting population trends in different UK management units (*Thompson et al., 2019*) also highlight the need for conservation managers to identify regional demographic drivers. Crucially, potential drivers include both anthropogenic stressors (which might require conservation interventions) and others such as competition with recovering grey seal populations (*Halichoerus grypus*) (*Wilson & Hammond, 2019*); where intervention is less likely to be appropriate. Previous studies in the Baltic Sea indicate that populations of harbour and grey seal have co-varied over historical time-scales (*Härkönen & Johannesson, 2005*). Our work illustrates how parallel coalescent studies on material from sympatric grey and harbour seal populations in different regions now provides an opportunity to inform conservation practices by exploring potential interactions between these two species at much larger temporal and spatial scales.

ACKNOWLEDGEMENTS

The authors wish to thank Shaneve Tripp (NYU School of Law) and Wendy West (DAFF) for their english corrections. Ludovic Hoarau (IFREMER) for his help on ArcGis. Katia Feve (INRAE) for her help with the DNA extraction protocol. DNA samples were extracted

at INRAE and genotyped at the Toulouse Genopole Platform (<http://www.genotoul.fr/>). Anonymous reviewers provided many helpful comments on an earlier version of the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by INRAE (FRANCE), Genotoul platform (FRANCE), and University of Aberdeen. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

INRAE (FRANCE).

Genotoul platform (FRANCE),.

University of Aberdeen.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Natacha Nikolic conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Paul Thompson performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Mark de Bruyn authored or reviewed drafts of the paper, and approved the final draft.
- Matthias Macé and Claude Chevalet performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

Blood samples were collected under Home Office licence issued to the University of Aberdeen under the Animal (Scientific Procedures) Act 1986 (PPL number 60/01351).

Data Availability

The following information was supplied regarding data availability:

Data is available at INRAE:

Nikolic, Natacha; Thompson, Paul; De Bruyn, Mark; Macé, Matthias; Chevalet, Claude, 2020, "Microsatellite data from: Evolutionary history of a Scottish harbour seal population", <https://doi.org/10.15454/AOZ7JI>, Portail Data INRAE, V2.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.9167#supplemental-information>.

REFERENCES

- Aarts G, Brasseur S, Poos JJ, Schop J, Kirkwood R, van. Kooten T, Mul E, Reijnders P, Rijnsdorp AD, Tulp I. 2019. Top-down pressure on a coastal ecosystem by harbor seals. *Ecosphere* 10:e02538.
- Allen PJ, Amos W, Pomedoy PP, Twiss SD. 1995. Microsatellite variation in grey seals (*Halichoerus grypus*) shows evidence of genetic differentiation between two British breeding colonies. *Molecular Ecology* 4:653–662
DOI 10.1111/j.1365-294X.1995.tb00266.x.
- Allendorf F, Ryman N. 2002. The role of genetics in population viability analysis. In: Beissinger SR, McCullough DR, eds. *Population viability analysis*. Chicago: University of Chicago Press, 50–85.
- Andersen LW, Lydersen C C, Frie AK, Rosing-Asvid A. 2011. A population on the edge: genetic diversity and population structure of the world's northernmost harbour seals (*Phoca vitulina*). *Biological Journal of the Linnean Society* 102:420–439
DOI 10.1111/j.1095-8312.2010.01577.x.
- Andersson G. 1902. Hasseln i Sverige fordom och nu. *Sveriges Geologiska Undersö kning Serie Ca* 3:1–168.
- Andersson G. 1909. The climate of Sweden in the Late-Quaternary period. *Sveriges geologiska undersö kning Serie Ca* 3:1–88.
- Bailleul D, Mackenzie A, Sacchi O, Poisson F, Bierne N, Arnaud-Haond S. 2018. Large-scale genetic panmixia in the blue shark (*Prionace glauca*): a single worldwide population, or a genetic lag-time effect of the grey zone of differentiation? *Evolutionary Applications* 11(5):614–630 DOI 10.1111/eva.12591.
- Baxter JM. 2001. Establishing management schemes on marine special areas of conservation in Scotland. *Aquatic Conservation: Marine and Freshwater Ecosystems* 11:261–265 DOI 10.1002/aqc.465.
- Beaumont MA. 1999. Detecting population expansion and decline using microsatellites. *Genetics* 153:2013–2029.
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F. 1998. *GENETIX, Logiciel sous Windows TM pour la Génétique des Populations*. Montpellier: Laboratoire Génome et Populations, CNRS UPR 9060, Université de Montpellier II.
- Berthier P, Beaumont MA, Cornuet J-M, Luikart G. 2002. Likelihood-based estimate of the effective population size using temporal changes in allele frequency: a genealogical approach. *Genetics* 160:741–751.
- Boveng PL, Bengtson JL, Withrow DE, Cesarone JC. 2003. The abundance of harbor seals in the Gulf of Alaska. *Marine Mammal Science* 19:111–127
DOI 10.1111/j.1748-7692.2003.tb01096.x.

- Bowen WD, Ellis SL, Iverson SJ, Boness DJ. 2003.** Maternal and newborn life-history traits during periods of contrasting population trends, implications for explaining the decline of harbour seals (*Phoca vitulina*), on Sable Island. *Journal of Zoology* 261:155–163 DOI [10.1017/S0952836903004047](https://doi.org/10.1017/S0952836903004047).
- Bradwell T, Fabel D, Stoker M, Mathers H, McHargue L, Howe J. 2008.** Ice caps existed throughout the Lateglacial Interstadial in northern Scotland. *Journal of Quaternary Science* 23:401–407 DOI [10.1002/jqs.1181](https://doi.org/10.1002/jqs.1181).
- Brook BW, Tonkyn D, O’Grady JJ, Frankham R. 2002.** Contribution of inbreeding to extinction risk in threatened species. *Conservation Ecology* 6(1):16.
- Buchanan FC, Maiers LD, Thue TD, De March BG, Stewart RE. 1998.** Microsatellites from the Atlantic walrus *Odobenus rosmarus rosmarus*. *Molecular Ecology* 7(8):1083–1085 DOI [10.1046/j.1365-294X.1998.00401.x](https://doi.org/10.1046/j.1365-294X.1998.00401.x).
- Burg TM, Trites AW, Smith JM. 1999.** Mitochondrial and microsatellite DNA analyses of harbour seal population structure in the Northeast Pacific Ocean. *Canadian Journal of Zoology* 77(6):930–943.
- Busch JD, Waser PM, DeWoody JA. 2007.** Recent demographic bottlenecks are not accompanied by a genetic signature in banner-tailed kangaroo rats (*Dipodomys spectabilis*). *Molecular Ecology* 16:2450–2462 DOI [10.1111/j.1365-294X.2007.03283.x](https://doi.org/10.1111/j.1365-294X.2007.03283.x).
- Chevalet C, Nikolic N. 2010.** Distribution of coalescent times and distances between microsatellite alleles with changing effective population size. *Theoretical Population Biology* 77(3):152–163 DOI [10.1016/j.tpb.2010.01.001](https://doi.org/10.1016/j.tpb.2010.01.001).
- Chybicki IJ, Burczyk J. 2009.** Simultaneous estimation of null alleles and inbreeding coefficients. *Journal of Heredity* 100:106–113 DOI [10.1093/jhered/esn088](https://doi.org/10.1093/jhered/esn088).
- Coltman DW, Bowen WD, Wright JM. 1996.** PCR primers for harbour seal (*Phoca vitulina concolour*) microsatellites amplify polymorphic loci in several pinniped species. *Molecular Ecology* 5:161–163 DOI [10.1111/j.1365-294X.1996.tb00303.x](https://doi.org/10.1111/j.1365-294X.1996.tb00303.x).
- Cordes LS, Duck CD, Mackey BL, Hall AJ, Thompson PM. 2011.** Long-term patterns in harbour seal site-use and the consequences for managing protected areas. *Animal Conservation* 14:430–438 DOI [10.1111/j.1469-1795.2011.00445.x](https://doi.org/10.1111/j.1469-1795.2011.00445.x).
- Cordes LS, Thompson PM. 2014.** Mark-recapture modeling accounting for state uncertainty provides concurrent estimates of survival and fecundity in a protected harbor seal population. *Marine Mammal Science* 30:691–705 DOI [10.1111/mms.12070](https://doi.org/10.1111/mms.12070).
- Cornuet J-M, Luikart G. 1997.** Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* 144:2001–2014.
- Curtis C, Stewart BS, Karl SA. 2009.** Pleistocene population expansions of Antarctic seals. *Molecular Ecology* 18(10):2112–2121 DOI [10.1111/j.1365-294X.2009.04166.x](https://doi.org/10.1111/j.1365-294X.2009.04166.x).
- Curtis C, Stewart BS, Karl SA. 2011.** Genetically effective population sizes of Antarctic seals estimated from nuclear genes. *Conservation Genetics* 12:1435–1446 DOI [10.1007/s10592-011-0241-x](https://doi.org/10.1007/s10592-011-0241-x).
- Davis CS, Gelatt TS, Siniff D, Strobeck C. 2002.** Dinucleotide microsatellite markers from the Antarctic seals and their use in other pinnipeds. *Molecular Ecology Resources* 2:203–208.

- De Bruyn M Hall, BL, Chauke LF, Baroni C Koch, PL, Hoelzel RA. 2009.** Rapid response of a marine mammal species to Holocene climate and habitat change. *PLOS Genetics* 5:e1000554 DOI [10.1371/journal.pgen.1000554](https://doi.org/10.1371/journal.pgen.1000554).
- Dishman DL. 2011.** Genetic substructure of Pacific harbor seal (*Phoca vitulina richardsi*) populations along the coasts of Oregon and Washington. In: *Dissertations and Theses (Open Access)*. Paper 312.
- Duck CD, Morris CD. 2014.** Surveys of harbour (common) and grey seals on the east, north and north-west coast of Scotland and in Orkney, including the Moray Firth and the Firth of Tay, in 2013. Research report (759). Scottish Natural Heritage..
- Earl Dent A, VonHoldt Bridgett M. 2012.** Structure harvester: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* 4(2):359–361 DOI [10.1007/s12686-011-9548-7](https://doi.org/10.1007/s12686-011-9548-7).
- Evanno G, Goudet J, Regnaut Sand. 2005.** Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Molecular Ecology* 14:2611–2620 DOI [10.1111/j.1365-294X.2005.02553.x](https://doi.org/10.1111/j.1365-294X.2005.02553.x).
- Excoffier L, Laval G, Schneider S. 2005.** Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1:47–50.
- Excoffier L, Lischer HEL. 2010.** Arlequin suite ver 3.5: A new series of programs to perform population genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10:564–567 DOI [10.1111/j.1755-0998.2010.02847.x](https://doi.org/10.1111/j.1755-0998.2010.02847.x).
- Fagan WF, Holmes EE. 2006.** Quantifying the extinction vortex. *Ecology Letters* 9(1):51–60.
- Faubet P, Waples R, Gaggiotti OE. 2007.** Evaluating the performance of a multilocus Bayesian method for the estimation of migration rates. *Molecular Ecology* 16:1149–1166 DOI [10.1111/j.1365-294X.2007.03218.x](https://doi.org/10.1111/j.1365-294X.2007.03218.x).
- Frankham R. 1995.** Effective population size/adult population size ratios in wildlife: a review. *Genetical Research* 66:95–106 DOI [10.1017/S0016672300034455](https://doi.org/10.1017/S0016672300034455).
- Franklin I. 1980.** Evolutionary changes in small populations. In: Soule M M, Wilcox B, eds. *Conservation biology: an evolutionary-ecological perspective*. Sunderland: Sinauer Associates, 135–150.
- Gelatt TS, Davis CS, Siniff DB, Strobeck C. 2001.** Molecular evidence for twinning in Weddell seals (*Leptonychotes weddellii*). *Journal of Mammalogy* 82(2):491–499 DOI [10.1644/1545-1542\(2001\)082<0491:MEFTIW>2.0.CO;2](https://doi.org/10.1644/1545-1542(2001)082<0491:MEFTIW>2.0.CO;2).
- Gelman A, Rubin DB. 1992.** Inference from iterative simulation using multiple sequences. *Statistical Science* 7:457–511 DOI [10.1214/ss/1177011136](https://doi.org/10.1214/ss/1177011136).
- Gemmell NJ, Allen PJ, Goodman SJ, Reed JZ. 1997.** Interspecific microsatellite markers for the study of pinniped populations. *Molecular Ecology* 6:661–666 DOI [10.1046/j.1365-294X.1997.00235.x](https://doi.org/10.1046/j.1365-294X.1997.00235.x).
- Gilpin ME, Soulé ME. 1986.** Minimum viable populations: processes of species extinction. In: Soulé ME, ed. *Conservation biology: the science of scarcity and diversity*. Sunderland: Sinauer Associates, 19–34.

- Girod C, Vitalis R, Lebois R, Freville H. 2011.** Inferring population decline and expansion from microsatellite data: a simulation-based evaluation of the MSvar methods. *Genetics* **188**:165–179 DOI [10.1534/genetics.110.121764](https://doi.org/10.1534/genetics.110.121764).
- Goodman SJ. 1997.** Dinucleotide repeat polymorphisms at seven anonymous microsatellite loci cloned from the European harbour seal (*Phoca vitulina vitulina*). *Animal Genetics* **28**(4):310–311.
- Goodman SJ. 1998.** Patterns of extensive genetic differentiation and variation among European Harbour seals (*Phoca vitulina vitulina*) revealed using microsatellite DNA polymorphisms. *Molecular Biology and Evolution* **15**:104–118 DOI [10.1093/oxfordjournals.molbev.a025907](https://doi.org/10.1093/oxfordjournals.molbev.a025907).
- Grove C. 2003.** *Drafting a conservation blueprint: a practitioner's guide to planning for biodiversity*. Arlington: Nature Conservancy, 188.
- Hall AJ, Mackey B, Kershaw JL, Thompson P. 2019.** Age–length relationships in UK harbour seals during a period of population decline. *Aquatic Conservation: Marine and Freshwater Ecosystems* **29**:61–70 DOI [10.1002/aqc.3104](https://doi.org/10.1002/aqc.3104).
- Han JB, Sun FY, Gao X, He C-B, Wang P-L, Ma Z-Q, Wang Z-H. 2010.** Low microsatellite variation in spotted seal (*Phoca largha*) shows a decrease in population size in the Liaodong Gulf colony. *Annales Zoologici Fennici* **47**:15–27 DOI [10.5735/086.047.0102](https://doi.org/10.5735/086.047.0102).
- Hanson N, Thompson D, Duck C, Baxter J, Lonergan M. 2015.** Harbour seal (*Phoca vitulina*) abundance within the Firth of Tay and Eden Estuary Scotland: recent trends and extrapolation to extinction. *Aquatic Conservation: Marine and Freshwater Ecosystems* **27**:268–281 DOI [10.1002/aqc.2609](https://doi.org/10.1002/aqc.2609).
- Härkönen T, Dietz R, Reijnders P, Teilmann J. 2006.** A review of the 1988 and 2002 phocine distemper virus seal epidemics in European harbour seals. *Diseases of Aquatic Organisms* **68**:115–130 DOI [10.3354/dao068115](https://doi.org/10.3354/dao068115).
- Härkönen T, Harding KC. 2001.** Spatial structure of harbour seal populations and the implications thereof. *Canadian Journal of Zoology* **79**:2115–2127 DOI [10.1139/z01-172](https://doi.org/10.1139/z01-172).
- Härkönen T, Harding KC, Goodman SJ, Johannesson K. 2005.** Colonization history of the Baltic harbour seals: integrating archaeological behavioral and genetic data. *Marine Mammal Science* **21**(4):695–716 DOI [10.1111/j.1748-7692.2005.tb01260.x](https://doi.org/10.1111/j.1748-7692.2005.tb01260.x).
- Hastings A. 1993.** Complex interactions between dispersal and dynamics—lessons from coupled logistic equations. *Ecology* **74**:1362–1372 DOI [10.2307/1940066](https://doi.org/10.2307/1940066).
- Hayes SA, Pearse DE, Costa DP, Harvey JT, Le Boeuf BJ, Garza JC. 2006.** Mating system and reproductive success in eastern Pacific harbour seals. *Molecular Ecology* **15**:3023–3034 DOI [10.1111/j.1365-294X.2006.02984.x](https://doi.org/10.1111/j.1365-294X.2006.02984.x).
- Hedrick P. 2005.** Large variance in reproductive success and the Ne/N ratio. *Evolution* **59**(7):1596–1599 DOI [10.1111/j.0014-3820.2005.tb01809.x](https://doi.org/10.1111/j.0014-3820.2005.tb01809.x).
- Heide-Jørgensen MP, Härkönen T, Aberg P. 1992.** Long term effects of epizootic in harbour seals in the Kattegat–Skagerrak and adjacent areas. *Ambio* **21**:511–516.

- Herreman JK, Blundell GM, Ben-David M. 2008.** False sex-linked microsatellite primer for *Phoca vitulina*. *Marine Mammal Science* **24**:411–413
[DOI 10.1111/j.1748-7692.2008.00186.x](https://doi.org/10.1111/j.1748-7692.2008.00186.x).
- Herreman JK, Blundell GM, McDonald DB, Ben-David M. 2009.** Asymmetrical male-mediated gene flow between Harbor seal (*Phoca vitulina*) populations in Alaska. *Canadian Journal of Zoology* **87**:498–507 [DOI 10.1139/Z09-033](https://doi.org/10.1139/Z09-033).
- Hoban S, Gaggiotti O, Bertorelle G. 2013.** Sample Planning Optimization Tool for conservation and population Genetics (SPOTG): a software for choosing the appropriate number of markers and samples. *Methods in Ecology and Evolution* **4**:299–303 [DOI 10.1111/2041-210x.12025](https://doi.org/10.1111/2041-210x.12025).
- Hoelzel AR, Goldsworthy SD, Fleischer RC. 2002.** Population genetics. In: Hoelzel AR, ed. *Marine mammal biology; an evolutionary approach*. Oxford: Blackwell Science, 325–352.
- Hoelzel AR, Halley JM, O'Brien SJ, Campagna C. 1993.** Elephant seal genetic variation and the use of simulation models to investigate historical population bottlenecks. *Journal of Heredity* **84**:443–449 [DOI 10.1093/oxfordjournals.jhered.a111370](https://doi.org/10.1093/oxfordjournals.jhered.a111370).
- Jamieson A. 1994.** The effectiveness of using co-dominant polymorphic allelic series for (1) checking pedigrees and (2) distinguishing full-sib pair members. *Animal Genetics* **25**(Suppl. 1):37–44 [DOI 10.1111/j.1365-2052.1994.tb00401.x](https://doi.org/10.1111/j.1365-2052.1994.tb00401.x).
- Jamieson A, Taylor SCS. 1997.** Comparisons of three probability formulae for parentage exclusion. *Animal Genetics* **28**:397–400 [DOI 10.1111/j.1365-2052.1997.00186.x](https://doi.org/10.1111/j.1365-2052.1997.00186.x).
- Jeffries S, Huber H, Calambokidis J, Laake J. 2003.** Trends and status of harbor seals in Washington State: 1978–1999. *The Journal of Wildlife Management* 207–218.
- Johnson PCD, Haydon DT. 2007.** Maximum likelihood estimation of allelic dropout and false allele error rates from microsatellite genotypes in the absence of reference data. *Genetics* **175**:827–42 [DOI 10.1534/genetics.106.064618](https://doi.org/10.1534/genetics.106.064618).
- Jombart T. 2008.** adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics* **24**:1403–1405 [DOI 10.1093/bioinformatics/btn129](https://doi.org/10.1093/bioinformatics/btn129).
- Jombart T, Ahmed I. 2011.** adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. *Bioinformatics* **27**(21):3070–3071.
- Jombart T, Devillard S, Balloux F. 2010.** Discriminant analysis of principal components: a new method for the analysis of genetically structured populations. *BMC Genetics* **11**:94.
- Kalinowski ST, Taper ML, Marshall TC. 2007.** Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Molecular Ecology* **16**:1099–1106 [DOI 10.1111/j.1365-294X.2007.03089.x](https://doi.org/10.1111/j.1365-294X.2007.03089.x).
- King TL, Kalinowski ST, Schill WB, Spidle AP, Lubinski BA. 2001.** Population structure of Atlantic salmon (*Salmo salar* L.): a range-wide perspective from microsatellite DNA variation. *Molecular Ecology* **10**:807–821 [DOI 10.1046/j.1365-294X.2001.01231.x](https://doi.org/10.1046/j.1365-294X.2001.01231.x).
- Kopelman NM, Mayzel J, Jakobsson M, Rosenberg NA, Mayrose I. 2015.** Clumpak: a program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources* **15**:1179–1191
[DOI 10.1111/1755-0998.12387](https://doi.org/10.1111/1755-0998.12387).

- Lande R. 1995.** Mutation and conservation. *Conservation Biology* **9**:782–791
[DOI 10.1046/j.1523-1739.1995.09040782.x](https://doi.org/10.1046/j.1523-1739.1995.09040782.x).
- Lonergan M, Duck Duck C, Thompson D, Mackey B. 2007.** Using sparse survey data to investigate the declining abundance of British harbour seals. *Journal Zoology* **271**:261–269 [DOI 10.1111/j.1469-7998.2007.00311.x](https://doi.org/10.1111/j.1469-7998.2007.00311.x).
- Lonsinger RC, Adams JR, Waits L. 2018.** Evaluating effective population size and genetic diversity of a declining kit fox population using contemporary and historical specimens. *Ecology and Evolution* **8**(23):12011–12021 [DOI 10.1002/ece3.4660](https://doi.org/10.1002/ece3.4660).
- Lotze HK, Worm B. 2009.** Historical baselines for large marine animals. *Trends in Ecology and Evolution* **24**:254–262 [DOI 10.1016/j.tree.2008.12.004](https://doi.org/10.1016/j.tree.2008.12.004).
- Luikart G, Allendorf FW, Cornuet J-M, Sherwin WB. 1997.** Distortion of allele frequency distributions provides a test for recent population bottlenecks. *Journal of Heredity* **89**(3):238–247.
- Marshall LG, Webb SD, Sepkoski Jr JJ, Raup DM. 1982.** Mammalian evolution and the great american interchange. *Science* **215**(4538):1351–1357
[DOI 10.1126/science.215.4538.1351](https://doi.org/10.1126/science.215.4538.1351).
- Matthiopoulos J, Cordes L, Mackey B, Thompson D, Duck C, Smout S, Caillat M, Thompson P. 2014.** State-space modelling reveals proximate causes of harbour seal population declines. *Oecologia* **174**:151–162 [DOI 10.1007/s00442-013-2764-y](https://doi.org/10.1007/s00442-013-2764-y).
- McClenachan L, Ferretti F, Baum JK. 2012.** From archives to conservation: why historical data are needed to set baselines for marine animals and ecosystems. *Conservation Letters* **5**:349–359 [DOI 10.1111/j.1755-263X.2012.00253.x](https://doi.org/10.1111/j.1755-263X.2012.00253.x).
- Merritt JW, Auton CA, Firth CR. 1995.** Ice-proximal glaciomarine sedimentation and sea-level change in the Inverness Area, Scotland: a review of the deglaciation of a major ice stream of the British Late Devensian Ice Sheet. *Quaternary Science Reviews* **14**:289–329 [DOI 10.1016/0277-3791\(95\)00008-D](https://doi.org/10.1016/0277-3791(95)00008-D).
- Mossman CA, Waser PM. 1999.** Genetic detection of sex-biased dispersal. *Molecular Ecology* **8**:1063–1067 [DOI 10.1046/j.1365-294x.1999.00652.x](https://doi.org/10.1046/j.1365-294x.1999.00652.x).
- Murray MS. 2008.** Zooarchaeology and arctic marine mammal biogeography, conservation, and management. *Ecological Applications* **18**(2):41–55 [DOI 10.1890/06-0818.1](https://doi.org/10.1890/06-0818.1).
- Nei M. 1978.** Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* **89**:583–590.
- Newman D, Pilson D. 1997.** Increased probability of extinction due to decreased genetic effective population size: experimental populations of *Clarkia pulchella*. *Evolution, the International Journal of Organic Evolution* **51**:354–362
[DOI 10.1111/j.1558-5646.1997.tb02422.x](https://doi.org/10.1111/j.1558-5646.1997.tb02422.x).
- Nikolic N, Butler JR, Baglinière JL, Laughton R, McMyn IA, Chevalet C. 2009.** An examination of genetic diversity and effective population size in Atlantic salmon populations. *Genetical Research* **91**(6):395–412 [DOI 10.1017/S0016672309990346](https://doi.org/10.1017/S0016672309990346).
- Nikolic N, Chevalet C. 2014a.** Detecting past changes of effective population size. *Evolutionary Applications* **7**:663–681 [DOI 10.1111/eva.12170](https://doi.org/10.1111/eva.12170).

- Nikolic N, Chevalet C. 2014b.** VarEff. Variation of Effective size. Software VAREFF (package R in file.zip) and the documentation. Available at <https://archimer.ifremer.fr/doc/00177/28781/>.
- Olsen MT, Islas V, Graves JA, Onoufriou A, Vincent C, Brasseur S, Frie AK, Hall AJ.** Microsatellite analysis of population structure in Canadian polar bears. *Aquatic Conservation: Marine and Freshwater Ecosystems* 27:839–845 DOI 10.1002/aqc.2760.
- Oosterhout CV, Hutchinson WF, Wills DPM, Shipley P. 2004.** Micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes* 4:535–538 DOI 10.1111/j.1471-8286.2004.00684.x.
- Paetkau D, Calvert W, Stirling I, Strobeck C. 1995.** Microsatellite analysis of population structure in Canadian polar bears. *Molecular Ecology* 4:347–354 DOI 10.1111/j.1365-294X.1995.tb00227.x.
- Paetkau D, Slade R, Burden M, Estoup A. 2004.** Direct real-time estimation of migration rate using assignment methods: a simulation-based exploration of accuracy and power. *Molecular Ecology* 13:55–65 DOI 10.1046/j.1365-294X.2004.02008.x.
- Palo JU, Mäkinen HS, Helles E, Väinölä R, Stenman O. 2001.** Microsatellite variation in ringed seals (*Phoca hispida*): genetic structure and history of the Baltic sea population. *Heredity* 86:609–617 DOI 10.1046/j.1365-2540.2001.00859.x.
- Palo JU, Hyvärinen H, Helle E, Mäkinen HS, Väinölä R. 2003.** Postglacial loss of microsatellite variation in the landlocked Lake Saimaa ringed seal. *Conservation Genetics* 4:117–128 DOI 10.1023/A:1023303109701.
- Palsbøll PJ, Heide-Jørgensen MP, Dietz R. 1997.** Population structure and seasonal movements of narwhals. *Monodon monoceros*, determined from mtDNA analysis. *Heredity* 78:284–292 DOI 10.1038/hdy.1997.43.
- Palsbøll PJ, Peery MZ, Olsen MT, Beissinger SR, Bérubé M. 2013.** Inferring recent historic abundance from current genetic diversity. *Molecular Ecology* 22:22–40 DOI 10.1111/mec.12094.
- Palstra FP, Ruzzante DE. 2008.** Genetic estimates of contemporary effective population size: what can they tell us about the importance of genetic stochasticity for wild population persistence? *Molecular Ecology* 17(15):3428–3447 DOI 10.1111/j.1365-294X.2008.03842.x.
- Pastor T, Garza JC, Allen P, Amos W, Aguilar A. 2004.** A low genetic variability in the highly endangered Mediterranean monk seal. *Journal of Heredity* 95:291–300 DOI 10.1093/jhered/esh055.
- Peakall R, Smouse PE. 2006.** GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295 DOI 10.1111/j.1471-8286.2005.01155.x.
- Peakall R, Smouse PE. 2012.** GenALEX 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28:2537–2539 DOI 10.1093/bioinformatics/bts460.
- Peery MZ, Kirby R, Reid BN, Stoelting R, Doucet-Béer E, Robinson S, Vásquez-Carrillo C, Pauli JN, Palsbøll PJ. 2012.** Reliability of genetic bottleneck tests

- for detecting recent population declines. *Molecular Ecology* **21**:3403–3418
[DOI 10.1111/j.1365-294X.2012.05635.x](https://doi.org/10.1111/j.1365-294X.2012.05635.x).
- Pichler FB, Baker CS. 2000.** Loss of genetic diversity in the endemic Hector's dolphin due to fisheries-related mortality. *Proceedings of the Royal Society B: Biological Sciences* **267**:97–102 [DOI 10.1098/rspb.2000.0972](https://doi.org/10.1098/rspb.2000.0972).
- Piry S, Luikart G, Cornuet J-M. 1999.** Bottleneck: a program for detecting recent effective population size reductions from allele data frequencies. *Journal of Heredity* **90**:502–503 [DOI 10.1093/jhered/90.4.502](https://doi.org/10.1093/jhered/90.4.502).
- Piry S, Alapetite A, Cornuet JM, Paetkau D, Baudouin L, Estoup A. 2004.** GeneClass2: a software for genetic assignment and first-generation migrant detection. *Journal of Heredity* **95**:536–539 [DOI 10.1093/jhered/esh074](https://doi.org/10.1093/jhered/esh074).
- Pritchard JK, Stephens P, Donnelly P. 2000.** Inference of population structure using multilocus genotype data. *Genetics* **155**:945–959.
- Puechmaille SJ. 2016.** The program structure does not reliably recover the correct population structure when sampling is uneven: subsampling and new estimators alleviate the problem. *Molecular Ecology Resources* **16**:608–627 [DOI 10.1111/1755-0998.12512](https://doi.org/10.1111/1755-0998.12512).
- Rannala B, Mountain JL. 1997.** Detecting immigration by using multilocus genotypes. *Proceedings of the National Academy of Sciences of the United States of America* **94**:9197–9221 [DOI 10.1073/pnas.94.17.9197](https://doi.org/10.1073/pnas.94.17.9197).
- Raymond M, Rousset F. 1995.** GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity* **86**:248–249
[DOI 10.1093/oxfordjournals.jhered.a111573](https://doi.org/10.1093/oxfordjournals.jhered.a111573).
- Reed DH, O'Grady JJ, Brook BW, Ballouc JD, Frankhama R. 2003.** Estimates of minimum viable population sizes for vertebrates and factors influencing those estimates. *Biological Conservation* **113**:23–24 [DOI 10.1016/S0006-3207\(02\)00346-4](https://doi.org/10.1016/S0006-3207(02)00346-4).
- Rick TC, Lockwood R. 2013.** Integrating paleobiology, archeology, and history to inform biological conservation. *Conservation Biology* **27**:45–54
[DOI 10.1111/j.1523-1739.2012.01920.x](https://doi.org/10.1111/j.1523-1739.2012.01920.x).
- Roman J, Palumbi SR. 2003.** Whales before whaling in the North Atlantic. *Science* **301**:508–510.
- Rousset F. 2008.** Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. *Molecular Ecology Resources* **8**:103–106
[DOI 10.1111/j.1471-8286.2007.01931.x](https://doi.org/10.1111/j.1471-8286.2007.01931.x).
- Rozen S, Skaletsky HJ. 2000.** Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S, Totowa N, eds. *Bioinformatics methods and protocols: methods in molecular biology*. Totowa: Humana Press, 365–386.
- Seppä H, Bjune AE, Telford RJ, Birks HJB, Veski S. 2009.** Last nine-thousand years of temperature variability in Northern Europe. *Climate of the Past* **5**:523–535
[DOI 10.5194/cp-5-523-2009](https://doi.org/10.5194/cp-5-523-2009).
- Swart JAA, Reijnders PJH, Van Delden W. 1996.** Absence of genetic variation in harbor seals (*Phoca vitulina*) in the Dutch Wadden Sea and the British Wash. *Conservation Biology* **10**:289–293 [DOI 10.1046/j.1523-1739.1996.10010289.x](https://doi.org/10.1046/j.1523-1739.1996.10010289.x).

- Thioulouse J, Chessel D, Dolédec J, Olivier JM. 1997.** ADE-4: a multivariate analysis and graphical display software. *Statistics and Computing* 7:75–83 DOI [10.1023/A:1018513530268](https://doi.org/10.1023/A:1018513530268).
- Thompson PM, Hall AJ. 1993.** Seals and epizootics—what factors might affect the severity of mass mortalities? *Mammal Review* 23:149–154 DOI [10.1111/j.1365-2907.1993.tb00427.x](https://doi.org/10.1111/j.1365-2907.1993.tb00427.x).
- Thompson PM, Kovacs KM, McConnell BJ. 1994.** Natal dispersal of harbour seals (*Phoca vitulina*) from breeding sites in Orkney. *Scotland. J. Zool. London* 234:668–673 DOI [10.1111/j.1469-7998.1994.tb04873.x](https://doi.org/10.1111/j.1469-7998.1994.tb04873.x).
- Thompson D, Duck CD, Morris CD, Russell DJF. 2019.** The status of harbour seals (*Phoca vitulina*) in the UK. *Aquatic Conservation Marine and Freshwater Ecosystems* 29:40–60 DOI [10.1002/aqc.3110](https://doi.org/10.1002/aqc.3110).
- Thompson PM, Tollit DJ, Wood DC, Corpe HM. 1997a.** Estimating harbour seal abundance and status in an estuarine habitat in north-east Scotland. *Journal of Applied Ecology* 34:43–52 DOI [10.2307/2404846](https://doi.org/10.2307/2404846).
- Thompson P, Tollit D, Corpe H, Reid R, Ross H. 1997b.** Changes in haematological parameters in relation to prey switching in a wild population of harbour seals. *Functional Ecology* 11:743–750 DOI [10.1046/j.1365-2435.1997.00148.x](https://doi.org/10.1046/j.1365-2435.1997.00148.x).
- Waples RS, England PR. 2011.** Estimating contemporary effective population size on the basis of linkage disequilibrium in the face of migration. *Genetics* 189:633–644 DOI [10.1534/genetics.111.132233](https://doi.org/10.1534/genetics.111.132233).
- Weir BS, Cockerham CC. 1984.** Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358–1370.
- Whitehead H. 1998.** Cultural selection and genetic diversity in matrilineal whales. *Science* 282:1708–1711 DOI [10.1126/science.282.5394.1708](https://doi.org/10.1126/science.282.5394.1708).
- Wilson GA, Rannala B. 2003.** Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* 163:1177–1191.
- Wilson LJ, Hammond PS. 2019.** The diet of harbour and grey seals around Britain: examining the role of prey as a potential cause of harbour seal declines. *Aquatic Conservation: Marine and Freshwater Ecosystems* 29:71–85.
- Wright S. 1931.** Evolution in Mendelian populations. *Genetics* 16:97–159.
- Wright S. 1969.** Evolution and genetics of populations. In: *The theory of gene frequencies*. Vol.2. Chicago: University of Chicago Press.