

REVIEW ARTICLE

COVID-19 and cutaneous manifestations: a systematic review

Qing Zhao, Xiaokai Fang, Zheng Pang, Bowen Zhang, Hong Liu, Furen Zhang

Shandong Provincial Hospital for Skin Diseases & Shandong Provincial Institute of Dermatology and Venereology, Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan, Shandong, China

*Correspondence: F. Zhang. E-mail: zhangfuren@hotmail.com

Abstract

The cutaneous manifestations of COVID-19 patients have been increasingly reported, but not summarized, and the potential mechanisms remain to be investigated. Herein, we performed a comprehensive review of literatures (from inception to 30 May 2020) using PubMed, CNKI, medRxiv and bioRxiv with the terms “((novel coronavirus) OR (2019 novel coronavirus) OR (2019-nCoV) OR (Coronavirus disease 2019) OR (COVID-19) OR (SARS-CoV-2)) AND ((Dermatology) OR (skin) OR (rash) OR (cutaneous))” and “((ACE2) OR (Angiotensin-converting enzyme)) AND ((skin) OR (epidermis) OR (dermis)).” Totally, 44 articles met the inclusion criteria. A total of 507 patients with cutaneous manifestations were summarized, and 96.25% patients were from Europe. The average age of the patients was 49.03 (range: 5–91) with a female ratio of 60.44%. The skin lesions were polymorphic, and erythema, chilblain-like and urticarial lesions were most common, occurring on an average of 9.92 days (range: 1–30) after the onset of systemic symptoms. The receptor of SARS-CoV-2, ACE2, was found to be expressed on skin, mainly on keratinocytes. Our review systematically presented the clinical characteristics of 507 patients and showed that skin might be the potential target of the infection according to ACE2 expression. More work should be done to better understand the underlying pathogenesis.

Received: 14 May 2020; Accepted: 8 June 2020

Conflicts of interest

None declared.

Funding sources

The authors acknowledge the financial support by the grants from Academic promotion programme of Shandong First Medical University (2019LJ002), the Innovation Project of Shandong Academy of Medical Sciences and the Shandong Province Taishan Scholar Project (tsqn201812124, tspd20150214).

Introduction

In December 2019, a novel coronavirus, severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), was firstly reported in Wuhan, China, as an etiological agent causing a new infectious respiratory disease (coronavirus disease 2019, COVID-19).¹ SARS-CoV-2 spread rapidly worldwide through human-to-human transmission and COVID-19 has been declared as a pandemic emergency by World Health Organization (WHO) on March, 2020. By 30 May 2020, 5 701 337 patients were confirmed as COVID-19 cases and 357 688 patients were dead.

The main clinical manifestations of COVID-19 were fever, cough, fatigue, sputum production, dyspnoea and muscle aches.^{2,3} The diarrhoea, olfactory and gustatory impairments caused by COVID-19 were also reported but less common.^{4,5} An increasing number of literatures on cutaneous manifestations of

COVID-19 patients have been recently reported, suggesting that skin lesions could be a potential clinical characteristic of COVID-19. Various viewpoints about the mechanisms inducing the skin rash in COVID-19 patients have been proposed^{6,7}: (i) Whether the virus could infect through an open wound of skin; (ii) Whether skin manifestations were related to immune responses; (iii) Whether skin manifestations were caused by a new prescribed medication. However, the researches on the pathogenesis of cutaneous manifestations of COVID-19 have not been performed yet. Given that angiotensin-converting enzyme 2 (ACE2) is a crucial functional receptor of SARS-CoV-2,^{8–12} exploration of the ACE2 expression in skin tissues could facilitate to clarify the mechanisms involved in cutaneous manifestations of COVID-19. Herein, to gain an in-depth understanding of the COVID-19, we summarized the publications related to the cutaneous manifestations of COVID-19 cases and the ACE2 expression in skin tissues.

Zhao and Fang contributed equally to the manuscript.

Methods/literature search

A systematic review was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. We searched in the Medline database (PubMed), CNKI (<https://www.cnki.net/>), medRxiv (<https://www.medrxiv.org/>) and bioRxiv (<https://www.biorxiv.org/>) with the combined terms “((novel coronavirus) OR (2019 novel coronavirus) OR (2019-nCoV) OR (Coronavirus disease 2019) OR (COVID-19) OR (SARS-CoV-2)) AND ((Dermatology) OR (skin) OR (rash) OR (cutaneous))” and “((ACE2) OR (Angiotensin-converting enzyme)) AND ((skin) OR (epidermis) OR (dermis)).” Searches were limited to the publications before 30 May 2020. Case reports, clinical analysis and fundamental research discussing the cutaneous manifestations and expression of ACE2 in skin were included. Unavailable articles, review articles, expert consensus, medical hypothesis articles and the articles focusing on the below aspects were excluded: the skin problems of medical staff caused by using protective equipment, the effects of the epidemic on the use of immunosuppressive agents and other drugs in dermatology, the effects of dermatological instruments, and the effects resulted from the treatment of COVID-19 transmission. The cases with clinical diagnosis of the disease (suspected cases) were included in addition to the cases confirmed of COVID-19 by viral nucleic acid detection, viral gene sequencing and serum antibody tests (Fig. S1).

Results

Study selection and study population

Data from 164 papers were collected. After screening, 120 of them were excluded. We recruited 44 articles, including 38 case reports (Table S1) and six publications of ACE2 expression in skin tissues (Table S2).

Cutaneous manifestations of COVID-19 patients

Lu *et al.*¹³ reported a COVID-19 case with urticaria as the first cutaneous manifestation on March 19, 2020. Subsequently, Recalcati *et al.*¹⁴ from Italy, Zhang *et al.*¹⁵ from China and Bouaziz *et al.*¹⁶ from France summarized 18, 7 and 14 cases, respectively. In Spain, Fernandez-Nieto *et al.*¹⁷ and Galván *et al.*¹⁸ conducted a nationwide case collection survey and reported 19 and 375 COVID-19 cases with cutaneous manifestations, respectively. However, 17 and 141 of the cases were clinically confirmed rather than diagnosed without etiological evidence, respectively. In addition, 73 sporadic cases have been reported.^{7,19–49} By 30 May 2020, a total of 507 cases of COVID-19 accompanied by skin manifestations were reported. And 488 out of 507 (96.25%) patients were from Europe.

Basic information

Among the 450 patients reported the basic information, the ratio of male to female was 178/272. The mean age of patients was

49.03 (range: 5–91). According to a previous study of COVID-19 in China, male patients accounted for about 53%, and the average age was 49.54.⁵⁰ Thus, we assumed that the incidence of skin lesions in COVID-19 patients was regardless of ages and genders.

The onset of skin manifestations

Eighty-eight of the 507 patients provided the information on the onset of skin manifestations. Among them, lesions occurred on average 9.92 days (range: 1–30) after the onset of systemic symptoms including fever, cough, dyspnoea, diarrhoea and fatigue. In addition, the longest incubation of lesions was 30 days,¹⁷ and it was noteworthy that 13 patients (14.77%) had skin lesions as the first symptom.^{17,25,26,29,31,45,47,49}

Clinical manifestations

All 507 patients had descriptions on their clinical manifestations. We found that the skin symptoms of COVID-19 patients were polymorphic. The most common skin lesion was erythema, which was observed in 224 patients (44.18%) and distributed on patients' trunk, extremities, flexural regions, face and mucous membranes. Moreover, the erythema lesions were also confined to specific sites, such as the heels¹⁴ without other triggers such as exposure. Chilblain-like lesions were described in 100 (19.72%) patients. Urticaria-like lesions were presented in 83 patients (16.37%) and distributed on patients' trunks or dispersed widely on their bodies.¹⁸ In addition, other manifestations such as vesicular (66, 13.02%), livedo/necrosis (31, 6.11%) and petechiae (8, 1.58%) were described. Significantly, 227 patients (44.77%) complained of significant pruritus.

Among the 507 patients, 431 had the information of systemic symptoms. Fever was the most common and presented in 327 patients (75.87%). Other symptoms including cough (299, 69.37%), fatigue or asthenia (252, 58.47%), dyspnoea (201, 46.64%), headache (131, 30.39%), gastrointestinal symptoms (116, 26.91%) and anosmia or ageusia (96, 22.27%) were also described.

Skin manifestations and COVID-19 severity

The severity of pneumonia in COVID-19 patients presenting cutaneous manifestations varied. Zhang *et al.*¹⁵ reported seven COVID-19 patients with critical symptoms. All the patients had extremity ischaemia with varied severity, which was manifested as cyanosis, purpura, hemophysallia, and dry gangrene. In addition, the skin of two patients become darker than usual after treatment, which might be related to severe symptoms, multiple organ damage and drug treatment.⁵¹ We speculate that the skin changes in these patients might be caused by multiple organ damages or systemic ischaemia, suggesting a severe condition.

For the remaining cases, Only five patients had clear information about the severity of COVID-19, including one mild, two ordinary and two severe cases, according to the diagnostic

criteria.⁵² Furthermore, Recalcati *et al.*¹⁴ argued that the skin manifestations of COVID-19 patients were similar to other skin diseases caused by viral infection, and not related to the severity of pneumonia. According to a study by Galván *et al.*,¹⁸ pseudo-chilblain, vesicular lesions, urticarial lesions, maculopapular lesions and livedoid/necrotic lesions were associated with progressively increasing disease severity.

Laboratory examinations

Due to the harsh conditions and equipment constraints during the COVID-19 epidemic, only 23 patients performed histopathological examination. All the 23 patients showed inflammatory features without specific pathological changes, such as lymphocyte infiltration.^{20,24,40,41} Amatore *et al.*²⁵ described that no virally induced cytopathic alterations or intranuclear inclusions were present. Sanchez *et al.*,³⁴ Fernandez-Nieto *et al.*³⁹ and Ahouach *et al.*³⁵ performed RT-PCR on skin biopsy specimen, and results were negative for SARS-CoV-2. None of the remaining patients underwent virus detection for skin lesions.

The peripheral blood examination was performed on 39 patients. The most common outcomes were lymphocytopenia (30 patients, 76.92%), C-reaction protein elevation (13 patients, 33.33%), and lactate dehydrogenase (LDH) elevation (18 patients, 46.15%). In the seven critical patients reported by Zhang *et al.*,¹⁵ D-dimer was significantly increased, and platelet count was normal. Fibrinogen and fibrin degradation product (FDP) were increased in six patients, and prothrombin time (PT) was prolonged in four patients. Two patients showed thrombocytopenia, corresponding to a skin lesion that includes petechiae.²¹

Complications

Ten patients had the complication of hypertension, and six patients had diabetes. In a study by Galván *et al.*,¹⁸ the number of smokers reached 44, accounting for 11.73% of a total of 375 cases.

Prognosis

For the prognosis of all 507 patients, only two cured and 13 dead cases were clearly described. In the seven critical patients, five patients died and the two others were still under treatment after 26 days of follow-up, and the causes of death were circulatory failure resulted from severe hypoxaemia and respiratory acidosis.¹⁵ In addition, a 71-year-old male patient died due to his advanced age and multiple underlying diseases.²⁶ In a study by Galván *et al.*,¹⁸ there were 369 survival and six dead cases. According to the description of the symptoms, we speculated that most of the patients had a relatively good prognosis and the reported death rate was 2.56% (13/507).

For skin lesions, 39 patients resolved spontaneously or with some basic medication within a week, and other cases did not

provide details data. It remains to be discussed whether seven patients develop skin lesions after taking a drug that targets COVID-19. Although some drugs, such as paracetamol, hydroxychloroquine and lopinavir/ritonavir, were commonly used with rare adverse reactions, it is still debatable whether the skin lesions were caused by viral infections or adverse drug reactions.^{20,33,48,53}

All the information of the 507 COVID-19 patients was summarized in Table 1.

SARS-CoV-2 entry factors expressed in skin

To our acknowledgement, no studies regarding the mechanisms of devolvement of cutaneous rash in COVID-19 patients have been carried out. Thus, we only summarized the studies linked to the expression of ACE2 in skin tissues. Sungnak *et al.*¹¹ reported that expression of ACE2 in skin tissues was not detectable in scRNA-seq data sets. However, the ARCHS4 database (<https://amp.pharm.mssm.edu/archs4/gene/ACE2>) showed that ACE2 was expressed mainly in keratinocytes and basal cells in skin tissues. Furthermore, the expression of ACE2 has been investigated by bulk samples of human skin⁵⁴ and in epidermal stem cells.⁵⁵ ACE2 immunoreactivity was detected by immunohistochemistry (IHC) in the cells of the basal layer of the epidermis extending to the basal cell layer of hair follicles, sebaceous gland cells as well as the smooth muscle cells surrounding the sebaceous glands, and the cells of the eccrine glands in normal skin.^{56,57} Recently, scRNA-seq was applied to analyse the ACE2 mRNA expression in different cell types in skin tissues.⁵⁸ Higher ACE2 expression was identified in keratinocytes mainly in differentiating keratinocytes and basal cells compared with the other cells of skin tissues. The results were further confirmed by IHC, which showed the ACE2-positive keratinocytes in the stratum basal, the stratum spinosum and the stratum granulosum of epidermis. In addition, Li *et al.*⁵⁹ found that CD8⁺ T-cell enrichment had significant positive correlations with the ACE2 expression in the skin.

Discussion

Infection with SARS-CoV-2 causes a broad spectrum of clinical syndromes, ranging from mild pneumonia to acute respiratory distress syndrome. Although cutaneous manifestations are rare, the number of COVID-19 patients with skin involvement has been reported to be increased. However, up to date, no *in vitro* or *in vivo* experiments have been performed to confirm whether SARS-Cov-2 could spread through skin. In this review, we found that ACE2 was expressed in the skin, especially in keratinocytes. This provides an evidence for percutaneous infection or the entry of virus into patients through skin tissue. Furthermore, SARS-CoV-2 RNA was detected in blood samples from 22% patients.⁶⁰ Thus, it is possible that the ACE2 expression in skin cells could be induced by viral infection via hematogenous dissemination, especially for the patients who were in a state of

Table 1 Summary of the demographics and clinical characteristics

Basic information	
Total number of cases	507
Female sex, n (%)	272/450 (60.44%)
Mean age (range), years	49.03 (5–91)
Country (n = 507), n (%)	
Spain	434 (85.60%)
Italy	33 (6.51%)
France	19 (3.75%)
China	8 (1.58%)
American	5 (0.99%)
Canada	3 (0.59%)
Belgium	2 (0.39%)
Thailand	1 (0.20%)
Indonesia	1 (0.20%)
Japan	1 (0.20%)
Mean time of appearance of skin lesions after onset of COVID-19 (range), days	9.92 (1–30)
Cutaneous manifestations (n = 507), n (%)	
Erythema	224 (44.18%)
Chilblain-like lesions	100 (19.72%)
Urticaria-like lesions	83 (16.37%)
Vesicular	66 (13.02%)
Livedo/necrosis	31 (6.11%)
Petechiae	8 (1.58%)
Accompanied by pruritus	227 (44.77%)
Mean duration of skin lesions, days	9.10
Information of COVID-19 (n = 507), n (%)	
Confirmed cases	343 (67.65%)
Suspected cases	164 (32.35%)
Systemic symptom (n = 431), n (%)	
Fever	327 (75.87%)
Cough	299 (69.37%)
dyspnoea	201 (46.64%)
Fatigue or asthenia	252 (58.47%)
Anosmia or ageusia	96 (22.27%)
Headache	131 (30.39%)
Gastrointestinal symptom	116 (26.91%)
Complications and medical history, n (%)	
Smoking	44/375 (11.73%)
Hypertension	10/18 (55.56%)
Diabetes	6/18 (33.33%)
Laboratory examinations, n (%)	
Histopathology	23/507 (4.54%)
Lymphocytopenia	30/39 (76.92%)
Elevated CRP	13/39 (33.33%)
Elevated LDH	18/39 (46.15%)
Case fatality, n (%)	13/507 (2.56%)

viraemia. More biopsy specimens should be collected to detect whether the presence of virus in skin lesions to confirm it.

According to our review, erythematous rash is the most common feature among 507 patients. However, the mechanisms of

erythema development in the COVID-19 patients remain unclear. It has been found that herpes simplex virus (HSV) could cause a subset of erythema multiforme (EM) lesions (herpes-associated EM (HAEM)).⁶¹ After infection by HSV, the DNA of HSV would be fragmented and the DNA fragments could be transported to distant skin sites by peripheral blood mononuclear cells (PBMC). The HSV-specific CD4⁺ Th1 cells could be recruited by the viral antigen to skin and initiate an inflammatory cascade, which induces the development of HAEM. For SARS-CoV-2 infection, although no studies have reported virus in skin lesions, hyperactivated CD4⁺ and CD8⁺ T cells were found in peripheral blood from patients by flow cytometry. Moreover, the concentration of CCR4- and CCR6-positive Th17 in CD4 T cells was increased. CCR4 has been implicated in the migration of T cells into skin, and its ligand, CCL17, is expressed on cutaneous endothelial cells.⁶² Thus, the virus-specific T cells in circulation could migrate into skin tissues. In addition, lymphocyte infiltration has also been observed in the skin lesions of COVID-19 patients. These findings suggested that the skin rash might be caused by the virus-specific T cells.

Urticaria is another common feature among those COVID-19 patients who presented rashes. Previous studies hypothesized that viral IgM and IgG could cross-react with mast cell IgE and cause the mast cell degranulation subsequently.⁶³ SARS-CoV-2-specific IgM and IgG can be detected in patients.⁶⁴ Thus, the cross-reaction between SARS-CoV-2 antibodies with mast cell IgE might result in degranulation of mast cell and wheal growth.

Chilblain-like lesions, ischaemic lesions and ecchymotic acral lesions were also observed in COVID-19 patients. It has been reported that blood coagulation is altered in COVID-19 patients and the acro-ischaemia might be associated with the hypercoagulation status in severe COVID-19 patients.¹⁵ Recently, these types of skin lesions were also found in asymptomatic or mildly symptomatic patients, probably caused by a coagulation disorder or a hypersensitivity reaction.^{17,65} Unfortunately, the mechanisms of cutaneous manifestations are only hypothetical at present. Interestingly, it is likely that the incidence of skin rashes among COVID-19 patients varied in different countries. Only eight cases were reported in China whereas 488 cases were from Europe. A study in China reported that only two of 1,099 patients showed skin lesions,⁶⁶ whereas Recalcati *et al.*,¹⁴ an Italian dermatologist team, described that skin changes occurred in 18 of the 88 confirmed COVID-19 patients, and Hedou *et al.*⁶⁷ reported that the incidence of cutaneous manifestations associated with COVID-19 infections in France was 4.9% based on a prospective study of 103 cases. It is unclear that whether the difference was caused by missed diagnoses and unreported cases, or regional differences.

It is noteworthy that skin rashes are also observed in general population. For example, in a population-based study, up to 1% of the general population in the USA and Europe suffered from

chronic urticarial at some point in their lifetime.⁶⁸ The ratio was significantly higher in Asia, which was about 9–20%.⁶⁹ Urticaria-like lesions were presented in 16.37% COVID-19 patients with skin manifestation in our study. Therefore, we could not rule out that the cutaneous lesions of COVID-19 patients were merely an incidental phenomenon based on the population incidence.

Conclusions

In conclusion, our review summarized the clinical characteristics of skin lesions in COVID-19 patients and the possible mechanisms by reviewing the ACE2 expression in skin tissues. However, we should acknowledge that lots of questions regarding COVID-19 remain unexplored. More clinical data should be collected, and more researches should be carried out for a better understanding of the cutaneous manifestations caused by COVID-19.

References

- Huang C, Wang Y, Li X *et al.* Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; **395**: 497–506.
- Lovato A, Filippis C. Clinical presentation of COVID-19: a systematic review focusing on upper airway symptoms. *Ear Nose Throat J* 2020. <https://doi.org/10.1177/0145561320920762>. [Epub ahead of print]
- Fang Z, Yi F, Wu K *et al.* Clinical characteristics of coronavirus pneumonia 2019 (COVID-19): an updated systematic review. *medRxiv* 2020. <https://doi.org/10.1101/2020.03.07.20032573>. [Epub ahead of print]
- Liang F, Rao S, Xiao C *et al.* Diarrhoea may be underestimated: a missing link in 2019 novel coronavirus. *Gut* 2020; **69**: 1141–1143.
- Xydakis M, Dehgani-Mobaraki P, Holbrook E *et al.* Smell and taste dysfunction in patients with COVID-19. *Lancet Infect Dis* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30293-0](https://doi.org/10.1016/S1473-3099(20)30293-0). [Epub ahead of print]
- Cavanagh G, Wambier C. Reply to: "Personal protective equipment recommendations based on COVID-19 route of transmission". *J Am Acad Dermatol* 2020; **83**: e47.
- Estebanez A, Perez-Santiago L, Silva E, Guillen-Climent S, Garcia-Vazquez A, Ramon M. Cutaneous manifestations in COVID-19: a new contribution. *J Eur Acad Dermatol Venereol* 2020; **34**: e250–e251. <https://doi.org/10.1111/jdv.16474>
- Lu R, Zhao X, Li J *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet* 2020; **395**:565–574.
- Wrapp W, Corbett K, Goldsmith J *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020; **367**: 1260–1263.
- Zhou P, Wang X, Hu B *et al.* Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *bioRxiv* 2020. <https://doi.org/10.1101/2020.01.22.914952>. [Epub ahead of print]
- Sungnak W, Huang N, Becavin C *et al.* SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat Med* 2020; **26**: 681–687.
- Lukassen S, Chua RL, Trefzer T *et al.* SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial transient secretory cells. *EMBO J* 2020; **39**: e105114.
- Lu S, Lin J, Zhang Z *et al.* Alert for non-respiratory symptoms of Coronavirus Disease 2019 (COVID-19) patients in epidemic period: a case report of familial cluster with three asymptomatic COVID-19 patients. *J Med Virol* 2020. <https://doi.org/10.1002/jmv.25776>. [Epub ahead of print]
- Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol* 2020; **34**: e212–e213.
- Zhang Y, Cao W, Xiao M *et al.* Clinical and coagulation characteristics of 7 patients with critical COVID-2019 pneumonia and acro-ischemia. *Zhonghua Xue Ye Xue Za Zhi* 2020; **41**: E006.
- Bouaziz J, Duong T, Jachiet M *et al.* Vascular skin symptoms in COVID-19: a French observational study. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16544>. [Epub ahead of print]
- Fernandez-Nieto D, Jimenez-Cauhe J, Suarez-Valle A *et al.* Characterization of acute acro-ischemic lesions in non-hospitalized patients: a case series of 132 patients during the COVID-19 outbreak. *J Am Acad Dermatol* 2020; **83**: e61–e63. <https://doi.org/10.1016/j.jaad.2020.04.093>
- Galván C, Català A, Carretero H *et al.* Classification of the cutaneous manifestations of COVID-19: a rapid prospective nationwide consensus study in Spain with 375 cases. *Br J Dermatol* 2020; **183**: 71–77. <https://doi.org/10.1111/bjd.19163>
- Mahe A, Birckel E, Krieger S, Merklen C, Bottlaender L. A distinctive skin rash associated with Coronavirus Disease 2019? *J Eur Acad Dermatol Venereol* 2020; **34**: e246–e247. <https://doi.org/10.1111/jdv.16471>
- Fernandez-Nieto D, Ortega-Quijano D, Segurado-Miravalles G, Pindado-Ortega C, Prieto-Barrios M, Jimenez-Cauhe J. Comment on: Cutaneous manifestations in COVID-19: a first perspective. Safety concerns of clinical images and skin biopsies. *J Eur Acad Dermatol Venereol* 2020; **34**: e252–e254. <https://doi.org/10.1111/jdv.16470>
- Joob B, Wiwanitkit V. COVID-19 can present with a rash and be mistaken for dengue. *J Am Acad Dermatol* 2020; **82**: e177.
- Tammaro A, Adebajo G, Parisella F, Pezuto A, Rello J. Cutaneous manifestations in COVID-19: the experiences of Barcelona and Rome. *J Eur Acad Dermatol Venereol* 2020; **34**: e306–e307. <https://doi.org/10.1111/jdv.16530>
- Landa N, Mendieta-Eckert M, Fonda-Pascual P, Aguirre T. Chilblain-like lesions on feet and hands during the COVID-19 pandemic. *Int J Dermatol* 2020; **59**: 739–743.
- Gianotti R, Veraldi S, Recalcati S *et al.* Cutaneous clinico-pathological findings in three COVID-19-positive patients observed in the metropolitan area of Milan, Italy. *Acta Derm Venereol* 2020; **100**: adv00124.
- Amatore F, Macagno N, Mailhe M *et al.* SARS-CoV-2 infection presenting as a febrile rash. *J Eur Acad Dermatol Venereol* 2020; **34**: e304–e306. <https://doi.org/10.1111/jdv.16528>
- van Damme C, Berlingin E, Saussez S, Accaputo O. Acute urticaria with pyrexia as the first manifestations of a COVID-19 infection. *J Eur Acad Dermatol Venereol* 2020; **34**: e300–e301. <https://doi.org/10.1111/jdv.16523>
- Avellana M, Villa E, Avellana M, Estela V, Aparicio M, Fontanella A. Cutaneous manifestation of COVID-19 in images: a case report. *J Eur Acad Dermatol Venereol* 2020; **34**: e307–e309. <https://doi.org/10.1111/jdv.16531>
- Najarjan DJ. Morbilliform exanthem associated with COVID-19. *JAAD Case Repo* 2020; **6**: 493–494.
- Hunt M, Koziatek C. Case of COVID-19 pneumonia in a young male with full body rash as a presenting symptom. *Clin Pract Cases Emerg Med* 2020; **4**: 219–221. <https://doi.org/10.5811/cpcem.2020.3.47349> [Epub ahead of print].
- Manalo I, Smith M, Cheeley J, Jacobs R. A dermatologic manifestation of COVID-19: transient livedo reticularis. *J Am Acad Dermatol* 2020; **83**: 700–. <https://doi.org/10.1016/j.jaad.2020.04.018>
- Henry D, Ackerman M, Sancelme E, Finon A, Esteve E. Urticarial eruption in COVID-19 infection. *J Eur Acad Dermatol Venereol* 2020; **34**: e244–e245. <https://doi.org/10.1111/jdv.16472>
- Genovese G, Colonna C, Marzano A. Varicella-like exanthem associated with COVID-19 in an 8-year-old girl: a diagnostic clue? *Pediatr Dermatol* 2020; **37**: 435–436. <https://doi.org/10.1111/pde.14201>
- Jimenez-Cauhe J, Ortega-Quijano D, Prieto-Barrios M, Moreno-Arrones OM, Fernandez-Nieto D. Reply to "COVID-19 can present with a rash and be mistaken for dengue": petechial rash in a patient with COVID-19 infection. *J Am Acad Dermatol* 2020; **83**: e141–e142. <https://doi.org/10.1016/j.jaad.2020.04>

- 34 Sanchez A, Sohler P, Benganem S et al. Digitate papulosquamous eruption associated with severe acute respiratory syndrome coronavirus 2 infection. *JAMA Dermatol* 2020. <https://doi.org/10.1001/jamadermatol.2020.1704>. [Epub ahead of print]
- 35 Ahouach B, Harant S, Ullmer A et al. Cutaneous lesions in a patient with COVID-19: are they related? *Br J Dermatol* 2020; **183**: e31–. <https://doi.org/10.1111/bjd.19168>
- 36 Diaz-Guimaraens B, Dominguez-Santas M, Suarez-Valle A et al. Petchial skin rash associated with severe acute respiratory syndrome coronavirus 2 infection. *JAMA Dermatol* 2020. <https://doi.org/10.1001/jamadermatol.2020.1741>. [Epub ahead of print]
- 37 Martin C, Amaro J, Lopez-Sanchez A, Jane-Salas E, Somacarrera M. Oral vesiculobullous lesions associated with SARS-CoV-2 infection. *Oral Dis* 2020. <https://doi.org/10.1111/odi.13382>. [Epub ahead of print]
- 38 Locatelli AG, Robustelli Test E, Vezzoli P et al. Histologic features of long lasting chilblain-like lesions in a pediatric COVID-19 patient. *J Eur Acad Dermatol Venereol* 2020; **34**: e365–e368. <https://doi.org/10.1111/jdv.16617>
- 39 Fernandez-Nieto D, Ortega-Quijano D, Jimenez-Cauhe J et al. Clinical and histological characterization of vesicular COVID-19 rashes: a prospective study in a tertiary care hospital. *Clin Exp Dermatol* 2020. <https://doi.org/10.1111/ced.14277>. [Epub ahead of print].
- 40 Jimenez-Cauhe J, Ortega-Quijano D, Carretero-Barrio I et al. Erythema multiforme-like eruption in patients with COVID-19 infection: clinical and histological findings. *Clin Exp Dermatol* 2020. <https://doi.org/10.1111/ced.14281>. [Epub ahead of print]
- 41 Colonna C, Monzani N, Rocchi A, Gianotti R, Boggio F, Gelmetti C. Chilblains-like lesions in children following suspected Covid-19 infection. *Pediatr Dermatol* 2020; **37**: 437–440. <https://doi.org/10.1111/pde.14210>
- 42 Gunawan C, Angela A, Widysanto A. Urticarial eruption in Coronavirus Disease 2019 (COVID-19) infection: a case report in Tangerang, Indonesia. *J Eur Acad Dermatol Venereol* 2020; **34**: e372–e373. <https://doi.org/10.1111/jdv.16622>
- 43 Robustelli E, Vezzoli P, Carugno A et al. Acute generalized exanthematous pustulosis with erythema multiforme-like lesions in a COVID-19 woman. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16613>. [Epub ahead of print]
- 44 Sachdeva M, Gianotti R, Shah M et al. Cutaneous manifestations of COVID-19: report of three cases and a review of literature. *J Dermatol Sci* 2020; **98**: 75–81.
- 45 Kalner S, Vergilis IJ. Periorbital erythema as a presenting sign of COVID-19. *JAAD Case Rep* 2020; <https://doi.org/10.1016/j.jdc.2020.05.001>. [Epub ahead of print]
- 46 Rodríguez-Jiménez P, Chicharro P, De A, Muñoz-Hernández P, Llamas-Velasco M. Reply to "Acute urticaria with pyrexia as the first manifestations of a COVID-19 infection": urticaria-like lesions in COVID-19 patients are not really urticaria. A case with clinicopathologic correlation. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16618>. [Epub ahead of print]
- 47 Balestri R, Termine S, Rech G, Girardelli C. Late onset of acral necrosis after SARS-CoV-2 infection resolution. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16668>. [Epub ahead of print]
- 48 Skroza N, Bernardini N, Balduzzi V et al. A late onset widespread skin rash in a previous Covid-19 infected patient: viral or multidrug effect? *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16633>. [Epub ahead of print]
- 49 Sakaida T, Isao T, Matsubara A, Nakamura M, Morita A. Unique skin manifestations of COVID-19: Is drug eruption specific to COVID-19? *J Dermatol Sci* 2020. <https://doi.org/10.1016/j.jdermsci.2020.05.002>. [Epub ahead of print]
- 50 Wu Z, McGoogan J. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China. *JAMA* 2020; **323**: 1239.
- 51 Zhu Y. The skin of COVID-19 patients become black, do not ignore pharmacological causes. *Health News* 2020. https://www.sohu.com/a/389573042_377345 (last accessed 20 April 2020).
- 52 Cheng Z, Shan J. 2019 Novel coronavirus: where we are and what we know. *Infection* 2020; **48**: 155–163.
- 53 Mahé A, Birckel E, Krieger S, Merklen C, Bottlaender L. A distinctive skin rash associated with Coronavirus Disease 2019? *J Eur Acad Dermatol Venereol* 2020; **34**: e246–e247. <https://doi.org/10.1111/jdv.16471>
- 54 Harmer G, Borman R, Clark K. Quantitative mRNA expression profiling of ACE 2, a novel homologue of angiotensin converting enzyme. *FEBS Lett* 2002; **532**:107–110.
- 55 Liao X, Xiao J, Li S et al. Critical role of the endogenous renin-angiotensin system in maintaining self-renewal and regeneration potential of epidermal stem cells. *Biochim Biophys Acta Mol Basis Dis* 2019; **1865**: 2647–2656.
- 56 Grzegorzolka S, Pula B, Zamirska A et al. ACE and ACE2 expression in normal and malignant skin lesions. *Folia Histochem Cytobiol* 2013; **51**:232–238.
- 57 Hamming I, Timens W, Bulthuis M, Lely A, Navis G, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol* 2004; **203**: 631–637.
- 58 Xue X, Mi Z, Wang Z, Pang Z, Liu H, Zhang F. High expression of ACE2 on the keratinocytes reveals skin as a potential target for SARS-CoV-2. *J Invest Dermatol* 2020. <https://doi.org/10.1016/j.jid.2020.05.087>. [Epub ahead of print]
- 59 Li M, Li L, Zhang Y, Wang X. Expression of the SARS-CoV-2 cell receptor gene ACE2 in a wide variety of human tissues. *Infect Dis Poverty* 2020; **9**: 45.
- 60 To K, Tsang O, Leung W et al. Temporal profiles of viral load in posterior oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-2: an observational cohort study. *Lancet Infect Dis* 2020; **20**: 565–574.
- 61 Aurelian L, Ono F, Burnett J. Herpes simplex virus (HSV)-associated erythema multiforme (HAEM): a viral disease with an autoimmune component. *Dermatol Online J* 2003; **9**: 1.
- 62 Soler D, Humphreys T, Spinola S, Campbell J. CCR4 versus CCR10 in human cutaneous TH lymphocyte trafficking. *Blood* 2003; **101**: 1677–1682.
- 63 Imbalzano E, Casciaro M, Quartuccio S et al. Association between urticaria and virus infections: a systematic review. *Allergy Asthma Proc* 2016; **37**: 18–22.
- 64 Cao X. COVID-19: immunopathology and its implications for therapy. *Nat Rev Immunol* 2020; **20**: 269–270.
- 65 Landa N, Mendieta-Eckert M, Fonda-Pascual P, Aguirre T. Chilblain-like lesions on feet and hands during the COVID-19 Pandemic. *Int J Dermatol* 2020; **59**: 739–743.
- 66 Guan WJ, Ni ZY, Hu Y et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med* 2020; **382**:1 708–1720.
- 67 Hedou M, Carsuzaa F, Chary E, Hainaut E, Cazenave-Roblot F, Masson Regnault M. Comment on "Cutaneous manifestations in COVID-19: a first perspective" by Recalcati S. *J Eur Acad Dermatol Venereol* 2020; **34**: e299–e300. <https://doi.org/10.1111/jdv.16519>
- 68 Hon K, Leung A, Ng W, Loo S. Chronic urticaria: an overview of treatment and recent patents. *Recent Pat Inflamm Allergy Drug Discov* 2019; **13**: 27–37.
- 69 Introduction to "urticaria" special issue of dermatology bulletin. *Chin J Dermatol Venereol* 2020; **34**: 328.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Table S1. General information of the identified case reports.

Table S2. General information of the identified literatures related to ACE2 expression on skin.

Fig. S1. PRISMA diagram.