



Published in final edited form as:

Biogeosciences. 2018 July ; 15(14): 4515–4532. doi:10.5194/bg-15-4515-2018.

A multi-method autonomous assessment of primary productivity and export efficiency in the springtime North Atlantic

Nathan Briggs¹, Kristinn Guðmundsson², Ivona Cetini^{3,4}, Eric D'Asaro⁵, Eric Rehm⁶, Craig Lee⁵, Mary Jane Perry⁷

¹National Oceanography Centre, Southampton SO14 3ZH, UK ²Marine Research Institute, PO Box 1390, 121 Reykjavík, Iceland ³GESTAR/Universities Space Research Association, 7178 Columbia Gateway Drive, Columbia, MD 21046, USA ⁴Ocean Ecology Laboratory, NASA Goddard Space Flight Center Code 616, Greenbelt, MD 20771, USA ⁵Applied Physics Laboratory and School of Oceanography, University of Washington, Seattle, WA 98105, USA ⁶Département de Biologie et Québec-Océan, Université Laval, Québec, QC G1V 0A6, Québec, Canada ⁷Darling Marine Center, School of Marine Sciences, University of Maine, Walpole, ME 04573, USA

Abstract

Fixation of organic carbon by phytoplankton is the foundation of nearly all open-ocean ecosystems and a critical part of the global carbon cycle. But quantification and validation of ocean primary productivity at large scale remains a major challenge, due to limited coverage of ship-based measurements and the difficulty of validating diverse measurement techniques. Accurate primary productivity measurements from autonomous platforms would be highly desirable, due to much greater potential coverage. In pursuit of this goal we estimate gross primary productivity over two months in the springtime North Atlantic from an autonomous Lagrangian float using diel cycles of particulate organic carbon derived from optical beam attenuation. We test method precision and accuracy by comparison against entirely independent estimates from a locally parameterized model based on chlorophyll *a* and light measurements from the same float. During nutrient replete conditions (80% of the study period), we obtain strong relative agreement between the independent methods across an order of magnitude of productivities ($r^2=0.97$), with slight under-estimation by the diel cycles method ($-19\pm 5\%$). At the end of the diatom bloom, this relative difference increases to -58% for a six-day period, likely a response to SiO_4 limitation, which is not included in the model. In addition, we estimate gross oxygen productivity from O_2 diel cycles and find strong correlation with diel cycles-based gross primary productivity over the entire deployment, providing further qualitative support to both methods. Finally, simultaneous estimates of net community productivity, carbon export and particle size suggest that bloom growth is halted by a combination of reduced productivity due to SiO_4 limitation and increased export efficiency due to rapid aggregation. After the diatom bloom, high chlorophyll *a* normalized productivity indicates that low net growth during this period is due to increased heterotrophic respiration and not nutrient limitation. These findings represent a significant advance in the accuracy and completeness of upper ocean carbon cycle measurements from an autonomous platform.

1 Introduction

Measurement of ocean primary productivity (PP), the origin of nearly all organic carbon available to marine organisms, is essential to the study of marine ecosystems and predicting how they might respond to human activities. Because human influences such as climate change and fishing have global impact, improvements in global, mechanistic understanding of both the drivers of PP and its effects on ecosystems and their services should be of great value. However, progress is limited by the difficulty of measuring PP, which traditionally involves incubation experiments and/or radio- or stable isotope analysis, requiring cost, expertise, and ship availability. Understanding is further limited by the difficulty in validating PP, as each method has potential sources of bias, but generally no two methods measure the exact same quantity at the same temporal scale. Therefore, it is often unclear whether discrepancies between independent measurements are caused by biases or real differences. Satellite PP algorithms and global models can achieve the desired coverage, but these products still must be validated, ideally using an in situ dataset of confirmed accuracy that spans many years, in all seasons and in all oceans. Autonomous platforms can achieve such in situ coverage at a fraction of the cost of ship-based sampling, so the ability to estimate PP from an autonomous platform and validate these estimates using independent methods is highly desirable, both for directly enhancing understanding of ocean ecosystems and validating the models and satellite products that can approach true continuous global coverage.

Methods for estimating PP from diel cycles in particulate beam attenuation c_p (Siegel et al. 1989; Claustre et al. 1999; Cullen et al. 1992; Kinkade et al. 1999; J. Marra 2002; Dall'Olmo et al. 2011; Gernez et al., 2011; Omand et al., 2017; White et al. 2017) or O_2 (Caffrey 2003; Hamme et al. 2012; Nicholson et al. 2015) are suited for application to autonomous platforms, many of which already carry O_2 sensors and/or transmissometers. These methods rely on the light-dependence of PP, which causes a diel cycle in O_2 and in c_p (due to its correlation with particulate organic carbon, POC). However, other factors such as zooplankton vertical migrations, mixing events, O_2 air-sea flux, and POC/ c_p ratios may have diel cycles that introduce bias in these PP estimates, so they cannot be relied upon without validation. Comparisons so far between diel cycles and independent PP estimates have been encouraging, generally agreeing within a factor of two to three (Cullen et al. 1992; Walsh et al. 1995; Kinkade et al. 1999; Hamme et al. 2012; Nicholson et al. 2015), but the independent estimates have not been of the same quantity at the same temporal scale, so these comparisons do not provide strong constraints on the accuracy of this method.

In this study we take three significant steps towards the goal of enhancing our understanding of ocean ecosystems by increasing coverage of accurate in situ PP estimates using autonomous platforms. First, we use diel cycles in measurements of c_p and O_2 to simultaneously estimate two related quantities, gross primary productivity of particulate organic carbon (GPP) and gross oxygen productivity (GOP), in the surface mixed layer over a two-month period from an autonomous Lagrangian float. To our knowledge, this is the first time that c_p -based GPP and GOP have been simultaneously calculated using diel cycles from any platform, let alone autonomously. Second, we compare our diel cycles-based GPP

estimates with entirely independent estimates of the same quantity at the same spatial and temporal scale across a wide dynamic range of productivities. Again, to our knowledge, this represents the most rigorous validation of the diel cycles method to date. Third, we apply our mixed layer PP estimates, in conjunction with mixed layer O₂, NO₃, and POC budgets, to better understand how PP, heterotrophic respiration, and sinking flux all interact to regulate mixed-layer biomass in our study system: the spring diatom bloom in the Iceland Basin.

2 Methods

2.1 Study area and Platforms

The data presented here were collected by an autonomous Lagrangian mixed-layer float, two ships, and three autonomous Seagliders during the North Atlantic Bloom 2008 (NAB08) project. All data used here are available online at <http://www.bco-dmo.org/project/2098>. The float was deployed on April 4 in the Iceland Basin at 59°N, 20.5°W, near the 60°N site of the 1989 Joint Global Ocean Flux Study (JGOFS). The NAB08 project centered around the float, which was designed to drift in the surface mixed layer, mimicking the movement of plankton, except for daily profiles to 250 m (D'Asaro 2003). The float gathered data for two months, drifting northwest towards the Reykjanes Ridge, and ceased collecting data on May 25 at 61.8°N, 26.7°W (Fig. 1; black line) and was recovered on June 3. The timing of the daily float profiles was irregular until April 14, after which the float profiled each day between 15:00 GMT and dusk. The float carried an array of sensors, including two SBE-43-CTs for temperature and salinity, a WET Labs C-Star transmissometer for particulate organic carbon (POC), via particulate beam attenuation c_p , a WET Labs FLNTU (fluorescence and turbidity meter) for chlorophyll a fluorescence and POC, via particulate optical backscattering b_{bp} , a Seabird SBE-43 and an Aanderaa optode for oxygen, an ISUS (In Situ Ultraviolet Spectrophotometer) for NO₃, a LICOR LI-192SA for planar photosynthetically active radiation (PAR). See Table 1 for a list of abbreviations used in more than one subsection. Three cruises provided calibration data for the float's sensors as well as more detailed biological and chemical measurements: a deployment cruise by the R.V. *Bjarni Saemundsson* (April 3–5), a process cruise by the R.V. *Knorr* (May 2–21) and a float “rescue” cruise by the R.V. *Bjarni Saemundsson* (June 4–5). The ships collected both in situ measurements and discrete water samples via an overboard CTD package, which profiled to 600 m. Both ships carried the same array of in situ sensors as the float, minus the ISUS NO₃ sensor and the Aanderaa optode. In addition, the R.V. *Knorr* carried a second CTD and an above-water PAR sensor. Unlike the float, both of the ship's PAR sensors measured scalar PAR. The Seagliders were deployed together with the float and piloted to follow it throughout the experiment. Over the deployment, the distance between the float and individual gliders ranged from 175 km to < 1 km. However, at least one glider was within 50 km of the float for almost the entire deployment, and starting on May 6, all gliders remained within 50 km. Seagliders carried an array of sensors, but here we only discuss Seaglider estimates of sinking flux, derived in Briggs et al. (2011) using spikes caused by large particles in b_{bp} , measured by a WET Labs BB2F.

2.2 Discrete sampling

Discrete samples from all three cruises were analyzed at depths ranging from near surface (3–5 m) to 600 m for particulate organic carbon (POC; n=343), chlorophyll *a* (*Chl*; n=935), SiO₄ and NO₃ (n=1001), and phytoplankton pigments (n=80). Detailed methodology for these analyses can be found in the following technical report: http://data.bco-dmo.org/NAB08/Laboratory_analysis_report-NAB08.pdf. Briefly, *Chl* samples were filtered onto GFF 0.7 μ m filters and analyzed onboard using a Turner Designs Model 10-AU fluorometer. Following JGOFS protocols, POC samples were filtered onto pre-combusted GFF 0.7 μ m filters, sealed in foil packets and stored at –20 °C until analysis onshore using a Perkin Elmer 2400 CHN analyzer. For nutrients, 60 mL samples were immediately frozen and stored at –20°C until analysis onshore using a Lachat Quickchem 8000 Flow Injection Analysis System. In addition, discrete samples on the May process cruise were analyzed for dissolved oxygen concentration via Winkler titrations (n=131) and for bacterial counts and phytoplankton community composition using a FACScan flow cytometer and a FlowCAM automated microscopic imager. Phytoplankton particles were divided into several groups based on optical properties, size, and morphology as described in Cetini et al. (2012), with more detailed methods in a technical report accompanying the dataset: http://data.bco-dmo.org/NAB08/Phytoplankton_Carbon-NAB08.pdf.

2.3 Calibration of in situ sensors

The ship's profiler was held at constant depth for 60 s prior to closing each bottle to capture a water sample. In situ measurements from the 30 s prior to bottle closing were averaged to obtain a single value for matchups with discrete samples. Ship in situ sensors were calibrated via linear regression against discrete measurements. Float in situ sensors were calibrated using data from ten calibration casts, in which the ship was brought to the float's location and both ship and float profiled simultaneously. Float NO₃ and oxygen sensors were calibrated directly against the discrete measurements taken during the calibration casts. All other float sensors were calibrated against the matching ship in situ sensors, in order to maximize the number of matchups. Individual calibration details for each float sensor are listed below.

2.3.1 Temperature and Salinity—The duplicate temperature (T) and salinity (S) sensors aboard the ship's profiler during the May process cruise agreed closely (median S difference = 0.0018 and a median T difference = 0.0006 °C for each of 134 profiles). The ship TS sensors were therefore used as standards, after de-spiking and averaging (more details at http://data.bco-dmo.org/NAB08/Ship_TS_despiking-NAB08.pdf). Duplicate T sensors aboard the float also agreed closely and were therefore combined into a single record without adjusting to match the ship. After reconciliation of duplicate S measurements on each platform, a small mismatch between float and ship salinity was identified from the calibration casts and corrected by subtracting 0.0075 from the float S (more details at http://data.bco-dmo.org/NAB08/CTD_float_Calibration-NAB08.pdf).

2.3.2 Oxygen—Comparison between the SBE-43 and optode oxygen sensors aboard the float revealed differing sources of bias in each sensor. Bias in SBE-43 oxygen was introduced by changes in pumping rate during different modes of float operation and by

wave action near the surface. Bias in optode oxygen arose from its factory calibration, T and pressure effects, and a slower time response. After reconciliation of the two sensors to reduce these biases, the SBE-43 oxygen was brought in line with the discrete oxygen samples on the best six calibration casts by subtracting a constant offset of $0.9 \mu\text{Mol kg}^{-1}$. We conclude that the accuracy of the corrected in situ oxygen estimates is better than $2 \mu\text{Mol kg}^{-1}$, based on agreement with discrete samples (Winkler titrations). More details of the float's oxygen calibration can be found at http://data.bco-dmo.org/NAB08/Oxygen_Calibration-NAB08.pdf.

2.3.3 POC from optical beam attenuation—Previous work has shown that measurements of light scattering by particles, including beam attenuation c_p and particulate backscattering b_{bp} correlate strongly with POC in the open ocean (Cetinic et al. 2012 and references therein). Calibration of our c_p and b_{bp} measurements and conversion to POC estimates are described in the next two subsections. Raw output from the float optical beam transmissometer was aligned with raw ship transmissometer output using matchups from eight of the calibration casts. Agreement was very good ($r^2 = 0.99$), showing no evidence of sensor drift. Intercalibrated raw transmissometer output was converted to particulate optical beam attenuation c_p using the mean of factory calibrations performed on the ship's transmissometer before and after deployment. More details can be found at http://data.bco-dmo.org/NAB08/C-Star_Calibration-NAB08.pdf. We estimated c_p -derived POC (POC_{cp}) following Cetinic et al. (2012), but with a time-dependent adjustment in POC/c_p ratio to account for community changes. After subtracting the POC/c_p regression offset of 0.015 m^{-1} (Cetinic et al. 2012) from our c_p measurements, we computed the POC/c_p ratio for all ship POC and c_p samples where $c_p > 0.2 \text{ m}^{-1}$ in the upper 30 m during the May process cruise (Fig. 2; gray points). Samples whose T, S, c_p , and b_{bp} matched the float ML measurements within 0.25°C , salinity of 0.01, 0.1 m^{-1} , and 0.001 m^{-1} , respectively are shown as black circles (Fig. 2). Three inflection points were fit by eye at 370, 310 and 450 mg m^{-2} on May 6, 11, and 15, respectively. A continuous estimate of POC/c_p at the float patch was obtained by interpolating between these points and assuming constant POC/c_p before May 6 and after May 15 (Fig. 2, red line). This continuous estimate of POC/c_p was multiplied by float c_p (minus offset of 0.015 m^{-1}) to yield a c_p -based float POC estimate (POC_{cp}).

2.3.4 POC from optical backscattering—An average of pre- and post-deployment calibrations were used to convert raw backscattering output from both the float and the ship to the volume scattering function at the angle (140°) and wavelength (700 nm) of the sensors. The volume scattering function of seawater was then calculated following Zhang et al. (2009) and subtracted to yield scattering due to particles. The result was multiplied by $2\pi\chi$ to yield the particulate backscattering coefficient b_{bp} , where the angle-dependent scale factor χ is 1.132 for the FLNTU scattering sensors used in this study (M. Twardowski, pers. comm.). Float b_{bp} was aligned with ship b_{bp} using matchups from eight calibration casts ($r^2 = 0.96$). More details can be found at http://data.bco-dmo.org/NAB08/Backscatter_Calibration-NAB08.pdf. Glider b_{bp} was calibrated against the ship FLNTU in a similar fashion to the float (Briggs et al. 2011). We estimated b_{bp} -derived POC (POC_{bbp}) following Cetinic et al. (2012) via the equation $\text{POC}_{bbp} [\text{mg C m}^{-3}] = 37500 b_{bp} [\text{m}^{-1}] - 14$,

derived from a linear regression between co-located measurements of POC and b_{bp} within the mixed layer from the May process cruise.

2.3.5 Chlorophyll a—Raw chlorophyll fluorometer output from the ship was converted to an initial Chl estimate $Chl_{factory}$ using the factory calibrated scale factor and a dark offset derived from the minimum of all per-cast deep values (defined as the median between 550–580 m). An empirical fit between $Chl_{factory}$, T, PAR, and ship discrete Chl measurements was used to derive an in-situ Chl product (Eq. 1), which was strongly correlated with discrete Chl ($r^2 = 0.87$). Float $Chl_{factory}$ was aligned with ship

$$Chl = Chl_{factory} * \frac{2.1 * 10^{(T - 9.2) * 0.8} + 0.44}{10^{(T - 9.2) * 0.8} + 1} * \frac{(\log_{10}(PAR) * 0.05 + 1.02) * \tanh\left(\frac{PAR}{95} * 0.55\right)}{0.55 * \frac{PAR}{95}}, \quad (1)$$

$Chl_{factory}$ using the matchups from eight calibration casts ($r^2 = 0.95$), allowing calculation of Chl via Eq. (1), at the float as well. The FLNTU sensor was located at the bottom of the float, facing down, so Chl data was removed whenever the float was moving upward at >1.7 cm s^{-1} , due to possible entrainment of deeper water. Chl measurements where $PAR > 75$ were also removed to eliminate non-photochemical quenching. In order to obtain a continuous, depth-resolved record of Chl for calculation of primary productivity, the remaining Chl estimates, from both mixed-layer mode and profiles, were filtered using a 5-point running median, averaged in one-hour, one-meter bins, and then interpolated in depth and time via triangulation-based 2-D linear interpolation, with distance calculated as $\sqrt{(dz[m]/30[m])^2 + dt[days]^2}$ (i.e. a 30 m vertical interval and a 1 day time interval were considered equidistant).

2.3.6 Nitrate—A post-deployment laboratory calibration, including temperature and salinity corrections, was used to obtain initial NO_3 estimates from the float's ISUS NO_3 sensor. An additional scale factor of 1.15 and offset of $+2.6 \mu\text{M}$ were required to bring these initial estimates in line with discrete samples taken during calibration casts. More details can be found in Alkire et al. (2012) and at http://data.bco-dmo.org/NAB08/ISUS_Nitrate_Calibration-NAB08.pdf.

2.3.7 Silicate— SiO_4 was not measured by the float, but discrete shipboard SiO_4 measurements from the top 15 m were considered to represent mixed-layer SiO_4 at the float location if the corresponding temperature, salinity and NO_3 measurements matched concurrent float ML measurements to within 0.25°C , salinity of 0.01, and 0.8 mmol m^{-3} respectively.

2.3.8 PAR—The factory calibration of the float PAR sensor was used “as is.”

2.4 Mixed layer depth

Mixed layer depth (MLD) was calculated at hourly intervals from float potential density anomaly estimates via the following steps: 1) Smooth density timeseries using a 5-point running median. 2) Average density into one-hour, one-meter bins. 3) Fill in the gaps with 2D linear interpolation, such that a 30 m vertical interval and a one-day time interval are considered equidistant. 4) For each hour, find the minimum potential density anomaly. 5) The MLD for each hour is defined as the shallowest depth where the potential density anomaly exceeds this minimum by 0.01 kg m^{-3} . The MLD was calculated twice, once excluding data when downward velocity exceeded 1 m min^{-1} and once excluding upward velocity exceeding 1 m min^{-1} . We use the average of these two estimates as the final MLD estimate, reducing the influence of single active profiles, which could differ from mean conditions due to entrainment or internal waves. When the float was close to neutral buoyancy, this MLD(t) estimate followed the lower limit of the vertical movement of the float during its ML mode. However, during periods of positive buoyancy, MLD(t) occasionally exceeded the maximum depth of the float during its ML mode (Fig. 3). Hourly MLD depth was also calculated using the same criteria from the output of the Bagniewski et al. (2011) data-assimilation model (see red line in Fig. 3) to permit testing of diel cycles methods within the model itself.

2.5 K_{PAR}

2.5.1 Instantaneous K_{PAR} estimates—The diffuse attenuation coefficient of PAR K_{PAR} was calculated from each pair of consecutive PAR measurements made at times t_1 and t_2 via Eq. (2), where z is depth, \bar{z} is the mean of $z(t_2)$ and $z(t_1)$, and \bar{t} is the mean of t_2 and t_1 .

$$K_{PAR(measured)}(\bar{z}, \bar{t}) = \frac{\ln(PAR(t_1)) - \ln(PAR(t_2))}{z(t_2) - z(t_1)} \quad (2)$$

2.5.2 K_{PAR} fit method—The uncertainty of individual $K_{PAR(measured)}$ estimates was high and depended strongly on dz , which ranged from 0.2 – 30 m with a mean of 1.3 m. These 14000 $K_{PAR(measured)}$ estimates were therefore fit to Chl and z using a non-linear least-squares multiple regression weighted by dz to obtain Eq. (3):

$$K_{PAR(modeled)}(Chl, z) = 0.064 * Chl^{0.51} + 0.20 * \max(z, 2.5)^{-0.63} + 0.0031. \quad (3)$$

In order to evaluate the performance of this fit, $K_{PAR(measured)}$ precision was increased by eliminating estimates with $dz < 2$ m and combining the remaining estimates into 21-point medians, yielding a total of 118 independent in-situ K_{PAR} estimates. A type- II linear regression of these estimates against 21-point medians of K_{PAR} estimated via Eq. (3) yielded an r^2 of 0.85, a root mean square error of 0.014 m^{-1} , and a mean bias of -0.004 m^{-1} . The residual error was not significantly correlated with depth, time, solar zenith angle or the ratio of in situ Chl to b_{bp} , a proxy for plankton community in this system (Cetini et al. 2015).

2.6 Depth-resolved PAR

In order to calculate PAR at all depths, PAR was extrapolated from a reference depth z_{ref} via Eq. (4):

$$PAR_{extrapolated}(z) = PAR(z_{ref}) * \exp\left(\int_z^{z_{ref}} K_{PAR} dz\right), \quad (4)$$

using $K_{PAR(modelled)}$ calculated via Eq. (3) from the float's continuous *Chl*. When the float was within the top 50 m, z_{ref} was the depth of the float and $PAR(z_{ref})$ was the float's PAR measurement. The performance of this extrapolation was evaluated by comparing $PAR_{extrapolated}(0-)$ (just below the surface) calculated via Eq. (4) with scalar $PAR(0+)$ measured by the ship's underway system. For all measurements where the ship was within 1 km of the float, the float was in the top 50 m, and $PAR(0+)$ was greater than $1 \mu \text{ mol m}^{-2} \text{ s}^{-1}$, $PAR(0+)$ and $PAR_{extrapolated}(0-)$ were highly correlated ($r^2 = 0.96$ on a linear scale and $r^2 = 0.99$ on a logarithmic scale). The geometric mean of the ratio of $PAR_{extrapolated}(0-)$ to $PAR(0+)$ was 0.92 and the geometric (multiplicative) standard deviation was a factor of 1.19. For several hours each afternoon, while the float profiled to 250 m, float PAR measurements were not available, so $PAR(0-)$ was estimated using an empirical function of solar zenith angle and an empirical index of cloud cover. First, a double exponential was fit to 36000 PAR measurements obtained in the top 1 m over a range of solar zenith angle θ from -6° to 90° by a global network of 100 "Biogeochemical Argo" type profiling floats to obtain $PAR_{modelled}(0-)$, an estimate of $PAR(0-)$ under mean cloud and atmospheric conditions:

$$\log_{10}(PAR_{modeled}(\theta)) = 2.5 * \exp(0.0030 * \theta) - 1.7 * \exp(-0.10 * \theta) \quad (5)$$

To adjust for clouds, $PAR_{extrapolated}(0-)$ from the Lagrangian float (via Eq. 4) was divided by corresponding estimates $PAR_{modelled}(0-)$ from to obtain an index of sunniness, which was averaged into 15 min bins to remove noise from wave focusing. This sunniness index ranged from 0.1 to 3.6 over the entire float deployment. Sunniness index at time t was estimated using a ± 1 -day running mean of these sunniness index estimates, weighted by the inverse square of $t-t_i$, where t_i is the time of each measurement. This running mean sunniness index was then multiplied by $PAR_{modelled}(0-)$ to obtain $PAR_{adjusted}(0-)$, which was used as $PAR(z_{ref})$ in Eq. (4) during the afternoon gaps.

2.7 O₂ air-sea flux

O₂ air-sea flux was calculated following Alkire et al. (2012). Briefly, wind speeds were taken from the NCEP WW3 Global Reanalysis product, except during the May cruise, when ship wind measurements were used. O₂ saturation was calculated following García and Gordon (1992). Air-sea flux was calculated following (Wanninkhof 1992), modified to account for bubble injection following Woolf and Thorpe (1991). Hourly dO_2/dt in the ML due to air-sea flux was estimated by dividing hourly flux estimates by hourly MLD.

2.8 Primary Productivity estimates

2.8.1 Diel cycles of O₂ and POC—“Typical” diel cycles (minimum near dawn and maximum near dusk) were observed in mixed layer records of O₂ (Fig. 4), consistent with previous studies (Caffrey 2003; Hamme et al. 2012; Nicholson et al. 2015). We estimated mixed layer gross oxygen productivity (GOP) at half-day intervals from these diel cycles. To estimate morning GOP, ML O₂ concentrations were smoothed with a 3-point running median and a type I linear regression (O₂ vs time) was fit to data from dusk to dawn (Fig. 4a; solid black line). The regression fit was projected forward to provide an estimate of noontime mixed-layer O₂ in the absence of GOP. Measured noontime O₂ was calculated from a type I linear regression of O₂ data taken within 1 h of local noon. Morning mixed layer GOP was calculated as the difference between measured and projected concentration (Fig. 4; blue vertical bar) and divided by 0.5 d to convert to units of mmol m⁻³ d⁻¹. Afternoon GOP was calculated in a similar fashion, by subtracting noontime mixed-layer O₂ from the noontime extrapolation of a linear fit of the following night’s data. Similar diel cycles were observed in mixed layer POC_{cp}, and the same method was used to calculate mixed layer gross primary productivity of POC (GPP_{cp}) from these cycles (Fig. 4b). Diel cycles in POC_{bbp} were less regular and usually out of phase with O₂ and POC_{cp} cycles, but GPP_{bbp} was calculated in the same way as GOP and GPP_{cp} for comparison. Note that this diel cycles method assumes homogeneous mixing to a constant depth and that any gain or loss terms other than GOP (or GPP) are constant day-to-night over the period of a single calculation (~18 h). However, we find a clear diel cycle in MLD (Fig. 3), which amplifies the diel cycle in O₂ (and *c_p* and *b_{bp}*), causing PP calculated from diel cycles to exceed mean PP within the daily mean MLD. This is because night-time ML deepening enhances loss of ML concentration relative to daytime mixing losses. Analysis of the output of a coupled physical-biological model assimilating data from the Lagrangian float (Bagniewski et al. 2011), which accurately reproduced the diel cycle in mixing (Fig. 3, black line) shows that the mixing-amplified diel cycles of O₂ in the ML yield daily GOP estimates that correspond approximately to the mean GOP above the daily *minimum* MLD. Regression of diel GOP, calculated from ML O₂ timeseries output by the model, as a function of “true” model GOP, forced through zero, yields a slope ±95 % confidence interval of 0.91±0.12 and a RMSE of 0.12 mmol m⁻³ d⁻¹. We therefore interpret our daily GOP and GPP_{cp} estimates as representing daily mean productivity between the surface and daily minimum MLD. Bias in GOP due to day-night differences in air-sea flux was also estimated using the difference between mean morning (or afternoon) dO₂/dt due to air-sea flux and that of the previous (or next) nighttime. Mean bias was small (<5% of GOP), and linked primarily to the MLD diel cycle, so a separate correction was not deemed necessary. Other potential biases are discussed in sections 4.1.2 to 4.1.4.

2.8.2 ¹⁴C incubations—During the April and May cruises, daily two-hour ¹⁴C incubation experiments were conducted (n=28) to estimate photosynthetic parameters. Each day, a water sample was taken from the *Chl* maximum, as determined by in situ fluorescence, and duplicate 2 h ¹⁴C incubations were carried out at 7 different PAR levels ranging from 0–400 μ mol m⁻² s⁻¹. The dark incubation ¹⁴C activities were weakly but significantly correlated with *Chl* (type II linear regression; r² = 0.19; p<0.05; apparent NPP = *Chl* * 0.036±0.026 mg C mg Chl⁻¹ h⁻¹ + 0.049±0.035 mg C m⁻³ h⁻¹); dark activities were

treated as sample-specific blanks and subtracted from the light incubation activities of the corresponding water sample. The resulting productivity estimates were interpreted as net primary productivity (NPP), based on findings that most phytoplankton do not respire old carbon when newly fixed carbon is available (John Marra and Barber 2004; Pei and Laws 2013). Note that if, contrary to our assumptions, phytoplankton did respire old carbon at all light levels during these incubations, then our calculations below overestimate NPP and underestimate phytoplankton respiration (R_{ϕ}), but GPP is unbiased. On the other hand, if old carbon is respired only in the low light incubations, then we under-estimate R_{ϕ} and GPP, but little bias is introduced in NPP. Seven of the 350 individual NPP estimates (all from the April cruise) were judged to be positive outliers and were manually removed before further analysis. In ~60% of the incubation experiments, NPP decreased with increasing PAR for $PAR > 200 \mu \text{mol quanta m}^{-2}\text{s}^{-1}$. We conclude that this apparent photoinhibition is likely not representative of most field conditions, because in situ measurements of *Chl*-normalized dO_2/dt showed no consistent relationship with PAR between PAR values of 100 and 1000 $\mu \text{mol m}^{-2}\text{s}^{-1}$. We therefore removed values of NPP where $PAR > 200 \mu \text{mol m}^{-2}\text{s}^{-1}$ if they were lower than the second-highest NPP observed where $PAR < 200 \mu \text{mol m}^{-2}\text{s}^{-1}$ (56 of 110 high light points removed). Remaining NPP vs PAR data were fit to an empirical “PvE” model represented in Eqs. (6–8):

$$\lambda = PAR \frac{\alpha}{P_M} \epsilon, \quad (6)$$

$$GPP = P_M \left(1 + \frac{1}{\epsilon} \sum_{i=0}^{\epsilon-1} e^{-\lambda} \frac{\lambda^i}{i!} - \sum_{i=0}^{\epsilon-1} e^{-\lambda} \frac{\lambda^i}{i!} \right), \quad (7)$$

$$NPP = GPP - R_{\phi}, \quad (8)$$

based on four parameters: maximum GPP (P_M), the initial slope of GPP/PAR (α), R_{ϕ} , and an efficiency factor (ϵ) representing “sharpness” of the transition between light-limited and light-saturated photosynthesis. This parameterization is based on a conceptual model of photosynthesis in which there is a rate-limiting step that can receive and “store” up to ϵ “packets” of energy at once at above the limiting rate without wasting any of these packets. We used a single ϵ value of six, which provided the best overall least squared fit across all incubation experiments. This ϵ value yields a NPP vs PAR relationship that is “sharper” than the commonly used “tanh” model (Harrison and Platt 1986) and more linear at low PAR, leading to smaller y- offset (smaller R_{ϕ} estimate). See Fig. 5 for example fits. A power law was then fit between in situ *Chl* estimates from the ship’s profiling package (calculated via Eq. 1) and each of the three parameters obtained from each NPP vs PAR fit (P_M ; Fig. 6a; α ; Fig. 6b; and R_{ϕ} ; Fig. 6c). Fits with P_M and α used data from all cruises, but the fit with R_{ϕ} included only data from the process cruise (Fig. 6c; solid circles), as signals were too low to constrain R_{ϕ} in April and R_{ϕ} appeared consistently higher during the June cruise, possibly due to higher temperature.

2.8.3 Chl-based GPP and NPP—The relationships in Fig. 6 were used to estimate photosynthetic parameters $P_M(t,z)$, $\alpha(t,z)$, and $R_\phi(t,z)$ and their uncertainty intervals at the float location from $Chl(t,z)$ (Section 2.3.5). We estimated gross primary productivity $GPP_{Chl}(t,z)$ and net primary productivity $NPP_{Chl}(t,z)$ via Eqs. (6–8) using the above photosynthetic parameters, $PAR_{extrapolated}(t,z)$ (Section 2.6), and $\epsilon=6$ as input. Uncertainties were propagated from $P_M(t,z)$, $\alpha(t,z)$, and $R_\phi(t,z)$ using the conservative assumption that they covary (i.e. upper bound of NPP_{Chl} was derived from upper bounds of P_M and α and lower bound of R_ϕ).

2.9. Area-weighted mean particle diameter

Area-weighted mean particle diameter D_{bbp} 10–50 m depth bin was estimated following Briggs et al. (2013) via Eqs. (9–11):

$$D_{bbp} = 2\sqrt{\frac{Var[b_{bp}(t)]}{E[b_{bp}(t)]} \frac{v}{Q_{bb}} \frac{1}{\gamma(\tau)} \frac{1}{\pi}}, \quad (9)$$

$$\gamma(\tau) = \begin{cases} 1 - (3\tau)^{-1}, & \text{if } \tau \geq 1 \\ \tau - \tau^2/3, & \text{if } \tau \leq 1 \end{cases}, \quad (10)$$

$$\tau = \left(\frac{t_{res}}{t_{samp}} \right), \quad (11)$$

where $Var[b_{bp}(t)]$ is the variance in b_{bp} due to random distribution of particles in space, $E[b_{bp}(t)]$ is mean b_{bp} , V is sensor sample volume, Q_{bb} is the backscattering efficiency, and γ and τ are functions of residence time in the sample volume t_{res} and sample integration time t_{samp} . $Var[b_{bp}(t)]$ and $E[b_{bp}(t)]$ were calculated once per profile (ascent or descent) using all data between 10–50 m. Prior to calculation of $Var[b_{bp}(t)]$, the b_{bp} timeseries was de-trended by subtracting a 7-point running median and large outliers (greater than five times the interquartile range) were removed before the variance was calculated on the residuals. A V of 0.62 ml was used (Briggs et al. 2013) and a Q_{bb} of 0.02 was assumed (based on empirical b_{bp}/c_p ratio of ~0.01 and theoretical value of $Q_c=2$ for diameter \gg wavelength; Bohren and Huffman, 1983). A t_{res} of 0.02 s was chosen based on a 6 mm path through the sample volume and a platform velocity of 30 cm s⁻¹ and t_{samp} was 1 s.

2.10 Sinking POC flux

POC_{bbp} profiles from both gliders and the float were divided into a “small” particle baseline (7-point running minimum followed by running maximum) and a “large” particle “spike” signal (residuals above the baseline). This approach, developed by Briggs et al. (2011), is based on the finding that large, fast sinking particles, owing to their rarity and light-scattering characteristics, can create individual large spikes in mesopelagic bbp clearly distinguishable from background concentrations (Briggs et al. 2011). Large particle POC_{bbp} was multiplied by a bulk sinking speed of 75 m d⁻¹ to estimate large POC flux (Briggs et al.

2011). A broad plausible range of bulk sinking speeds $5 \pm 5 \text{ m d}^{-1}$ was used to estimate small POC sinking flux, which was added to large POC flux to yield total sinking POC flux. Sinking POC flux was bin averaged in 50 m vertical bins and either running 2-day bins (gliders) or longer discrete bins to match bloom stages (float).

3 Results

3.1 Evolution of the spring bloom

From float deployment through April 17, MLD was variable (often $>200 \text{ m}$ but occasionally $< 50 \text{ m}$; Fig. 3), mixed-layer nutrients were high ($\text{NO}_3 \approx 12 \text{ mmol m}^{-3}$; $\text{SiO}_4 \approx 4 \text{ mmol m}^{-3}$), biomass was low ($\text{Chl} \approx 0.35 \text{ mg m}^{-3}$; $\text{POC}_{\text{cp}} \approx 35 \text{ mg m}^{-3}$), and O_2 was undersaturated by $\sim 10 \text{ mmol m}^{-3}$ (Fig. 7). Mixed-layer biomass concentrations increased over the next month, peaking in mid-May. This broad increase was punctuated by several 1–2 day periods of decrease, most associated with clear mixed-layer deepening (Fig. 7). SiO_4 was depleted to its lowest level on May 11, Chl concentration peaked on May 12, and NO_3 depletion and POC_{cp} and O_2 concentrations peaked on May 13. From bloom peak to May 16, Chl decreased dramatically (77%), POC_{cp} and O_2 decreased moderately (by 9 and 13 mmol m^{-3} , respectively), and NO_3 and SiO_4 concentrations recovered slightly (by 0.8 and 0.4 mmol m^{-3} , respectively).

3.2 Primary productivity estimates

All GPP and GOP estimates were averaged into 3-day bins to improve precision of the diel-cycles-based estimates. To first order, GPP_{Chl} followed Chl , low in early April ($0.5\text{--}1.0 \text{ mmol m}^{-3} \text{ d}^{-1}$), peaking near $10 \text{ mmol m}^{-3} \text{ d}^{-1}$ between May 7 and 13, then decreasing to near $3 \text{ mmol m}^{-3} \text{ d}^{-1}$ or below after the bloom (Fig. 8). But increases in GPP_{Chl} led increases in Chl by 1–2 days during ML shoaling (and high growth) events on April 24–27 and May 6–8 (Fig. 8; pale vs. dark green) due to higher ML averaged PAR (not shown). For the entire “bloom growth” phase from early April through May 9, GPP_{Chl} was strongly correlated with both cycle-based estimates of both GOP (Fig. 8 and Fig. 9b; blue) and GPP_{cp} (Fig. 8 and Fig. 9c; blue). GOP was a factor of 2.1 higher than GPP_{Chl} on a molar basis, while GPP_{cp} was slightly lower (factor of 0.81). GPP_{bbp} was poorly correlated with GPP_{Chl} and significantly lower (by 60%; Fig. 9d). From noon May 10 to noon May 11, diel cycles could not be calculated, because the float was trapped at the surface, due to high stratification and slight positive buoyancy. At peak biomass (May 11–13), and the bloom decline (May 13–16), both $\text{GOP}/\text{GPP}_{\text{Chl}}$ and $\text{GPP}_{\text{cp}}/\text{GPP}_{\text{Chl}}$ were substantially lower than during bloom growth (Fig. 8, pink highlighted region, and Fig. 9; pink symbols). In the post-bloom period (May 16–24), $\text{GOP}/\text{GPP}_{\text{Chl}}$ and $\text{GPP}_{\text{cp}}/\text{GPP}_{\text{Chl}}$ increased again, similar to the bloom growth ratios (Fig. 8 and Fig. 9; red symbols). When all bloom phases are combined, best-fit ratios of GOP and GPP_{cp} to GPP_{Chl} are 1.7 and 0.6, respectively and correlations are considerably less strong (r^2 of 0.67 and 0.49, respectively). However, the estimates of productivity from diel cycles (GOP and GPP_{cp}) remained strongly correlated for the entire deployment. Over the entire study period, morning estimates of GOP and GPP_{cp} were not significantly different from the afternoon estimates, while morning GPP_{bbp} estimates were significantly lower than afternoon estimates (80% lower overall). However, morning-afternoon patterns appear to change starting on May 13, when the bloom decline starts (e.g.

Fig. 4). From May 13–24, there is no significant difference between morning and afternoon GPP_{bbp} , but afternoon estimates of GPP_{cp} and GOP , were lower than morning estimates by 70% and 43%, respectively. These differences were near the threshold of statistical significance: mean afternoon-morning difference ± 2 standard errors was $-2.3 \pm 2.3 \text{ mmol m}^{-3} \text{ d}^{-1}$ for GPP_{cp} and $-3.0 \pm 2.5 \text{ mmol m}^{-3} \text{ d}^{-1}$ for GOP .

3.3 Depth integrated GPP, NPP, and NCP and carbon export

Alkire et al. (2012) estimated net community productivity (NCP) integrated within the top 50–60 m and carbon export from 50–60 m at the float location for four periods of stable stratification: the “early bloom” (April 23–27), “main bloom” (May 6–13), “decline” (May 13–14) and “post bloom” (May 20–24). We integrated GPP_{Chl} and NPP_{Chl} to the same depth and time ranges in order to assemble detailed organic carbon budgets for these periods (Fig. 10). Each budget term carries considerable uncertainty, but based on the central estimates, the partitioning of fixed carbon appeared to change substantially over the course of the bloom. Note that these NCP estimates include net production of dissolved organic carbon (DOC), while NPP_{Chl} excludes any photosynthetic DOC production. NPP_{Chl} and NCP estimates were similar during the early and main bloom, suggesting moderate-to low heterotrophic respiration. During the early bloom period, export was also low (~22–28% of GPP_{Chl}), allowing rapid accumulation of biomass. During the main bloom, GPP_{Chl} nearly doubled as biomass increased, but a larger fraction (~50%) was exported, leaving ~25% to accumulate. During the bloom decline, apparent community respiration (defined as difference between GPP_{Chl} and NCP) was 156% of GPP_{Chl} and export was an additional 50–80%. In the post-bloom period, community respiration was again high (~100% of GPP), and export was much lower (0–15% of GPP). Our NPP_{Chl} estimates and b_{bp} “spike”-based sinking flux estimates provide a continuous, high-resolution picture of the link between productivity and export at 125 m for the entire study period (Fig. 11a). Float and glider-based POC export estimates agree broadly at this depth (red lines), suggesting that the higher-resolution glider timeseries are representative of the float patch as well. While export at 125 m is coupled with NPP_{Chl} (Fig. 11a), there is a rapid increase in export efficiency between May 3–6 from ~20% to 40%. Area-weighted mean particle diameter (D_{bbp}) ranged from 90–150 μm during April, peaked at 250 μm on May 7–8 (Fig. 11b), coincident with peak biomass as measured by both Chl and POC_{bbp} from the gliders (not shown). D_{bbp} fell rapidly on May 9, coincident with a ML deepening event. Post-bloom D_{bbp} ranged from 150–190 μm (Fig. 11b).

4 Discussion

4.1 Accuracy of PP estimates

The combination of three estimates of primary productivity and one estimate of community productivity, all from the same platform at comparable temporal and horizontal scales, provides a unique opportunity to evaluate the accuracy of all methods. Each of our PP methods is discussed in turn in sections 4.1.1–4.1.4.

4.1.1 GPP_{Chl} — GPP_{Chl} and GPP_{cp} are estimates of the same quantity, obtained independently. GPP_{Chl} is derived from PAR and Chl estimates using robust local

parameterizations obtained from ^{14}C incubations. GPP_{cp} is derived entirely from c_p measurements, converted to POC using another robust, local empirical relationship. The averaging depth (daily minimum MLD) for GPP_{chl} was chosen to match the diel cycles method based on results of a model tuned to match local conditions (Bagniewski et al. 2011). In this context, the combination of strong correlation and absolute agreement between GPP_{chl} and GPP_{cp} (Fig. 9c; within 19%) provides confidence in both methods during the bloom growth and post bloom periods. The $\text{GOP}/\text{GPP}_{\text{chl}}$ slope of 2.1 (Fig. 9b) is at the upper end of the expected range, providing additional first-order support for GPP_{chl} accuracy. Neither GPP method includes DOC production, so the range of expected photosynthetic quotients ($\sim 1\text{--}1.45$; Laws 1991; Robertson et al. 1993), combined with the fraction of GPP released as DOC in marine/estuarine environments (2–50%; Baines and Pace, 1991) imply a possible GOP/GPP range of 1–2.9. During the main bloom observed by the float in this study, Alkire et al. (2012) estimate that DOC accounts for 22–40% of NCP in the mixed layer during the main bloom. If these estimates apply to GPP as well, our expected GOP/GPP range narrows to 1.3–2.4 (Fig. 9a,b; gray dashed lines). Thus, our GOP estimates suggest either that both the photosynthetic quotient and phytoplankton DOC production are high during bloom growth (and GPP is accurate) or that both GPP estimates are biased low. As mentioned in section 2.8.2, a negative bias in $\text{GPP}_{\text{model}}$ could be explained if phytoplankton respired substantial old, unlabelled carbon in our low light incubations, but not in the high light incubations. In this case a separate explanation (see next section) is needed for the high $\text{GOP}/\text{GPP}_{\text{cp}}$ slope of 2.6 (Fig. 9a).

During bloom peak and decline, the strong discrepancies between GPP_{cp} and GPP_{chl} imply either an underestimate by GPP_{cp} (discussed in the next section) or an over-estimate by GPP_{chl} . If diatoms reduce GPP in response to sustained SiO_4 limitation then we expect GPP_{chl} over-estimation at peak biomass, given that GPP_{chl} is only a function of Chl and PAR, without any nutrient limitation term. Twelve mixed-layer SiO_4 samples were collected in the vicinity of the float on May 11–13, and the mean and maximum measured concentrations were 0.3, and 0.6 mmol m^{-3} , respectively, suggesting that diatom growth was most likely severely limited (Fig. 7a). This does not necessarily imply that diatom carbon fixation rates were reduced, but previous studies have indeed observed a large and reversible reduction in apparent diatom photosynthetic efficiency under multi-day SiO_4 limitation (Lippemeier, Hartig, and Colijn 1999; Lippemeier et al. 2001). Both FlowCAM microscopy and HPLC pigments indicate that diatoms accounted for $\sim 50\%$ of phytoplankton biomass at bloom peak (Cetini et al. 2015), so we expect a substantial reduction in bulk phytoplankton growth (and likely GPP) under these conditions. This expectation, combined with the observed reduction in both GPP_{cp} and GOP at bloom peak, leads us to conclude that GPP_{chl} is most likely over-estimated at bloom peak. This conclusion agrees with the coupled physical-biological model of Bagniewski et al. (2011), which assimilated float biogeochemical measurements and achieved optimal fit when diatom GPP was limited by SiO_4 with a half-saturation constant of 1 $\mu\text{mol m}^{-3}$. GPP inferred from this model closely matches our observed GPP_{cp} during SiO_4 limitation (Fig. 8, gray line vs black circles), even though Bagniewski et al. (2011) assimilated daily binned data, removing any diel cycle information. On the other hand, three ^{14}C incubations were conducted between May 10–14 using water with $\text{SiO}_4 < 0.5 \text{ mmol m}^{-3}$ although not at the float location, and there was not a

substantial reduction in measured P . These samples may not be representative of the water sampled by the float, despite similar Chl and SiO_4 concentrations, or it is possible that a bottle effect enhanced GPP. But we cannot rule out the alternative hypothesis that the Si-limited community continued to fix carbon at a constant rate, and that GPP_{cp} and GOP estimates were reduced for another reason (discussed in next sections).

Apart from Si limitation, possible explanations for GPP_{Chl} over-estimation include over-estimation of Chl due to increased fluorescence, underestimation of the MLD, or photoinhibition. Again, Chl was calculated the same way for the floats and ship, so if the in-situ fluorometric method over-estimated Chl at bloom peak, we would expect to see a deviation from the observed relationships between photosynthetic parameters and Chl (Fig. 6). So this explanation, while plausible given the high Chl/b_{bp} ratio at bloom peak (Cetini et al. 2015), also requires that none of our low SiO_4 bottle samples were representative of the bloom peak at the float location. The density-based MLD estimates appear quite robust during this period, consistently shallow and stable at 10–20 m and matched by the vertical motion of the float. And the daytime increases in O_2 (Fig. 4) and c_p on May 11 and 12 show no sign of photoinhibition, despite a peak hourly-averaged PAR of $>750 \mu\text{mol m}^{-2}\text{s}^{-1}$; increases are smooth throughout the morning and appear to continue at the same rate in the afternoon (Fig. 4). This result supports our decision to exclude photoinhibited bottle incubation data from our productivity vs. PAR fits, and we suggest that photoinhibition terms from bottle incubations should be applied with caution, if at all, in future studies. In this system, photoinhibition is likely reduced during deeper mixing events due to the shorter time phytoplankton is exposed to high light. During the stratified conditions, reduction in photoinhibition can be attributed to photoadaptation.

SiO_4 limitation may also explain some of the discrepancy between GPP_{cp} and GPP_{Chl} during the bloom decline (May 13–16), at least when afternoon estimates are excluded (see Fig. 9b,c; open pink circles). Lower afternoon GPP_{cp} and GOP, combined with very shallow (<5 m) MLDs at noon on May 13 and 15, also raise the possibility of significant photodamage inhibiting afternoon productivity. Mean ML PAR exceeded $500 \mu\text{mol m}^{-2}\text{s}^{-1}$ for at least two hours on both days. However, the negative afternoon GPP_{cp} estimates at this time suggest a bias in the diel cycles method as well (see next section).

4.1.2 GPP_{cp} —Potential sources of bias unique to GPP_{cp} include a diel cycle in grazing (e.g. due to diel migration of zooplankton), a diel cycle in export loss, or a diel cycle in the POC/ c_p ratio. The tight correlations between GPP_{cp} and GOP throughout the entire study period ($r^2=0.95$; Fig. 9a) and between GPP_{cp} and GPP_{Chl} during bloom growth ($r^2=0.96$; Fig. 9b) provide encouraging support for GPP_{cp} as a measure of *relative* primary productivity at the very least. Furthermore, the quantitative agreement between GPP_{cp} and GPP_{Chl} during both bloom growth and post-bloom (Fig. 9c; slope: 0.82 ± 0.06) is very close to our expected slope of 0.93 from model results (reanalysis of Bagniewski et al. 2011), suggesting that GPP_{cp} accuracy is comparable to other methods across most of the conditions encountered. These findings agree closely with those of White et al. (2017), who find a GPP_{cp}/NPP ratio of 1.1 across a factor of three dynamic range of productivities in the subtropical North Pacific, suggesting that GPP_{cp} is either accurate, or slightly under-estimates true GPP. Taken together, our results are highly encouraging regarding the

widespread applicability and accuracy of the GPP_{cp} method. However, it should be noted that our results still may not apply to certain other systems, where different phytoplankton size and/or timing of cell division could alter the diel POC/ c_p relationship (Dall'Olmo, et al. 2011).

If the SiO_4 limitation hypothesis is correct, then GPP_{cp} during the bloom peak and morning GPP_{cp} during the bloom decline may be accurate as well. On the other hand, if GPP_{chl} is accurate during this time, then GPP_{cp} is biased low by ~50% at this time. It is unclear what might cause such a low bias, especially at bloom peak. Between the afternoon of May 11 and the morning of May 13, there is no anomaly in the diel cycles of POC $_{cp}$ or O_2 (Fig. 10) indicative of daytime mixing, advection, or possible photoinhibition, and there is no change in the relationship between GPP_{cp} and GOP (Fig. 9a; rightmost pink symbol). Without grazing data, we cannot rule out enhanced daytime grazing as a possible explanation, although grazing is generally expected to be higher at night. Alternatively, particularly high photo-oxidation could potentially dampen O_2 diel cycles during this period and perhaps c_p diel cycles as well. This hypothesis is supported by laboratory measurements of diatom productivity under nutrient limitation (Spilling et al. 2015), although again we would need to explain why reduced P_M was not observed in our bottle incubations. On the other hand, the afternoon GPP_{cp} estimates during the bloom decline period show a clear example of negative bias in the diel cycles method. On May 13, 14, and 15 (bloom decline), rates of net POC $_{cp}$ (and O_2) accumulation are positive or near zero in the morning but negative each afternoon (e.g. Fig. 4; May 13). One plausible explanation is horizontal advection of the float relative to the ML during its afternoon profile, causing it to resurface in water with lower biomass. During this period, comparison with ship, autonomous glider, and satellite measurements (Alkire et al. 2012) shows that the float was at the edge of a high biomass (and O_2) patch, so advection during this time would most likely cause loss in POC $_{cp}$ and O_2 . Note that the afternoon reductions in GPP_{cp} during bloom decline are greater than the afternoon reductions in GOP (e.g. Fig. 4). This result is possible with the horizontal advection/mixing hypothesis alone, but high export, combined with a shallow afternoon MLD may also play a role. Shallow MLD enhances the loss of ML concentration for a given export rate, and night-time mixing can re-entrain some of this export, reducing the ML POC diel cycle relative to the O_2 diel cycle.

4.1.3 GOP—The tight fit between GOP and both GPP estimates over most of the study period provides important support for the O_2 diel cycles method as a measure of relative primary production in this region. Again, because all estimates were independent and taken at the same scale, and because the two-month deployment allowed 14 independent matchups at 3-day timescale, spanning a wide range of productivities, this dataset represents the most extensive validation to date of the O_2 diel cycles method as a measure of *relative* primary productivity. Additionally, the overall *accuracy* of our GOP estimates may be assessed indirectly through comparison with independent ship-based GOP estimates made during the May process cruise (Quay et al., 2012) and through comparison of our GOP/GPP and GOP/NPP ratio estimates with previous estimates from this region. Quay et al. (2012) estimated ML-integrated GOP using measurements of three oxygen isotopes: ^{16}O , ^{17}O , and ^{18}O , taken daily between May 3 and 21 during the process cruise. Mean GOP calculated by

this method was $245 \text{ mmol O}_2 \text{ m}^{-2} \text{ d}^{-1}$. This method integrates over several weeks, so we interpret their estimate to correspond roughly to mean ML depth-integrated GOP between April 19 (2 weeks before the first sample) and May 21. For comparison, we multiply each half-day GOP estimate by MLD to obtain ML-integrated GOP and obtain an average from April 19 to May 21 of $149 \text{ mmol m}^{-2} \text{ d}^{-1}$, 40% lower than Quay et al. (2012)'s estimate. However, our estimate integrates to the daily minimum MLD, and while the triple O_2 isotope method assumes constant MLD, we expect it to more closely approximate daily maximum MLD in the presence of diel MLD fluctuations, given its long integration time. Mean GPP_{Chl} during this period, integrated to the bottom of the daily *minimum* MLD is 30% lower than mean GPP_{Chl} integrated to the daily *maximum* MLD. If we assume the same relative difference for GOP, we obtain a revised ML-integrated GOP estimate of $213 \text{ mmol m}^{-2} \text{ d}^{-1}$, 13% lower than Quay et al. (2012)'s estimate. Given the uncertainties associated with the GOP methods as well as the differing spatio-temporal scales, this result provides first-order support for the accuracy of both methods. Our findings reinforce those of Hamme et al. (2012), who, in the Southern Ocean in March/April, found that mean ML integrated GOP calculated via O_2/Ar diel cycles (similar to our method) was 18% lower than GOP calculated via triple oxygen isotope method (similar to Quay et al., 2012).

Bender et al. (1992) calculated a GOP/NPP ratio of 2.5 during the spring bloom in the Northeast Atlantic, using in situ ^{18}O incubations and 24h ^{14}C incubations. We calculate $\text{GOP}/\text{NPP}_{\text{Chl}}$ as shown in Fig. 9b, but replace GPP_{Chl} with NPP_{Chl} and obtain a best-fit ratio and 95% confidence interval of 2.4 ± 0.2 for the bloom growth period and 1.7 ± 0.4 for the entire deployment. These fits appear to support the accuracy of both our NPP_{Chl} and GOP estimates during the bloom growth phase, consistent with our other findings. However, our GOP/GPP ratio estimates of 2.6 ± 0.2 (Fig. 9a) and 2.1 ± 0.2 (Fig. 9b) are near or above the high end of our expected range of 1.3–2.4 (see section 4.1.1). As discussed in previous sections, these ratios may be the result of high photosynthetic quotient and high DOC production, combined with a small negative bias in GPP_{cp} . Our GOP estimates may also be too high, but we cannot think of a plausible mechanism that would cause a substantial overestimate of diel-based GOP (but not of GPP_{cp}). Regardless of the source of our high GOP/GPP ratios, they are also consistent with Hamme et al. (2012), who also estimated GPP from on-deck PvE incubations as well as GOP via O_2/Ar diel cycles, providing a very close methodological comparison in a different environment (autumn, Southern Ocean). They obtain an even higher GOP/GPP ratio of 3.6. However, Hamme et al. (2012) assumed that 1–2h ^{14}C incubations represent GPP, while we assume that these same incubations represent NPP (when $\text{NPP} > 0$). If our assumption is correct, then their method provides a quantity closer to daytime NPP than GPP. However, even in this case, assuming moderate daytime phytoplankton respiration rates (30% of GPP), GOP/GPP during their study was > 2.5 , in agreement with our estimates. It is also worth noting that related studies comparing diel cycles in O_2 and pCO_2 measurements (Johnson 2010; Merlivat, et al., 2015), both of which include the effects of DOC production, have found ratios of daytime oxygen production to carbon production that are within the expected range of 1–1.45. These results provide further support for diel cycles-based O_2 production and the hypothesis that DOC production may drive the high GOP/GPP observed in this and other studies (Bender et al. 1992; Hamme et al. 2012).

In total, the available evidence provides first-order support for the accuracy of our diel cycles-based GOP estimates. Our findings build on important recent work in diverse environments showing that diel cycles in O_2/Ar ratio yield ML GOP estimates that are consistent with independent GOP estimates (Hamme et al. 2012), and that diel cycles in O_2 measurements from autonomous gliders in the subtropical Pacific provide GOP estimates that are a reasonable multiple of independent NPP results (Nicholson et al. 2015). Our results add a third ocean region (springtime North Atlantic) and a third platform (Lagrangian mixed-layer float), in addition to new comparisons with c_p and b_{bp} diel cycles.

4.1.4 GPP_{bbp}—Because diurnal variability in b_{bp} can be estimated from geostationary satellites (Neukermans et al. 2012), the ability to accurately estimate GPP from b_{bp} diel cycles would be extremely valuable. While ship-based measurements from NAB08 show that b_{bp} and c_p were equally well correlated with POC over the May cruise (Cetini et al. 2012), the poor matchups we find between GPP_{chl} and GPP_{bbp} suggest that diel changes are present in POC/ b_{bp} and can cause strong, consistent bias in GPP_{bbp}. Our results agree with previous findings that while beam attenuation and forward scattering by phytoplankton increase immediately after they begin to photosynthesize, b_{bp} and side scattering often do not, both in the lab (Ackleson et al. 1993; Poulin et al., 2018) and in the ocean (Kheireddine and Antoine 2014). These results caution against the use of b_{bp} diel cycles to estimate GPP without further research. However, it is worth noting that our afternoon GPP_{bbp} estimates are reasonably well correlated with GPP_{chl} ($r^2=0.63$, $m = 0.75\pm 0.23$; data not shown) during the bloom growth period. If this result is found to be robust in other times and places, then a useful estimate of GPP from satellite (and other) b_{bp} timeseries may be possible. However, even if the b_{bp} diel cycle cannot be used to estimate GPP, it likely contains other useful information, especially in combination with c_p and/or O_2 . If robust relationships between plankton community and/or physiology and b_{bp} diel cycles can be established (and, ideally, understood mechanistically), then measurements of b_{bp} diel cycles may still provide valuable oceanographic information, whether from in situ platforms or satellite.

4.2 Combined upper layer carbon budgets

Taken together with Alkire et al.'s (2012) NCP and carbon export estimates and our adaptation of Briggs et al.'s (2011) depth-resolved carbon fluxes, our productivity and bulk particle estimates provide a remarkably detailed, high-resolution picture of carbon flows over the entire spring bloom. From April 4–17, ML *Chl*, POC, and O_2 concentrations changed little, despite large fluctuations in MLD, while NO_3 increased slightly during deep-mixing, presumably due to entrainment, but was stable during shallow (<100 m) mixing. Consistent positive 125 m integrated NPP_{chl} (Fig. 11a) was therefore likely balanced by heterotrophic respiration. From April 18 to May 7, ML shoaling events coincided with several pulses of high net growth in POC_{cp} and *Chl* (Fig. 7), and the close match between NPP_{chl} and NCP during these periods (Fig. 10) suggests minimal role of grazing in regulating this growth. From May 6–7, all four gliders observed a rapid, aggregation event (Fig. 11b) that triggered a dramatic pulse in carbon export, both from the float patch and the broader (~30 km) glider survey area (Fig. 11a; blue and red lines). This pulse sank through the mesopelagic at ~75 m d⁻¹ and was composed primarily of fragile aggregates containing live phytoplankton including *Chaetoceros sp.* resting spores (Martin et al. 2011; Briggs et al.

2011; Rynearson et al. 2013). This aggregate export was the largest loss term of surface POC during the “main bloom”, reducing biomass accumulation rate by ~50% (Fig. 10). While SiO₄ limitation has been proposed as a cause of this rapid sinking event (Bagniewski et al. 2011), this aggregation commenced when SiO₄ concentrations were still >2 mmol m⁻³ (Fig. 7a) and five days prior to the ~35% reduction in GOP and GPP_{cp} that we attribute to SiO₄ limitation (pink band in Fig. 11b). The exact cause of this rapid aggregation event is unknown, but likely involves a combination of moderately high particle concentration (POC_{cp} > 10 mmol m⁻³), weakening of mixing (which could break fragile aggregates), and production of transparent exopolymer particles (P. Martin et al. 2011; Alkire et al. 2012). The combination of high export and reduced productivity at the end of the diatom bloom (May 12–14) appears to end the ML biomass accumulation. However, we conclude that the subsequent, sharp decline in ML Chl, POC_{cp}, and O₂ from May 14–15 (Fig. 7) was probably not the result of a dramatic increase in heterotrophic respiration, as implied by the strong negative NCP estimate (Fig. 10) of Alkire et al. (2012). Our conclusion stems from the night-time ML O₂ loss rates, which do not increase at all between the bloom peak the bloom decline (see Fig. 4a). Instead, the ML O₂ decline appears to be caused by further GPP decreases, due to continued SiO₄ limitation and a decline in *Chl* (Fig. 7b,d), likely enhanced by export of phytoplankton from the shallow ML. The O₂ decline (and accelerating *Chl* decline) may have been enhanced by advection of the float relative to the thin surface ML during afternoon profiles (see section 4.1.2), or perhaps an additional, light-dependent process, such as photoinhibition or photorespiration (Spilling et al. 2015), nearly eliminated GPP during this time, but only in the afternoons. After the decline of the diatom bloom, the different productivity estimates again provide a consistent picture, this time of top-down control. GOP and GPP_{cp} again show no sign of nutrient limitation (Fig. 9b,c, red symbols), and NPP_{Chl} is apparently balanced by heterotrophic respiration. Glider estimates of sinking POC export were low, but higher than early bloom export, despite similar NPP (Fig. 11a) and higher respiration. This result highlights the de-coupling between NCP and export on weekly-monthly timescales in this dynamic system and suggests that biomass and particle size are better predictors of sub-seasonal export dynamics. The changing export efficiencies that we observed (<15% through most of April, to ~57% during the main bloom to ~33% in the post-bloom period), provide a complex picture of “the spring bloom”, but still agree broadly with the export ratio of 45% calculated by Buesseler and Boyd (2009) in the North Atlantic spring bloom using JGOFS data, among the highest export efficiencies observed in the open ocean. However, unlike Buesseler and Boyd (2009), and in line with the conclusions of Martin et al. (1993), we see significant flux attenuation in the 100 m below the euphotic zone. For example, 35–48% of flux is lost between 60 m (Fig. 10) and 125 m (Fig. 11a) during the main bloom).

5 Conclusions

Our results, placed in the context of previous studies, provide strong support for the diel cycles method as a means to obtain estimates of GOP (from O₂) and GPP (from *c_p*) with reasonable accuracy relative to existing methods and enough precision on 3-day timescales to clearly resolve a spring diatom bloom. The range of biomass, mixing regimes and phytoplankton communities in this study, combined with previous results from the

subtropics, suggest that these methods are not overly dependent on particular ocean conditions. Because the diel cycles method is well suited for autonomous platforms, it has the potential to greatly increase our coverage of in-situ productivity estimates, providing both direct knowledge of this critical biological rate and greatly enhanced validation datasets for satellite-derived and modelled productivity. Our results also support the use of short-term ^{14}C incubations to parameterize simple PvE models for application to autonomous measurements, at least in the absence of strong nutrient limitation. We find high GOP/GPP ratios of 2.1–2.6 through most of the study, suggesting high DOC production and/or a possible moderate under-estimation of GPP by both methods. Finally, combined high-resolution estimates of NPP, particle size and sinking flux during the North Atlantic spring bloom show strong coupling between the three, modulated by a dramatic increase in export efficiency at bloom peak, apparently due to rapid aggregation.

Acknowledgements

Collection of data for this study was funded by the US National Science Foundation (Grants OCE-0628107 and OCE-0628379) and NASA (Grants NNX-08AL92G and NNX-10AP29H). Analysis and writing was further funded by a University of Maine Doctoral Research Fellowship, National Science Foundation grant OCE-1420929 and European Research Council grant. The authors would also like to thank Andrew Thomas and Emmanuel Boss for valuable comments and feedback as PhD committee members and the crew and technicians of the *R. V. Knorr* and *R. V. Bjarni Saemundsson* for making this entire study possible. We would also like to thank two anonymous reviewers for their feedback, which has substantially improved this manuscript.

7 References

- Ackleson SG, Cullen JJ, Brown J, and Lesser Michael. 1993 “Irradiance-Induced Variability in Light Scatter from Marine Phytoplankton in Culture.” *Journal of Plankton* 15 (7): 737–59. <http://plankt.oxfordjournals.org/content/15/7/737.short>.
- Alkire Matthew B., D’Asaro Eric, Lee Craig, Perry Mary Jane, Gray Amanda, Cetini Ivona, Briggs Nathan, et al. 2012 “Estimates of Net Community Production and Export Using High-Resolution, Lagrangian Measurements of O₂, NO₃⁻, and POC through the Evolution of a Spring Diatom Bloom in the North Atlantic.” *Deep Sea Research Part I: Oceanographic Research Papers* 64 (June): 157–74. doi:10.1016/j.dsr.2012.01.012.
- Bagniewski W, Fennel K, Perry MJ, and D’Asaro EA. 2011 “Optimizing Models of the North Atlantic Spring Bloom Using Physical, Chemical and Bio-Optical Observations from a Lagrangian Float.” *Biogeosciences* 8: 1291–1307. doi:10.5194/bg-8-1291-2011.
- Baines SB, and Pace ML. 1991 “The Production of Dissolved Organic Matter by Phytoplankton and Its Importance to Bacteria: Patterns across Marine and Freshwater Systems.” *Limnology and Oceanography* 36 (6): 1078–90. doi:10.4319/lo.1991.36.6.1078.
- Bender Michael, Ducklow Hugh, Kiddon John, Marra John, and Martin John. 1992 “The Carbon Balance during the 1989 Spring Bloom in the North Atlantic Ocean, 47°N, 20°W.” *Deep Sea Research Part I: Oceanographic Research Papers* 39 (10): 1707–25. <http://www.sciencedirect.com/science/article/pii/0198014992900250>.
- Bohren Craig F., and Huffman Donald R.. 1983 *Absorption and Scattering of Light by Small Particles*. New York: John Wiley & Sons.
- Briggs Nathan, Perry Mary Jane, Cetini Ivona, Lee Craig, D’Asaro Eric, Gray Amanda M., and Rehm Eric. 2011 “High-Resolution Observations of Aggregate Flux during a Sub-Polar North Atlantic Spring Bloom.” *Deep Sea Research Part I: Oceanographic Research Papers* 58 (10): 1031–39. doi:10.1016/j.dsr.2011.07.007.
- Briggs Nathan T, Slade Wayne H, Boss Emmanuel, and Perry Mary Jane. 2013 “Method for Estimating Mean Particle Size from High-Frequency Fluctuations in Beam Attenuation or Scattering Measurements.” *Applied Optics* 52 (27): 6710–25. <http://www.ncbi.nlm.nih.gov/pubmed/24085170>. [PubMed: 24085170]

- Buesseler KO, and Boyd PW. 2009 “Shedding Light on Processes That Control Particle Export and Flux Attenuation in the Twilight Zone of the Open Ocean.” *Limnology and Oceanography* 54 (4): 1210–32.
- Caffrey Jane M. 2003 “Production, Respiration and Net Ecosystem Metabolism in U.S. Estuaries” In *Coastal Monitoring through Partnerships*, edited by Melzian Brian D., Engle Virginia, McAlister Malissa, Sandhu Shabeg, and Eads Lisa Kay, 81:207–19. Springer. doi:10.1007/978-94-017-0299-7_19.
- Cetini I, Perry MJ, Asaro ED, Briggs N, Poulton N, Sieracki ME, and Lee CM. 2015 “A Simple Optical Index Shows Spatial and Temporal Heterogeneity in Phytoplankton Community Composition during the 2008 North Atlantic Bloom Experiment,” 2179–94. doi:10.5194/bg-12-2179-2015.
- Cetinic Ivona, Perry Mary Jane, Briggs Nathan T, Kallin Emily, D’Asaro Eric a, and Lee Craig M.. 2012 “Particulate Organic Carbon and Inherent Optical Properties during 2008 North Atlantic Bloom Experiment.” *Journal of Geophysical Research* 117. doi:10.1029/2011JC007771.
- Claustre H, Morel A, Babin M, Cailliau C, Marie D, Marty JC, Tailliez D, and Vaultot D. 1999 “Variability in Particle Attenuation and Chlorophyll Fluorescence in the Tropical Pacific: Scales, Patterns, and Biogeochemical Implications.” *Journal of Geophysical Research-Oceans* 104 (C2): 3401–22. doi:10.1029/98jc01334.
- Cullen John J, Lewis Marlon R, Davis Curtiss O, and Barber Richard T. 1992 “Photosynthetic Characteristics and Estimated Growth Rates Indicate Grazing Is the Proximate Control of Primary Production in the Equatorial Pacific.” *Journal of Geophysical Research* 97 (C1): 639. doi:10.1029/91JC01320.
- D’Asaro EA. 2003 “Performance of Autonomous Lagrangian Floats.” *Journal of Atmospheric and Oceanic Technology* 20 (6): 896–911.
- Dall’Olmo G, Boss E, Behrenfeld MJ, Westberry TK, Courties C, Prieur L, Pujo-Pay M, Hardman-Mountford N, and Moutin T. 2011 “Inferring Phytoplankton Carbon and Eco-Physiological Rates from Diel Cycles of Spectral Particulate Beam-Attenuation Coefficient.” *Biogeosciences* 8 (11): 3423–39. doi:10.5194/bg-8-3423-2011.
- Dall’Olmo G, Westberry TK, Behrenfeld MJ, Boss E, Courties C, Prieur L, Hardman-Mountford N, and Moutin T. 2011 “Inferring Phytoplankton Carbon and Eco-Physiological Rates from Diel Cycles of Spectral Particulate Beam- Attenuation Coefficient.” *Biogeosciences Discussions* 8 (2): 3009–50. doi:10.5194/bgd-8-3009-2011.
- García Hernán E., and Gordon Louis I. 1992 “Oxygen Solubility in Seawater: Better Fitting Equations.” *Limnology and Oceanography*. doi:10.4319/lo.1992.37.6.1307.
- Gernez Pierre, Antoine David, and Huot Yannick. 2011 “Diel Cycles of the Particulate Beam Attenuation Coefficient under Varying Trophic Conditions in the Northwestern Mediterranean Sea: Observations and Modeling.” *Limnology and Oceanography* 56 (1): 17–36. doi:10.4319/lo.2011.56.1.0017.
- Hamme Roberta C., Cassar Nicolas, Lance Veronica P., Vaillancourt Robert D., Bender Michael L., Strutton Peter G., Moore Tommy S., et al. 2012 “Dissolved O₂/Ar and Other Methods Reveal Rapid Changes in Productivity during a Lagrangian Experiment in the Southern Ocean.” *Journal of Geophysical Research* 117 (January): C00F12. doi:10.1029/2011JC007046.
- Harrison WG, and Platt T. 1986 “Photosynthesis-Irradiance Relationships in Polar and Temperate Phytoplankton Populations.” *Polar Biology*, 153–64. <http://link.springer.com/article/10.1007/BF00441695>.
- Johnson KS. 2010 “Simultaneous Measurements of Nitrate, Oxygen, and Carbon Dioxide on Oceanographic Moorings: Observing the Redfield Ratio in Real Time.” *Limnology and Oceanography* 55 (2): 615–27. doi:10.4319/lo.2009.55.2.0615.
- Kheireddine M, and Antoine D. 2014 “Diel Variability of the Beam Attenuation and Backscattering Coefficients in the Northwestern Mediterranean Sea (BOUSSOLE Site).” *Journal of Geophysical Research*: ..., 1–18. doi:10.1002/2014JC010007.Received.
- Kinkade CS, Marra J, Dickey TD, Langdon C, Sigurdson DE, and Weller R. 1999 “Diel Bio-Optical Variability Observed from Moored Sensors in the Arabian Sea.” *Deep Sea Research Part II: Topical Studies in Oceanography* 46 (8–9): 1813–31. doi:10.1016/S0967-0645(99)00045-4.

- Laws Edward A. 1991 “Photosynthetic Quotients, New Production and Net Community Production in the Open Ocean.” *Deep Sea Research Part A, Oceanographic Research Papers* 38 (1): 143–67. doi:10.1016/0198-0149(91)90059-O.
- Lippemeier Sebastian, Hartig Peter, and Colijn Franciscus. 1999 “Direct Impact of Silicate on the Photosynthetic Performance of the Diatom *Thalassiosira weissflogii* Assessed by On- and off-Line PAM Fluorescence Measurements.” *Journal of Plankton Research* 21 (2): 269–83. doi:10.1093/plankt/21.2.269.
- Lippemeier Sebastian, Hintze Ralf, Vanselow Klaus, Hartig Peter, and Colijn Franciscus. 2001 “In-Line Recording of PAM Fluorescence of Phytoplankton Cultures as a New Tool for Studying Effects of Fluctuating Nutrient Supply on Photosynthesis.” *European Journal of Phycology* 36 (1): 89–100. doi:10.1080/09670260110001735238.
- Marra J 2002 “Approaches to the Measurement of Plankton Production” In *Phytoplankton Productivity and Carbon Assimilation in Marine and Freshwater Ecosystems*, edited by Williams PJ, Thomas DR, and Reynolds CS, 222–64. London: Blackwell.
- Marra John, and Barber Richard T.. 2004 “Phytoplankton and Heterotrophic Respiration in the Surface Layer of the Ocean.” *Geophysical Research Letters* 31 (9): n/a-n/a. doi:10.1029/2004GL019664.
- Martin JH, Fitzwater SE, Gordon RM, Hunter CN, and Tanner SJ. 1993 “Iron, Primary Production and Carbon-Nitrogen Flux Studies during the JGOFS North Atlantic Bloom Experiment.” *Deep-Sea Research Part II-Topical Studies in Oceanography* 40 (1–2): 115–34.
- Martin P, Lampitt RS, Perry MJ, Sanders R, Lee C, and D’Asaro EA. 2011 “Export and Mesopelagic Particle Flux during a North Atlantic Spring Diatom Bloom.” *Deep-Sea Research Part I- Oceanographic Research Papers* 58 (4): 338–49.
- Merlivat L, Boutin J, and D’Ovidio F. 2015 “Carbon, Oxygen and Biological Productivity in the Southern Ocean in and out the Kerguelen Plume: CARIOCA Drifter Results.” *Biogeosciences* 12 (11): 3513–24. doi:10.5194/bg-12-3513-2015.
- Neukermans Griet, Loisel Hubert, Mériaux Xavier, Astoreca Rosa, and McKee David. 2012 “In Situ Variability of Mass-Specific Beam Attenuation and Backscattering of Marine Particles with Respect to Particle Size, Density, and Composition.” *Limnology and Oceanography* 57 (1): 124–44.
- Nicholson David P., Wilson Samuel T., Doney Scott C., and Karl David M.. 2015 “Quantifying Subtropical North Pacific Gyre Mixed Layer Primary Productivity from Seaglider Observations of Diel Oxygen Cycles.” *Geophysical Research Letters* 42: 4032–39. doi:10.1002/2015GL063065.
- Omand Melissa, Cetini Ivona, and Lucas Andrew. 2017 “Using Bio-Optics to Reveal Phytoplankton Physiology from a Wirewalker Autonomous Platform.” *Oceanography* 30 (2): 128–31. doi:10.5670/oceanog.2017.233.
- Pei Shaofeng, and Laws Edward a.. 2013 “Does the 14C Method Estimate Net Photosynthesis? Implications from Batch and Continuous Culture Studies of Marine Phytoplankton.” *Deep Sea Research Part I: Oceanographic Research Papers* 82 (December). Elsevier: 1–9. doi:10.1016/j.dsr.2013.07.011.
- Poulin Carina, Antoine David, and Huot Yannick. 2018 “Diurnal Variations of the Optical Properties of Phytoplankton in a Laboratory Experiment and Their Implication for Using Inherent Optical Properties to Measure Biomass.” *Optics Express* 26 (2): 711. doi:10.1364/OE.26.000711. [PubMed: 29401953]
- Quay P, Stutsman J, and Steinhoff T. 2012 “Primary Production and Carbon Export Rates across the Subpolar N. Atlantic Ocean Basin Based on Triple Oxygen Isotope and Dissolved O₂ and Ar Gas Measurements.” *Global Biogeochemical Cycles* 26 (2): 1–13. doi:10.1029/2010GB004003.
- Robertson JE, Watson AJ, Langdon C, Ling RD, and Wood JW. 1993 “Diurnal Variation in Surface PCO₂ and O₂ at 60°N, 20°W in the North Atlantic.” *Deep-Sea Research Part II* 40 (1–2): 409–22. doi:10.1016/0967-0645(93)90024-H.
- Rynearson TA, Richardson K, Lampitt RS, Sieracki ME, Poulton AJ, Lyngsgaard MM, and Perry MJ. 2013 “Major Contribution of Diatom Resting Spores to Vertical Flux in the Sub-Polar North Atlantic.” *Deep-Sea Research Part I: Oceanographic Research Papers* 82 Elsevier: 60–71. doi:10.1016/j.dsr.2013.07.013.

- Siegel DA, Dickey TD, Washburn L, Hamilton MK, and Mitchell BG. 1989 “Optical Determination of Particulate Abundance and Production Variations in the Oligotrophic Ocean.” *Deep Sea Research, Part A* 36: 211–22.
- Spilling Kristian, Ylöstalo Pasi, Simis Stefan, and Seppälä Jukka. 2015 “Interaction Effects of Light, Temperature and Nutrient Limitations (N, P and Si) on Growth, Stoichiometry and Photosynthetic Parameters of the Cold-Water Diatom *Chaetoceros Wighamii*.” *PLoS ONE* 10 (5): 1–18. doi:10.1371/journal.pone.0126308.
- Walsh ID, Chung SP, Richardson MJ, and Gardner WD. 1995 “The Diel Cycle in the Integrated Particle Load in the Equatorial Pacific - a Comparison with Primary Production.” *Deep-Sea Research Part II-Topical Studies in Oceanography* 42 (2–3): 465–77. doi:10.1016/0967-0645(95)00030-t.
- Wanninkhof Rik. 1992 “Relationship Between Wind Speed and Gas Exchange” 97 (92): 7373–82.
- White Angelique E., Barone Benedetto, Letelier Ricardo M., and Karl David M.. 2017 “Productivity Diagnosed from the Diel Cycle of Particulate Carbon in the North Pacific Subtropical Gyre.” *Geophysical Research Letters* 44 (8): 3752–60. doi:10.1002/2016GL071607.
- Wolf David K., and Thorpe SA. 1991 “Bubbles and the Air-Sea Exchange of Gases in near-Saturation Conditions.” *Journal of Marine Research* 49 (3). KLINE GEOLOGY LABORATORY, YALE UNIV, NEW HAVEN, CT 06520: 435–66. doi:10.1357/002224091784995765.
- Zhang XD, Hu LB, and He MX. 2009 “Scattering by Pure Seawater: Effect of Salinity.” *Optics Express* 17 (7): 5698–5710. [PubMed: 19333338]




Fig. 1. Study area with tracks of autonomous Lagrangian mixed-layer float and autonomous Seagliders.




Fig. 2.
 POC/c_p from the May cruise in upper 30 m, and fit used to calculate POC_{cp} .




Fig. 3. Hourly mixed-layer depth estimates calculated directly from float density measurements and from the Bagniewski et al. (2011) data assimilation model (black line), along with the depth of the float in mixed-layer mode (red line). Inset shows mean MLD diel cycle over the entire duration of the model (April 21-May 24). All MLD estimates use a density threshold of 0.01 kg m^{-3} to better approximate active mixing on an hourly timescale.




Fig. 4. Calculation of gross production of O₂ (a) and POC_{cp} (b) in the ML from their diel cycles.




Fig. 5. Example NPP vs PAR relationship from ¹⁴C incubations, with best fit “PvE” curve.




Fig. 6. Photosynthetic parameters P_M (a), α (b), and R_ϕ (c) vs in situ *Chl* with least squares power law fits and 95% confidence intervals. R_ϕ estimates from April and June cruises are excluded from fit (panel c, open circles).




Fig. 7. Float patch mixed layer timeseries of NO_3^- (a), SiO_4 (a), MLD (b), Chl (b), POC_{cp} (c), O_2 (d), and the concentration of O_2 saturation (d).





Fig. 8. Primary productivity estimates within the daily minimum ML. GPP_{ChI} , GOP, GPP_{cp} and GPP from Bagniewski et al. (2011), along with ML Chl. Diel cycles-based estimates are 3-day means; other productivity estimates are daily and Chl is continuous.

**Fig. 9.**

Relationships between primary productivity estimates: GOP vs GPP_{cp} (a), GOP vs GPP_{ChI} (b), GPP_{cp} vs GPP_{ChI} (c), and GPP_{bbp} vs GPP_{ChI} (d). Type I linear regressions are forced through the origin and include all data except the SiO_4 -depleted period (pink circles).

Expected range of GOP/GPP (a,b; dashed lines) assumes a photosynthetic quotient between 1–1.45 and 22–40% of fixed carbon released as DOC (see text).




Fig. 10. Estimates of sources and sinks of organic carbon integrated over the top 60 m: GPP_{Chl} and NPP_{Chl} and sinking particle export (this study), as well as NCP and loss due to the sum of sinking particle export and net DOC production and sinking particle export only (Alkire et al. 2012). Bloom periods follow Alkire et al. (2012) and are defined in the text (Section 3.3).




Fig. 11.

(a) Continuous productivity and export from the autonomous float and gliders, to/from the top 125 m over the entire study period. Productivity and glider export are 2-day running means while float export is averaged over longer periods denoted by the width of the bars. Bar height denotes uncertainty bounds. (b) Near surface glider D_{bbp} estimates from 10–50 m.

Table 1.

Abbreviations used in more than one subsection of the text

Abbreviation	Description
b_{bp}	particulate optical backscattering coefficient
chl	Chlorophyll <i>a</i> concentration
c_p	particulate optical beam attenuation coefficient
D_{bbp}	area-weighted mean particle diameter from optical backscattering
DOC	dissolved organic carbon concentration
GOP	Gross O ₂ productivity from O ₂ diel cycles
GPP	Gross primary productivity
GPP _{bbp}	GPP from optical backscattering diel cycles
GPP _{chl}	GPP from in situ chlorophyll and light measurements
GPP _{cp}	GPP from optical beam attenuation diel cycles
JGOFS	Joint Global Ocean Flux Study
K_{PAR}	Diffuse attenuation coefficient of PAR
ML	Mixed layer
MLD	Mixed layer depth
NPP	Net primary productivity
NPP _{chl}	NPP from in situ chlorophyll and light measurements
PAR	photosynthetically available radiation
P_m	Maximum GPP (light saturated)
POC	Particulate organic carbon concentration
POC _{bbp}	POC from optical backscattering
POC _{cp}	POC from optical beam attenuation
PP	Primary productivity
R_ϕ	Phytoplankton respiration rate
S	Salinity
T	Temperature
a	Initial slope GPP/PAR (light limited)
e	coefficient representing “sharpness” of NPP vs PAR relationship