



Published in final edited form as:

Qual Life Res. 2020 August ; 29(8): 2299–2310. doi:10.1007/s11136-020-02489-3.

Assessing Meaning & Purpose in Life: Development and Validation of an Item Bank and Short Forms for the NIH PROMIS®

John M. Salsman, PhD¹, Benjamin D. Schalet, PhD², Crystal L. Park, PhD³, Login George, PhD⁴, Michael F. Steger, PhD⁵, Elizabeth A. Hahn, MA^{2,6}, Mallory A. Snyder, MS, MPH⁷, David Cella, PhD^{2,6}

¹Department of Social Sciences and Health Policy, Wake Forest School of Medicine & the Wake Forest Baptist Comprehensive Cancer Center, Winston Salem, NC, USA

²Department of Medical Social Sciences, Northwestern University Feinberg School of Medicine, Chicago, IL, USA

³Department of Psychological Sciences, University of Connecticut, Storrs, CT, USA

⁴Department of Psychiatry and Behavioral Sciences, Memorial Sloan Kettering Cancer Center, New York, NY, USA

⁵Department of Psychology, Colorado State University, Fort Collins, CO, USA

⁶The Robert H. Lurie Comprehensive Cancer Center of Northwestern University, Chicago, IL, USA

⁷Office of Research and National Laboratories, The University of Chicago, Chicago, IL, USA

Abstract

Purpose: A sense of meaning and purpose is important for people living with acute and chronic illness. It can buffer the effects of stress and facilitate adaptive coping. As part of the Patient-Reported Outcomes Measurement Information System (PROMIS), we developed and validated an item response theory (IRT)-based measure of meaning and purpose in life (M&P).

Methods: Informed by a literature review and patient and content expert input, we wrote 52 items to assess M&P and administered them to a general population sample ($n=1000$) along with the Meaning in Life Questionnaire-Presence of Meaning Subscale (MLQ-Presence) and the Life Engagement Test (LET). We split the sample in half for exploratory factor analysis (EFA)

Corresponding Author: John M. Salsman, Ph.D., Department of Social Sciences and Health Policy, Wake Forest School of Medicine, Medical Center Boulevard, Winston-Salem, NC 27157, 336.713.3613 (phone), 336.716.7554 (fax), jsalsman@wakehealth.edu.

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

All procedures performed involving human participants were in accordance with the ethical standards of the Institutional Review Board of Northwestern University and with the 1964 Helsinki declaration and its later amendments.

Informed consent

Informed consent was obtained from all participants in the study.

followed by confirmatory factor analysis (CFA). IRT analyses included assessments of differential item functioning (DIF).

Results: Participants had a mean age of 47.8 years and 50.3% were male. EFA revealed one dominant factor and CFA yielded a good fitting model for a 37-item bank (CFI=0.962, TLI=0.960, RMSEA=0.085). All items were free of gender, age, education and race DIF. Internal consistency reliability estimates ranged from $\alpha=0.90$ (4-item short form) to $\alpha=0.98$ (37-item bank). The 8-item M&P short form was correlated with the MLQ-Presence ($r=0.89$), the LET ($r=0.79$), and the full PROMIS M&P item bank ($r=0.98$).

Conclusions: The PROMIS M&P measures demonstrated sufficient unidimensionality and displayed good internal consistency, model fit and convergent validity. Further psychometric testing of the PROMIS M&P item bank and short forms in people with chronic diseases will help evaluate the generalizability of this new tool.

Keywords

Meaning; life purpose; patient-reported outcomes; cancer; item-response theory; psychological well-being

INTRODUCTION

The Patient-Reported Outcomes Measurement Information System (PROMIS®) is an NIH Roadmap/Common Fund initiative to improve and standardize patient-reported outcomes across a range of conditions and demographic characteristics [1, 2]. It is the most ambitious attempt to date to apply models from item response theory (IRT) to health-related assessments across domains of physical, mental, and social health, yielding measures that are flexible, efficient, and precise [3]. The PROMIS approach involves iterative steps of comprehensive literature searches, development of conceptual frameworks, item pooling, qualitative assessment of items using focus groups and cognitive interviewing, and quantitative evaluation of items [4, 5].

The PROMIS initiative has primarily focused on developing instruments to assess health status for chronic conditions. Consequently, item banks developed thus far focus on symptoms and function, such as emotional distress, pain, fatigue, and social function [6]. However, many individuals with chronic conditions experience themselves as more than symptomatic or disabled, having learned to cope with their conditions in positive and adaptive ways [7]. Existing measures of health status often neglect psychological well-being and positive adjustment to illness. Most conceptualizations of psychological well-being include both hedonic (positive affect) and eudaimonic (life satisfaction, meaning and purpose) components [8, 9]. Psychometrically robust, IRT-informed measures of psychological well-being for healthy and ill adults are sparse. The NIH Toolbox initiative developed measures to assess meaning and purpose (an 18-item bank), but the raw score distributions tended to be negatively skewed, and precision estimates at the high and low ends of the information function continuum were less precise [10].

To address these limitations, we aimed to develop and validate an IRT-based patient-reported outcome tool of meaning and purpose for inclusion in PROMIS. Meaning in life refers to the

feeling that one's life and experiences make sense and matter [11]. Life purpose is characterized by the extent to which one experiences life as being directed, organized, and motivated by important goals [12]. The presence of meaning and purpose in life is considered a core component of mental health [13] and is a protective factor in health outcomes such as morbidity and mortality from cardiovascular disease [14, 15], quality of life among rheumatoid arthritis patients [16] and prevention of depressive symptoms, including suicidality [17, 18]. Meaning and purpose in life provide important perspectives through which we may better understand patients' experiences of illness.

We report the development, calibration and validation of the PROMIS Meaning and Purpose Item Bank. We aimed to: (1) refine a patient-reported outcome assessment tool of meaning and purpose for PROMIS and evaluate assumptions for IRT consistent with PROMIS Scientific Standards (e.g., unidimensionality, local independence) [5]; (2) examine item-level properties to support computer adaptive testing and evaluate possible differential item functioning (DIF); and (3) create short forms and examine convergent validity of the PROMIS Meaning & Purpose short forms and item bank.

METHODS

Participants and Procedures

We partnered with Opinions for Good (Op4G), an online research panel, to recruit a demographically diverse, general population sample from the United States ($n=1000$). Representativeness of data from internet samples is comparable to data from probability-based general population samples [19]. The internet is an efficient and low-cost means of data collection widely accessible to diverse groups [20]. The Institutional Review Board of Northwestern University approved this study. All interested and eligible participants provided informed consent electronically.

Op4G recruited participants by sending email invitations to a random selection of English-speaking panel members. Target distributions for age and gender were pre-specified (minimum $n=300$ in each of three age strata "18–39," "40–59," and "60–85" with a minimum $n=120$ men and 120 women in each age group), race and ethnicity (minimum $n=200$ participants who self-identify as Hispanic or Latino and minimum $n=200$ participants who self-identify as Black or African American), and educational attainment (minimum $n=400$ for high school graduate/GED and minimum $n=400$ for some college).

Following screening to ensure eligibility, participants provided informed consent and then completed a demographic survey and other self-report measures (described below). To reduce the potential for order effects, all measures were administered in random, thematic blocks, and order of measures within the blocks were also randomized. Participants who completed questionnaires were eligible for incentive-based compensation and donations made to a charity of their choice by Op4G.

Study Measures

PROMIS Meaning and Purpose Item Pool.—Informed by a literature review and qualitative input from patients and content experts, a pool of 52 items was created [21]. The

item pool comprised 18 items from the NIH Toolbox® Meaning and Purpose Item Bank [10], 8 items from the PROMIS Pediatric Meaning and Purpose Short Form [22], and 26 newly-written items to ensure adequate content coverage across the meaning and purpose continuum [23]. Each item was rated on a 5-point Likert scale with responses ranging from “strongly disagree” to “strongly agree” and from “not at all” to “very much.”

NIH Toolbox Meaning and Purpose Short Form.—The goal of the NIH Toolbox was to identify, create and validate brief comprehensive assessment tools to measure cognition, emotion, motor, and sensory function in longitudinal, epidemiological and intervention studies [24]. Within the emotional health domain, item banks and short forms were developed to assess positive affect, life satisfaction, and meaning and purpose, representing the first effort to develop IRT-informed measures of these important aspects of psychological well-being [25]. The NIH Toolbox Meaning and Purpose Short Form is an 8-item, calibrated short form that assesses the degree to which participants feel their lives matter or make sense [10]. Each item is rated on a 5-point Likert scale with responses ranging from “strongly disagree” to “strongly agree” and from “not at all” to “very much.” Cronbach’s alpha for this study was .92.

PROMIS Pediatric Meaning and Purpose Short Form.—The PROMIS Pediatric Meaning and Purpose Short Form is an 8-item, calibrated short form that assesses children’s evaluation of life as having purpose, goals to pursue, and a positive future [26]. Each item is rated on a 5-point Likert scale with responses ranging from “not at all” to “very much.” Cronbach’s alpha for this study was .95.

Life Engagement Test (LET).—The LET is a self-report measure of purpose in life or the extent to which a person engages in activities that are personally valued [27]. It includes six items rated on a 5-point Likert scale that ranges from “strongly disagree” to “strongly agree.” Three of the items are framed positively (e.g., “I value my activities a lot”), and three of the items are framed negatively and reverse-scored (e.g., “There is not enough purpose in my life”). Cronbach’s alpha for this study was .86.

Meaning in Life Questionnaire - Presence of Meaning subscale (MLQ-Presence).—The MLQ-Presence is a 5-item, self-report subscale used to evaluate how much participants feel their lives have meaning [28]. Each item is rated on a seven-point Likert scale with responses ranging from “absolutely untrue” to “absolutely true.” Sample items include, “My life has a clear sense of purpose” and “I understand my life’s meaning.” Cronbach’s alpha for this study was .87.

Positive and Negative Affective States (PANAS).—The PANAS is a 20-item, self-report measure that yields separate scores for positive affect (e.g., interested, excited, enthusiastic) and negative affect (e.g., distressed, irritable, afraid) [29]. Participants rate the extent they have felt “this” way over the past week. Each item is rated on a five-point Likert scale from “very slightly or not at all” to “extremely.” Cronbach’s alpha for this study was .92 for both positive and negative affect scales.

Satisfaction with Life Scale (SWLS).—The SWLS is a 5-item, self-report measure that captures the degree to which participants are content with or believe they have a good life [30]. Participants are asked to indicate how much they agree or disagree with statements using a seven-point Likert scale ranging from “strongly disagree” to “strongly agree.” High life satisfaction, along with high positive affect and low negative affect, is considered a key component of subjective well-being and closely related to meaning and purpose in life [31]. Cronbach’s alpha for this study was .87.

PROMIS Global-10.—The PROMIS Global is a 10-item short form that assesses general domains of health and functioning, including overall physical, mental, and social health, as well as pain, fatigue and overall perceived quality of life [32]. Participants respond using 5-point Likert scales or an 11-point Likert scale (i.e., pain) to indicate the quality of their health or the frequency or severity of their symptoms. We used the physical and mental health summary scores for this project. Cronbach’s alphas for the summary scores were .77 for Global Physical Health and .80 for Global Mental Health.

Statistical Analysis

We followed the general guidelines used in the PROMIS Scientific Standards for item bank development [4, 5, 33] and grouped them into three stages: (1) Testing assumptions for IRT modeling; unidimensionality and local independence of items; (2) Estimating item parameters using IRT, IRT-based local dependence analysis, evaluating items for DIF; and (3) Selecting items for static short forms and examining preliminary validity. After reviewing item content and analytic results, we used group consensus to decide the final composition of the static short forms.

During the first stage, we examined items for sparse data within any rating scale response category (i.e., $n < 5$). Data were randomly divided into two datasets ($n=500$ each), one for exploratory factor analysis (EFA) and the other for confirmatory factor analysis (CFA). We used the psych package in R for exploratory analyses [34, 35], and MPlus 7.2 [36] for confirmatory analysis. We applied EFAs of the polychoric correlation matrix with oblique rotation to identify potential factors among items; CFA was used to confirm final factor structure. In the EFAs, we examined the scree plot and parallel analysis as criteria to estimate meaningful factors. Parallel analysis compares the succession of factors of the observed data with that of a random data of the same size [37]. Items representing secondary factors or with loadings $< .4$ on the primary factor were considered for exclusion. Next, we estimated the proportion of total variance attributable to a general factor with omega hierarchical (omega-h) using the psych package [34]. This method estimates omega-h from the general factor loadings derived from an exploratory factor analysis and a Schmid-Leiman transformation [38]. Values of .70 or higher suggest that the item set is sufficiently unidimensional [39]. Finally, arriving at a single factor model, we examined residual correlations to identify any remaining locally dependent item pairs ($> .20$).

For CFA, we evaluated the final selection of items in a single factor model with fit statistics. We used the weighted least squares estimator with adjustments for the mean and variance (WLSMV) in Mplus, based on a polychoric correlation matrix, as appropriate for the

ordered categorical data [40]. We selected the commonly used indices for item banking as recommended by PROMIS Scientific Standards: Comparative Fit Index (CFI), Tucker-Lewis index (TLI), and Root Mean Square Error of Approximation (RMSEA). We used the following model fit indices as guidelines: $RMSEA < .08$; $CFI > .95$; $TLI > .95$. [4].

In the second stage, the total sample ($n=1000$) was used and items that met unidimensionality assumptions were analyzed using Samejima's Graded Response Model (GRM) [41] as implemented in IRTPRO software [42, 43]. The GRM is one of the most commonly used IRT models in health-related quality of life research [44]. Item threshold parameters represent items along the measured trait and show the coverage across the meaning and purpose continuum. The item slope parameter represents the discriminative ability of the items, with higher slope values indicating better ability to discriminate between adjoining values on the construct. Items displaying poor IRT fit (criterion: significant $S-X^2$ fit statistic, $p < 0.01$ [45, 46]) and poorly discriminating items (i.e., those with unacceptable IRT slopes; criterion: slope < 1) were candidates for exclusion at this stage. To ensure that parameter estimates are not unduly distorted by pairs of associated items, we assessed local dependence in the IRT framework with the chi-square (LD X^2) statistic; values of 10 or greater are considered large and unexpected [43].

We used the lordif package in R to conduct DIF analyses on the basis of age ("18–39" versus "40–59," "18–39" versus "60–85," "40–59" versus "60–85"), sex ("male" versus "female"), education ("high school" versus "some college") and race ("White" versus "non-White," "Black" versus "non-Black") for groups with a minimum of 150–200 participants per subgroup [47]. An item has significant DIF if the item exhibits different measurement properties between subgroups, which is similar to "item bias." We tested for DIF using an ordinal logistic regression procedure [48] with Chi-square to detect items ($p < 0.01$), and McFadden pseudo $R^2 > 0.02$ as the threshold for substantial DIF [49]. Items that demonstrated DIF greater than $R^2 > 0.02$ were considered for removal.

In the third and final stage, a fixed-length short form was determined by consensus. Our team of content-expert consultants, psychometricians, and measurement scientists reviewed item content, threshold and slopes for all meaning and purpose items in the newly calibrated bank to identify optimal 4-, 6-, and 8-item short forms. Finally, the convergent validity of the PROMIS Meaning & Purpose Item Bank and 8-item Short Form were examined using bivariate Pearson correlations with comparable constructs. For measures that provided item content for the development of the PROMIS Meaning & Purpose Item Pool and served as comparison measures (e.g., NIH Toolbox), we examined correlations with and without overlapping items. We hypothesized that the PROMIS Meaning & Purpose Item Bank and Short Forms would demonstrate the largest correlations with the NIH Toolbox Meaning and Purpose Short Form but would also be significantly correlated with the LET and the MLQ-Presence. We also expected PROMIS Meaning & Purpose scores to be significantly correlated with the PROMIS Global Mental Health scores and less strongly correlated with the PROMIS Global Physical Health scores.

RESULTS

Sample Characteristics

Our sample comprised approximately equal numbers of older (ages 60 to 85), middle-aged (ages 40 to 59), and young (ages 18 to 39) adults. It was primarily non-Hispanic, White (62.1%) but had good representation from racial and ethnic minorities. Approximately equal numbers of participants had received a high school education or less and greater than a high school education. The most common comorbidities reported were high blood pressure (39.2%), anxiety (27.7%), depression (27.0%), arthritis (26.6%), and migraines (24.2%). Additional demographic characteristics are shown in Table 1.

IRT assumptions

We examined frequencies for the 52-item PROMIS Meaning and Purpose Item Pool to ensure adequate numbers of responses for each category for all items. None of the candidate items had sparse data (i.e., $n < 5$) within any response category. Item-total correlations ranged from $r = 0.52$ “I understand the world around me” to $r = 0.86$ “My life has purpose.” To establish the relative unidimensionality of the PROMIS Meaning and Purpose Item Pool, we randomly split the sample into halves and conducted EFAs on the first half ($n = 500$) and a CFA on the second half to confirm a single model fit for the final item set. The EFAs were conducted (with the psych package in R) by generating a polychoric correlation matrix, followed by weighted least squares estimation. We first examined the unidimensionality of the item data with a scree plot, parallel analysis, and the residual correlation matrix of the single factor model. Results suggest that a second factor is formed by the reverse-scored items (nearly all showed residual correlations $> .20$). The two-factor EFA model (oblimin rotation) showed a dominant factor (eigenvalue = 26.5; 51% variance explained) with a second distinguishable factor (eigenvalue = 8.6; 17% variance explained) defined by the 10 negatively-worded items (e.g., “Most of what I do seems trivial and unimportant to me”). The output of this two-factor model is presented in Appendix 1 along with each item. Given the potential for the multidimensionality introduced by negatively-worded items to distort the interpretation and reliability of our final instrument’s scores, we opted to remove these items from further consideration.

During the exploratory phase of our analysis, we also removed five additional items based on conceptual and content grounds. We excluded three conceptually weaker items (“I feel grateful for each day”, “I expect to enjoy my future life”, “I feel hopeful about my future”). Finally, we excluded two additional items that were redundant with other item content (“I have a reason for living” and “I know where I am going in my life.”)

Next, we investigated distribution and unidimensionality of the remaining 37 item set. The frequency response distribution of these 37 items revealed a distribution with small level of skew (Mean = 134.3, $SD = 33.4$, Median = 140, Range = 38 to 185; Figure 1). Turning to unidimensionality, we produced a combined scree and parallel analysis plot of these items (Figure 2). This plot shows that all secondary factors have eigenvalues below 1 and close to the eigenvalues produced by random data. Consistent with these findings, the omega-

hierarchical index (based on the polychoric correlations) produced a high value (.87) suggesting the presence of a dominant general factor.

Finally, we conducted a single-factor CFA on a polychoric correlation matrix of the other half of the sample ($n=500$). Acceptable fit indices were obtained CFI=0.96, TLI=0.96, RMSEA=0.085). Table 2 shows the relatively high factor loadings of this model, ranging from 0.65 to 0.92. Thus, the proposed meaning and purpose bank was essentially unidimensional for purposes of scaling with IRT models.

Estimating Item Parameters and Evaluating DIF

Once we established essential unidimensionality, the next step was to calibrate the new meaning and purpose bank using estimated IRT parameters from a GRM to inform item slope (discrimination) and threshold (location) parameters. All item slopes were >1.0 , which met our inclusion criteria with the average slope = 2.28. The location parameters ranged from -5.59 to 1.38. However, two items suggested a poor fit ($S-X^2 < .01$) and were candidates for exclusion (“I have a reason for living,” “My life matters”).

Next we examined local dependency statistics. Out of 666 possible pairs, 60 pairs showed X^2 LD values of 10 or higher, affecting 16 items (out of 37). Because local dependencies may inflate discrimination parameter estimates, we estimated additional models. First, we identified 21 items that were relatively free of local dependencies. We then re-ran each of the 16 LD items with this 21 item set, and compared the resulting parameters with those that were generated from the full 37 item set. Discrimination parameter estimates from the 21+1 calibration runs were very similar to those obtained as part of the 37-item set. The average difference was 0.06 (range -0.29 to 0.30). The average discrimination parameter value for the 21+1 item runs was 2.79 (range 1.91 to 3.90). The average for those same items in the 37-item calibration was only slightly higher, 2.85 (range 1.88 to 4.06). We concluded that local dependencies did not meaningfully bias parameter estimates.

None of the 37 items exceeded the McFadden pseudo R^2 threshold of 0.02 in any of the DIF comparisons (sex, age, education and race). Since the two items with poor fit had good slopes, were free of DIF, and provided important and conceptually congruent content for meaning and purpose, they were retained for the final bank.

Next, IRT parameters were estimated using a GRM and linked to the NIH Toolbox metric, such that T-scores ($M=50$ and $SD=10$) are comparable and representative of the United States 2010 census [50, 51]. This was accomplished by following the multi-method linking procedure described by PROsetta Stone investigators [52]. Briefly, we obtained the official Toolbox item parameters from the investigators, and used these previously established values to fix the 10 overlapping items to anchor our analyses. In a co-calibration of the 10 Toolbox items, the new 27 Meaning and Purpose PROMIS items were freely estimated. As a second method, we used the Stocking-Lord procedure [53] to estimate linking constants defined by the difference of Toolbox item parameters we obtained from our sample compared to those we received from the Toolbox developers. The resulting linking constants were as follows: $A = 1.314$ and $B = -0.525$. They were then applied uniformly to the 27 new PROMIS item parameters to place them on the Toolbox metric. Both the fixed co-calibration and the

Stocking-Lord methods lead to similar test characteristic curves, with a maximum expected score difference of 2.25 points on a raw summed score range of 148 (37×4) at very low levels of the trait (< 2 SDs below the mean). The resulting Stocking-Lord linking constants ($A = 1.314$ and $B = -0.525$) were applied to the PROMIS item parameters to place them on the Toolbox metric.

Identifying a Short Form and Examining Preliminary Validity

Of particular relevance for identifying the “best” items for short forms was the information accounted for by each item across the meaning and purpose continuum. These calibrations and content considerations (identifying a conceptual range of meaning and purpose concepts) guided the selection of 4-, 6-, and 8-item short forms (Table 3) to go along with the 37-item bank. The 4-, 6-, and 8-item short forms and item bank demonstrated excellent internal consistency reliability, with coefficient alphas=0.90, 0.91, 0.93, and 0.98, respectively. In addition, the short forms were all positively correlated with the item bank ($r_s=0.94$ to 0.96). Table 4 presents bivariate correlations among the PROMIS Meaning & Purpose Short Forms and Item Bank with related constructs (life engagement, presence of meaning, positive affect, life satisfaction), the NIH Toolbox Meaning and Purpose Short Form and the PROMIS Global 10. All correlations between the PROMIS Meaning and Purpose short forms and the legacy measures, the MLQ-Presence and the LET were significant ($p<.001$) with $r_s=0.75$ or higher. Similarly, high correlations were found between the PROMIS Meaning and Purpose Short Forms and the comparable Toolbox and Pediatric short forms ($r_s=0.87$ to 0.96). Not surprisingly, correlations with the PROMIS Global measure were stronger with the Mental score ($r_s=0.65$ to 0.67) than with the Physical score ($r_s=0.37$ to 0.39).

CONCLUSIONS

The PROMIS Meaning and Purpose measure demonstrated sufficient unidimensionality and good internal consistency, model fit, and convergent validity. This is the first report summarizing the psychometric properties of this important component of psychological well-being for PROMIS and one of only three studies of which we are aware that applied a systematic, rigorous, and state-of-the-art measurement development approach to create a patient-reported outcome measure of meaning and purpose [10, 26, 54]. Of those three studies, only one measure (NIH Toolbox Meaning and Purpose Item Bank) was designed for use among healthy and ill adults [10]. The PROMIS Meaning and Purpose Item Bank builds on and extends the work of the NIH Toolbox in order to refine and strengthen the assessment of this important domain and further our understanding of healthy adaptation to illness.

The content of our Meaning and Purpose Bank was represented by 37 items that cover the conceptual breadth of the construct and yet remain sufficiently unidimensional. Recent work in the measurement of meaning in life suggests it comprises distinct but related concepts of mattering, purpose, and comprehension [23]. Other measurement approaches have focused on the search for meaning as well as the presence of meaning [28]. Within the scope of PROMIS, we prioritized the presence of meaning while also intentionally capturing the range of the construct, identifying existing and writing new item content [21]. One dominant

factor that included items from the tripartite approach to meaning in life emerged in our large general population sample. While not necessarily precluding a tripartite understanding of meaning, this finding does suggest the presence of an underlying, general meaning in life factor.

Our calibration testing further supported the potential utility of the PROMIS Meaning and Purpose Item Bank. In contrast to the NIH Toolbox Meaning and Purpose Item Bank, we obtained a normal distribution of scores from a similar general population sample. Although both measurement approaches can be administered as computer adaptive tests, the PROMIS Meaning and Purpose Item Bank includes 12 of the Toolbox Meaning and Purpose Bank items (all but the 5 negatively worded items and the item “I feel grateful for each day”) as well as an additional 24 items. Thus, administration of the full 37 items of the PROMIS Meaning and Purpose Bank or flexible administration of the Bank as a computer adaptive test should yield greater precision than the NIH Toolbox measure, over the range of the latent meaning and purpose continuum.

The newly developed PROMIS Meaning and Purpose short forms and item bank all had excellent internal consistency reliability and evidence of convergent validity. Although there are no true “gold standards” for assessing meaning in life within health-related research, the measures we included as indices of convergent validity are some of the more commonly used and psychometrically sound, brief measures of meaning and purpose [11, 27, 28] as well as the most commonly used measures of related well-being concepts of positive affect [29] and life satisfaction [30]. Although our item bank included overlapping content, the convergent validity correlations remained quite strong even after excluding the redundant items from the PROMIS measures. Similarly, the correlations with the existing NIH Toolbox and Pediatric PROMIS measures of meaning and purpose were quite large, suggesting considerable overlap in the construct. Lastly, the positive associations with global mental and global physical quality of life underscore the relationship between meaning in life and positive health [55–58].

Study limitations should be acknowledged: The cross-sectional design precludes examining potential responsiveness of the PROMIS Meaning in Life measures. A robust body of work focuses on meaning-making within the context of acute and chronic illnesses [7, 59–62] and the mutability of meaning is an important, patient-centered outcome. Further, psychosocial interventions to promote meaning have demonstrated efficacy [63] and psychometrically sound indices of meaning in life that capture change over time with minimal participant burden and maximal measurement precision are inherently valuable. A related concern is that the current calibration and validation testing did not include a clinical sample. Since PROMIS measures are designed for patients with a range of acute and chronic illnesses, it is not yet known how these new measures will perform among patients. To establish useful *T*-scores, it is important to calibrate and validate these new measures with a general population sample to serve as a meaningful reference group as a first step. Subsequent work will extend and increase the psychometric evidence for the PROMIS Meaning and Purpose measures.

In summary, the work described here provides initial and strong psychometric support for the PROMIS Meaning and Purpose item bank and short forms. These assessment tools were

designed to aid clinicians and researchers to better evaluate and understand the potential role of positive psychological processes for individuals with chronic health conditions. Further psychometric testing to examine criterion validity and responsiveness alongside commonly used measures of psychological well-being and in patients with chronic diseases will help evaluate the added benefit and generalizability of these new measures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Funding

This study was funded by the National Cancer Institute of the NIH (K07CA158008).

Funding Source: Research reported in this publication was supported by the National Cancer Institute of the NIH under award number K07CA158008. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

REFERENCES

1. Cella D, et al., Initial adult health item banks and first wave testing of the Patient-Reported Outcomes Measurement Information System (PROMIS™) Network: 2005–2008. *Journal of Clinical Epidemiology*, 2010 63(11): p. 1179. [PubMed: 20685078]
2. Cella D, et al., The Patient-Reported Outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005–2008. *Journal of Clinical Epidemiology*, 2010 63(11): p. 1179–1194. [PubMed: 20685078]
3. Segawa E, Schalet B, and Cella D, A comparison of computer adaptive tests (CATs) and short forms in terms of accuracy and number of items administered using PROMIS profile. *Qual Life Res*, 2020 29(1): p. 213–221. [PubMed: 31595451]
4. Reeve BB, et al., Psychometric Evaluation and Calibration of Health-Related Quality of Life Item Banks: Plans for the Patient-Reported Outcomes Measurement Information System (PROMIS). *Medical Care*, 2007 45(5 Suppl 1): p. S22–S31. [PubMed: 17443115]
5. PROMIS Health Organization and PROMIS Cooperative Group. PROMIS® Instrument Development and Validation: Scientific Standards Version 2.0 (revised May 2013) 2013 September 19, 2013; Available from: http://www.nihpromis.org/Documents/PROMISStandards_Vers2.0_Final.pdf.
6. Cook KF, et al., PROMIS measures of pain, fatigue, negative affect, physical function, and social function demonstrated clinical validity across a range of chronic conditions. *J Clin Epidemiol*, 2016 73: p. 89–102. [PubMed: 26952842]
7. Linley PA and Joseph S, Positive change following trauma and adversity: a review. *Journal of Traumatic Stress*, 2004 17(1): p. 11–21. [PubMed: 15027788]
8. Ryan RM and Deci EL, On happiness and human potentials: a review of research on hedonic and eudaimonic well-being. *Annual Review of Psychology*, 2001 52: p. 141–66.
9. Samman E, Psychological and subjective wellbeing: a proposal for internationally comparable indicators. *Oxford Development Studies*, 2007 35(4): p. 459–486.
10. Salsman J, et al., Assessing psychological well-being: self-report instruments for the NIH Toolbox. *Quality of Life Research*, 2014 23(1): p. 205–15. [PubMed: 23771709]
11. Steger MF, Oishi S, and Kashdan TB, Meaning in life across the life span: Levels and correlates of meaning in life from emerging adulthood to older adulthood. *Journal of Positive Psychology*, 2009 4(1): p. 43–52.
12. McKnight PE and Kashdan TB, Purpose in life as a system that creates and sustains health and well-being: An integrative, testable theory. *Review of General Psychology*, 2009 13(3): p. 242–251.

13. Fusar-Poli P, et al., What is good mental health? A scoping review. *Eur Neuropsychopharmacol*, 2020 31: p. 33–46. [PubMed: 31901337]
14. Koizumi M, et al., Effect of having a sense of purpose in life on the risk of death from cardiovascular diseases. *Journal of Epidemiology*, 2008 18(5): p. 191–6. [PubMed: 18753736]
15. Kim ES, Delaney SW, and Kubzansky LD, Sense of Purpose in Life and Cardiovascular Disease: Underlying Mechanisms and Future Directions. *Curr Cardiol Rep*, 2019 21(11): p. 135. [PubMed: 31673815]
16. Verduin PJ, et al., Purpose in life in patients with rheumatoid arthritis. *Clinical Rheumatology*, 2008 27(7): p. 899–908. [PubMed: 18214572]
17. Mascaro N and Rosen DH, Existential meaning's role in the enhancement of hope and prevention of depressive symptoms. *Journal of Personality*, 2005 73(4): p. 985–1013. [PubMed: 15958142]
18. Costanza A, Prelati M, and Pompili M, The Meaning in Life in Suicidal Patients: The Presence and the Search for Constructs. A Systematic Review. *Medicina (Kaunas)*, 2019 55(8).
19. Liu H, et al., Representativeness of the PROMIS Internet Panel. *Journal of Clinical Epidemiology*, 2010 63(11): p. 1169–78. [PubMed: 20688473]
20. Roster CA, et al., A comparison of response characteristics from web and telephone surveys. *International Journal of Market Research*, 2004 46(3): p. 359–374.
21. Salsman JM, et al., Refining and supplementing candidate measures of psychological well-being for the NIH PROMIS(R): qualitative results from a mixed cancer sample. *Qual Life Res*, 2018.
22. Ravens-Sieberer U, et al., Subjective well-being measures for children were developed within the PROMIS project: presentation of first results. *Journal of Clinical Epidemiology*, 2014 67(2): p. 207–218. [PubMed: 24295987]
23. George LS and Park CL, Meaning in life as comprehension, purpose, and mattering: Toward integration and new research questions. *Review of General Psychology*, 2016 20(3): p. 205.
24. Gershon RC, et al., Assessment of neurological and behavioural function: the NIH Toolbox. *Lancet Neurology*, 2010 9(2): p. 138–9. [PubMed: 20129161]
25. Salsman JM, et al., Emotion assessment using the NIH Toolbox. *Neurology*, 2013 80(11 Supplement 3): p. S76–S86. [PubMed: 23479549]
26. Forrest CB, et al., Assessing Children's Eudaimonic Well-Being: The PROMIS Pediatric Meaning and Purpose Item Banks. *J Pediatr Psychol*, 2019.
27. Scheier M, et al., The Life Engagement Test: Assessing Purpose in Life. *Journal of Behavioral Medicine*, 2006 29(3): p. 291–298. [PubMed: 16565785]
28. Steger MF, et al., The Meaning in Life Questionnaire: Assessing the Presence of and Search for Meaning in Life. *Journal of Counseling Psychology*, 2006 53(1): p. 80–93.
29. Watson D, Clark LA, and Tellegen A, Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 1988 54(6): p. 1063–1070. [PubMed: 3397865]
30. Diener E, et al., The Satisfaction with Life Scale. *Journal of Personality Assessment*, 1985 49(1): p. 71–75. [PubMed: 16367493]
31. Diener E, et al., Subjective Well-Being: Three Decades of Progress. *Psychological Bulletin*, 1999 125(2): p. 276–302.
32. Hays RD, et al., Development of physical and mental health summary scores from the Patient Reported Outcomes Measurement Information System (PROMIS) global items. *Quality of Life Research*, 2009 18(7): p. 873–80. [PubMed: 19543809]
33. Pilkonis PA, et al., Item banks for measuring emotional distress from the Patient-Reported Outcomes Measurement Information System (PROMIS): depression, anxiety, and anger. *Assessment*, 2011 18(3): p. 263–83. [PubMed: 21697139]
34. Revelle W, Psych: procedures for personality and psychological research (R Package Version 1.8.12) [Computer software] Evanston, IL: Northwestern University, 2019 Retrieved from: <http://cran.r-project.org/web/packages/psych/index.html>.
35. R Core Development Team. R: A language and environment for statistical computing [Computer software]. Vienna, Austria: R Foundation for Statistical Computing, 2018 Retrieved from: <http://www.r-project.org/>.

36. Muthen LK and Muthen BO, Mplus User's Guide. Seventh Edition 1998–2012, Los Angeles, CA: Muthen & Muthen.
37. Humphreys LG and Montanelli RG Jr, An investigation of the parallel analysis criterion for determining the number of common factors. *Multivariate Behavioral Research*, 1975 10(2): p. 193–205.
38. Schmid J and Leiman J, The development of hierarchical factor solutions. *Psychometrika*, 1957 22(1): p. 53–61.
39. Reise SP, et al., Multidimensionality and Structural Coefficient Bias in Structural Equation Modeling: A Bifactor Perspective. *Educational and Psychological Measurement*, 2013 73(1): p. 5–26.
40. Muthén B, et al., Robust inference using weighted least squares and quadratic estimating equations in latent variable modeling with categorical and continuous outcomes. 1997.
41. Samejima F, Estimation of latent ability using a response pattern of graded scores, in *Psychometrika Monograph Supplement, No. 17* 1969, Psychometric Society: Richmond, VA.
42. Thissen D, MULTILOG. 2003, Scientific Software International, Inc.: Lincolnwood, IL.
43. Cai L, Thissen D, and de Toit S, IRTPRO 2.01 2011, Scientific Software International: Lincolnwood, IL.
44. Preston K, et al., Using the nominal response model to evaluate response category discrimination in the PROMIS emotional distress item pools. *Educational and Psychological Measurement*, 2011 71(3): p. 523–550.
45. Orlando M and Thissen D, Further examination of the performance of S-X 2, an item fit index for dichotomous item response theory models. *Applied Psychological Measurement*, 2003 27: p. 289–298.
46. Kang T and Chen TT, Performance of the generalized S-X2 item fit index for the graded response model. *Asia Pacific Education Review*, 2011 12(1): p. 89–96.
47. Scott NW, et al., A simulation study provided sample size guidance for differential item functioning (DIF) studies using short scales. *Journal of Clinical Epidemiology*, 2009 62(3): p. 288–95. [PubMed: 18774693]
48. Zumbo BD, A handbook on the theory and methods of differential item functioning (DIF): Logistic regression modeling as a unitary framework for binary and Likert-type (ordinal) item scores. 1999, Ottawa, ON: Directorate of Human Resources Research and Evaluation, Department of National Defense.
49. Choi SW, Gibbons LE, and Crane PK, lordif: An R Package for Detecting Differential Item Functioning Using Iterative Hybrid Ordinal Logistic Regression/Item Response Theory and Monte Carlo Simulations. *Journal of Statistical Software*, 2011 39(8): p. 1–30.
50. Beaumont JL, et al., Norming plans for the NIH Toolbox. *Neurology*, 2013 80(11 Supplement 3): p. S87–S92. [PubMed: 23479550]
51. Babakhanyan I, et al., National Institutes of Health Toolbox Emotion Battery for English- and Spanish-speaking adults: normative data and factor-based summary scores. *Patient Relat Outcome Meas*, 2018 9: p. 115–127. [PubMed: 29588623]
52. Choi SW, et al., Establishing a common metric for depressive symptoms: linking the BDI-II, CES-D, and PHQ-9 to PROMIS depression. *Psychol Assess*, 2014 26(2): p. 513–27. [PubMed: 24548149]
53. Stocking ML and Lord FM, Developing a Common Metric in Item Response Theory. *Applied Psychological Measurement*, 1983 7(2): p. 201–210.
54. Carlozzi NE, et al., New measures to capture end of life concerns in Huntington disease: Meaning and Purpose and Concern with Death and Dying from HDQLIFE (a patient-reported outcomes measurement system). *Qual Life Res*, 2016 25(10): p. 2403–2415. [PubMed: 27393121]
55. Zika S and Chamberlain K, On the relation between meaning in life and psychological well-being. *British Journal of Psychology*, 1992 83 (Pt 1): p. 133–145. [PubMed: 1559144]
56. Ryff CD, Psychological well-being revisited: advances in the science and practice of eudaimonia. *Psychother Psychosom*, 2014 83(1): p. 10–28. [PubMed: 24281296]
57. Steptoe A, Deaton A, and Stone AA, Subjective wellbeing, health, and ageing. *Lancet*, 2015 385(9968): p. 640–648. [PubMed: 25468152]

58. Czekierda K, et al., Meaning in life and physical health: systematic review and meta-analysis. *Health Psychol Rev*, 2017 11(4): p. 387–418. [PubMed: 28488471]
59. Lee V, The existential plight of cancer: meaning making as a concrete approach to the intangible search for meaning. *Support Care Cancer*, 2008 16(7): p. 779–85 [PubMed: 18197427]
60. Park CL, Making sense of the meaning literature: An integrative review of meaning making and its effects on adjustment to stressful life events. *Psychological Bulletin*, 2010 136(2): p. 257–301. [PubMed: 20192563]
61. Martz E and Livneh H, Psychosocial Adaptation to Disability Within the Context of Positive Psychology: Findings from the Literature. *J Occup Rehabil*, 2016 26(1): p. 4–12. [PubMed: 26283187]
62. Leonhardt BL, et al., Recovery and serious mental illness: a review of current clinical and research paradigms and future directions. *Expert Rev Neurother*, 2017 17(11): p. 1117–1130. [PubMed: 28885065]
63. Park CL, et al., Effects of psychosocial interventions on meaning and purpose in adults with cancer: A systematic review and meta-analysis. *Cancer*, 2019 125(14): p. 2383–2393. [PubMed: 31034600]




Figure 1.
PROMIS Meaning and Purpose raw score distributions
Note: Scree and parallel analysis plot of final 37 items.




Figure 2. PROMIS Meaning and Purpose scree and parallel analysis plot

Table 1.Demographic Characteristics ($n=1000$)

Age	47.8 (M)	16.2 (SD)
	<i>N</i>	%
18–39	341	34.1
40–59	335	33.5
60–85	324	32.4
Sex		
Female	497	49.7
Ethnicity		
Hispanic Origin	199	19.9
Race		
White	683	68.3
Black/African American	200	20.0
Asian or Pacific Islander	41	4.1
Native American or Alaskan Native	37	3.7
Native Hawaiian or Other Pacific Islander	14	1.4
Other	50	5.0
Education		
8 th grade or less	50	5.0
Some high school	114	11.4
High school grad/GED	334	33.4
Some college/Technical degree/AA	217	21.7
College degree (BA/BS)	203	20.3
Graduate school	82	8.2
Country of Origin		
United States	897	89.7
Other	103	10.3
Relationship Status		
Married/Living with partner in a committed relationship	548	54.8
Never married	267	26.7
Divorced/Separated	154	15.4
Widowed	31	3.1
Times stayed in hospital overnight or longer in the past 12 months		
0	672	67.2
1–2	206	20.6
3–4	54	5.4
5–6	31	3.1
7–10	22	2.2
11–50	13	1.3

Age	47.8 (M)	16.2 (SD)
	N	%
51–75	2	0.2
Days that poor physical or mental health kept you from doing normal activities in the past 30 days		
0	448	44.8
1–2	193	19.3
3–4	96	9.6
5–6	77	7.7
7–10	75	7.5
11–30	111	11.1
Comorbidities		
None	236	23.6
1 or 2	349	34.9
3 or more	415	41.5
Used/taken prescription medication in the past 30 days		
Yes	646	64.6
No	340	34.0
Don't Know	14	1.4

Table 2 -

Confirmatory Factor Analyses Results

Item	Factor 1	Communality	Uniqueness
My life has purpose.	0.92	0.84	0.16
My life has significance.	0.91	0.82	0.18
I feel that my life has meaning.	0.91	0.82	0.18
My life is filled with meaning.	0.90	0.81	0.19
I feel a sense of purpose in my life.	0.89	0.79	0.21
My life has value.	0.89	0.79	0.21
My life has meaning.	0.89	0.78	0.22
The things I do in my life are of significance.	0.87	0.76	0.24
The things I do in my life are of value.	0.87	0.76	0.24
My life matters.	0.87	0.76	0.24
I can make sense of my life.	0.87	0.75	0.25
My life is fulfilling.	0.86	0.75	0.25
I have a clear sense of direction in life.	0.86	0.74	0.26
I can make sense of my existence.	0.86	0.74	0.26
My life has a clear sense of purpose.	0.86	0.73	0.27
I generally feel that what I do in my life is valuable and worthwhile.	0.85	0.72	0.28
I am positive about my future.	0.84	0.71	0.29
I can understand my life.	0.84	0.70	0.30
I have a reason for living.	0.83	0.69	0.31
My life makes sense to me.	0.83	0.69	0.31
I have discovered a satisfying life purpose.	0.83	0.69	0.31
I know where I am going in life.	0.83	0.69	0.31
I experience deep fulfillment in my life.	0.83	0.69	0.31
I understand that there is a reason for my life.	0.82	0.68	0.32
I understand my life's meaning.	0.82	0.67	0.33
My life as a whole has meaning.	0.81	0.66	0.34
My life has been productive.	0.80	0.63	0.37
I realize my life has a great deal of personal meaning to me.	0.79	0.62	0.38
I have a good sense of what makes my life meaningful.	0.78	0.61	0.39
I can reach my goals in life.	0.78	0.60	0.40
I have lots of reasons for living.	0.77	0.60	0.40
To me, the things I do are all worthwhile.	0.76	0.58	0.42
I have a clear understanding of what life is about.	0.76	0.58	0.42
My daily life is full of things that are interesting to me.	0.74	0.55	0.45
I have very clear goals and aims for my life.	0.74	0.55	0.45
I value my activities a lot.	0.73	0.53	0.47
I understand the world around me.	0.65	0.43	0.57

Note: Pattern matrix of factor loadings, based on polychoric correlations and weighted least squares estimation with adjustments for the mean and variance (WLSMV).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

PROMIS Meaning and Purpose Short Forms

PROMIS Meaning & Purpose Short Forms	4	6	8
My life has meaning.	*	*	*
I have a clear sense of direction in life.	*	*	*
I experience deep fulfillment in my life.	*	*	*
My life has purpose.	*	*	*
I have a good sense of what makes my life meaningful.		*	*
I have very clear goals and aims for my life.		*	*
I generally feel that what I do in my life is valuable and worthwhile.			*
My life has significance.			*

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4

Construct validity of PROMIS Meaning and Purpose measures

PROMIS Meaning & Purpose Measures	Meaning in Life Questionnaire-Presence Subscale	Life Engagement Test	Toolbox Meaning & Purpose SF	PROMIS Pediatric Meaning & Purpose SF	PROMIS Global Mental Health	PROMIS Global Physical Health	PANAS Positive	PANAS Negative	SWLS
4-item SF	0.84	0.75	0.87	0.91 (0.87)	0.65	0.37	0.65	-0.42	0.71
6-item SF	0.88 (0.85)	0.76	0.89	0.92 (0.89)	0.65	0.37	0.66	-0.43	0.71
8-item SF	0.89 (0.87)	0.79	0.91 (0.89)	0.92 (0.90)	0.66	0.38	0.66	-0.45	0.71
37-item Bank	0.91 (0.89)	0.82 (0.81)	0.95 (0.92)	0.94 (0.92)	0.67	0.39	0.67	-0.47	0.72

Note: All Pearson correlations are significant at the $p < 0.001$ level. Correlations in () indicate the association between the PROMIS Meaning and Purpose measure and the comparison measures without the overlapping item(s). SF= Short Form. SWLS = Satisfaction with Life Scale.