

# Potential adverse effects of ciprofloxacin and tetracycline on ARPE-19 cell lines

Nasim Salimiaghdam,<sup>1</sup> Lata Singh,<sup>1</sup> Kevin Schneider,<sup>1</sup> Angele Nalbandian,<sup>1</sup> Marilyn Chwa,<sup>1</sup> Shari R Atilano,<sup>1</sup> Andrea Bao,<sup>1</sup> M Cristina Kenney <sup>2,3</sup>

**To cite:** Salimiaghdam N, Singh L, Schneider K, *et al*. Potential adverse effects of ciprofloxacin and tetracycline on ARPE-19 cell lines. *BMJ Open Ophthalmology* 2020;**5**:e000458. doi:10.1136/bmjophth-2020-000458

Received 20 February 2020  
Revised 20 May 2020  
Accepted 1 June 2020

## ABSTRACT

**Background** We aim to determine the possible adverse effects of ciprofloxacin (CPF) and tetracycline (TETRA), as examples of bactericidal and bacteriostatic agents, respectively, on cultured human retinal pigment epithelial cells (ARPE-19).

**Methods** Cells were treated with 30, 60 and 120 µg/mL of CPF and TETRA. Cell metabolism was measured by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay. JC-1 dye (5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolyl carbocyanine iodide) assay was conducted to measure the mitochondrial membrane potential (MMP). The level of reactive oxygen species (ROS) was measured using the -2',7'-dichlorodihydrofluorescein diacetate assay (H2DCFDA). Quantitative real-time PCR was performed to analyse the gene expression levels associated with apoptosis (*BAX*, *BCL2-L13*, *BCL2*, *Caspase 3*, *Caspase 7* and *Caspase 9*), inflammatory (interleukin-1β (*IL-1β*), *IL-6*, *IL-33*, transforming growth factor-α (*TGF-α*), *TGF-β1* and *TGF-β2*) and antioxidant pathways (*SOD2*, *SOD3*, *GPX3* and *NOX4*), along with the mitochondrial DNA (mtDNA) copy numbers.

**Results** Results illustrated that while all three concentrations of CPF decreased cellular viability of ARPE-19 during all incubation periods, the 120 µg/mL TETRA resulted in increased cellular viability. At 48 and 72 hours, levels of MMP and ROS decreased significantly with each antibiotic. *BAX*, *BCL2-L13*, *CASP-7*, *CASP-9*, *SOD2* and *GPX3* genes overexpressed by either antibiotics. There was higher expression of *IL-6* and *IL-1B* with TETRA treatment. The level of mtDNA decreased using both treatments.

**Conclusions** Clinically relevant concentrations of CPF and TETRA have detrimental impacts on ARPE-19 cell lines in vitro, including upregulation of genes related to apoptosis, inflammation and antioxidant pathways. Additional studies are warranted to investigate if these harmful effects might be seen in retinal degeneration models in vivo.

## INTRODUCTION

Fluoroquinolones (ie, ciprofloxacin (CPF), ofloxacin, levofloxacin, fleroxacin, lomefloxacin, gatifloxacin) are currently a popular group of bactericidal antibiotics used to treat (1) skin, urinary tract, joint, sinus and lung infections; (2) ocular infections, such as endophthalmitis and bacterial keratitis<sup>1,2</sup> and (3) traumatic injuries

## Key messages

### What is already known about this subject?

- ▶ This subject is relatively new, this study examines the possible adverse effects of ciprofloxacin (CPF) and tetracycline (TETRA).
- ▶ Bactericidal and bacteriostatic antibiotics might induce detrimental impacts on retinal cell lines, notably in the population suffering from age-related disease.

### What are the new findings?

- ▶ Our new findings include clinically adjusted dosages of CPF and TETRA may have detrimental impacts human retinal pigment epithelial (ARPE-19) cells.

### How might these results change the focus of research or clinical practice?

- ▶ We speculate that if CPF can have deleterious effects on the wild type, healthy ARPE-19 cells, then it may also have deleterious effects on damaged mitochondria of patients with age-related disease such as age-related macular degeneration who are exposed to treatments with the fluoroquinolones.

prophylactically. Fluoroquinolones preliminarily inhibit the DNA gyrase (topoisomerase II) enzyme, which is involved in supercoiling, separation and replication of circular bacterial DNA.<sup>3</sup> Fluoroquinolones are very effective against intracellular, gram negative and positive organisms. Moreover, the fluoroquinolone penetrate through ocular barriers increases its impact for eye diseases.<sup>4</sup>

Bacteriostatic tetracyclines (TETRA) are efficient in treating skin, urinary, respiratory and chlamydia/trachoma infections. These antibiotics impede bacterial protein synthesis, which eventually prevent further bacterial growth and replication. TETRA bind to the bacterial 30S ribosome, preventing aminoacyl tRNA from interacting with the ribosome RNA complex. Furthermore, these antibiotics may alter the bacterial cytoplasmic membrane, causing leakage of cell material, thereby facilitating cell death.<sup>5</sup> While antibiotics have significant preventative and therapeutic effectiveness clinically, studies suggest that some antibiotics have significant



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Ophthalmology, University of California, Irvine, California, USA

<sup>2</sup>Ophthalmology, University of California School of Medicine, Irvine, California, USA

<sup>3</sup>Department of Pathology and Laboratory Medicine, University of California Irvine School of Medicine, Irvine, California, USA

## Correspondence to

Dr M Cristina Kenney; mkenney@hs.uci.edu

adverse effects. For instance, with respect to the eye, there are increased risks of retinal detachment, optic neuritis and retinal haemorrhage associated with administrations of fluoroquinolones.<sup>6</sup> Other adverse effects of antibiotics include permanent damage to the inner ear cells (auditory and vestibular), tendon damage and rupture, arthropathy, destruction of kidney cells and psychosis.<sup>7</sup> Human mammary epithelial cells (HMEC) in vitro show elevated levels of reactive oxygen species (ROS), protein carbonylation, lipid peroxidation and 8-hydroxy-2'-deoxyguanosine (marker for DNA damage) after treatment with fluoroquinolones.<sup>8</sup> These features were also present in mice treated 16 weeks with the antibiotics.<sup>8</sup> Primary human osteoblasts (PHO), osteosarcoma and HeLa cells were impaired by fluoroquinolone treatments.<sup>9</sup> High-content screening for mitochondrial proteins showed that several fluoroquinolones damaged the mitochondria of human liver cells.<sup>10</sup>

In 2015, over 32 million prescriptions for fluoroquinolones were given to patients in the USA for various medical conditions. Anticancer studies report that CPFX can halt cell cycle and cause double-strand DNA breaks that lead to increased apoptosis.<sup>11</sup> Exposure to CPFX causes lower cell viability and induces apoptosis in lung, melanoma and hepatocellular cancer cell lines.<sup>12</sup> CPFX blocks topoisomerase II inhibition in malignant cells but not normal cells<sup>13</sup> and induces G2 cell cycle arrest,<sup>14</sup> ultimately suggesting that a course of CPFX may be a reasonable adjunct therapy for some cancers.<sup>15</sup> While most people do not have any serious side effects, a small percentage have progressive, severe complications. Case reports of serious damage to multiple systems, including peripheral neuropathies, muscle weakness, pain in joints and tendons, cognitive impairment, along with gastrointestinal and respiratory disturbances, were shown for individuals treated with levofloxacin.<sup>16</sup> More recently, an excellent review on the negative impact of fluoroquinolones for a small number of individuals was published.<sup>17</sup> In 2015, the Food and Drug Administration (FDA) recognised a syndrome called fluoroquinolone-associated disability (FQAD) to describe otherwise healthy subjects that took fluoroquinolones and subsequently developed irreversible, severe side effects.

Mitochondria originated from ancestral aerobic bacteria,<sup>18 19</sup> and present-day bacteria and mitochondria possess many structural and biological similarities, such as similar outer membrane proteins and genomic sequence.<sup>20 21</sup> Therefore, it is not surprising that both would be detrimentally impacted by antibiotics. In particular, it is critical to determine if these antibiotics have negative influence on mitochondria from elderly patients that already have compromised mitochondrial functions. For example, studies have shown that patients with age-related macular degeneration (AMD) possess damaged and dysfunctional mitochondria that have increased susceptibility to stressors.<sup>22 23</sup> In AMD retinas, the retinal pigment epithelial (RPE) cells are the first cell type affected in this disease. In this study, we investigated

how CPFX and TETRA affect mitochondrial and cellular health in human ARPE-19 cells.

## MATERIALS AND METHODS

### Cell culture

ARPE-19 cells were purchased from the American Type Culture Collection (Manassas, Virginia, USA) and grown in a mixture of Dulbecco's Modified Eagle's medium/nutrient mixture F-12 (Invitrogen, Carlsbad, California, USA), 10% fetal bovine serum, 0.37% sodium bicarbonate, 0.58% L-glutamine, antibiotics (streptomycin sulphate 0.1 mg/mL, amphotericin-B 2.5 mg/mL, penicillin G 100 U/mL and gentamycin 10 mg/mL) and 10 mM non-essential amino acids. Cells were incubated in standard conditions (95% humidity, 5% CO<sub>2</sub> at 37°C).

Cells were treated with either CPFX (Cat#17850, Sigma-Aldrich, St. Louis, Missouri, USA) or TETRA (Cat# 87128, Sigma-Aldrich) at a of 0, 30, 60 and 120 µg/mL and cultured for 24, 48 and 72 hours. We used a hydrochloric acid (HCl) solution with 0.1 normality and methanol (Meth) as the vehicles for CPFX and TETRA, respectively.

### Cell metabolism (MTT assay)

Cell metabolism levels were measured with the MTT assay. ARPE-19 cells were cultured in 96-well plates (10<sup>4</sup>/well) and 10 µL MTT assay reagent (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) (Catalogue# 30006, Biotium, California, USA) was added to each well and plates were incubated at 37°C for 2 hours. Then, 100 µL/well DMSO was added to each well and plates were read in an absorbance reader (signal at 570 nm and reference at 630 nm) (Biotek Elx808 Absorbance Reader, Winooski, Vermont, USA). Experiments were performed three times. There were 12 replicate wells for each treatment modality.

### Reactive oxygen species (ROS assay)

Cells were cultured at a density of 10<sup>4</sup>/well in 96-well plates. Subsequently, 2', 7'-dichlorodihydrofluorescein diacetate; Catalogue# D399, Thermo Fisher Scientific, Waltham, MA solution, which is converted into a fluorescent molecule in the presence of ROS, was added to each well and plates were read on a fluorescent plate reader (SoftMax Pro, V.6.4, Catalogue# 94089, Sunnyvale, California, USA) at excitation (EX, 492 nm) and emission (EM, 520 nm) wavelengths. There were 12 wells for each treatment modality. Experiments were performed three times. There were 12 replicate wells for each treatment modality.

### Mitochondria membrane potential ( $\Delta\Psi_m$ )

Cells were seeded in 96-well plates (10<sup>4</sup>/well). After the treatment periods, the JC-1 reagent (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethyl-benzimidazolylcarbocyanine iodide; Catalogue# 30001, Biotium, California, USA) was added to each well and plates were incubated at 37°C for 15 min. Finally, plates were read on a

fluorescent plate reader (SoftMax Pro, V.6.4, Catalogue# 94089,) at red (EX 550 nm and EM 600 nm) and green (EX 485 nm and EM 535 nm) emissions to determine the ratios of red to green fluorescence. Experiments were performed three times. There were 12 replicate wells for each treatment modality.

### RNA/DNA isolation and cDNA amplification

ARPE-19 cells were treated with 120 µg/mL treatment concentrations of CPFEX and TETRA. Subsequently, DNA and RNA were isolated from the cellular lysate via application of Pure Genomic DNA Mini Kit (Thermo Fisher Scientific; Cat#K1820-01,) and RNeasy Mini-Extraction kit (Qiagen), according to manufacturer's protocol. The Nano Drop 1000 (Thermo-scientific) was used to determine the RNA/DNA concentration and purity from 15 samples of the ARPE-19 cells. Each RNA sample (100 ng) was reverse transcribed into cDNA using the QuantiTect reverse transcription kit (Qiagen).

### Quantitative real time polymerase chain reaction (qRT-PCR)

We evaluated the expression levels of genes related to the apoptosis (*BAX*, *BCL2-L13*, *BCL2*, *Caspase 3*, *Caspase 7* and *Caspase 9* (mitochondria specific)) (QuantiTect Primer Assay, Qiagen), inflammatory markers (interleukin-1β (*IL-1β*), *IL-6*, *IL-33*, *transforming growth factor-α* (*TGF-α*), *TGF-β1* and *TGF-β2*) and antioxidant enzymes (*SOD2*, *SOD3*, *GPX3* and *NOX4*) (table 1) by quantitative real-time-PCR (qRT-PCR). The total RNA was isolated from cultured treated cells and vehicle-control cells. Then QuantiFast SYBR Green PCR Kit (Qiagen, USA) on a Bio-Rad iCycler detection system was used for the Q-PCR. The HPRT1 primer was used as the housekeeping gene and standardisation of expression levels for all primers. Analyses were done in triplicates and no template control wells were used to assess the contamination. For all the experiments, HPRT1 gene was selected as the housekeeping gene.  $\Delta\Delta Ct$  method used for analysing the obtained data of qRT-PCR, which  $\Delta Ct = [Ct \text{ (threshold value) of the target gene}] - [Ct \text{ for HPRT1}]$ , and  $\Delta\Delta Ct = \Delta Ct \text{ of the treatment condition} - \Delta Ct \text{ of the untreated condition}$ . The fold changes of treated conditions compared with untreated condition were calculated as:  $\text{fold change} = 2^{-\Delta\Delta Ct}$ .

### Mitochondrial DNA copy number assay

The relative levels of mitochondrial DNA (mtDNA) copy numbers were measured for all samples by comparing the levels of mtDNA (MT-ND2) versus nuclear DNA (18S). The total DNA was isolated from the cultured treated cells, untreated and vehicle-control cells. TaqMan Gene Expression assay (Thermo Fisher Scientific, USA) was performed for the quantitative measurement of mtDNA. Analyses were done in triplicates.

### Statistical analyses

Using GraphPad Prism (V.5.0, GraphPad Software., and San Diego, California, USA) was used for all statistical analyses. The data was analysed by two-way analysis of

variance with the BONFERRONI test.  $P < 0.05$  was considered statistically significant.

## RESULTS

### Cell viability

Performed MTT assay revealed the mean percentages of cellular metabolism in cultures treated for 24 hours with 60 and 120 µg/mL concentrations of CPFEX decreased by 21.33% ( $p=0.04$ ) and 28.4% ( $p=0.014$ ), respectively, compared with the vehicle-control cells. After 48 hours of incubation, viability was reduced by 20.79% ( $p=0.025$ ) with 30 µg/mL, 27.85% ( $p=0.018$ ) with 60 µg/mL and 34.57% ( $p=0.011$ ) with 120 µg/mL CPFEX in comparison with vehicle-control samples. Moreover, in the 72 hours treatment period, the mean percentages declined by 27.46% ( $p=0.03$ ) with 30 µg/mL, 40.68% ( $p=0.014$ ) with 60 µg/mL and 45.55% ( $p=0.01$ ) with 120 µg/mL compared with vehicle-control samples (figure 1A).

In 24 hours cultures with TETRA, the mean percentage of cell viability of ARPE-19 cells treated with 120 µg/mL TETRA increased by 22.1% ( $p=0.001$ ) compared with vehicle-control wells (figure 1B). After a 48-hour treatment period at 120 µg/mL TETRA, there was a 17.7% ( $p=0.049$ ) increase in the mean percentage of cellular metabolism compared with the vehicle-control cells.

Altogether, these findings demonstrated that there is a negative association between the higher concentrations of CPFEX and cellular viability of ARPE-19 cells, notably in prolonged exposure periods. However, high concentrations of TETRA may induce positive influential effects on the viability of ARPE-19 cells.

### ROS production

The ROS levels in CPFEX-treated cultures at 24 and 48 hours showed no significant changes compared with vehicle-control samples (figure 1C). However, after 72 hours culture with either CPFEX 60 or 120 µg/mL, the mean percentages of ROS levels declined by 11.27% ( $p=0.048$ ) and 22.59% ( $p=0.028$ ), respectively, in comparison with vehicle-control cells.

TETRA-treated ARPE-19 cultures showed similar results on ROS levels (figure 1D). After 48 hours, cells treated with TETRA 120 µg/mL had a 32.6% ( $p=0.031$ ) decrease in ROS levels compared with untreated samples. Notably, at 72 hours, the mean percentage of ROS levels decreased by 15.62% ( $p=0.023$ ) with TETRA 60 µg/mL and 44.36% ( $p=0.032$ ) with TETRA 120 µg/mL compared with vehicle-control cells. These results suggest that higher concentrations of both antibiotics decrease ROS levels, most notably after longer incubation times.

### Changes in mitochondrial membrane potential

Compared with vehicle-control cells, the mean percentage of mitochondrial membrane potential (MMP) diminished by 36.81% ( $p=0.031$ ) in samples treated 24 hours with 120 µg/mL CPFEX. The MMP was reduced by 12.08% ( $p=0.037$ ) with CPFEX 30 µg/mL, by 25.27% ( $P=0.032$ ) with CPFEX 60 µg/mL and by

**Table 1** Information of the genes related to apoptotic, inflammatory and antioxidant pathways

Symbol	Gene name	GenBank accession no	Function
BAX	BCL2-associated X	NM_001291429 NM_001291428 NM_001291430 NM_001291431 NM_004324 NM_138761 NM_138763 NM_138764	This gene encodes a mitochondrially localised protein with conserved B-cell lymphoma two homology motifs. Overexpression of the encoded protein induces apoptosis.
BCL2-L13	BCL2 like 13	NM_015367 NM_001270729 NM_001270731 NM_001270732 NM_001270734 NM_001270735	Encodes a mitochondrially localised protein, apoptosis inducer.
BCL2	BCL2 apoptosis regulator	NM_000633	Encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells (eg, lymphocytes).
CASP-3	Caspase 3, apoptosis-related cysteine peptidase	NM_004346 NM_032991	Encodes protein as a cysteine-aspartic acid protease that plays a central role in the execution phase of cell apoptosis.
CASP-7	Caspase 7, apoptosis-related cysteine peptidase	NM_145248, XM_006725153, XM_006725154, XM_005268295, XM_006725155, XM_005268294, XM_006719962	This gene encodes a member of the cysteine-aspartic acid protease (Caspase) family. Sequential activation of caspases plays a central role in the execution phase of cell apoptosis.
CASP-9	Caspase 9, apoptosis-related cysteine peptidase	NM_001229 NM_032996	Encodes a member of the cysteine aspartic acid protease (caspase) family, which is involved in the execution phase of cell apoptosis.
IL-1 $\beta$	IL-1, beta	NM_000576	Produced by activated macrophages, IL-1 stimulates thymocyte proliferation. The protein encoded by this gene is a member of the IL-1 cytokine family. This cytokine is a pleiotropic cytokine involved in various immune responses, inflammatory processes and haematopoiesis
IL-6	IL-6	NM_000600	This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections.
IL-33	IL-33	NM_033439 NM_001199640 NM_001127180	The protein encoded by this gene is a cytokine that binds to the IL1RL1/ST2 receptor. The encoded protein is involved in the maturation of Th2 cells and the activation of mast cells, basophils, eosinophils and natural killer cells.

Continued

Table 1 Continued

Symbol	Gene name	GenBank accession no	Function
TGF- $\alpha$	TGF alpha	NM_003236 NM_001099691	This gene encodes a growth factor that is a ligand for the epidermal growth factor receptor, which activates a signalling pathway for cell proliferation, differentiation and development. This protein may act as either a transmembrane-bound ligand or a soluble ligand.
TGF- $\beta$ 1	TGF beta-1-like	NM_003238	This gene is a polypeptide member of the TGF beta superfamily of cytokines. It is a secreted protein that performs many cellular functions, including the control of cell growth, cell proliferation, cell differentiation and apoptosis.
TGF- $\beta$ 2	TGF beta 2	NM_001135599	This gene encodes a secreted ligand of the TGF-beta superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression.
SOD2	Superoxide dismutase 2	NM_000636	This gene is a member of the iron/manganese superoxide dismutase family. It encodes a mitochondrial protein that forms a tetramer and binds one manganese ion per subunit. This protein binds to the superoxide byproducts of oxidative phosphorylation and converts them to hydrogen peroxide and diatomic oxygen.
SOD3	Superoxide dismutase 3	NM-003102	This gene encodes a member of the SOD protein family, which catalyses the conversion of superoxide radicals into hydrogen peroxide and oxygen, effective in protection of the brain, lungs and other tissues from oxidative stress.
GPX3	Glutathione peroxidase 3	NM_002084	The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyse the reduction of organic hydroperoxides and hydrogen peroxide (H <sub>2</sub> O <sub>2</sub> ) by glutathione, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity.
SOD2	Superoxide dismutase 2	NM_000636	This gene is a member of the iron/manganese superoxide dismutase family. It encodes a mitochondrial protein that forms a homotetramer and binds one manganese ion per subunit. This protein binds to the superoxide byproducts of oxidative phosphorylation and converts them to hydrogen peroxide and diatomic oxygen.
SOD3	Superoxide dismutase 3	NM-003102	This gene encodes a member of the SOD protein family, which catalyses the conversion of superoxide radicals into hydrogen peroxide and oxygen, effective in protection of the brain, lungs and other tissues from oxidative stress.

Continued


Table 1 Continued

Symbol	Gene name	GenBank accession no	Function
GPX3	Glutathione peroxidase 3	NM_002084	The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyse the reduction of organic H <sub>2</sub> O <sub>2</sub> by glutathione, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity.
NOX4	NADPH oxidase 4	NM_001143836 NM_016931 NM_001143837 NR_026571 NM_001291926 NM_001300995 XM_006718848 NM_001291927 XM_006718852 XM_006718853 NM_001291929 XM_006718849	This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localised to non-phagocytic cells where it acts as an oxygen sensor and catalyses the reduction of molecular oxygen to various reactive oxygen species.

IL-1, interleukin-1; SOD, superoxide dismutase; TGF, transforming growth factor.

39.68% ( $p=0.013$ ) with CPFX 120  $\mu\text{g}/\text{mL}$  treatment after 48 hours culture compared with untreated samples. The results showed that cells treated with 60 and 120  $\mu\text{g}/\text{mL}$  CPFX for 72 hours had 11.27% ( $p=0.035$ ) and 22.59% ( $p=0.025$ ) decrease of mean percentages in comparison with vehicle-control cells (figure 1E). The MMP mean percentages of cells treated with 60 and 120  $\mu\text{g}/\text{mL}$  TETRA for 24 hours were reduced by 21.61% ( $p=0.021$ ) and 43.58% ( $p=0.014$ ), respectively, compared with vehicle-control cultures. Furthermore, cells cultured for

48 hours showed a reduction by 18.6% with 30  $\mu\text{g}/\text{mL}$  ( $p=0.026$ ), 26.75% with 60  $\mu\text{g}/\text{mL}$  ( $p=0.021$ ) and 44.26% with 120  $\mu\text{g}/\text{mL}$  ( $p=0.014$ ) compared with vehicle-control cultures. There was also a decrease in  $\Delta\Psi\text{m}$  after 72 hours of TETRA treatment; with mean percentages decline by 17.44% ( $p=0.003$ ) with 30  $\mu\text{g}/\text{mL}$ , 44.51% ( $p=0.004$ ) with 60  $\mu\text{g}/\text{mL}$  and 67.69% ( $p=0.001$ ) with 120  $\mu\text{g}/\text{mL}$ , respectively, compared with vehicle-control wells (figure 1F). Altogether, these in vitro findings show



**Figure 1** Treatment effects of CPFX and TETRA on cellular metabolism (figure 1A,B), ROS levels (figure 1C,D) and mitochondrial membrane potential (MMP, figure 1E,F) in ARPE-19 cells after 24, 48 and 72 hours as measured with the MTT, H<sub>2</sub>DCFDA and JC-1 assays. \* $P<0.05$ , \*\* $P<0.01$ . ARPE-19, human retinal pigment epithelial cells; CPFX, ciprofloxacin; H<sub>2</sub>DCFDA, 2',7'-dichlorodihydrofluorescein diacetate; ns, not significant; ROS, reactive oxygen species; TETRA, tetracycline.

that higher concentrations of both CFX and TETRA decreased MMP over time.

### Expression levels for pro-apoptosis, antioxidant and pro-inflammation genes

In all experiments of the qRT-PCR, the vehicle-control cells were considered at a value of 1 (table 2). After treatment with CPF, the relative gene expression of *BAX* (1.977-fold,  $p=0.002$ ) (figure 2A), *BCL2-L13* (1.414-fold,  $p=0.014$ ) (figure 2B), *Caspase 7* (1.712-fold,  $p=0.003$ ) (figure 2E), *Caspase 9* (2.319-fold,  $p=0.013$ ) (figure 2F), *SOD2* (1.26-fold,  $p=0.01$ ) (figure 3A) and *GPX3* (1.24-fold,  $p=0.018$ ) (figure 3D) were increased as compared with the HCl-treated control cells. However, the relative gene expression levels of NOX4 in CPF exposed cells decreased to 0.84-fold ( $p=0.031$ ) compared with HCl-treated controls (figure 3C).

The TETRA-treated cells had relative gene expression levels of *BAX* (1.465-fold,  $p=0.04$ ) (figure 2A), *BCL2-L13* (1.142-fold,  $p=0.026$ ) (figure 2B), *BCL2* (2.82-fold,  $p=0.024$ ) (figure 2C), *Caspase-7* (1.712-fold,  $p=0.002$ ) (figure 2E), *Caspase-9* (1.434-fold,  $p=0.007$ ) (figure 2F) and *SOD2* (1.27-fold,  $p=0.048$ ) (figure 3A) compared with the Meth-treated control group. Moreover, the relative gene expression levels of SOD3 and NOX4 in 120 µg/mL TETRA-treated cells decreased to 0.41-fold ( $p=0.041$ ) (figure 3B) and 0.37-fold, ( $p=0.045$ ) (figure 3C) in comparison with the Meth-treated control cells.

Although the relative gene expression of inflammatory marker IL-6 in CPF-treated cells increased to 1.31-fold ( $p=0.048$ ) (figure 4A), the relative gene expression of IL-33 declined to 0.34-fold ( $p=0.047$ ) (figure 4C) compared with HCl-treated control cells. Moreover, in TETRA-treated cells, the relative gene expression of TGF- $\alpha$ 1 (0.41-fold,  $p=0.014$ ) (figure 4D), TGF $\beta$ 1 (0.54-fold,  $p=0.043$ ) (figure 4E) and IL-33 (0.11-fold,  $p=0.011$ ) (figure 4C) compared with Meth-treated control samples. Also, in 120 µg/mL TETRA-treated cells, the relative gene expression of IL-6 and IL-1 $\beta$  increased to 5.1-fold ( $p=0.001$ ) (figure 4A) and 2.29-fold ( $p=0.023$ ) (figure 4B), compared with the Meth-treated control cells.

Therefore, it is possible that exposure of ARPE-19 cells with CPF and TETRA facilitated higher expression of some apoptotic, inflammatory and antioxidant enzyme-related genes.

### Significant reduction of mtDNA copy number

In CPF-treated ARPE-19 cultures, the level of mtDNA decreased to 0.23-fold ( $p=0.008$ ) compared with HCl-treated control cells (figure 5). In addition, treatment with TETRA reduced the relative level of mtDNA to 0.28-fold ( $p=0.041$ ) in comparison with Meth-treated control cells (figure 5). The untreated cultures were assigned a value of 1. Our findings show that treatment of ARPE-19 with CPF and TETRA led to significant reduction of mtDNA copy numbers.

**Table 2** Expression levels in ARPE-19 Treated cells with CPF and TETRA compared with vehicle-treated controls

Symbol	Treated cells with CPF versus vehicle-control* P value, fold	Treated cells with TETRA versus vehicle-control* P value, fold
<b>Apoptosis-regulator genes</b>		
BAX	0.002 1.977±0.03	0.04 1.465±0.09
BCL2-L13	0.014 1.41±0.02	0.026 1.14±0.03
BCL2	0.061 1.76±0.04	0.024 2.82±0.56
CASP3	0.246 1.39±0.09	0.250 1.25±0.30
CASP7	0.003 1.71±0.02	0.002 1.71±0.02
CASP9	0.013 2.13±0.19	0.007 1.43±0.03
<b>Antioxidant enzymes related genes</b>		
SOD2	0.014 1.26±0.06	0.048 1.27±0.07
SOD3	0.24 0.80±0.05	0.041 0.41±0.05
NOX4	0.031 0.84±0.06	0.045 0.37±0.02
GPX3	0.018 1.24±0.06	0.249 0.69±0.01
<b>Inflammatory pathway genes</b>		
IL-6	0.048 1.31±0.03	0.001 5.11±0.16
IL-1 $\beta$	0.86 0.31±0.05	0.023 2.29±0.19
IL-33	0.047 0.34±0.04	0.011 0.11±0.008
TGF- $\alpha$ 1	0.211 0.67±0.06	0.014 0.41±0.02
TGF- $\beta$ 1	0.061 1.003±0.08	0.043 0.54±0.02
TGF- $\beta$ 2	0.410 0.62±0.04	0.177 0.74±0.07


Fold values greater than 1 indicate upregulation of the gene. Fold values less than 1 indicate downregulation of the gene.

\*Are assigned a value of 1.

ARPE-19, human retinal pigment epithelial cells; CPF, ciprofloxacin; IL-6, interleukin-6; TETRA, tetracycline; TGF- $\beta$ 1, transforming growth factor- $\beta$ 1.

### DISCUSSION


In this study, we evaluated the effects of CPF and TETRA on ARPE-19 cells in vitro. We assessed the outcomes



**Figure 2** Expression of apoptotic genes (A) BAX, (B) BCL2-L13, (C) BCL-2, (D) CASP-3, (E) CASP-7 and (F) CASP-9 in CPFX and TETRA treated ARPE19 cells. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ . ARPE19, human retinal pigment epithelial cells; Cont, untreated control; CPFX, ciprofloxacin; HCl, hydrochloric acid vehicle control; Meth, methanol vehicle control; ns, not significant; TETRA, tetracycline.


of these treatments on levels of cellular viability, ROS, MMP and expression levels of genes involved in inflammatory, apoptotic and antioxidant enzymes pathways. Our findings showed that treatment with clinically relevant dosages of CPFX resulted in significantly decreased ARPE-19 cell viability, metabolic activity and MMP that were more prominent with longer incubation and higher concentrations of CPFX. Four of the proapoptosis genes

(*BAX*, *BCL2-L13*, *Casp-7* and *Casp-9*) were upregulated after CPFX treatment, which is consistent with the lower cell numbers. Additionally, IL-6 levels were higher in the CPFX-treated cells, indicating that elevated levels of inflammation would be found associated with CPFX treatment. None of the other inflammation-related genes (*TGF- $\alpha$* , *TGF- $\beta$ 1*, *TGF- $\beta$ 2* or *IL1 $\alpha$* ) changed expression levels on CPFX exposure.



**Figure 3** Expression of antioxidant genes (A) SOD2, (B) SOD3, (C) NOX4 (D) GPX3 in CPFX and TETRA treated ARPE19 cells. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ . ARPE19, human retinal pigment epithelial cells; Cont, untreated control; CPFX, ciprofloxacin; HCL, hydrochloric acid vehicle control; Meth, methanol vehicle control; ns, not significant; TETRA, tetracycline.






**Figure 4** Expression of inflammatory pathway genes (A) IL-6, (B) IL-1 $\beta$ , (C) IL-33, (D) TGF- $\alpha$ 1, (E) TGF- $\beta$ 1 and (F) TGF- $\beta$ 2 in CPF and TETRA treated ARPE19 cells. \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ . ARPE19, human retinal pigment epithelial cells; Cont, untreated control; CPF, ciprofloxacin; HCl, hydrochloric acid vehicle control; IL-6, interleukin-6; Meth, methanol vehicle control; ns, not significant; TETRA, tetracycline; TGF- $\beta$ 1, transforming growth factor- $\beta$ 1.

In the ARPE-19 cultures, ROS levels remained unchanged at the 24 and 48 hours time points with the 60 and 120  $\mu\text{g}/\text{mL}$  CPF dosages by 72 hours. These data suggest that the ARPE-19 response to CPF was via the mitochondria (lower MMP) and the cellular metabolic activity through the Nicotinamide adenine dinucleotide phosphate (NADPH) flux (lower MTT) and not through the generation of ROS. The lower ROS levels observed in the CPF and TETRA treated ARPE-19 cells may also in part be due to the upregulation of SOD2, and the downregulation of NOX4, which is responsible for the production of ROS. Moreover, other reports have shown enhanced lactate production due to stimulated glycolysis,<sup>24</sup> which would also lead to lower electron transport chain (ETC) activity and decreased ROS production with reduced mitochondrial activity.

Figure 5



**Figure 5** Effects of TETRA and CPF on the of mtDNA copy number level compared with nDNA. \* $P < 0.05$ , \*\* $P < 0.01$ . Cont, untreated control; CPF, ciprofloxacin; HCL, hydrochloric acid vehicle control; Meth, methanol vehicle control; mtDNA, mitochondrial DNA; ns, not significant; TETRA, tetracycline.

In contrast to our findings, Kalghatgi *et al* evaluated CPF-treated non-cancer mammary epithelial cells (MCF10A), primary human aortic endothelial cells (PAEC), primary HMEC, human gut epithelial cells and normal human diploid skin fibroblasts and reported higher levels of ROS, lipid peroxidation and oxidative damage to DNA and proteins. Usually bactericidal antibiotics disrupt the ETC and tricarboxylic cycle which would lead to lower formation of the endogenously generated ROS.<sup>8</sup>

Many of the published data related to CPF have been conducted using various cancer cell lines so it might be expected that our findings with ARPE-19 cells would be different. Cancer-related studies have illustrated CPF has inhibitory effects on leukaemia cells, glioblastoma cells, colorectal and bladder cancer cells that act via diminished MMP and increased apoptosis.<sup>25–28</sup> Duetzelhenke *et al* evaluated the effects of 20 antibiotics on PHO, the osteosarcoma cell lines (MG63) and Hela epithelial cell lines demonstrating fluoroquinolones, including CPF, decreased the metabolic activity and proliferation of PHO by 20% and inhibited mitochondrial oxidative phosphorylation, but the cells underwent a metabolic shift to use glycolysis, as measured by lactate concentrations.<sup>9</sup> Similarly, Holtom *et al* analysed murine osteoblast-like cell lines (MC3T3-E1) and reported a 50% inhibitory concentration of 40  $\mu\text{g}/\text{mL}$  CPF at both 48 and 72 hours and decreased 5-bromo-2'-deoxyuridine at 80  $\mu\text{g}/\text{mL}$  CPF.<sup>29</sup> In this study, we illustrated that similar harmful effects are found in normal, non-cancer RPE cells.

Fluoroquinolones, including CPF, can induce considerable harmful influences on a variety of cancer cells. For instance, bladder cancer cells exposed to 100  $\mu\text{g}/$



mL CPFX showed an upregulation of BAX, the proapoptotic regulator, and arrested S/G2-M phase of the cell cycle.<sup>27</sup> Kloskowski *et al* reported similar results when they exposed three different cancer cell lines (lung cancer, melanoma and glioblastoma) to a range of concentrations (12–1092 µg/mL) of CPFX for 24–48 hours and they found that CPFX induced cell cycle arrest at the G<sub>2</sub>/M phase.<sup>13</sup> This was supported in CPFX-treated melanoma cells (COLO829) that showed inhibited DNA polymerase II and arrest of the S checkpoint of cell cycle.<sup>30</sup>

In the ARPE-19 cells, the long-term incubation with higher dosages of TETRA resulted in increased cell viability and metabolism along with lower ROS levels, suggesting positive effects on cellular health with decreased MMP. Our data suggest that the overall effect of TETRA on the ARPE-19 cells is less disruptive than CPFX. Similarly, a study by Kalghatgi *et al* reported significant CPFX-induced damaged mitochondrial morphology, along with lower levels of MMP, ATP production and metabolic activity in treated cells.<sup>8</sup> However, the same dosage of TETRA did not induce significant alterations of these mitochondrial-related features or affect ROS production<sup>31</sup> compared with untreated group.

Our gene expression studies showed that ARPE-19 cells had a differential response to TETRA with upregulation of the antiapoptosis gene BCL2, downregulation of TGF-β1 (growth and differentiation) and higher levels of IL-1β, a proinflammatory gene compared with the CPFX-treated cells. However, in a Zebrafish model (*Danio rerio*), Ding *et al* found that TETRA induced damage at several concentrations (45, 60 and 90 mg/L) after 7, 14 and 21 days with mitochondrial damage, including diminished mitochondrial cristae and mitochondrial swelling.<sup>32</sup> Another study in hepatocytes showed significant negative influences on mitochondrial calcium uniporter with treatment after TETRA derived compounds (50 µM, 22.22 g). TETRA prevented Ca<sup>2+</sup> uptake by mitochondria and then inhibited the inducement of mitochondrial permeability transmission by Ca<sup>2+</sup>.<sup>33</sup>

## CONCLUSIONS

Future investigations including in vivo studies and clinical trials are necessary in the discovery of effective treatments with antibiotics with nominal adverse effects. In conclusion, clinically adjusted dosages of CPFX and TETRA may have detrimental impacts ARPE-19 cells. We speculate that if CPFX can have these deleterious effects on the wildtype, healthy ARPE-19 cells, then it may also have deleterious effects on the older, damaged mitochondria of the AMD subjects are exposed to treatments with the fluoroquinolones.

**Acknowledgements** This work was supported by the Discovery Eye Foundation, Polly and Michael Smith, Edith and Roy Carver, Iris and B. Gerald Cantor Foundation, Max Factor Family Foundation, and NEI R01 EY0127363 (MCK). Supported in part by an Unrestricted Departmental Grant from Research to Prevent Blindness. We acknowledge the support of the Institute for Clinical and Translational Science (ICTS) at University of California Irvine.

**Contributors** NS planned the study, designed experiments, wrote the manuscript, and analysed data. LS and AN edited the manuscript. MCK, the PI, developed the concepts, edited the manuscript, provided resources for the study. KS, SA, MC and AB helped with the experimental designs and performed some of the experiments. NS is an Arnold and Mabel Beckman Retinal Degeneration Fellow. KS is a Genentech AMD Research Fellow.

**Funding** This work was supported by the Discovery Eye Foundation, Polly and Michael Smith, Edith and Roy Carver, Iris and B. Gerald Cantor Foundation, Max Factor Family Foundation, and NEI R01 EY0127363 (MCK). Supported in part by an Unrestricted Departmental Grant from Research to Prevent Blindness. We acknowledge the support of the Institute for Clinical and Translational Science (ICTS) at University of California Irvine.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All data relevant to the study are included in the article.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID ID

M Cristina Kenney <http://orcid.org/0000-0003-1765-1750>

## REFERENCES

- Mather R, Karenchak LM, Romanowski EG, *et al*. Fourth generation fluoroquinolones: new weapons in the arsenal of ophthalmic antibiotics. *Am J Ophthalmol* 2002;133:463–6.
- Nishida T, Kuse Y, Mochizuki K, *et al*. Protective effects of fluoroquinolones on UV-induced damage of cultured ocular cell lines. *Eur J Pharmacol* 2017;806:59–66.
- LeBel M. Ciprofloxacin: chemistry, mechanism of action, resistance, antimicrobial spectrum, pharmacokinetics, clinical trials, and adverse reactions. *Pharmacotherapy* 1988;8:3–30.
- Blondeau JM. Fluoroquinolones: mechanism of action, classification, and development of resistance. *Surv Ophthalmol* 2004;49 Suppl 2:S73–8.
- Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiol Mol Biol Rev* 2001;65:232–60.
- Etminan M, Forooghian F, Brophy JM, *et al*. Oral fluoroquinolones and the risk of retinal detachment. *JAMA* 2012;307:1414–9.
- Patterson DR. Quinolone toxicity: methods of assessment. *Am J Med* 1991;91:S35–7.
- Kalghatgi S, Spina CS, Costello JC, *et al*. Bactericidal antibiotics induce mitochondrial dysfunction and oxidative damage in mammalian cells. *Sci Transl Med* 2013;5:192ra85.
- Duwelhenke N, Krut O, Eysel P. Influence on mitochondria and cytotoxicity of different antibiotics administered in high concentrations on primary human osteoblasts and cell lines. *Antimicrob Agents Chemother* 2007;51:54–63.
- Nadanaciva S, Dillman K, Gebhard DF, *et al*. High-Content screening for compounds that affect mtDNA-encoded protein levels in eukaryotic cells. *J Biomol Screen* 2010;15:937–48.
- Herbold BA, Brendler-Schwaab SY, Ahr HJ. Ciprofloxacin: in vivo genotoxicity studies. *Mutat Res* 2001;498:193–205.
- Kloskowski T, Gurtowska N, Nowak M, *et al*. The influence of ciprofloxacin on viability of A549, HepG2, A375.S2, B16 and C6 cell lines in vitro. *Acta Pol Pharm* 2011;68:859–65.
- Kloskowski T, Gurtowska N, Olkowska J, *et al*. Ciprofloxacin is a potential topoisomerase II inhibitor for the treatment of NSCLC. *Int J Oncol* 2012;41:1943–9.
- Smart DJ, Halicka HD, Traganos F, *et al*. Ciprofloxacin-Induced G2 arrest and apoptosis in TK6 lymphoblastoid cells is not dependent on DNA double-strand break formation. *Cancer Biol Ther* 2008;7:113–9.
- Mohammed HHH, Abuo-Rahma GE-DAA, Abbas SH, *et al*. Current trends and future directions of fluoroquinolones. *Curr Med Chem* 2019;26:3132–49.

- 16 Golomb BA, Koslik HJ, Redd AJ. Fluoroquinolone-induced serious, persistent, multisymptom adverse effects. *BMJ Case Rep* 2015;2015:bcr2015209821.
- 17 Marchant J. When antibiotics turn toxic. *Nature* 2018;555:431–3.
- 18 Kurland CG, Andersson SG. Origin and evolution of the mitochondrial proteome. *Microbiol Mol Biol Rev* 2000;64:786–820.
- 19 Andersson SGE, Karlberg O, Canbäck B, *et al.* On the origin of mitochondria: a genomics perspective. *Philos Trans R Soc Lond B Biol Sci* 2003;358:165–79.
- 20 Tommassen J. Assembly of outer-membrane proteins in bacteria and mitochondria. *Microbiology* 2010;156:2587–96.
- 21 Andersson SG, Kurland CG. Ancient and recent horizontal transfer events: the origins of mitochondria. *APMIS Suppl* 1998;84:5–14.
- 22 Udar N, Atilano SR, Memarzadeh M, *et al.* Mitochondrial DNA haplogroups associated with age-related macular degeneration. *Invest Ophthalmol Vis Sci* 2009;50:2966–74.
- 23 Kenney MC, Atilano SR, Boyer D, *et al.* Characterization of retinal and blood mitochondrial DNA from age-related macular degeneration patients. *Invest Ophthalmol Vis Sci* 2010;51:4289–97.
- 24 D'Aurelio M, Merlo Pich M, Catani L, *et al.* Decreased Pasteur effect in platelets of aged individuals. *Mech Ageing Dev* 2001;122:823–33.
- 25 Somekh E, Douer D, Shaked N, *et al.* In vitro effects of ciprofloxacin and pefloxacin on growth of normal human hematopoietic progenitor cells and on leukemic cell lines. *J Pharmacol Exp Ther* 1989;248:415–8.
- 26 Beberok A, Rzepka Z, Respondek M, *et al.* Gsh depletion, mitochondrial membrane breakdown, caspase-3/7 activation and DNA fragmentation in U87MG glioblastoma cells: new insight into the mechanism of cytotoxicity induced by fluoroquinolones. *Eur J Pharmacol* 2018;835:94–107.
- 27 Aranha O, Wood DP, Sarkar FH. Ciprofloxacin mediated cell growth inhibition, S/G2-M cell cycle arrest, and apoptosis in a human transitional cell carcinoma of the bladder cell line. *Clin Cancer Res* 2000;6:891–900.
- 28 Herold C, Ocker M, Ganslmayer M, *et al.* Ciprofloxacin induces apoptosis and inhibits proliferation of human colorectal carcinoma cells. *Br J Cancer* 2002;86:443–8.
- 29 Holtom PD, Pavkovic SA, Bravos PD, *et al.* Inhibitory effects of the quinolone antibiotics trovafloxacin, ciprofloxacin, and levofloxacin on osteoblastic cells in vitro. *J Orthop Res* 2000;18:721–7.
- 30 Beberok A, Wrześniok D, Minecka A, *et al.* Ciprofloxacin-mediated induction of S-phase cell cycle arrest and apoptosis in COLO829 melanoma cells. *Pharmacol Rep* 2018;70:6–13.
- 31 Kohanski MA, Dwyer DJ, Hayete B, *et al.* A common mechanism of cellular death induced by bactericidal antibiotics. *Cell* 2007;130:797–810.
- 32 Ding L, Zang L, Zhang Y, *et al.* Joint toxicity of fluoroquinolone and tetracycline antibiotics to zebrafish (*Danio rerio*) based on biochemical biomarkers and histopathological observation. *J Toxicol Sci* 2017;42:267–80.
- 33 Kholmukhamedov A, Czerny C, Hu J, *et al.* Minocycline and doxycycline, but not tetracycline, mitigate liver and kidney injury after hemorrhagic shock/resuscitation. *Shock* 2014;42:256–63.