



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Growth and photosynthetic characteristics of sweet potato (*Ipomoea batatas*) leaves grown under natural sunlight with supplemental LED lighting in a tropical greenhouse

Jie He*, Lin Qin

Natural Sciences and Science Education Academic Group, National Institute of Education, Nanyang Technological University, 1 Nanyang Walk, 637 616, Singapore

ARTICLE INFO

Keywords:
LED lighting
Leaf growth
Photosynthetic performance
Sweet potato leaves
Tropical greenhouse

ABSTRACT

Leaf growth and photosynthetic characteristics of sweet potato (*Ipomoea batatas* var. Biru Putih) grown under different light quantities were studied in a tropical greenhouse. The stem cuttings of *I. batatas* with adventitious roots were grown hydroponically under (1) only natural sunlight (SL); (2) SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$ (SL + L-LED); and (3) SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ (SL + H-LED). One week after emergence, all leaves had similar area and water content. However, leaf fresh weight and dry weight were significantly higher in plants grown under SL+L-LED and SL + H-LED than under SL due to their thicker leaves reflected by the lower specific leaf area. Plants grown under SL had significantly lower concentrations of total chlorophyll (Chl) and total carotenoids (Car) but higher Chl a/b ratio than under SL + L-LED and SL + H-LED. However, all plants had similar Chl/Car ratios. Although midday F_v/F_m ratio was the lowest in leaves grown under SL+ H-LED followed by SL + L-LED and SL, predawn F_v/F_m ratios of all leaves were higher than 0.8. Increasing growth irradiance with supplemental LED resulted in higher electron transport rate and photochemical quenching but lower non-photochemical quenching compared to those of plants grown under SL. Measured under their respective growth irradiance in the greenhouse, attached leaves grown under SL + L-LED and SL+H-LED had significantly higher photosynthetic CO_2 assimilation rate and stomatal conductance than under SL. However, measuring the detached leaves at 25°C in the laboratory, there were no significant differences in PS II and Cyt b_6/f concentrations although light- and CO_2 -saturated photosynthetic O_2 evolution rates were slightly higher in leaves grown under SL + H-LED than under SL. Impacts of supplemental LED on leaf growth and photosynthetic characteristics were discussed.

1. Introduction

Sweet potato (*Ipomoea batatas* L.) is widely grown in developing countries due to its low production cost and its high adaptability (Lin et al., 2007; Mekonen et al., 2015). Most people grow sweet potato as food and mainly consume their modified root tubers (Yoshimoto et al., 2002). Although the leaves of sweet potato have been neglected, they are consumed as fresh vegetables in tropical areas in Southeast Asia (Nwinyi, 1992) because both leaf blades and petioles are rich in protein, dietary fiber, vitamins, antioxidants, essential fatty acids and minerals (Ishida et al., 2000; Johnson and Pace, 2010). Several studies

have demonstrated that sweet potato leaves inhibit mutagenicity, diabetes, leukemia and viruses and the growth of colon and stomach cancer cells (Yoshimoto et al., 2002; Kurata et al., 2007; Ludvik et al., 2008).

Due to limited land in Singapore, local farming currently accounts for only 10 per cent of the leafy vegetables consumed. The maintenance of food security, especially the supply of vegetable is an increasing challenge for Singapore where natural resources are limited. Furthermore, the disconnect between supply and demand is the result of the global food supply chains being disrupted in unprecedented ways due to the COVID-19 pandemic. Back to March 2019, the Environment

Abbreviations: A, Photosynthetic CO_2 assimilation rate; Car, carotenoids; Chl, Chlorophyll; C_i , Internal CO_2 concentration; Cyt b_6/f , Cytochrome b_6/f complex; DW, Dry weight; ETR, Electron transport rate; FW, Fresh weight; gs, Stomatal conductance; LED, Light emitting diode; NPQ, Non-photochemical quenching; P_{max} , Photosynthetic capacity; P_N , Net photosynthetic O_2 evolution rate; PPFD, Photosynthetic photon flux density; qp, Photochemical quenching; RuBP, ribulose-1,5-bisphosphate; Rubisco, ribulose 1,5-bisphosphate carboxylase/oxygenase; SL, natural sunlight; SL + L-LED, natural sunlight with supplemental LED lighting at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, natural sunlight with supplemental LED lighting at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$; SLA, specific leaf area; T_r , Transpiration

* Corresponding author.

E-mail address: jie.he@nie.edu.sg (J. He).

<https://doi.org/10.1016/j.jplph.2020.153239>

Received 27 May 2020; Received in revised form 4 July 2020; Accepted 9 July 2020

Available online 24 July 2020

0176-1617/ © 2020 Elsevier GmbH. All rights reserved.

and Water Resources Minister, Singapore announced the ambitious “30 by 30” goal to produce 30 per cent of Singapore’s nutritional needs locally by 2030 (Ai-Lien, 2019). Sweet potato leaves are considered an indigenous and tropical leafy vegetable, which could play an important role in alleviating the shortage of leafy vegetables as they grow quickly in the tropics under warm temperature and humid conditions (An et al., 2003). They have much higher annual yield than that of other green vegetables and could be harvested several times a year. Sweet potato are mainly grown over a broad range of environment. In Singapore sweet potato leaves are commonly cultured in outdoor soil farms. It was reported that high light intensity enhanced growth of sweet potato plants and thus, shade conditions should be avoided (Oswald et al., 1994). The sweet potato plant is also considered to be a drought tolerant crop (Ghuman and Lal, 1983). However, drought is a major environmental constraint for sweet potato production in the tropical area as its growth and development are significantly influenced by soil moisture (Yooyongwecha et al., 2013).

Soilless cultures such as hydroponics and aeroponics are increasingly adopted as major technological components in the modern greenhouses to replace community gardens and traditional outdoor soil farms in Singapore (He, 2015). Today, in Singapore, all kinds of leafy vegetables are grown all year round in the greenhouse using soilless culture systems with adequate water and nutrient supplies, which is not affected by drought conditions. Apart from water, light intensity critically affects plant growth. In the past two decades, Singapore has been frequently experiencing increasingly unpredictable cloudy and hazy weather (Nobre et al., 2016), resulting in lowered light intensity which reduced crop productivity (Jones, 2006). We have previously reported that in Singapore, when lettuce (He et al., 2011) and *Brassica alboglabra* (Chinese broccoli) plants (He et al., 2019b) were grown under low light during the haze episodes in the greenhouse, lower photosynthetic rate, stomatal conductance and productivity were measured. To circumvent the problem of insufficient sunlight, in another study with lettuce plants, light emitting diode (LED) lighting was supplemented to low sunlight intensity in the greenhouse (Choong et al., 2018; He et al., 2019a). Leaf expansion rate was faster in both heat resistant and heat sensitive recombinant inbred lines (RILs) of lettuce grown under natural sunlight with supplemental LED. However, impacts of supplementary LED lighting on shoot and root productivity and photosynthetic performance such as photosynthetic light use efficiency and maximal oxygen evolution rate seemed to be genotype-dependent due to their different sensitivities to heat (Choong et al., 2018). In the study with *B. alboglabra* (Chinese broccoli), we have found that plants grown under shade with supplemental LED lighting improved photosynthetic CO₂ assimilation, stomatal conductance and productivity (He et al., 2019). As mentioned earlier, sweet potato leaves are tropical leafy vegetables and have fast growth rates in the tropics under warm and humid conditions (An et al., 2003). The nutrient-rich leaf blade and petiole make it as one of the suitable vegetable crops to achieve the Singapore’s goal to produce 30 per cent of nutritional needs locally by 2030. However, little is known about its photosynthetic characteristics when grown under different light intensities using soilless culture such as hydroponics. We hypothesize that plant growth should increase with increasing of growth irradiance, with leaf traits adjustment that enhance light capture and carbon fixation in the tropical greenhouse. Therefore, this study aimed to investigate the effects of two different supplementary LED lightings with photosynthetic photon flux density (PPFD) of 150 and 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ respectively on the leaf growth of sweet potato leaves, *I. batatas* (var. Biru Putih) in the tropical greenhouse. Impacts of supplementary LED quantity on photosynthetic light use efficiency measured by chlorophyll fluorescence, the functions of PS II and Cyt b₆f, and photosynthetic gas exchanges were also studied.

2. Materials and methods

2.1. Plant materials and experimental design

Sweet potato leaves, *I. batatas* (var. Biru Putih), were purchased from one of the local farms (KOK FAH Technology Farm Pte Ltd). Apical stems were cut into 15 cm pieces. Roots were induced from cuttings grown hydroponically under modified half strength Netherlands Standard Composition nutrient solution (details are given below) for 1 week. After root induction, the stem cuttings with adventitious roots were grown hydroponically in greenhouse under three different quantities of lights: (1) only natural sunlight (SL) with average maximum PPFD of 800 $\mu\text{mol m}^{-2} \text{s}^{-1}$; (2) SL with supplemental LED lighting at a PPFD of 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (SL + L-LED); and (3) SL with supplemental LED lighting at a PPFD of 300 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (SL + H-LED). The photoperiod of supplemental LED lighting (Dissis LED Lighting Technology, Singapore) was 12-h (from 0700 h to 1900 h) provided by a combination of red- (633 nm and 656 nm) and blue-LED (463.5 nm) lightings in the ratio of 9:1. All the light intensities were measured by holding PAR quantum sensor with a reading unit (SKP 215 and 200, Skye Instruments Ltd, Llandrindod Wells, UK). All treatments were supplied with modified full strength Netherlands Standard Composition nutrient solution (Douglas, 1985) with $2.0 \pm 0.2 \text{ mS cm}^{-1}$ conductivity and pH 6.0 ± 0.2 . The modified full strength Netherlands Standard Composition solution had the following composition: N, 191.89 ppm; P, 33.29 ppm; K, 309.00 ppm; Ca, 210.29 ppm; Mg, 60.04 ppm; S, 125.45 ppm; Fe, 9.36 ppm; B, 0.105 ppm; Mn, 0.238 ppm; Zn, 0.014 ppm; Cu, 0.015 ppm; Mo, 0.408 ppm; Na, 3.860 ppm. The fluctuating ambient temperatures of 24–43 °C and relative humidity of 26–96 % were recorded using DataHog2 (Skye Instruments Ltd, UK).

2.2. Measurements of leaf fresh weight (FW) and dry weight (DW), leaf area, specific leaf area (SLA) and leaf water content

The just-emerged leaves of *I. batatas* plants were labeled 27 days after transplanting and they were harvested one week after emergence from 0730 h to 0800 h. The cut leaves were kept in sealed plastic bags with sheets of damp paper towel and brought back to lab. The FW of leaf (blade only) was first recorded before measuring their areas using a leaf area meter (WinDIAS3 Image Analysis system). All leaves were then wrapped individually in pre-weighed aluminium foil, dried at 80 °C for at least four days, before re-weighing them to obtain DW. Specific leaf area (SLA) was determined as L_A/L_{DW} where L_A = leaf area (cm²) and L_{DW} = leaf DW(g) (Hunt et al., 2002). Leaf water content (LWC) was determined as $(L_{FW} - L_{DW})/L_{FW}$ where L_{FW} = leaf FW (g).

2.3. Measurements of Chl and carotenoids (Car) pigments

For each measurement, two leaf discs (1 cm in diameter) were cut from the youngest fully expanded leaves 15 days after transplanting and soaked in 2.5 ml of *N,N*-dimethylformamide (*N,N*-DMF, Sigma Chemical Co.) in darkness for 48 h at 4 °C. The absorption of pigments were measured using a spectrophotometer (UV-2550 Shimadzu, Japan) at 647 nm, 664 nm and 480 nm respectively. Chl a, Chl b and Car concentrations were calculated as described by Wellburn (1994).

2.4. Measurements of predawn and midday F_v/F_m ratio

The maximum photochemical efficiency of PS II was estimated in dark-adapted samples by the F_v/F_m ratio. Predawn and midday F_v/F_m ratios were measured from the attached youngest fully expanded leaves in the greenhouse before photoperiod and during mid-photoperiod on sunny days after transplanting for 34 days using the Plant Efficiency Analyser (Hansatech Instruments, UK) according to He et al. (2001).




Fig. 1. Leaf area (A) and SLA (B) of the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 5$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.

2.5. Measurements of electron transport rate (ETR), photochemical quenching (qP) and non-photochemical quenching (NPQ)

After 30 days of transplanting, the youngest fully expanded leaves were harvested between 0900 h–1000 h and ETR, qP and NPQ were determined at 25°C in the laboratory. Prior to measurements, the leaves were pre-darkened for 15 min. By using the IMAGING PAM MAXI (Walz, Effeltrich, Germany), images of fluorescence emission were digitized within the camera and transferred via ethernet interface (GigEVision®) to the PC for storage and analysis. Measurements and calculations of ETR, qP and NPQ were described previously (He et al., 2017b).

2.6. Measurements of light response curves of net photosynthetic O_2 evolution rate (P_N), PS II concentration and Cyt b_6/f concentration

These parameters were measured according to He and Chow (2003), and Zhu et al. (2017). O_2 evolution from leaf discs was measured in a gas-phase oxygen electrode (Hansatech, King's Lynn, UK) chamber maintained at 25°C . Each leaf disc was 3.4 cm^2 in area, punched from the similar part of the youngest fully expanded *I. batatas* leaves grown under different light conditions. The sample chamber contained 1% CO_2 supplied by fabric matting moistened with 1 M $\text{NaHCO}_3/\text{Na}_2\text{CO}_3$ (pH 9). Two illumination regimes were used: (1) repetitive flash illumination with saturating, single-turnover flashes, or (2) continuous white light from light emitting diodes. First, repetitive flash illumination of the leaf sample with saturating, single-turnover xenon flashes (at 10 Hz) was performed to obtain a net rate of O_2 evolution on a leaf area basis. Following an initial dark equilibration for 10 min, the repetitive flash illumination was applied for 4 min, followed by 4 min darkness. This was followed by a second cycle of flashes and darkness. The average dark drift in the signal before and after repetitive-flash illumination was subtracted algebraically from the net rate of O_2 evolution during flash illumination to obtain the gross rate of flash-induced O_2 evolution. A small heating artefact signal due to flash illumination was obtained by substituting a green paper disc for a leaf disc, and was corrected for. The limitation of linear electron transport by PS I was minimized by the use of background far-red light. The ratio of the gross rate of O_2 evolution to the flash frequency was used to derive the PS II concentration on a leaf area basis (p), assuming that after four flashes, each active PS II evolves one O_2 molecule (Chow et al., 1991). Second, after repetitive-flash illumination, a light response curve of P_N was measured under continuous white light. The leaf disc was illuminated at 15 different light intensities, starting from the lowest PPFD of $0\text{--}1870 \mu\text{mol m}^{-2} \text{s}^{-1}$. The leaf disc was illuminated at each PPFD over several minutes until steady-state of photosynthetic O_2 evolution rate was obtained. The saturating, continuous irradiance ($1870 \mu\text{mol m}^{-2} \text{s}^{-1}$) was used to determine the photosynthetic capacity (P_{max}). The post-illumination drift was subtracted algebraically from the steady-state net O_2

evolution rate at PPFD of $1870 \mu\text{mol m}^{-2} \text{s}^{-1}$ to yield the gross O_2 evolution rate, P_{max} . For calibration of the oxygen signals, 1 ml of air at 25°C (taken to contain $8.584 \mu\text{mol O}_2$) was injected into the gas-phase O_2 electrode chamber.

Calculation of the Cyt b_6/f concentration: After measurements of p and P_{max} , the Cyt b_6/f concentration (f) was calculated from the equation, $P_{\text{max}} = 1/[(0.022/f) + (0.004/p)]$, all parameters being on a leaf area basis. The Cyt b_6/f concentration, calculated from the two activity measurements, represents the functional Cyt b_6/f concentration in leaves (Zhu et al., 2017)

2.7. Measurements of photosynthetic CO_2 assimilation rate (A), stomatal conductance (g_s), internal CO_2 concentration (C_i) and transpiration (T_r)

A, g_s , C_i and T_r were measured from the youngest fully expanded attached leaves under growth irradiances in the greenhouse using the LI-COR Portable Photosynthetic System (LI-6400, Bioscience, USA) from 1030 h to 1230 h. During the measurement, the average ambient CO_2 concentration in the greenhouse was $383 \pm 6 \mu\text{mol mol}^{-1}$, the relative humidity was $40 \pm 10\%$ and the temperature was $39 \pm 4^\circ\text{C}$. The measurement was performed twice on both the 44th and 45th days after transplanting. The results presented were the means of data collected in these two days.

2.8. Statistical analysis

One-way analysis of variance (ANOVA) was used to test for significant differences of variances crossed with the three different treatments. LSD multiple comparison tests were used to discriminate between means of the different treatments, where means with $p < 0.05$ has significant differences (IBM, SPSS Version 25).

3. Results

3.1. Leaf growth, leaf productivity and leaf water content

It was observed that all leaves were fully expanded after one week emergence regardless of growth irradiances. Fig. 1 shows an average leaf area and a SLA of fully-expanded leaves developed under different growth irradiances. Although the average leaf area of sweet potato leaves grown under SL + H-LED (56 cm^2) was slightly larger than those grown under SL + L-LED (50 cm^2) and SL (51 cm^2), statistically, there were no significant differences in their values (Fig. 1A). However, sweet potato leaves grown under SL had significantly higher SLA compared to those grown under SL + L-LED and SL + H-LED (Fig. 1B). For leaf FW and leaf DW, they were significantly higher in sweet potato plants grown under SL + L-LED and SL + H-LED than under only natural SL (Fig. 2A and B). However, all leaves had similar water content (Fig. 2C).




Fig. 2. Leaf FW (A), leaf DW (B) and leaf water content of the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 5$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.

3.2. Photosynthetic pigments

Sweet potato leaves grown under SL + H-LED had the highest concentrations of Chl a, Chl b, total Chl and total Car followed by those grown under SL + L-LED. For plants grown under SL, their leaves had the lowest Chl a, Chl b, total Chl and total Car concentrations (Fig. 3A–D). However, Chl a/b ratio of sweet potato leaves grown under SL was significantly higher than those grown under SL + L-LED and SL + H-LED (Fig. 3E). There were no significant differences in Chl/Car

ratios among the leaves grown under different light conditions (Fig. 3F).

3.3. Photosynthetic light utilization efficiency measured by F_v/F_m ratio, ETR, qP and NPQ

Fig. 4 shows the predawn and midday F_v/F_m ratios measured from the attached leaves on the same sunny day in the greenhouse. The measurements were repeated once on another sunny day and similar




Fig. 3. Chl a concentration (A), Chl b concentration (B), total Chl concentration (C), total Car concentration (D), Chl a/b ratio (E) and Chl/Car ratio (F) of the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 6$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.




Fig. 4. Pre-dawn (A) and midday (B) F_v/F_m ratios on a sunny day measured from the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 8$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.

results were obtained. All leaves had their predawn F_v/F_m ratios greater than 0.8 although leaves grown under SL had F_v/F_m ratio significantly higher than those grown under SL + L-LED and SL + H-LED (Fig. 4A). Leaves grown under SL + H-LED had significantly lower midday F_v/F_m ratio (0.689) than those grown under SL (0.750) and SL + L-LED (0.744) (Fig. 4B).

The youngest fully expanded leaves were also harvested to measure the light response curves of ETR, qP and NPQ, starting from the lowest PPFD of $0\text{--}1501 \mu\text{mol m}^{-2} \text{s}^{-1}$. The measurements were repeated once on a different day and similar results were obtained. The readings of ETR increased as PPFDs increased from $0\text{--}1501 \mu\text{mol m}^{-2} \text{s}^{-1}$ for all leaves (Fig. 5A). All leaves had similar values of ETR when they were measured under PPFDs $< 111 \mu\text{mol m}^{-2} \text{s}^{-1}$. However, leaves grown under SL had significantly lower ETR compared to those grown under SL + L-LED and SL + H-LED when they were measured under PPFDs from 146 to $1501 \mu\text{mol m}^{-2} \text{s}^{-1}$. Measured under the two highest PPFDs of 1076 and $1501 \mu\text{mol m}^{-2} \text{s}^{-1}$, leaves grown under SL + H-LED had the highest ETR, followed by those grown under SL + L-LED and the leaves grown under SL had the lowest ETR under these two PPFDs (Fig. 5A). The values of qP decreased with increasing PPFDs from $0\text{--}1501 \mu\text{mol m}^{-2} \text{s}^{-1}$ for all leaves (Fig. 5B). All leaves had similar decrease rates of qP when they were measured under PPFDs from 0 to $56 \mu\text{mol m}^{-2} \text{s}^{-1}$. However, decreases in qP were significantly faster in leaves grown under SL than under SL + L-LED and SL + H-LED when measured under PPFDs above $81 \mu\text{mol m}^{-2} \text{s}^{-1}$ except for the higher PPFD of $1501 \mu\text{mol m}^{-2} \text{s}^{-1}$, under which all leaves had similar lowest level of qP (Fig. 5B). Similar to the changes of ETR, NPQ of all leaves increased with increasing PPFDs from $0\text{--}1501 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 5C). When measured under PPFDs $< 146 \mu\text{mol m}^{-2} \text{s}^{-1}$, all leaves had similar levels of NPQ. Leaves grown under SL + H-LED had significantly lower NPQ compared to those grown under SL + L-LED and SL under PPFDs from 146 to $611 \mu\text{mol m}^{-2} \text{s}^{-1}$. However, under the two highest PPFDs of 1076 and $1501 \mu\text{mol m}^{-2} \text{s}^{-1}$, NPQ values were similarly higher in leaves grown under SL + L-LED and SL + H-LED than under SL (Fig. 5C).

3.4. Light response curves of P_N , PS II and Cyt b_6/f concentrations of detached leaves measured in the laboratory

Fig. 6 shows light response curves P_N , PS II and Cyt b_6/f concentrations measured from the same detached leaves in the laboratory at 25°C . P_N increased with increasing PPFDs similarly in all leaves from 0 to $602 \mu\text{mol m}^{-2} \text{s}^{-1}$, and started to saturate between $602\text{--}808 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 6A), after which, slow increases of P_N were still observed from 808 to $1435 \mu\text{mol m}^{-2} \text{s}^{-1}$ in leaves grown under SL + H-LED. Thus, the light response curve of P_N measured from leaves grown under SL + H-LED was above those grown under SL + L-LED and SL. Measured under the two highest PPFDs of 1435 and $1870 \mu\text{mol m}^{-2} \text{s}^{-1}$, leaves grown under SL + H-LED had significantly higher P_N than those grown under SL + L-LED while leaves grown under SL + H-LED and SL had similar P_N (Fig. 6A). However, there were no significantly differences in




Fig. 5. Light response curves of ETR (A), qP (B) and NPQ (C) measured from the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 4$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.




Fig. 6. Light response curves of photosynthetic O_2 evolution (A), PS II concentration (B) and Cyt b_6f concentration (C) measured from the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 3$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$.

PS II (Fig. 6B) and Cyt b_6f (Fig. 6C) concentrations among the sweet potato leaves grown under different light conditions.

3.5. A , g_s , C_i , T_r of attached leaves measured in the greenhouse

Fig. 7A and B show the results of A and g_s of sweet potato leaves measured under their respective growth irradiance with similar but much higher values obtained from leaves grown under SL + L-LED and SL + H-LED than under SL. The C_i was significantly lower in leaves grown under SL + H-LED compared to those grown under SL + L-LED and SL (Fig. 7C). However, all leaves had similar T_r (Fig. 7D).

4. Discussion

Although sweet potato leaves have been consumed in Singapore as a high nutritional leafy vegetable, they are mainly cultivated in outdoor soil farms. The supply of sweet potato leaves to the vegetable markets is not secured due to the fact that its productivity is influenced by

variation in sunlight availability and intensity. To the best of our knowledge, this was the first project to grow sweet potato leaves in the tropical greenhouse using hydroponic systems under supplemental LED lighting to prevailing natural sunlight. Light is the most crucial factor for plant growth and development as well as the efficiency of the photosynthesis (Violet-Chabrand et al., 2017; Gommers, 2020). Plants can adjust their morphological and physiological traits such as leaf size, SLA, leaf mass and Chl content when they are subjected to changing light conditions (Liu et al., 2016; He et al., 2017a). This study was carried out in the tropical greenhouse from the middle of February to early April 2020, when the weather was warm and sunny. The average highest ambient temperature was about $34\text{--}36^\circ\text{C}$ under full sunlight and maximum PPFD in an open field on sunny days outside the greenhouse ranged from 1400 to $1600 \mu\text{mol m}^{-2} \text{s}^{-1}$ for at least 4 h from 1100 h to 1500 h. However, the average maximum PPFD inside the greenhouse was round $700\text{--}800 \mu\text{mol m}^{-2} \text{s}^{-1}$, which was 50 % of full sunlight but the average ambient temperature could be as high as $38\text{--}40^\circ\text{C}$ with the highest of 43°C during midday for at least 4 h. It was a surprise to observe that inside the hot greenhouse, the leaves of all sweet potato plants grew very fast and were very healthy, especially those grown under SL supplemented with constant PPFDs of either 150 (SL + L-LED) or $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ (SL + H-LED), for a 12 h photoperiod. These observations were supported by the results shown in Figs. 1 and 2. Although there was no significant difference in leaf area of the youngest fully expanded leaves grown under different light conditions one week after emergence (Fig. 1A), leaf FW and DW were significantly higher in leaves grown under SL + L-LED and SL + H-LED than under SL alone (Fig. 2A and B). All leaves had similar water content (Fig. 2C) and thus, higher leaf FW and DW resulted from the greater thickness of those grown under supplemental LED lighting, which was supported by the lower SLA (Fig. 1B). Low SLA is associated with greater leaf thickness as SLA is measured as the ratio of leaf area to leaf dry mass (Hunt et al. 2002). It was previously reported by our studies that both light quantity and quality affected SLA of leafy vegetables within the species (Choong et al., 2018; He et al., 2019).

Grown under low-light conditions, plants usually develop leaves with higher SLA compared to those grown under high-light conditions (He et al., 2017a; Violet-Chabrand et al., 2017; Zhang et al., 2019). According to Poorter et al. (2018), light-capture-related traits such as SLA and Chl content are strongly related to light, and therefore vary mostly within species. The higher SLA of sweet potato leaves grown under low-light conditions such as only SL (Fig. 1B) could be interpreted as a mechanism to optimize light harvesting. However, under high-light conditions such as SL + H-LED conditions, lower SLA (Fig. 1B) could help plants to increase the efficiency of light capture (Evans and Poorter, 2001; Liu et al., 2016). In the study with soybean, Fan et al. (2018) and Feng et al. (2019) reported that Chl a, Chl b, total Chl and Car contents increased with the increase in light intensity, which were directly associated with leaf thickness. The results of this study with potato leaves grown under SL + L-LED and SL + H-LED conditions having higher Chl a, Chl b, total Chl and Car concentrations on a per area basis (Fig. 3A–D) are consistent with those previous results (Fan et al., 2018; Feng et al., 2019). However, Anderson et al. (1988) demonstrated that it was not the Chl concentration per leaf area but the ratio of Chl a/b showed a close correlation to the growth irradiance. In this study, when growth irradiance was increased by supplementing LED lighting to SL, Chl a/b ratio of sweet potato leaves decreased (Fig. 3E). Increasing light intensity decreased Chl a/b ratio was also reported in soybean (Feng et al., 2019). These results are not in accordance with the usually increased Chl a/b ratio of plants grown under high-irradiance compared to those grown under low-irradiance (Evans and Poorter, 2001). Zivcak et al. (2014) suggested that effects of light intensity on Chl a/b ratio is not a universal phenomenon and the dependence of Chl a/b ratio on light intensity is strongly correlated to plant species. In this study, the supplemental LED-lighting was provided by a combination of red- and blue-LED lighting in the ratio of 9:1. The




Fig. 7. Photosynthetic CO_2 assimilation rate, A (A), stomatal conductance, g_s (B), internal CO_2 concentration, C_i (C) and transpiration, T_r (D) measured from the youngest fully expanded leaves of *I. batatas* grown under different light conditions. Means with different letters are statistically different ($P < 0.05$; $n = 6$) as determined by LSD multiple comparison test. SL, Sunlight; SL + L-LED, SL with supplemental LED at a PPFD of $150 \mu\text{mol m}^{-2} \text{ s}^{-1}$; SL + H-LED, SL with supplemental LED at a PPFD of $300 \mu\text{mol m}^{-2} \text{ s}^{-1}$.

effects of light conditions on Chl a/b ratio could result from both the light intensity and light quality, which merits our further study. Car concentration of sweet potato leaves increased by 8 and 27 %, under SL + L-LED and SL + H-LED respectively, compared to those grown under SL alone (Fig. 3D). Generally, Car concentration was higher for sun-acclimated leaves compared to that of shade-grown leaves (Demmig-Adams and Adam, 1996; Lichtenthaler, 2007). Carotenoids play important roles in photosynthesis (Pogson et al., 2005) and protect plants from the harmful effects of excess exposure to light by maintaining proper Chl/Car ratio (or Car/Chl ratio) for optimal photosynthesis and photoprotection (Hashimoto et al., 2016). All sweet potato leaves had similar Chl/Car ratios as those grown under higher growth irradiance increased the concentrations of both total Chl and total Car concentrations (Fig. 3C and D). In the study with barley leaves (*Hordeum vulgare* L.), Zivcak et al. (2014) also reported that no significant changes were observed in the Chl/Car ratio between sun and shade leaves.

Biosynthesis of Chl increases with increasing light intensity (Björkmann, 1981). However, under adverse environmental conditions such as high temperature, high light inhibits Chl formation due to photoinhibition (He et al., 1996). In this study, all leaves had predawn F_v/F_m ratio > 0.8 (Fig. 4A), indicating that no chronic photoinhibition occurred in any plants. However, dynamic photoinhibition occurred in all leaves during midday with F_v/F_m ratios < 0.8 . The greatest decrease of midday F_v/F_m ratio was observed in leaves grown under SL + H-LED, which were exposed to the highest PPFD about $1150 \mu\text{mol m}^{-2} \text{ s}^{-1}$. The midday F_v/F_m ratio remained higher for leaves of plants grown under SL with the lowest midday PPFD ($700\text{--}800 \mu\text{mol m}^{-2} \text{ s}^{-1}$). Grown under SL + L-LED subjected to a PPFD of $1000 \mu\text{mol m}^{-2} \text{ s}^{-1}$, the decrease of midday F_v/F_m ratio was in-between (Fig. 4B). Similar to the results of this study, grown under natural conditions with dynamic fluctuations in light, high PPFDs coupled with high temperature during midday caused dynamic photoinhibition but no sustained chronic photoinhibition were observed in certain native orchid species in Singapore (He et al., 2017a; Tay et al., 2019).

Chls are essential molecules that catch light energy to drive photosynthetic electron transfer (Fromme et al., 2003). Sweet potato leaves grown under high-irradiance had thicker leaves and higher Chl concentrations are associated with higher ETR compared to those grown under low-irradiance (Fig. 5A). When measurements were carried out from 801 to the highest $1501 \mu\text{mol m}^{-2} \text{ s}^{-1}$, which were within the maximum ranges of PPFD under which all leaves were developed, the ETR readings were 80–98 % and 55–75 % respectively higher in SL + H-LED and SL + L-LED leaves than that in leaves grown under SL (Fig. 5A). Not only ETR but also qP was modified by the light conditions, with slower declines in qP in leaves grown under SL + L-LED and SL + H-LED compared to that in leaves grown under SL except for the measurements at the higher PPFD of $1501 \mu\text{mol m}^{-2} \text{ s}^{-1}$ (Fig. 5B). The values of qP were significantly higher for the leaves grown under high-irradiance such as SL + L-LED and SL + H-LED than those under lower irradiance of SL inside the greenhouse, indicating that high-irradiance improve photosynthetic light use efficiency (Feng et al., 2019). It was reported that high growth irradiance increased ETR and qP while decreased the NPQ (Zivcak et al., 2014; Feng, 2019). These were also observed in this study with sweet potato leaves grown under SL + H-LED had low NPQ when measurements were made from 146 to $611 \mu\text{mol m}^{-2} \text{ s}^{-1}$ compared to those grown under SL. However, decreased NPQ was not observed in leaves grown under SL + L-LED (Fig. 5C).

High growth irradiance enhances ETR and qP that could increase photosynthetic rate by improving the energy transport from PSII to PSI in tobacco leaves (Yamori et al., 2010) and soybean leaves (Yang et al., 2018). It was also reported earlier that pea leaves grown under high-light had higher concentrations of PS II and Cyt b_6/f and thus a higher capacity for electron transport and photosynthetic oxygen evolution compared to those grown under low-light (Leong and Anderson, 1984; Chow and Anderson, 1987). Similar results were also reported in spinach (Chow and Hope, 1987), *Alocasia macrorrhiza* (Chow et al., 1988) and *Hordeum vulgare* (De la Torre and Burkey, 1990). However, in this study, sweet potato leaves grown under SL + L-LED and SL + H-LED increased ETR and qP while the concentrations of PS II (Fig. 6B) and Cyt

b_6f (Fig. 6C) remained unchanged compared to those grown under SL. Higher ETR and qP observed in this study may be due to the tuning of the amount of active PSII reaction centres and regulating the electron transfer by the Cyt b_6f complex (Tikkanen et al., 2012) rather than the modified concentrations. As all leaves in this study were developed under the maximum PPFDs about $700\text{--}800 \mu\text{mol m}^{-2} \text{s}^{-1}$ where P_N started to saturate (Fig. 6A), they may have produced maximum amount of PS II and Cyt b_6f complex. On the other hand, P_N was measured at the temperature (25°C) lower than the maximum growth temperature of sweet potato leaves, which may also be another factor affecting the values of P_N (to be discussed in the next section). In our future study, corrections among ETR, qP , PS II, Cyt b_6f and P_N in sweet potato leaves should be carried out under different light intensity and different leaf temperatures.

Many studies have shown that PS II and Cyt b_6f may be the site of the rate-limiting step in the electron transport (Heber et al., 1988; Eichelmann et al., 2000). The concentration of Cyt b_6f is the main rate-limited factor that determines light and CO_2 -saturated photosynthetic capacity (Tikkanen et al., 2012; Zhu et al., 2017). In this study, there was a slightly higher P_N measured from leaves grown under SL + H-LED at the highest PPFD of $1870 \mu\text{mol m}^{-2} \text{s}^{-1}$ compared to that of leaves grown under SL (Fig. 6A), which could be due to the similar concentrations of Cyt b_6f (Fig. 6B) among all leaves.

Although all detached leaves had similar P_N measured at 25°C in the laboratory under saturated CO_2 (Fig. 6A), there were significant differences in A , g_s , and C_i measured from attached leaves under their respective growth irradiance in the greenhouse (Fig. 7). Generally, leaves developed under high-irradiance are thicker than those grown under low-irradiance and thus enhance light use efficiency for carbon fixation (Terashima et al., 2006). In their paper, Terashima et al. (2006) confirmed that there were sufficient mesophyll surfaces in the thicker sun leaves for CO_2 dissolution and transport to the chloroplasts. Thicker sun leaves of C_3 plants could maintain the CO_2 concentration in the chloroplast as high as possible for ribulose 1,5-bisphosphate carboxylase/oxygenase (Rubisco) (Terashima et al., 2006). Furthermore, thicker leaves accumulated more photosynthetic enzymes on a leaf area basis and thus contributed to greater CO_2 fixation capacity of high-light grown leaves (Evans and Poorter, 2001). The results of this study support those earlier studies as photosynthetic CO_2 assimilation rate, A was 31 and 40 % higher in SL + L-LED and SL + H-LED, respectively than that of SL (Fig. 7A). Generally, plants grown at higher temperature have a higher optimal temperature of photosynthetic rate (Yamasaki et al., 2002). In many species, the optimal temperature of photosynthetic rate increases with increasing growth temperature (Hikosaka et al., 2006). These earlier studies explain why the values of A (Fig. 7A) measured from the attached leaves in the greenhouse with their leaf temperatures as high as $38\text{--}43^\circ\text{C}$ under ambient CO_2 were higher than those of P_N (Fig. 6A) measured from detached leaves at 25°C in the presence of a saturating CO_2 concentration in the laboratory. It has been reported that g_s is sensitive to temperature, humidity and light (Bunce, 2001; Lawson, 2009). Generally, high temperature promotes stomatal opening to facilitate leaf cooling. In this study, there were small variations in leaf temperature and humidity among sweet potato leaves as they were grown in the same area of the same greenhouse. However, the values of g_s were significantly higher in sweet potato leaves grown under SL + L-LED and SL + H-LED than that of leaves grown under SL (Fig. 7B). Different light conditions could be the main factors resulting in different g_s . Optimal combinations of blue- and red-LED used in this study could enhance g_s , which have been previously reported in cucumber (Hernández and Kubota, 2016) and lettuce (Wang et al., 2016; Choong et al., 2018). It is well known that stomata are responsible for balancing photosynthetic CO_2 uptake with water loss through transpiration. Although it was unable to measure the root biomass in this study, all sweet potato plants had well developed big root systems to ensure water and nutrient uptake (Poorter et al., 2012). Thus, all leaves had similar transpiration rate (Fig. 7D) although leaves

grown under SL had significantly lower g_s compared to those grown under SL supplemented with LED lighting.

In certain species, the greater A by plants grown under high-irradiance means that they have lower C_i than those grown under low-irradiance (Hanba et al., 2002; Yamori et al., 2010; Huang et al., 2014). For instance, Huang et al. (2014) reported that sun-grown tobacco leaves had greater A , leading to lower C_i . In this study, sweet potato leaves grown under the highest irradiance (SL + H-LED) also had the highest A which was as high as $38.1 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ (Fig. 7A) and the lowest C_i of $236.8 \mu\text{mol CO}_2 \text{ mol}^{-1}$ compared to those grown under low-irradiance (Fig. 7C). Lower C_i could result in higher ribulose-1,5-bisphosphate (RuBP) oxygenation rate in sun-grown-leaves than in shade-grown leaves and thus upregulate photorespiratory pathway. However, lower C_i of sun-grown leaves also increased RuBP regeneration resulting from higher electron transport capacity and thus RuBP oxygenation and regeneration were balanced (Foyer et al., 2012; Huang et al., 2014). In this study, sweet potato leaves grown under high-irradiance also had higher electron transport capacity (Fig. 5A). Instead of suppressing A , enhancement of photorespiratory pathway through enhanced RuBP regeneration potentially improved the A in *Arabidopsis thaliana* (Timm et al., 2012) and tobacco leaves (Huang et al., 2014). Sweet potato leaves grown in the tropical greenhouse under high-irradiance may also have higher photorespiration which enhanced photosynthetic CO_2 assimilation and at the same time prevented them from suffering sustained chronic photoinhibition supported by their predawn F_v/F_m ratios of greater than 0.8 (Fig. 3A). Changes in the light dependence of photosynthesis may be ascribed to changes in not only CO_2 concentration in the chloroplasts but also the activity and amount of photosynthetic components especially the activation and the amount of Rubisco (Simkin et al., 2015, 2019). Compared to sweet potato leaves grown under low-irradiance such as SL only, leaves grown under high-irradiance such as SL + H-LED probably have increased synthesis of Rubisco. Supplemental LED lighting to SL increased total leaf soluble protein and Rubisco protein of Cos lettuce (*Lactuca sativa* L.) grown in the same greenhouse (unpublished data). In this study, samples have been collected for the analysis of total leaf soluble protein and Rubisco protein. Unfortunately, we were unable to determine these parameters as we had to close our laboratory due to the COVID-19 pandemic. The effects of light on Rubisco protein would be studied in near future. We also hypothesize that supplementing LED to natural SL enhances not only leaf growth and photosynthetic characteristics of sweet potato leaves in a tropical greenhouse but also their nutritional quality, which also merits our further study.

5. Conclusion

The results of this study show that supplemental LED lighting to natural SL increased leaf fresh and dry weights of hydroponically grown sweet potato (*I. batatas* var. Biru Putih) in the hot tropical greenhouse. The enhancement of leaf biomass was due to the increased thickness of high-irradiance grown leaves, which improved electron transport capacity and photosynthetic CO_2 assimilation rate of attached leaves under their growth temperature and irradiances. Corrections among ETR, qP , PS II, Cyt b_6f , P_N and Rubisco protein in sweet potato leaves under different light intensities and different leaf temperatures merit our further study.

Author contributions

JH initiated and funded the expenses for the project, planned and carried out some parts of the experiments and wrote most part of the manuscript. LQ planned, carried out most experiments, analyzed the data and wrote some part of the manuscript.

CRedit authorship contribution statement

Jie He: Conceptualization, Methodology, Investigation, Writing - original draft, Writing - review & editing. **Lin Qin:** Formal analysis, Investigation, Methodology, Project administration, Writing - review & editing.

Declaration of Competing Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Acknowledgements

We thank Singapore Millennium Foundation (SMF-Farming System) and National Institute of Education, Nanyang Technological University, Singapore for providing us with the funding and facility support.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.jplph.2020.153239>.

References

- Ai-Lien, C., 2019. Singapore sets 30% goal for home-grown food by 2030. The Straits Times. Singapore Press Holdings, Singapore (Accessed 8 March 2019). <https://www.straitstimes.com/singapore/spore-sets-30-goal-for-home-grown-food-by-2030>.
- An, L.V., Frankow-Lindberg, B.E., Lindberg, J.E., 2003. Effect of harvesting interval and defoliation on yields and chemical composition of leaves, stems and tubers of sweet potato (*Ipomoea batatas* L. (Lan.)) plant parts. *Field Crops Res.* 82, 49–58. [https://doi.org/10.1016/S0378-4290\(03\)00018-2](https://doi.org/10.1016/S0378-4290(03)00018-2).
- Anderson, J.M., Chow, W.S., Goodchild, D.J., 1988. Thylakoid membrane organisation in sun/shade acclimation. *Aust. J. Plant Physiol.* 15, 11–26. <https://doi.org/10.1071/PP9880011>.
- Björkmann, O., 1981. Responses to different quantum flux densities. In: Lange, O.K., Nobel, P.S., Osmond, C.B., Ziegler, H. (Eds.), *Encyclopedia of Plant Physiology*, N.S., Vol. 12A, Physiological Plant Ecology I. Springer, Berlin Heidelberg, New York, pp. 57–107.
- Bunce, J.A., 2001. Responses of stomatal conductance to light, humidity and temperature in winter wheat and barley grown at three concentrations of carbon dioxide in the field. *Glob. Change Biol.* 6, 371–382. <https://doi.org/10.1046/j.1365-2486.2000.00314.x>.
- Choong, T.W., He, J., Qin, L., Lee, S.K., 2018. Quality of supplementary LED lighting effects on growth and photosynthesis of two different *Lactuca* recombinant inbred lines (RILs) grown in a tropical greenhouse. *Photosynthetica* 56, 1278–1286. <https://doi.org/10.1007/s11099-018-0828-2>.
- Chow, W.S., Anderson, J.M., 1987. Photosynthetic responses of *Pisum sativum* to an increase in irradiance during growth. II. Thylakoid membrane components. *Aust. J. Plant Physiol.* 14, 9–19. <https://doi.org/10.1071/PP9870009>.
- Chow, W.S., Hope, A.B., 1987. The stoichiometries of supramolecular complexes in thylakoid membranes from spinach chloroplasts. *Aust. J. Plant Physiol.* 14, 21–28. <https://doi.org/10.1071/PP9870021>.
- Chow, W.S., Qian, L., Goodchild, D.J., Anderson, J.M., 1988. Photosynthetic acclimation of *Alocasia macrorrhiza* (L.) G. Don to growth irradiance: structure, function and composition of chloroplasts. *Aust. J. Plant Physiol.* 15, 107–122. <https://doi.org/10.1071/PP9880107>.
- Chow, W.S., Hope, A.B., Anderson, J.M., 1991. Further studies on quantifying photosystem II *in vivo* by flash-induced oxygen yield from leaf discs. *Aust. J. Plant Physiol.* 18, 397–410. <https://doi.org/10.1071/PP910397>.
- De la Torre, W.R., Burkey, K.O., 1990. Acclimation of barley to changes in light intensity: photosynthetic electron transport activity and components. *Photosynth. Res.* 24, 127–136. <https://doi.org/10.1007/BF00032593>.
- Demmig-Adams, B., Adams, W.W., 1996. The role of xanthophyll cycle carotenoids in the protection of photosynthesis. *Trends Plant Sci.* 1, 21–26. [https://doi.org/10.1016/S1360-1385\(96\)80019-7](https://doi.org/10.1016/S1360-1385(96)80019-7).
- Douglas, J.S., 1985. *Advanced guide to hydroponics*. Pelham Books, London.
- Eichelmann, H., Price, D., Badger, M., Laisk, A., 2000. Photosynthetic parameters of wild-type and *Cytb_f* deficient transgenic tobacco studied by CO₂ uptake and transmittance at 800 nm. *Plant Cell Physiol.* 41, 432–439. <https://doi.org/10.1093/pcp/41.4.432>.
- Evans, J.R., Poorter, H., 2001. Photosynthetic acclimation of plants to growth irradiance: the relative importance of specific leaf area and nitrogen partitioning in maximizing carbon gain. *Plant Cell Environ.* 24, 755–767. <https://doi.org/10.1046/j.1365-3040.2001.00724.x>.
- Fan, Y., Chen, J., Cheng, Y., Raza, M.A., Wu, X., Wang, Z., et al., 2018. Effect of shading and light recovery on the growth, leaf structure, and photosynthetic performance of soybean in a maize-soybean relay-strip intercropping system. *PLoS One* 13, e0198159. <https://doi.org/10.1371/journal.pone.0198159>.
- Feng, L., Raza, M.A., Li, Z., Chen, Y., Khalid, M.H.B., Du, J., Liu, W., Wu, X., Song, C., Yu, L., Zhang, Z., Yuan, S., Yang, W., Yang, F., 2019. The influence of light intensity and leaf movement on photosynthesis characteristics and carbon balance of soybean. *Front. Plant Sci.* 9, 1952. <https://doi.org/10.3389/fpls.2018.01952>.
- Foyer, C.H., Neukermans, J., Queval, G., Noctor, G., Harbinson, J., 2012. Photosynthetic control of electron transport and the regulation of gene expression. *J. Exp. Bot.* 63, 1637–1661. <https://doi.org/10.1093/jxb/ers013>.
- Fromme, P., Melkozernov, A., Jordan, P., Krauss, N., 2003. Structure and function of photosystem I: interaction with its soluble electron carriers and external antenna systems. *FEBS Lett.* 555, 40–44. [https://doi.org/10.1016/S0014-5793\(03\)01124-4](https://doi.org/10.1016/S0014-5793(03)01124-4).
- Ghuman, B.S., Lal, R., 1983. Mulch and irrigation effects on plant-water relations and performance of cassava and sweet potato. *Field Crops Res.* 7, 13–29. [https://doi.org/10.1016/0378-4290\(83\)90003-5](https://doi.org/10.1016/0378-4290(83)90003-5).
- Gommers, C.M.M., 2020. Adapting to High Light: At a Different Time and Place? *Plant Physiol.* 182, 10–11. <https://doi.org/10.1104/pp.19.01445>.
- Hanba, Y.T., Kogami, H., Terashima, I., 2002. The effect of growth irradiance on leaf anatomy and photosynthesis in *Acer* species differing in light demand. *Plant Cell Environ.* 25, 1021–1030. <https://doi.org/10.1046/j.1365-3040.2002.00881.x>.
- Hashimoto, H., Uragami, C., Cogdell, R.J., 2016. Carotenoids and photosynthesis. *Subcell. Biochem.* 79, 111–139. https://doi.org/10.1007/978-3-319-39126-7_4.
- He, J., 2015. Farming of Vegetables in Space-limited Environments. 2015 Cosmos 11. pp. 21–36. <https://doi.org/10.1142/S0219607715500020>.
- He, J., Chow, W.S., 2003. The rate coefficient of repair of photosystem II after photo-inactivation. *Physiol. Plant.* 118, 297–304. <https://doi.org/10.1034/j.1399-3054.2003.00107.x>.
- He, J., Chee, C.W., Goh, C.J., 1996. "Photoinhibition" of Heliconia under natural tropical conditions- Importance of leaf orientation for light interception and leaf temperature. *Plant Cell Environ.* 19, 1238–1248. <https://doi.org/10.1111/j.1365-3040.1996.tb00002.x>.
- He, J., Cheok, L., Qin, L., 2011. Nitrate accumulation, productivity and photosynthesis of temperate butter head lettuce under different nitrate availabilities and growth irradiances. *Open Hortic. J.* 4, 17–24. <https://doi.org/10.2174/1874840601104010017>.
- He, J., Lim, R.M.P., Dass, S.H.J., Yam, T.W., 2017a. Photosynthetic acclimation of *Grammatophyllum speciosum* to growth irradiance under natural conditions in Singapore. *Bot. Stud.* 58, 58. <https://doi.org/10.1186/s40529-017-0210-x>.
- He, J., Qin, L., Chong, E.L.C., Choong, T.W., Lee, S.K., 2017b. Plant growth and photosynthetic characteristics of *Mesembryanthemum crystallinum* grown aeroponically under different blue- and red-LEDs. *Front. Plant Sci.* 8, 361. <https://doi.org/10.3389/fpls.2017.00361>.
- He, J., Qin, L., Chow, W.S., 2019a. Impacts of LED spectral quality on leafy vegetables: productivity closely linked to photosynthetic performance or associated with leaf traits? *Int. J. Agric. Biol. Eng.* 12, 16–25. <https://doi.org/10.25165/ij.ajibae.20191206.5178>.
- He, J., Qin, L., Teo, L.J.L., Choong, T.W., 2019b. Nitrate accumulation, productivity and photosynthesis of Brassica albobolabra grown under low light with supplemental LED lighting in the tropical greenhouse. *J. Plant Nutr.* 42, 1740–1749. <https://doi.org/10.1080/01904167.2019.1643367>.
- Heber, U., Neimanis, S., Dietz, K.J., 1988. Fractional control of photosynthesis by the Q_B protein, the cytochrome *f/b₆* complex and other components of the photosynthetic apparatus. *Planta* 173, 267–274. <https://doi.org/10.1007/BF00403020>.
- Hernández, R., Kubota, C., 2016. Physiological responses of cucumber seedlings under different blue and red photon flux ratios using LEDs. *Environ. Exp. Bot.* 121, 66–74. <https://doi.org/10.1016/j.envexpbot.2015.04.001>.
- Hikosaka, K., Ishikawa, K., Borjigidai, A., Muller, O., Onoda, Y., 2006. Temperature acclimation of photosynthesis: mechanisms involved in the changes in temperature dependence of photosynthetic rate. *J. Exp. Bot.* 57, 291–302. <https://doi.org/10.1093/jxb/erj049>.
- Huang, W., Zhang, S.B., Hu, H., 2014. Sun leaves up-regulate the photorespiratory pathway to maintain a high rate of CO₂ assimilation in tobacco. *Front. Plant Sci.* 5, 688. <https://doi.org/10.3389/fpls.2014.00688>.
- Hunt, R., Causton, D.R., Shipley, B., Askew, A.P., 2002. A modern tool for classical plant growth analysis. *Ann. Bot.* 90, 485–488. <https://doi.org/10.1093/aob/mcf214>.
- Ishida, H., Suzuno, H., Sugiyama, N., Innami, S., Tadokoro, T., Maekawa, A., 2000. Nutritive evaluation on chemical components of leaves, stalks and stems of sweet potatoes (*Ipomoea batatas* L. var. *bonjir*). *Food Chem.* 68, 359–367. [https://doi.org/10.1016/S0308-8146\(99\)00206-X](https://doi.org/10.1016/S0308-8146(99)00206-X).
- Johnson, M., Pace, R.D., 2010. Sweet potato leaves: properties and synergistic interactions that promote health and prevent disease. *Nutr. Rev.* 68, 604–615. <https://doi.org/10.1111/j.1753-4887.2010.00320.x>.
- Jones, D.S., 2006. ASEAN and transboundary haze pollution in Southeast Asia. *Asia Eur. J.* 4, 431–446. <https://doi.org/10.1007/s10308-006-0067-1>.
- Kurata, R., Adachi, M., Yamakawa, O., Yoshimoto, M., 2007. Growth suppression of human cancer cells by polyphenolics from sweetpotato (*Ipomoea batatas* L.) leaves. *J. Agric. Food Chem.* 55, 185–190. <https://doi.org/10.1021/jf0620259>.
- Lawson, T., 2009. Guard cell photosynthesis and stomatal function. *New Phytol.* 181, 13–34. <https://doi.org/10.1111/j.1469-8137.2008.02685.x>.
- Leong, T.Y., Anderson, J.M., 1984. Adaptation of the thylakoid membranes of pea chloroplasts to light intensities. II. Regulation of electron transport capacities, electron carriers, coupling factor (CF1) activity and rates of photosynthesis. *Photosynth. Res.* 5, 117–128. <https://doi.org/10.1007/BF00028525>.
- Lichtenthaler, H.K., 2007. Biosynthesis, accumulation and emission of carotenoids, lipo- tocopherol, plastoquinone, and isoprene in leaves under high photosynthetic irradiance. *Photosynth. Res.* 92, 163–179. <https://doi.org/10.1007/s11200-007-9204-y>.

- Lin, K.H., Lai, Y.C., Chang, K.Y., Chen, Y.F., Hwang, S.Y., Lo, H.F., 2007. Improving breeding efficiency for quality and yield of sweet potato. *Bot. Stud.* 48, 283–292.
- Liu, Y., Dawson, W., Prati, D., Haeuser, E., Feng, Y., van Kleunen, M., 2016. Does greater specific leaf area plasticity help plants to maintain a high performance when shaded? *Ann. Bot.* 118, 1329–1336. <https://doi.org/10.1093/aob/mcw180>.
- Ludvik, B., Hanefeld, M., Pacini, M., 2008. Improved metabolic control by *Ipomoea batatas* (Caiapo) is associated with increased adiponectin and decreased fibrinogen levels in type 2 diabetic subjects. *Diabetes Obes. Metab.* 10, 586–592. <https://doi.org/10.1111/j.1463-1326.2007.00752.x>.
- Mekonen, B., Tulu, S., Nego, J., 2015. Orange fleshed sweet potato (*Ipomoea batatas* L.) varieties evaluated with respect to growth parameters at Jimma in Southwestern Ethiopia. *J. Agron.* 14, 164–169. <https://doi.org/10.3923/ja.2015.164.169>.
- Nobre, A.M., Karthik, S., Liu, H., Yang, D., Martins, F.R., Pereira, E.B., R  ther, R., Reindl, T., Peters, I.M., 2016. On the impact of haze on the yield of photovoltaic systems in Singapore. *Renew. Energy* 89, 389–400. <https://doi.org/10.1016/j.renene.2015.11.079>.
- Nwinyi, S.C.O., 1992. Effect of age at short removal on tuber and shoot yields at harvest of five sweet potato (*Ipomoea batatas* (L) Lam) cultivars. *Field Crops Res.* 29, 47–54. [https://doi.org/10.1016/0378-4290\(92\)90075-K](https://doi.org/10.1016/0378-4290(92)90075-K).
- Oswald, A., Alk  mper, J., Midmore, D.J., 1994. The effect of different shade levels on growth and tuber yield of sweet potato: I. Plant development. *J. Agro. Crop Sci.* 173, 41–52. <https://doi.org/10.1111/j.1439-037X.1994.tb00572.x>.
- Pogson, B.J., Rissler, H.M., Frank, H.A., 2005. The roles of carotenoids in photosystem II of higher plants. In: Wydrzynski, T., Satoh, K. (Eds.), *Photosystem II: the Light-Driven Water: Plastoquinone Oxidoreductase*. Springer-Verlag, Dordrecht, The Netherlands, pp. 515–537.
- Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P., Mommer, L., 2012. Biomass allocation to leaves, stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytol.* 193, 30–50. <https://doi.org/10.1111/j.1469-8137.2011.03952.x>.
- Poorter, L., Castilho, C.V., Schiatti, J., Oliveira, R.S., Costa, F.R.C., 2018. Can traits predict individual growth performance? A test in a hyperdiverse tropical forest. *New Phytol.* 219, 109–121. <https://doi.org/10.1111/nph.15206>.
- Simkin, A.J., McAusland, L., Headland, L.R., Lawson, T., Raines, C.A., 2015. Multigene manipulation of photosynthetic carbon assimilation increases CO₂ fixation and biomass yield in tobacco. *J. Exp. Bot.* 66, 4075–4090. <https://doi.org/10.1093/jxb/erv204>.
- Simkin, A.J., L  pez-Calcagno, P.E., Raines, C.A., 2019. Feeding the world: improving photosynthetic efficiency for sustainable crop production. *J. Exp. Bot.* 70, 1119–1140. <https://doi.org/10.1093/jxb/ery445>.
- Tay, S., He, J., Yam, T.W., 2019. CAM plasticity in epiphytic tropical orchid species responding to environmental stress. *Bot. Stud.* 60, 7. <https://doi.org/10.1186/s40529-019-0255-0>. 2019.
- Terashima, I., Hanba, Y.T., Tazoe, Y., Vyas, P., Yano, S., 2006. Irradiance and phenotype: comparative eco-development of sun and shade leaves in relation to photosynthetic CO₂ diffusion. *J. Exp. Bot.* 57, 343–354. <https://doi.org/10.1093/jxb/erj014>.
- Tikkanen, M., Grieco, M., Nurmi, M., Rantala, M., Suorsa, M., Aro, E.M., 2012. Regulation of the photosynthetic apparatus under fluctuating growth light. *Philos. Trans. R. Soc. B* 367, 3486–3493. <https://doi.org/10.1098/rstb.2012.0067>.
- Timm, S., Florian, A., Arrivault, S., Stitt, M., Fernie, A.R., 2012. Glycine decarboxylase controls photosynthesis and plant growth. *FEBS Lett.* 586, 3692–3697. <https://doi.org/10.1016/j.febslet.2012.08.027>.
- Violet-Chabrand, Matthews, J.S.A., Simkin, A.J., Raines, C.A., Lawson, T., 2017. Importance of fluctuations in light on plant photosynthetic acclimation. *Plant Physiol.* 173, 2163–2179. <https://doi.org/10.1104/pp.16.01767>.
- Wang, J., Lu, W., Tong, Y., Yang, Q., 2016. Leaf morphology, photosynthetic performance, chlorophyll fluorescence, stomatal development of lettuce (*Lactuca sativa* L.) exposed to different ratios of red light to blue light. *Front. Plant Sci.* 7, 250. <https://doi.org/10.3389/fpls.2016.00250>.
- Wellburn, A.R., 1994. The spectral determination of chlorophylls a and b, as well as carotenoids, using various solvents with spectrophotometers of different resolution. *J. Plant Physiol.* 144, 307–313. [https://doi.org/10.1016/s0176-1617\(11\)81192-2](https://doi.org/10.1016/s0176-1617(11)81192-2).
- Yamasaki, T., Yamakawa, T., Yamane, Y., Koike, H., Satoh, K., Katoh, S., 2002. Temperature acclimation of photosynthesis and related changes in photosystem II electron transport in winter wheat. *Plant Physiol.* 128, 1087–1097. <https://doi.org/10.1104/pp.010919>.
- Yamori, W., Evans, J.R., Von Caemmerer, S., 2010. Effects of growth and measurement light intensities on temperature dependence of CO₂ assimilation rate in tobacco leaves. *Plant Cell Environ.* 33, 332–343. <https://doi.org/10.1111/j.1365-3040.2009.02067.x>.
- Yang, F., Feng, L., Liu, Q., Wu, X., Fan, Y., Ali Raza, M., et al., 2018. Effect of interactions between light intensity and red-to-far-red ratio on the photosynthesis of soybean leaves under shade condition. *Environ. Exp. Bot.* 150, 79–87. <https://doi.org/10.1016/j.envexpbot.2018.03.008>.
- Yooyongwecha, S., Theerawitayab, C., Samphumphuangb, T., Chaumb, S., 2013. Water-deficit tolerant identification in sweet potato genotypes (*Ipomoea batatas* (L.) Lam.) in vegetative developmental stage using multivariate physiological indices. *Sci. Hortic.* 162, 242–251. <https://doi.org/10.1016/j.scienta.2013.07.041>.
- Yoshimoto, M., Yahara, S., Okuno, S., Islam, M.S., Ishiguro, K., Yamakawa, O., 2002. Antimutagenicity of mono-, di-, and tricaffeoylquinic acid derivatives isolated from sweet potato (*Ipomoea batatas* L) leaf. *Biosci. Biotech. Biochem.* 66, 2336–2341. <https://doi.org/10.1271/bbb.66.2336>.
- Zhang, Y., Yu, T., Ma, W., Tian, C., Sha, Z., Li, J., 2019. Morphological and physiological response of *Acer catalpifolium* Rehd. Seedlings to water and light stresses. *Glob. Ecol. Conserv.* 19, e00660. <https://doi.org/10.1016/j.gecco.2019.e00660>.
- Zhu, H., Zeng, L.D., Yi, X.P., Peng, C.L., Zhang, W.F., Chow, W.S., 2017. The half-life of the cytochrome b_f complex in leaves of pea plants after transfer from moderately-high growth light to low light. *Funct. Plant Biol.* 44, 351–357. <https://doi.org/10.1071/FP16222>.
- Zivcak, M., Brestic, M., Kalaji, H.M., Govindjee, 2014. Photosynthetic responses of sun- and shade-grown barley leaves to high light: is the lower PSII connectivity in shade leaves associated with protection against excess of light? *Photosynth. Res.* 119, 339–354. <https://doi.org/10.1007/s11120-014-9969-8>.