



Published in final edited form as:

Horm Behav. 2019 May ; 111: 131–134. doi:10.1016/j.yhbeh.2018.11.005.

Glucocorticoids and resilience

Joanna L. Spencer-Segal^{1,2}, Huda Akil²

¹Department of Internal Medicine, Division of Metabolism, Endocrinology, and Diabetes, University of Michigan, Ann Arbor, MI

²Molecular and Behavioral Neuroscience Institute, University of Michigan, Ann Arbor, MI

Abstract

All organisms endure frequent challenges to homeostasis, or stressors, that require adaptation. Depending on the individual, the context, and the magnitude of stress, this active adaptation can lead to behavioral susceptibility or resilience. The latter is an under-appreciated consequence of stress, as the damaging effects of chronic stress and chronically elevated glucocorticoids have received much more attention. We suggest here that neural pathways promoting resilience are initiated at the time of stress, and that they involve glucocorticoid signaling. By focusing on the neurobiology of resilience induction and the identification of vulnerable individuals, we may be able to intervene in the future at the time of stress to promote positive adaptation.

Stress is the foundation of evolution. All organisms undergo stressors, defined as internal or external challenges to homeostasis that require adaptation (Karatsoreos et al., 2013). Organisms that adapt positively to stress will survive; thus, evolution has selected for resilience promoting traits. The classic idea of allostasis, or stability through change, argues that an organism can handle a certain amount of stress or allostatic load, adapting well to it in order to return to equilibrium (McEwen, 1998). Thus, in allostasis, individuals who recover successfully are resilient, generally not exhibiting negative physiological or behavioral effects after stress. However, beyond a certain level, the homeostatic compensation to accumulated stress breaks down, and this allostatic overload results in negative health effects and increased vulnerability to subsequent stressors (McEwen et al., 2015).

As implied in the concept of allostasis, the maintenance of biological stability in the face of change is an active process associated with distinct changes in gene expression, neuroanatomy, and neurogenesis involving many of the same brain regions that mediate stress susceptibility (Anacker et al., 2018; 2016; Bagot et al., 2017; Gray et al., 2014).

Correspondence: Joanna Spencer-Segal, MD, PhD 205 Zina Pitcher Place, Room 1033 Ann Arbor, MI 48109 (734) 936-0207
sjoanna@med.umich.edu.

The authors have no conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Moreover, individuals are not only resilient *in spite of* stress; rather, the inevitable stressors of life can promote positive adaptation, termed “good” stress (McEwen et al., 2015).

As the attention to resilience has recently increased, most studies have focused on what a resilient organism versus a susceptible one looks like based on specific behavioral criteria at a distal time point to a particular stress paradigm. This line of investigation has yielded fruitful insights into individual differences in neurobiological adaptation, including how genetic predisposition and early developmental events can bias an organism towards greater vulnerability or resilience (Aydin et al., 2015; Clinton et al., 2013; Meaney, 2001; Rana et al., 2016).

However, by only studying downstream resilient and vulnerable phenotypes, we miss information about resilience-promoting mechanisms. For example, gene expression changes in the brain of stress-recovered animals have very little overlap with the changes seen immediately following stress (Gray et al., 2014). Therefore, the mechanisms of resilience are distinct and not necessarily discoverable by studying a resilient organism in recovery. Here we argue for a renewed focus on the acute neurobiological adaptations to stress. We suggest that acute stress activates resilience-promoting pathways in part via glucocorticoids. Studying the mechanisms by which these resilience-promoting pathways are initiated in the brain in response to acute stress will yield critical insights into the neurobiology of adaptation that will have strong translational potential.

Glucocorticoids in short- and long-term adaptation to stress

Two concepts in the field, while important, have overshadowed the notion that glucocorticoids may promote positive neurobiological adaptation to stress. One is the idea that elevated glucocorticoids during acute stress serve mostly to promote physiologic mechanisms important for survival, such as increased vascular tone, alteration of metabolism, and a balanced enhancement and suppression of inflammation (Dhabhar, 2018; Hamrahian et al., 2017). While glucocorticoids do support these processes, very high levels of glucocorticoid are not clearly required for an organism to adequately withstand significant physical stressors (Udelsman et al., 1986). The second concept underscores the damaging effects of sustained high levels of glucocorticoids. A key notion of allostasis and allostatic load is that chronic elevation of glucocorticoids, which occurs in chronic stress and other situations, results in structural and functional alterations in the hippocampus and other brain regions leading to significant affective, cognitive, and metabolic problems (McEwen, 2008a; Starkman et al., 1992). However, as discussed below, acute and chronic glucocorticoids may activate distinct and often opposing neurobiological processes. The focus on the damaging effects of chronic glucocorticoids has unfortunately detracted attention from their short-term role in resilience.

Several human studies support the notion that glucocorticoids promote resilience via their actions in the brain during acute stress (Table 1). In humans, corticosteroid treatment during critical illness is associated with lower symptoms of PTSD and anxiety in survivors, suggesting a possible protective effect (Bienvenu et al., 2013; Schelling et al., 1999; Spencer-Segal et al., 2017). Indeed, in a few small trials hydrocortisone treatment improved

mental health outcomes after sepsis or cardiac surgery (Schelling et al., 2001; 2004; Weis et al., 2006). The association between high cortisol and lower PTSD symptoms has also been seen after other traumatic experiences such as a motor vehicle accident (Delahanty et al., 2000). Finally, in one small trial, treatment with high doses of hydrocortisone after a traumatic experience attenuated PTSD development in patients exhibiting acute stress reactions (Zohar et al., 2011).

The putative protective effect of glucocorticoids in acute stress has also been investigated in rats. Compared with other strains, Lewis rats showed a blunted corticosterone response to predator odor stress along with greater anxiety- and PTSD-like behaviors later on, which were attenuated by corticosterone treatment immediately following the stressor (Cohen et al., 2006). Similarly, treatment with high-dose corticosterone during or immediately following restraint or predator odor stress prevented the later emergence of anxiety- and PTSD-like behavior (Cohen et al., 2008; Rao et al., 2012). Thus, in both humans and rats undergoing varied acute stressors, glucocorticoids seem to prevent the development of the anxiety- and PTSD-like behaviors. The conservation of this effect between species suggests evolutionary importance to this phenomenon and implies that rodent studies may help elucidate the mechanism.

Neural mechanisms of acute and chronic glucocorticoid action

Acute and chronic glucocorticoids may promote resilience or vulnerability through direct actions on brain regions important for emotional behavior. Glucocorticoid receptors (GR) are expressed in neurons throughout the brain, while mineralocorticoid receptors (MR) are found in limbic areas, especially the hippocampus, commonly implicated in stress neurobiology (Reul and De Kloet, 1985). Due to the higher affinity of MR relative to GR for the main adrenal glucocorticoids, corticosterone (in rodents) and cortisol (in humans) (Reul and De Kloet, 1985), GR is considered a likely candidate to mediate the acute effects of stress levels of hormone. Rapid nongenomic actions via membrane MR are likely important, as well (Heegde et al., 2015). Acting through MR and GR, glucocorticoids alter many processes including dendritic branching, spine formation, apoptosis, neurogenesis, and memory formation (Heegde et al., 2015; McEwen, 2008b; Schoenfeld and Gould, 2012).

Acute and chronic glucocorticoids may have different and even opposite effects on neural processes. For example, chronic stress and glucocorticoids cause decreased dendritic branching and dendritic spine density in hippocampal and prefrontal cortical pyramidal cells (Anderson et al., 2016; Gourley et al., 2013; Woolley et al., 1990). In general, these changes are linked to negative behavioral adaptations such as impaired cognitive processing and anhedonia (Gourley et al., 2013; Liston et al., 2006). In contrast, during acute stress, corticosterone seems to prevent detrimental changes in neural architecture. For example, endogenous and exogenous corticosterone prevent specific changes to dendritic branching in the rat hippocampal dentate gyrus and basolateral amygdala after acute stress (Rao et al., 2012; Zohar et al., 2011). In both of those cases, corticosterone at the time of stress prevented not only the changes to dendritic architecture but also the increase in anxiety-like behavior. This evidence suggests that glucocorticoids during acute stress promote behavioral resilience in part via the preservation of limbic dendritic architecture.

A better understanding of the different effects of acute and chronic glucocorticoids on neural processes will be an important step toward identifying the mechanisms by which acute glucocorticoids may promote resilience. In regard to dendritic spine density, acute and chronic corticosterone were reported to have different effects on spine turnover (Liston and Gan, 2011). In that study, acute corticosterone increased turnover of new spines, while chronic corticosterone increased elimination of both old and new spines without affecting rates of spine formation. It is possible that rapid nongenomic mechanisms dominate the acute effects while genomic actions are more prominent chronically. Additionally, MR and GR can have opposing cellular effects; for example, GR seems to increase cell death in several paradigms while MR has the opposite effect (Heegde et al., 2015). Therefore, changes in receptor expression level, the ratio of MR to GR, or receptor sensitivity could alter the effect of glucocorticoids. In some cases, acute and chronic glucocorticoids may have similar outcomes that are perceived as adaptive in the short term but detrimental in the long term. For example, glucocorticoids may prevent PTSD by interfering with the formation or consolidation of traumatic memories (Roozendaal, 2003), but in the long term this impairment in cognitive processing would be undesirable.

Importantly, the downstream effects of acute glucocorticoids depend on context. For example, in the absence of stress, acute corticosterone treatment can actually promote long-term anxiety and dendritic hypertrophy in the amygdala, the opposite of its effect when paired with acute stress (Mitra and Sapolsky, 2008). Moreover, more “positively” viewed stimuli that activate the hypothalamic-pituitary-adrenal (HPA) axis such as environmental enrichment and exercise can stimulate neurogenesis, while more aversive stressors have the opposite effect (Schoenfeld and Gould, 2012). Furthermore, dose is crucial, as the effects of glucocorticoids often follow a U-shaped curve. For example, in one study, lower doses of acute corticosterone actually exacerbated the anxiety-like and startle behavior seen after predator odor stress, while high doses prevented these responses (Cohen et al., 2008). Of course, glucocorticoids will interact with other systems such as adrenergic, endocannabinoid, and estrogen signaling (Lorsch et al., 2018). All these factors need to be accounted for when considering the mechanisms by which glucocorticoids may promote resilience after acute stress.

Harnessing biology to promote resilience

Despite decades of studying glucocorticoid signaling and stress biology, no glucocorticoid-directed treatments are currently available for stress-susceptible individuals. Reasons for this likely include the complexity of acute and chronic glucocorticoid actions in stressed individuals, and individual variation in glucocorticoid signaling and stress susceptibility. More effective development and implementation of endocrine treatments could result from a renewed focus on the role of glucocorticoids in promoting resilience to acute stress. Genetic polymorphisms and epigenetic modifications of HPA axis components have been associated with PTSD and mood disorders, and they could influence vulnerability in part by affecting glucocorticoid signaling during stress (Bogdan et al., 2012; Davydow et al., 2014; Griffiths and Hunter, 2014; van Zuiden et al., 2012; Vogel et al., 2014; Yehuda et al., 2014). The identification of glucocorticoid-based markers of vulnerability and implementation of a

mechanistically appropriate intervention to enhance resilience would be a major success for precision mental health.

Conclusion

We suggest that when faced with a stressor, circulating endocrine and other signals initiate resilience-promoting pathways in the brain to protect against detrimental behavioral outcomes and to prepare for future challenges. Glucocorticoids in particular appear to be crucial components of resilience induction in the brain, subject to nuances of context, dose, and timing. In this way the acutely stressed brain is like a sore muscle, with glucocorticoids the physical activity which can, if calibrated correctly, trigger mechanisms that are beneficial to short-term healing (Søgaard and Sjøgaard, 2017). On the other hand, just as chronic overuse of any muscle or joint can be damaging, natural mechanisms of resilience may break down in the long term because they are unsustainable or have damaging side effects. Though intrinsic to the original allostasis model, the benefits of short-term stress and the stress response are often overshadowed by the attention paid to damaging effects of chronic stress and stress hormones. We therefore propose a renewed focus on short-term adaptation, or the initiation of resilience-promoting pathways by acute stress and glucocorticoids. In the future, we envision that individuals vulnerable to maladaptation to acute or chronic stress may be identified by their genetic, epigenetic, or phenotypic (e.g., neuroendocrine) signatures. Real-time interventions that promote natural mechanisms of resilience programming during stress could then be designed to promote positive adaptation.

Acknowledgments

Financial support was provided by the Hope for Depression Research Foundation (H.A.), Pritzker Neuropsychiatric Disorders Research Fund LLC (H.A.), Office of Naval Research Grant Nos. N00014-09-1-0598 and N00014-12-1-0366 (H.A.), National Institutes of Health grants R01 MH104261 (H.A.) and K08 MH116267 (J.S.S.), and the Brain and Behavior Research Foundation (J.S.S.).

References

- Anacker C, Luna VM, Stevens GS, Millette A, Shores R, Jimenez JC, Chen B, Hen R, 2018 Hippocampal neurogenesis confers stress resilience by inhibiting the ventral dentate gyrus. *Nature* 559, 98–102. doi:10.1038/s41586-018-0262-4 [PubMed: 29950730]
- Anacker C, Scholz J, O'Donnell KJ, Allemang-Grand R, Diorio J, Bagot RC, Nestler EJ, Hen R, Lerch JP, Meaney MJ, 2016 Neuroanatomic Differences Associated With Stress Susceptibility and Resilience. *Biol. Psychiatry* 79, 840–849. doi:10.1016/j.biopsych.2015.08.009 [PubMed: 26422005]
- Anderson RM, Glanz RM, Johnson SB, Miller MM, Romig-Martin SA, Radley JJ, 2016 Prolonged corticosterone exposure induces dendritic spine remodeling and attrition in the rat medial prefrontal cortex. *J. Comp. Neurol* 524, 3729–3746. doi:10.1002/cne.24027 [PubMed: 27113541]
- Aydin C, Frohmader K, Akil H, 2015 Revealing a latent variable: individual differences in affective response to repeated injections. *Behav. Neurosci* 129, 679–682. doi:10.1037/bne0000084 [PubMed: 26191946]
- Bagot RC, Cates HM, Purushothaman I, Vialou V, Heller EA, Yieh L, Labonté B, Peña CJ, Shen L, Wittenberg GM, Nestler EJ, 2017 Ketamine and Imipramine Reverse Transcriptional Signatures of Susceptibility and Induce Resilience-Specific Gene Expression Profiles. *Biol. Psychiatry* 81, 285–295. doi:10.1016/j.biopsych.2016.06.012 [PubMed: 27569543]
- Bienvenu OJ, Gellar J, Althouse BM, Colantuoni E, Sricharoenchai T, Mendez-Tellez PA, Shanholtz C, Dennison CR, Pronovost PJ, Needham DM, 2013 Post-traumatic stress disorder symptoms after

- acute lung injury: a 2-year prospective longitudinal study. *Psychol Med* 43, 2657–2671. doi:10.1017/S0033291713000214 [PubMed: 23438256]
- Bogdan R, Williamson DE, Hariri AR, 2012 Mineralocorticoid receptor Iso/Val (rs5522) genotype moderates the association between previous childhood emotional neglect and amygdala reactivity. *Am J Psychiatry* 169, 515–522. doi:10.1176/appi.ajp.2011.11060855 [PubMed: 22407082]
- Clinton SM, Watson SJ, Akil H, 2013 High novelty-seeking rats are resilient to negative physiological effects of the early life stress. *Stress* 17, 97–107. doi:10.3109/10253890.2013.850670 [PubMed: 24090131]
- Cohen H, Matar MA, Buskila D, Kaplan Z, Zohar J, 2008 Early post-stressor intervention with high-dose corticosterone attenuates posttraumatic stress response in an animal model of posttraumatic stress disorder. *Biol. Psychiatry* 64, 708–717. doi:10.1016/j.biopsych.2008.05.025 [PubMed: 18635156]
- Cohen H, Zohar J, Gidron Y, Matar MA, Belkind D, Loewenthal U, Kozlovsky N, Kaplan Z, 2006 Blunted HPA axis response to stress influences susceptibility to posttraumatic stress response in rats. *Biol. Psychiatry* 59, 1208–1218. doi:10.1016/j.biopsych.2005.12.003 [PubMed: 16458266]
- Davydov DS, Kohen R, Hough CL, Tracy JH, Zatzick D, Katon WJ, 2014 A pilot investigation of the association of genetic polymorphisms regulating corticotrophin-releasing hormone with posttraumatic stress and depressive symptoms in medical-surgical intensive care unit survivors. *J Crit Care* 29, 101–106. doi:10.1016/j.jcrc.2013.08.016 [PubMed: 24075295]
- Delahanty DL, Raimonde AJ, Spoonster E, 2000 Initial posttraumatic urinary cortisol levels predict subsequent PTSD symptoms in motor vehicle accident victims. *Biol. Psychiatry* 48, 940–947. [PubMed: 11074232]
- Dhabhar FS, 2018 The short-term stress response - Mother nature's mechanism for enhancing protection and performance under conditions of threat, challenge, and opportunity. *Front Neuroendocrinol* 49, 175–192. doi:10.1016/j.yfrne.2018.03.004 [PubMed: 29596867]
- Gourley SL, Swanson AM, Koleske AJ, 2013 Corticosteroid-induced neural remodeling predicts behavioral vulnerability and resilience. *J. Neurosci* 33, 3107–3112. doi:10.1523/JNEUROSCI.2138-12.2013 [PubMed: 23407965]
- Gray JD, Rubin TG, Hunter RG, McEwen BS, 2014 Hippocampal gene expression changes underlying stress sensitization and recovery. *Mol. Psychiatry* 19, 1171–1178. doi:10.1038/mp.2013.175 [PubMed: 24342991]
- Griffiths BB, Hunter RG, 2014 Neuroepigenetics of stress. *Neuroscience* 275, 420–435. doi:10.1016/j.neuroscience.2014.06.041 [PubMed: 24976514]
- Hamrahian AH, Fleseriu M, AACE Adrenal Scientific Committee, 2017 EVALUATION AND MANAGEMENT OF ADRENAL INSUFFICIENCY IN CRITICALLY ILL PATIENTS: DISEASE STATE REVIEW. *Endocr Pract* 23, 716–725. doi:10.4158/EP161720.RA [PubMed: 28332876]
- Heegde, Ter F, De Rijk RH, Vinkers CH, 2015 The brain mineralocorticoid receptor and stress resilience. *Psychoneuroendocrinology* 52C, 92–110. doi:10.1016/j.psyneuen.2014.10.022
- Karatsoreos IN, Karatsoreos IN, McEwen BS, 2013 Annual Research Review: The neurobiology and physiology of resilience and adaptation across the life course. *J Child Psychol Psychiatry* 54, 337–347. doi:10.1111/jcpp.12054 [PubMed: 23517425]
- Liston C, Gan WB, 2011 Glucocorticoids are critical regulators of dendritic spine development and plasticity in vivo. *Proceedings of the National Academy of Sciences* 108, 16074–16079. doi:10.1073/pnas.1110444108
- Liston C, Miller MM, Goldwater DS, Radley JJ, Rocher AB, Hof PR, Morrison JH, McEwen BS, 2006 Stress-induced alterations in prefrontal cortical dendritic morphology predict selective impairments in perceptual attentional set-shifting. *J. Neurosci* 26, 7870–7874. doi:10.1523/JNEUROSCI.1184-06.2006 [PubMed: 16870732]
- Lorsch ZS, Loh Y-HE, Purushothaman I, Walker DM, Parise EM, Salery M, Cahill ME, Hodes GE, Pfau ML, Kronman H, Hamilton PJ, Issler O, Labonté B, Symonds AE, Zucker M, Zhang TY, Meaney MJ, Russo SJ, Shen L, Bagot RC, Nestler EJ, 2018 Estrogen receptor α drives pro-resilient transcription in mouse models of depression. *Nat Commun* 9, 1116. doi:10.1038/s41467-018-03567-4 [PubMed: 29549264]

- McEwen BS, 2008a Central effects of stress hormones in health and disease: Understanding the protective and damaging effects of stress and stress mediators. *Eur. J. Pharmacol* 583, 174–185. doi:10.1016/j.ejphar.2007.11.071 [PubMed: 18282566]
- McEwen BS, 2008b Central effects of stress hormones in health and disease: Understanding the protective and damaging effects of stress and stress mediators. *Eur. J. Pharmacol* 583, 174–185. doi:10.1016/j.ejphar.2007.11.071 [PubMed: 18282566]
- McEwen BS, 1998 Stress, adaptation, and disease. Allostasis and allostatic load. *Annals of the New York Academy of Sciences* 840, 33–44. [PubMed: 9629234]
- McEwen BS, Bowles NP, Gray JD, Hill MN, Hunter RG, Karatsoreos IN, Nasca C, 2015 Mechanisms of stress in the brain. *Nature Publishing Group* 18, 1353–1363. doi:10.1038/nm.4086
- Meaney MJ, 2001 Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. *Annu. Rev. Neurosci* 24, 1161–1192. doi:10.1146/annurev.neuro.24.1.1161 [PubMed: 11520931]
- Mitra R, Sapolsky RM, 2008 Acute corticosterone treatment is sufficient to induce anxiety and amygdaloid dendritic hypertrophy. *Proc. Natl. Acad. Sci. U.S.A* 105, 5573–5578. doi:10.1073/pnas.0705615105 [PubMed: 18391224]
- Rana S, Nam H, Glover ME, Akil H, Watson SJ, Clinton SM, Kerman IA, 2016 Protective effects of chronic mild stress during adolescence in the low-novelty responder rat. *Stress* 19, 133–138. doi:10.3109/10253890.2015.1108304 [PubMed: 26473581]
- Rao RP, Anilkumar S, McEwen BS, Chattarji S, 2012 Glucocorticoids protect against the delayed behavioral and cellular effects of acute stress on the amygdala. *Biol. Psychiatry* 72, 466–475. doi:10.1016/j.biopsych.2012.04.008 [PubMed: 22572034]
- Reul JM, De Kloet ER, 1985 Two receptor systems for corticosterone in rat brain: microdistribution and differential occupation. *Endocrinology* 117, 2505–2511. doi:10.1210/endo-117-6-2505 [PubMed: 2998738]
- Roosendaal B, 2003 Systems mediating acute glucocorticoid effects on memory consolidation and retrieval. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 27, 1213–1223. doi:10.1016/j.pnpbp.2003.09.015 [PubMed: 14659476]
- Schelling G, Briegel J, Roosendaal B, Stoll C, Rothenhäusler HB, Kapfhammer HP, 2001 The effect of stress doses of hydrocortisone during septic shock on posttraumatic stress disorder in survivors. *Biol. Psychiatry* 50, 978–985. [PubMed: 11750894]
- Schelling G, Kilger E, Roosendaal B, de Quervain DJ-F, Briegel J, Dagge A, Rothenhäusler H-B, Krauseneck T, Nollert G, Kapfhammer H-P, 2004 Stress doses of hydrocortisone, traumatic memories, and symptoms of posttraumatic stress disorder in patients after cardiac surgery: a randomized study. *Biol. Psychiatry* 55, 627–633. doi:10.1016/j.biopsych.2003.09.014 [PubMed: 15013832]
- Schelling G, Stoll C, Kapfhammer HP, Rothenhäusler HB, Krauseneck T, Durst K, Haller M, Briegel J, 1999 The effect of stress doses of hydrocortisone during septic shock on posttraumatic stress disorder and health-related quality of life in survivors. *Critical Care Medicine* 27, 2678–2683. [PubMed: 10628609]
- Schoenfeld TJ, Gould E, 2012 Stress, stress hormones, and adult neurogenesis. *Exp. Neurol* 233, 12–21. doi:10.1016/j.expneurol.2011.01.008 [PubMed: 21281629]
- Spencer-Segal JL, Hyzy RC, Iwashyna TJ, Standiford TJ, 2017 Psychiatric Symptoms in Survivors of the Acute Respiratory Distress Syndrome: Effects of Age, Sex, and Immune Modulation. *Ann Am Thorac Soc* 14, AnnalsATS.201606–468OC–967. doi:10.1513/AnnalsATS.201606-468OC
- Starkman MN, Gebarski SS, Berent S, Scheingart DE, 1992 Hippocampal formation volume, memory dysfunction, and cortisol levels in patients with Cushing’s syndrome. *Biol. Psychiatry* 32, 756–765. [PubMed: 1450290]
- Sjøgaard K, Sjøgaard G, 2017 Physical Activity as Cause and Cure of Muscular Pain: Evidence of Underlying Mechanisms. *Exerc Sport Sci Rev* 45, 136–145. doi:10.1249/JES.000000000000112 [PubMed: 28418998]
- Udelsman R, Ramp J, Gallucci WT, Gordon A, Lipford E, Norton JA, Loriaux DL, Chrousos GP, 1986 Adaptation during surgical stress. A reevaluation of the role of glucocorticoids. *J. Clin. Invest* 77, 1377–1381. doi:10.1172/JCI112443 [PubMed: 3958189]

- van Zuiden M, Geuze E, Willems HLDM, Vermetten E, Maas M, Amarouchi K, Kavelaars A, Heijnen CJ, 2012 Glucocorticoid receptor pathway components predict posttraumatic stress disorder symptom development: a prospective study. *Biol. Psychiatry* 71, 309–316. doi:10.1016/j.biopsych.2011.10.026 [PubMed: 22137507]
- Vogel S, Gerritsen L, van Oostrom I, Arias-Vásquez A, Rijpkema M, Joëls M, Franke B, Tendolkar I, Fernández G, 2014 Linking genetic variants of the mineralocorticoid receptor and negative memory bias: interaction with prior life adversity. *Psychoneuroendocrinology* 40, 181–190. doi:10.1016/j.psyneuen.2013.11.010 [PubMed: 24485490]
- Weis F, Kilger E, Roozendaal B, de Quervain DJ-F, Lamm P, Schmidt M, Schmölz M, Briegel J, Schelling G, 2006 Stress doses of hydrocortisone reduce chronic stress symptoms and improve health-related quality of life in high-risk patients after cardiac surgery: a randomized study. *J. Thorac. Cardiovasc. Surg* 131, 277–282. doi:10.1016/j.jtcvs.2005.07.063 [PubMed: 16434254]
- Woolley CS, Gould E, McEwen BS, 1990 Exposure to excess glucocorticoids alters dendritic morphology of adult hippocampal pyramidal neurons. *Brain Res.* 531, 225–231. [PubMed: 1705153]
- Yehuda R, Pratchett LC, Elmes MW, Lehrner A, Daskalakis NP, Koch E, Makotkine I, Flory JD, Bierer LM, 2014 Glucocorticoid-related predictors and correlates of post-traumatic stress disorder treatment response in combat veterans. *Interface Focus* 4, 20140048–20140048. doi:10.1098/rsfs.2014.0048 [PubMed: 25285201]
- Zohar J, Yahalom H, Kozlovsky N, Cwikel-Hamzany S, Matar MA, Kaplan Z, Yehuda R, Cohen H, 2011 High dose hydrocortisone immediately after trauma may alter the trajectory of PTSD: interplay between clinical and animal studies. *Eur Neuropsychopharmacol* 21, 796–809. doi:10.1016/j.euroneuro.2011.06.001 [PubMed: 21741804]

Highlights

- Challenges or stressors can cause positive adaptations leading to behavioral resilience, and this positive effect of stress is under-appreciated.
- While chronic elevations in glucocorticoids have adverse consequences for the brain and behavior, acute elevations in glucocorticoids during stress may promote resilience in both rodents and humans.
- We encourage future research on the initiation of resilience-promoting pathways during and immediately following acute stressors and the role of glucocorticoids in this process.

Table 1.

Studies showing a role for glucocorticoids during acute stress in behavioral resilience.

Study	Organism	Challenge	Measurement/intervention	Follow-up	Findings
Delahanty et al. 2000	Human	Motor vehicle accident	Post-trauma urinary cortisol (15 hours)	1 month	Lower urinary cortisol predicted higher PTSD symptoms
Schelling et al. 2001	Human	Septic shock	Hydrocortisone during illness (observational)	31 months (mean)	Hydrocortisone was associated with lower PTSD incidence
Schelling et al. 2004	Human	Cardiac surgery	Perioperative hydrocortisone (randomized)	6 months	Hydrocortisone decreased PTSD symptoms
Weis et al. 2006	Human	Cardiac surgery	Hydrocortisone after surgery	6 months	Hydrocortisone decreased PTSD symptoms and increased HRQoL
Zohar et al. 2011	Human	Traumatic event	Hydrocortisone or placebo injection after trauma	Up to 3 months	Hydrocortisone decreased all symptoms
Bienvenu et al. 2013	Human	ARDS	ICU corticosteroid treatment days	2 years	More steroid days predicted fewer PTSD symptoms
Spencer-Segal et al. 2017	Human	ARDS	ICU corticosteroid treatment days	6 months	More steroid days predicted lower anxiety
Zohar et al. 2011	Rat	Predator scent	Corticosterone injection 1 hour after exposure	7 days	Corticosterone decreased anxiety and startle, and prevented dentate dendritic retraction
Cohen et al. 2006	Rat	Predator scent	Corticosterone response to predator scent/Preexposure corticosterone injection	7 days	Lower corticosterone predicted greater anxiety and startle, prevented with corticosterone
Cohen et al. 2008	Rat	Predator scent	Corticosterone injection 1 hour after exposure	30 days	High dose corticosterone decreased anxiety, startle, and freezing
Rao et al. 2012	Rat	Acute restraint	ADX/corticosterone treatment prerestraint	10 days	ADX exacerbated and corticosterone prevented anxiety and BLA dendritic spine increase

Abbreviations: PTSD, Post-traumatic stress disorder; ARDS, Acute Respiratory Distress Syndrome; HRQoL, Health-related quality of life; BLA, basolateral amygdala; ASD, acute stress disorder