



Corrigendum: Randomized, crossover questionnaire survey of acceptabilities of controlled-release mesalazine tablets and granules in ulcerative colitis patients

Keiji Yagisawa¹, Taku Kobayashi², Ryo Ozaki², Shinji Okabayashi², Takahiko Toyonaga², Miki Miura³, Mari Hayashida³, Eiko Saito², Masaru Nakano², Hajime Matsubara¹, Tadakazu Hisamatsu³, Toshifumi Hibi²

¹Department of Pharmacy and ²Center for Advanced IBD Research and Treatment, Kitasato University Kitasato Institute Hospital, Tokyo; ³The Third Department of Internal Medicine, Kyorin University School of Medicine, Tokyo, Japan

https://doi.org/10.5217/ir.2018.00078

Intest Res 2019;17(1):87-93

In the version of this article initially published, Figure 2 contains errors in the content of the questionnaire. On page 89, the figure should be corrected as following.

From:



<p>A Questionnaire 1 (about usability of the tablet)</p> <p>Q I. Did you feel the tablets were hard to take?</p> <p>Q II. In case you answered "yes" in Q I, why? (multiple answers are allowed)</p> <ol style="list-style-type: none">1. Size of the tablet2. Pill number3. Frequency4. I do not like tablets not limited to the CR mesalazine5. Tastes6. Portability7. The other reason <p>Q III. How much percentage of dose did you take correctly?</p>  <p>Q IV. Have you missed the doses once or more often because of the usability of the tablets?</p>	<p>C Questionnaire 3 (on comparison of the tablets and the granules)</p> <p>Q I. Which formulation did you feel easier to take?</p> <p>Q II. In case you answered "tablet" in Q I, why? (multiple answers are allowed)</p> <ol style="list-style-type: none">1. The volume of the granules was large2. The sachet number of the granules was large3. Taking granules is harder than tablets regardless of products4. The tablets taste or smell better than the granules.5. The tablets are easier to carry than the granules.6. The other reason <p>Q III. In case you answered "granules" in Q I, why?</p> <ol style="list-style-type: none">1. The size of the tablet is big2. The tablet number is large3. Taking granules is harder than tablets regardless of products4. The granules taste or smell better than the tablets5. The granules are easier to carry than the tablets6. The other reason <p>Q V. Which formulation did you feel more effective?</p>
<p>B Questionnaire 2 (on the usability of the granules)</p> <p>Q I. Did you feel the granules were hard to take?</p> <p>Q II. In case you answered "yes" in Q I, why? (multiple answers are allowed)</p> <ol style="list-style-type: none">1. Volume of the granules2. Sachet number3. Frequency4. I do not like granules not limited to the CR mesalazine5. Tastes6. Portability7. The other reason <p>Q III. How much percentage of dose did you take correctly?</p>  <p>Q IV. Have you missed the doses once or more often because of the usability of the granules?</p>	

Fig. 2. (A) Questionnaire 1: acceptability of tablets. (B) Questionnaire 2: acceptability of granules. (C) Questionnaire 3: comparison of tablets and granules. All the original questionnaires were written in Japanese. CR, controlled-release.

To:



<p>A Questionnaire 1 (about usability of the tablet)</p> <p>Q I. Did you feel the tablets were hard to take?</p> <p>Q II. In case you answered "yes" in Q I, why? (multiple answers are allowed)</p> <p>1. Size of the tablet 2. Pill number 3. Frequency</p> <p>4. I do not like tablets not limited to the CR mesalazine 5. Tastes</p> <p>6. Portability 7. The other reason</p> <p>Q III. How much percentage of dose did you take correctly?</p>  <p>Q IV. Have you missed the doses once or more often because of the usability of the tablets?</p>	<p>C Questionnaire 3 (on comparison of the tablets and the granules)</p> <p>Q I. Which formulation did you feel easier to take?</p> <p>Q II. In case you answered "tablet" in Q I, why? (multiple answers are allowed)</p> <p>1. The volume of the granules was large</p> <p>2. The sachet number of the granules was large</p> <p>3. Taking granules is harder than tablets regardless of products</p> <p>4. The tablets taste or smell better than the granules.</p> <p>5. The tablets are easier to carry than the granules.</p> <p>6. The other reason</p> <p>Q III. In case you answered "granules" in Q I, why?</p> <p>1. The size of the tablet is big</p> <p>2. The tablet number is large</p> <p>3. Taking tablets is harder than granules regardless of products</p> <p>4. The granules taste or smell better than the tablets</p> <p>5. The granules are easier to carry than the tablets</p> <p>6. The other reason</p> <p>Q V. Which formulation did you feel more effective?</p>
<p>B Questionnaire 2 (on the usability of the granules)</p> <p>Q I. Did you feel the granules were hard to take?</p> <p>Q II. In case you answered "yes" in Q I, why? (multiple answers are allowed)</p> <p>1. Volume of the granules 2. Sachet number 3. Frequency</p> <p>4. I do not like granules not limited to the CR mesalazine 5. Tastes</p> <p>6. Portability 7. The other reason</p> <p>Q III. How much percentage of dose did you take correctly?</p>  <p>Q IV. Have you missed the doses once or more often because of the usability of the granules?</p>	

Fig. 2. (A) Questionnaire 1: acceptability of tablets. (B) Questionnaire 2: acceptability of granules. (C) Questionnaire 3: comparison of tablets and granules. All the original questionnaires were written in Japanese. CR, controlled-release.