



Published in final edited form as:

Int J Cancer. 2020 August 01; 147(3): 675–685. doi:10.1002/ijc.32760.

Abdominal and Gluteofemoral Size and Risk of Liver Cancer: The Liver Cancer Pooling Project

Andrea A. Florio¹, Peter T. Campbell², Xuehong Zhang³, Anne Zeleniuch-Jacquotte⁴, Jean Wactawski-Wende⁵, Stephanie A. Smith-Warner^{6,7}, Rashmi Sinha¹, Tracey G. Simon⁸, Howard D. Sesso^{7,9}, Catherine Schairer¹, Lynn Rosenberg¹⁰, Thomas E. Rohan¹¹, Kim Robien¹², Andrew G. Renehan¹³, Mark P. Purdue¹, Jenny N. Poynter¹⁴, Julie R. Palmer¹⁰, Christina C. Newton², Yunxia Lu¹⁵, Martha S. Linet¹, Linda M. Liao¹, I-Min Lee^{7,9}, Jill Koshiol¹, Cari M. Kitahara¹, Victoria A. Kirsh¹⁶, Jonathan N. Hofmann¹, Barry I. Graubard¹, Edward Giovannucci⁷, J. Michael Gaziano^{9,17}, Susan M. Gapstur², Neal D. Freedman¹, Jane Demuth¹⁸, Dawn Q. Chong¹⁹, Andrew T. Chan^{3,20,8}, Julie E. Buring^{7,9}, Patrick T. Bradshaw²¹, Laura E. Beane Freeman¹, Katherine A. McGlynn¹, Jessica L. Petrick^{1,10}

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD

²Behavioral and Epidemiology Research Group, American Cancer Society, Atlanta, GA

³Channing Division of Network Medicine, Brigham and Women's Hospital, Boston, MA

⁴Department of Population Health, New York University School of Medicine, New York, NY

⁵Department of Epidemiology and Environmental Health, University at Buffalo, Buffalo, NY

⁶Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA

⁷Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, MA

⁸Division of Gastroenterology, Massachusetts General Hospital and Harvard Medical School, Boston, MA

⁹Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital, Boston, MA

¹⁰Slone Epidemiology Center, Boston University, Boston, MA

Corresponding author: Jessica L. Petrick, PhD, MPH, 72 East Concord Street, L-7, Boston, MA 02118 phone: 617-206-6179
jpetrick@bu.edu.

Authors' contributions

Study concept and design: All authors

Acquisition of data: PTC, HZ, AZ, JW, RS, TGS, HDS, CS, LR, TER, KR, MPP, JNP, JRP, CCN, YL, MSL, LML, IL, CMK, VAK, JNH, EG, JMG, SMG, NDF, DQC, ATC, JEB, LEBF, KAM, JLP

Analysis and interpretation of data: All authors

Drafting of the manuscript: JLP, KAM

Critical revision of the manuscript for important intellectual content: All authors

Statistical analysis: AAF, CCN, JD, JLP

Obtained funding: KAM

Study supervision: KAM

Revised the manuscript: AAF, JLP

Approved manuscript: All authors

Conflict of interest: The authors have declared that no conflicts of interest exist.

Consent for publication: N/A

¹¹Department of Epidemiology & Population Health, Albert Einstein College of Medicine, Bronx, NY

¹²Department of Exercise and Nutrition Sciences, Milken Institute School of Public Health, George Washington University, Washington, DC

¹³Faculty Institute of Cancer Sciences, University of Manchester, Manchester, United Kingdom

¹⁴Division of Pediatric Epidemiology and Clinical Research and Masonic Cancer Center, University of Minnesota, Minneapolis, MN

¹⁵Department of Population Health and Disease Prevention, Program in Public Health, University of California Irvine, Irvine, CA

¹⁶Epidemiology Division, Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada

¹⁷VA Boston Healthcare System, Boston, MA

¹⁸Information Management Services Inc., Silver Spring, MD

¹⁹Division of Medical Oncology, National Cancer Centre Singapore, Singapore

²⁰Clinical and Translational Epidemiology Unit, Massachusetts General Hospital, Boston, MA

²¹Division of Epidemiology and Biostatistics, School of Public Health, University of California, Berkeley, CA

Abstract

Obesity is known to be associated with primary liver cancer (PLC), but the separate effects of excess abdominal and gluteofemoral size are unclear. Thus, we examined the association between waist and hip circumference with risk of PLC overall and by histologic type – hepatocellular carcinoma (HCC) and intrahepatic cholangiocarcinoma (ICC). The Liver Cancer Pooling Project is a consortium of prospective cohort studies that includes data from 1,167,244 individuals (PLC n=2,208, HCC n=1,154, ICC n=335). Multivariable-adjusted hazards ratios (HRs) and 95% confidence intervals (CI) were estimated using proportional hazards regression. Waist circumference, per 5 cm increase, was associated with an 11% increased PLC risk (HR=1.11, 95%CI: 1.09–1.14), including when adjusted for hip circumference (HR=1.12, 95%CI: 1.08–1.17) and also when restricted to individuals in a normal body mass index (BMI) range (18.5–<25 kg/m²; HR=1.14, 95%CI: 1.07–1.21). Hip circumference, per 5 cm increase, was associated with a 9% increased PLC risk (HR=1.09, 95%CI: 1.06–1.12), but no association remained after adjustment for waist circumference (HR=0.99, 95%CI: 0.94–1.03). HCC and ICC results were similar. These findings suggest that excess abdominal size is associated with an increased risk of liver cancer, even among individuals considered to have a normal BMI. However, excess gluteofemoral size alone confers no increased risk. Our findings extend prior analyses, which found an association between excess adiposity and risk of liver cancer, by disentangling the separate effects of excess abdominal and gluteofemoral size through utilization of both waist and hip circumference measurements.

Keywords

hepatocellular carcinoma; intrahepatic cholangiocarcinoma; abdominal obesity; gluteofemoral obesity; epidemiology

Introduction

Obesity, most commonly characterized as having a body mass index (BMI, kg/m²) greater than 30, is an established risk factor for primary liver cancer (PLC),¹ including the predominant type of liver cancer, hepatocellular carcinoma (HCC),² and the second most common type, intrahepatic cholangiocarcinoma (ICC).³ Liver cancer is typically predated by oxidative stress and inflammation in the liver, which can result from hepatitis B or C virus (HBV or HCV) infection, aflatoxin consumption, smoking, excess alcohol consumption, diabetes, fatty liver, or obesity.^{2, 4} To examine obesity, many studies have utilized BMI as a proxy. However, the location of excess body size is critically important, as abdominal and gluteofemoral fat deposits have distinct physiologic properties. Abdominal visceral adipose tissue is hypothesized to predispose individuals to metabolic disorders based on its venous drainage directly into liver portal circulation, potentially leading to lipolysis, insulin resistance, and systematic inflammation.⁵⁻⁷ In contrast, gluteofemoral adiposity, which is subcutaneous fat deposited in the hip and thigh region, has venous drainage into the systemic circulation and is associated with a reduced risk of diabetes and dyslipidemia.^{8, 9} The reduced risk of metabolic complications arising as a result of gluteofemoral adiposity may be due to the delayed release of fatty acids and long-term protection to visceral organs compared to abdominal adiposity.⁸

Previous studies have found inverse associations of excess gluteofemoral size, as measured by hip circumference, with adverse outcomes, including cardiovascular disease^{10, 11} and total mortality.¹²⁻¹⁴ However, the association between excess gluteofemoral size and liver cancer is understudied. A recent meta-analysis of European prospective cohort studies examined BMI, waist circumference, hip circumference, and waist-to-hip ratio and reported that waist circumference was associated with an increased risk of all obesity-related cancers; however, the study did not examine excess body size in relation to liver cancer, specifically.¹⁵ Additionally, another study based in the Liver Cancer Pooling Project found that BMI and waist circumference were associated with increased risk of liver cancer but did not examine hip circumference, waist-to-hip ratio, or other classifications of body size.² Thus, the current study aimed to determine the association between excess abdominal and gluteofemoral body size and risk of liver cancer overall and by histologic type.

Methods

Study Population

As described previously,¹⁶ 15 North American-based cohort studies that are members of the National Cancer Institute's Cohort Consortium agreed to participate in the Liver Cancer Pooling Project. Of these, 12 studies contributed individual participant level data on waist and/or hip circumference: NIH-AARP Diet and Health Study (AARP),¹⁷ The Breast Cancer

Detection Demonstration Project (BCDDP),¹⁸ Women's Health Study (WHS),¹⁹ Physicians' Health Study (PHS),²⁰ Health Professionals Follow-Up Study (HPFS),²¹ New York University Women's Health Study (NYUWHS),²² Cancer Prevention Study–II Nutrition Cohort (CPS-II),²³ Iowa Women's Health Study (IWHS),²⁴ Black Women's Health Study (BWHS),²⁵ Women's Health Initiative (WHI),²⁶ Nurses' Health Study (NHS),²⁷ and the Canadian Study of Diet, Lifestyle, and Health (CSDLH)²⁸ (Supporting Table S1). All studies had complete information available on the entire cohort, with the exception of the CSDLH, which employed a case-cohort design. In the CSDLH, a sub-cohort of 6,127 participants was created by randomly selecting an age-stratified sample of participants from the total cohort at baseline (n=73,909).²⁸ There were 1,590,648 study participants that were eligible to be included in the current study. We excluded participants with missing age (n=9,813), no follow-up time (n=69,736), no waist or hip measurements (n=311,626), BMI <15 kg/m² (n=32,156), and prior diagnosis of stomach, colorectal, lung, pancreas, or breast cancer within 5 years preceding PLC diagnosis (i.e., potential primary cancer of another site that metastasized to the liver, n=73). Our final study population included 1,167,244 individuals (707,281 women and 457,755 men). The individual cohorts were approved by the institutional review boards of the participating institutions; LCPP was approved by the NIH Office of Human Subjects Research.

Outcomes

Linkage to state or national cancer registries or medical/pathology record review was used to ascertain incident PLC (defined as International Classification of Diseases, 10th edition [ICD-10] diagnostic code C22). Cases were further classified as HCC (International Classification of Diseases for Oncology, 3rd edition [ICD-O-3] histology codes of 8170–8175) or ICC (ICD-O-3 histology codes of 8032–8033, 8041, 8050, 8070–8071, 8140–8141, 8160, 8260, 8480, 8481, 8490, and 8560). The current study included 2,208 PLC cases, 1,154 HCC cases, 335 ICC cases, and 1,165,036 non-cases.

Exposure

Waist and hip circumference were available from all studies, with the exception of CPSII which only collected waist circumference. Measurements were taken by trained staff (BCDDP, NYUWHS, WHI) or self-measured by participants (AARP, WHS, PHS, HPFS, CPSII, IWHS, BWHS, NHS, CSDLH). Participants who self-measured were provided specific instructions, which usually included a measuring tape and an illustration demonstrating exactly where on the body to measure. Prior studies have reported self-measured waist and hip circumference to be accurate and reproducible.^{29–32} To examine possible effect modification by measurement method in the current analysis, results were stratified by method. Results were similar (data not shown). Waist circumference, in centimeters (cm), was categorized according to previously published literature (women: <70, 70–<80, 80–<90, and 90; men: <90, 90–<100, 100–<110, and 110),² World Health Organization categories (women: <80, 80–<88, and 90; men: <94, 94–<102, and 102),³³ and using sex- and study-specific quartiles. Similarly, waist-to-hip ratio (WHR) was classified according to previously published literature (women: <0.80, 0.80–<0.85, 0.85–<0.90, and 0.90; men: <0.85, 0.85–<0.90, 0.90–<0.95, and 0.95),³⁴ World Health Organization categories (women: <0.85 and 0.85; men: <0.90 and 0.90),³³ and using sex-

and study-specific quartiles. Hip circumference was categorized using sex- and study-specific quartiles. In addition, we created a cross-classification of sex- and study-specific categories based on waist and hip circumference quartiles (Category 1: Both hip and waist circumference below the 75th percentile; Category 2: Hip circumference above the 75th percentile and waist circumference below the 75th percentile; Category 3: Hip circumference below the 75th percentile and waist circumference above the 75th percentile; Category 4: Both hip and waist circumference above the 75th percentile).

Nested Case-Control Study of HBV/HCV

HBV and HCV are known risk factors for PLC but none of the participating cohorts had previously ascertained HBV/HCV status. Thus, a nested case-control study was conducted within LCPP to determine HBV/HCV status among a subset of participants. Serum was selected from all LCPP participants with samples available for determination of HBV and HCV status, which included 185 PLC cases and 419 controls. To determine HBV status, hepatitis B surface antigen (HBsAg) was assayed using the Bio-Rad GS HBsAg 3.0 enzyme immunoassay (Bio-Rad Laboratories, Redmond, WA, USA). To determine HCV status, antibody to hepatitis C virus (anti-HCV) was assessed using the Ortho HCV Version 3.0 ELISA test system (Ortho-Clinical Diagnostics, Inc.). For the current analysis, we examined whether HBV/HCV was a potential confounder for the association between hip or waist circumference and PLC.

Statistical Analysis

Data were harmonized and pooled for analysis. Between-study heterogeneity was assessed by individual participant data (IPD) random-effects meta-analysis using a chi-square test based on the Q statistic and the I² statistic (where 0% indicates no heterogeneity and larger values indicate increasing heterogeneity between studies).³⁵ Cox proportional hazard regression analysis, with follow-up time as the underlying time metric, was used to calculate adjusted hazard ratios (HRs) and 95% confidence intervals (CIs) for the associations of excess body size measurements with PLC, HCC, and ICC, with modification for the case-cohort design of CSDLH.³⁶ In CSDLH, all cases were weighted as one, and non-cases were weighted according to the inverse of their stratum-specific sampling fractions. All participants in the remaining cohorts were weighted as one. Follow-up of the analytic cohort was from time of waist or hip circumference measurement until the occurrence of an event (i.e., incident liver cancer) or right-censoring (i.e., death, other cancer diagnosis, loss to follow-up, or last date of follow-up), whichever occurred first. A cause-specific analysis was utilized, rather than a competing risk analyses, as it does not require independence of the outcome and competing events to estimate relative risk.³⁷ The proportional hazards assumption was tested using an interaction between waist circumference, hip circumference, or waist-to-hip ratio with log(time), as a continuous variable, in models that included confounders. No interactions were observed (p > 0.05). BMI-adjusted (15–<18.5 kg/m², 18.5–<25 kg/m², 25–<30 kg/m², and ≥ 30 kg/m²) models are presented for all body fat distribution measurements. Additionally, results for waist circumference were adjusted for hip circumference, while results for hip circumference were adjusted for waist circumference.

Based on existing literature, potential confounders³⁸ included age, race, sex, alcohol consumption, smoking, and history of diabetes diagnosis. Variables remained in the adjusted model if they were associated with the exposure and outcome and not a mediator between excess body size and liver cancer.³⁹ Final models included age (continuous), race (white, black, other), sex, alcohol consumption (nondrinker and 1.08, 1.09–3.58, 3.59–13.54, >13.54 g/day), and cigarette smoking (never, former, current). Parent study was also adjusted for in all models (AARP, BCDDP, WHS, PHS, HPFS, NYUWHS, CPS-II, IWHS, BWHS, WHI, NHS, and CSDLH). Diabetes is a potential mediator of the association between excess body size and liver cancer and was therefore not included as a potential confounder.

Effect measure modification by sex and menopausal hormone therapy (MHT) use was assessed. We also stratified by BMI category (18.5–<25 kg/m², 25–<30 kg/m², and ≥30 kg/m²). Departures from the null were assessed using likelihood ratio tests to compare regression models with and without a multiplicative term.³⁸ Tests of linear trend were conducted using continuous variables, per 5 cm increase for waist and hip circumference and per 0.05 unit increase for waist-to-hip ratio. All p-values are two-sided. Analyses were conducted using SAS version 9.4 (SAS Institute, Cary, NC).

Data availability

The authors confirm that some access restrictions apply to the data underlying the findings. Data are stored at NCI and initial requests for data may be directed to Katherine McGlynn (mcglynnk@mail.nih.gov).

Ethics approval and consent to participate

The individual cohorts were approved by the institutional review boards of the participating institutions; LCPP was approved by the NIH Office of Human Subjects Research.

Results

Table 1 summarizes participant characteristics. Compared with non-cases, individuals who developed PLC were more likely to be older (median baseline age of cases vs. non-cases: 64 years vs. 60 years), male (68% vs. 39%), heavy alcohol drinkers (quartile 4 of alcohol consumption: 24% vs. 18%), and current/former smokers (67% vs. 55%).

Waist circumference, per 5 cm increase, was associated with an 11% increased risk of PLC (HR=1.11, 95% CI: 1.09–1.14); results were similar by histologic type, HCC (HR=1.14, 95% CI: 1.10–1.17) and ICC (HR=1.11, 95% CI: 1.06–1.16; Table 2). Waist circumference remained associated with an increased risk of PLC after adjustment for BMI (HR=1.08, 95% CI: 1.04–1.12) and hip circumference (HR=1.12, 95% CI: 1.08–1.17), although the association was attenuated in the BMI-adjusted models. The highest sex- and study-specific quartile of waist circumference was associated with a 90% increased risk of PLC compared to the first quartile (HR=1.90, 95% CI: 1.61–2.23); results were similar when examined by *a priori* categories.

Hip circumference, per 5 cm increase, was associated with a 9% increased risk of PLC (HR=1.09, 95% CI: 1.06–1.12), but no association remained after adjustment for BMI (HR=1.01, 95% CI: 0.97–1.05) or waist circumference (HR=0.99, 95% CI: 0.94–1.03; Table 3).

The associations between WHR and liver cancer are shown in Table 4. Participants with roughly equivalent waist and hip circumference (0.95 in men and 0.90 in women) had a 61% increased risk of PLC (HR=1.61, 95% CI: 1.31–1.98). When adjusted for BMI, results were attenuated but remained significant. Estimates for HCC and ICC were similar.

To examine the separate effects of excess abdominal and gluteofemoral size, quartiles of waist and hip circumference were cross-classified (Table 4). Compared to individuals below the 75th percentile for both waist and hip circumference (Category 1), individuals that had an elevated hip circumference (above the 75th percentile) and low waist circumference (below the 75th percentile) had no increased risk of PLC (Category 2 HR=1.04, 95% CI: 0.82–1.32). However, the highest category of waist circumference was associated with an increased risk of PLC for individuals in the lower (Category 3 HR=1.46, 95% CI: 1.20–1.77) or the higher (Category 4 HR=1.67, 95% CI: 1.45–1.92) category of hip circumference.

There was no evidence of effect measure modification by sex (Tables 2–4 and Supplemental Table S2). Although the p-value for interaction by MHT was not statistically significant, the association between excess body size and PLC appeared stronger among women who used MHT (Supplemental Table S3). For example, the highest study-specific quartile of waist circumference was associated with a 3-fold increased risk of PLC among MHT users (HR=2.87, 95% CI: 1.86–4.42), while MHT non-users only had a 35% increased risk (HR=1.35, 95% CI: 0.89–2.03, p-interaction=0.09).

When stratified by BMI categories (Supplemental Table S4), waist circumference was associated with an increased risk of PLC in all BMI categories (18.5–<25 kg/m² HR=1.14, 95% CI: 1.07–1.21, 25–<30 kg/m² HR=1.06, 95% CI: 1.00–1.12, and ≥30 kg/m² HR=1.07, 95% CI: 1.02–1.12).

Low-to-moderate heterogeneity was observed between studies (waist circumference overall $I^2=38.8%$, $p=0.08$, hip circumference overall $I^2=59.2%$, $p=0.006$; Supplemental Figures 1–4).

We tested a subset of both PLC cases (n=185) and matched controls (n=419) for hepatitis B surface antigen (HBsAg) and antibody to HCV (anti-HCV). For HBV, 9 PLC cases (4.9%) and 7 controls (1.7%) were positive for HBsAg. For HCV, 32 PLC cases (17.3%) and 6 controls (1.4%) were positive for anti-HCV. There was no association between HCV or HBV status and waist or hip circumference (data not shown).

Discussion

In our pooled analysis of 12 North American-based prospective cohort studies, comprising over 1.16 million adults, excess abdominal size was associated with a significantly increased risk of PLC. Each 5 cm increase in waist circumference was associated with an 11%

increased risk of PLC. Waist circumference remained associated with an increased risk of PLC, even after adjustment for BMI or hip circumference. Further, when we examined the cross-classification of waist and hip circumference, the highest category of waist circumference was associated with a 46–67% increased risk of PLC, regardless of hip circumference. However, individuals in the highest category of hip, but not waist, circumference had no increased risk of PLC. Results were similar by sex, BMI, MHT use, and for the two main histologic types of PLC, HCC and ICC. Existing heterogeneity reported between studies may be explained by methodological variability in cohort study design (e.g., measurement method), though the estimated HRs were similar across stratifications (i.e., waist circumference per 5 cm increase: self-reported HR=1.10, 95% CI 1.07–1.12; direct measurement HR=1.13, 95% CI 1.07–1.20). Additionally, when we examined the influence of individual studies, whereby we excluded one study at a time and the summary effect estimates were re-estimated, results were similar.

Our findings extend prior analyses, that have found an association between excess adiposity (i.e., BMI ≥ 30 or increased waist circumference) and risk of liver cancer,^{1–3} by disentangling the separate effects of excess abdominal and gluteofemoral size through utilization of both waist and hip circumference measurements. This approach enabled us to identify the region of excess body size associated with liver cancer risk, specifically excess abdominal size. One previous study, utilizing data from seven European cohort studies, reported that one standard deviation (~11 cm) increase in waist circumference was associated with a 13% increased risk of *all* obesity-associated cancers, which included liver cancer.¹⁵ However, this study did not have a sufficient sample size to examine liver cancer as a distinct outcome.

Obesity is a heterogeneous condition, with abdominal and gluteofemoral fat deposits having distinct properties. Though waist circumference and the other anthropometric measurements used here are not exact measurements of body fat, excess abdominal or gluteofemoral size can be indicative of fat in those regions and may be good proxies for risk associated with these specific body fat distributions in both men and women.^{40–42} As reported in the current study, excess abdominal size conferred an increased risk of liver cancer, which was independent of excess generalized or gluteofemoral size. Conversely, we found that excess gluteofemoral size, as measured by hip circumference, conferred no increased risk after accounting for excess abdominal size. Specifically, among those in the group with high hip and low waist circumference, representing excess gluteofemoral size, there was no increased liver cancer risk. While among individuals with a low hip circumference and high waist circumference, representing excess abdominal size, we saw a significantly increased risk of liver cancer, which remained significant after BMI adjustment. Those with high hip and high waist circumference, representing the greatest general excess size, also had a significantly increased risk of liver cancer. This suggests that the obesity-liver cancer association is primarily driven by excess abdominal size. Moreover, excess abdominal size, indicative of abdominal adiposity, is often considered a marker of metabolic dysregulation in overweight or obese individuals. However, we report herein that increased waist circumference conferred an increased risk of liver cancer even among “normal” weight individuals (18.5– <25 kg/m²), which suggests that it could be important to target waist circumference for prevention strategies, even among individuals with a normal BMI.

Why excess abdominal and gluteofemoral size would have different effects on the risk of PLC is not certain, but several plausible mechanisms are hypothesized, presuming that excess size is indicative of adiposity. Visceral, or intraperitoneal, fat accounts for approximately 1/3 of abdominal fat,⁴³ which also includes intraabdominal retroperitoneal fat and subcutaneous fat.⁷ Visceral fat is metabolically active and releases substantial amounts of growth factors, inflammatory markers, free fatty acids (which contribute to insulin resistance), locally produced estrogen, and adipokines, which might contribute to cancer development.^{44, 45} Only visceral, and not subcutaneous, adipose tissue has venous drainage into the portal vein and therefore has a direct connection with the liver. Due to the higher lipolytic activity of visceral versus subcutaneous adipocytes, free fatty acids are more rapidly mobilized from visceral fat cells. Chronic exposure to free fatty acids contributes to β -cell failure and type 2 diabetes.⁴⁴ Additionally, as more fat is accumulated in the midsection, adipose tissue undergoes tissue remodeling to accommodate excess energy storage, resulting in chronic inflammation that can lead to severe hepatic injury.⁴⁶

In contrast to visceral fat, gluteofemoral fat, which is subcutaneous fat deposited in the hip and thigh region, is thought to act as a 'metabolic sink', entrapping fatty acids and preventing lipid overflow in tissues surrounding abdominal organs.^{8, 47} The primary function of subcutaneous adipose tissue is fatty acid storage, which are reintroduced into the circulation in the form of non-esterified fatty acids during periods of fasting, exercise, or starvation.⁸ Additionally, excess hip size is associated with lower risk of various metabolic dysfunctions, including diabetes.^{48–50} In persons with a greater hip circumference compared to waist circumference, excess hip size may offer long-term protection from inflammation and elevated lipid levels.

Liver cancer incidence has notable sex differences, with rates among males being two to three-fold higher than rates among females.⁴ However, sex differences in liver cancer rates are most pronounced around the menopause transition (45–60 years of age), with males having four-fold higher rates than females during this period.⁵¹ While estrone and estradiol production in visceral adipose tissue increases with increasing waist circumference, estradiol is produced more efficiently in subcutaneous fat.⁵² Men and post-menopausal women have more visceral fat deposits than pre-menopausal women,⁵³ but post-menopausal women still have a lower proportion of abdominal fat compared to men.⁵⁴ In the current study, we did not find effect measure modification by sex, which may be in part due to the female study population predominately including post-menopausal women. Additionally, MHT use did not modify the associations among post-menopausal women. However, the associations between excess abdominal size and liver cancer were slightly more pronounced among MHT users compared to non-users. While post-menopausal women are using MHT, they typically do not have adiposity gains seen in post-menopausal women not using MHT. However, we were only able to examine ever use of MHT. Thus, the women classified as using MHT may not have been currently using MHT when they had their waist and hip circumference measurements taken.

The current study has several strengths, including long follow-up time and wide geographic representation of North America. The large sample size of our study provided the statistical power to stratify by BMI, sex, and histologic type within each adiposity measure. Due to the

low incidence of liver cancer in North America and other western countries,⁵⁵ other studies conducted in these geographic locations have been underpowered to examine the association between excess body size distribution and liver cancer risk by BMI, sex, and histologic type. Additionally, we were able to identify HCC and ICC cases through cancer registries or medical record review, minimizing misclassification of metastatic tumors. Most importantly, we were able to mutually adjust our analyses for each body size measure when examining waist and hip circumference. The inclusion of this adjustment allowed us to isolate the effect of each body size distribution profile; without it, all three measures could simply be indicative of general adiposity, which does not offer insight to the potential biological mechanisms involved.

Our study was not without limitations. Waist circumference, hip circumference, and waist-to-hip ratio are proxies for fat deposition, and thus are imperfect measures of abdominal and gluteofemoral adiposity. However, studies have shown that anthropometric measures tend to reliably measure total fat and adiposity risk when compared to gold standard measurements.^{40–42} Several cohorts pooled for this analysis collect only self-measured data on waist and hip circumference, which could result in exposure misclassification. However, this potential limitation was lessened by the detailed instructions given to participants to take these measurements. Additionally, individuals with higher BMI or larger waist circumference tend to underestimate self-measured waist circumference,^{56, 57} which would bias our results towards the null. Another limitation was the lack of information on potential confounders such as physical activity, diet, and HBV and HCV infection status. Data on physical activity and diet either were not ascertained by the parent cohort or could not be harmonized in a meaningful way to examine as potential confounders. For individuals with HBV or HCV status available, there was no association between these potential covariates and the exposures of excess body size. While this suggests that HBV and HCV are not confounders of the association between excess body size and PLC, our sample size was limited to examine this. Lastly, further research is needed to determine whether our results are generalizable to individuals of non-European descent, due to the characteristics of the current study population.

In conclusion, these results suggest that excess abdominal size is associated with an increased risk of liver cancer, independent of excess generalized or gluteofemoral size, and that excess gluteofemoral size does not confer an increased risk of liver cancer independent of waist circumference. Further studies should investigate whether these associations exist in other populations, including groups at high risk for liver cancer. Additionally, in a study population with repeated measures data, it would be valuable to examine risk of liver cancer among those with changing excess adiposity over time (i.e., reduced or increased waist or hip circumference over time). Future studies should also utilize imaging techniques, rather than anthropometric measurements, to further disentangle the association between visceral versus subcutaneous fat and liver cancer. Overall, these results suggest that targeted prevention efforts, focused on reducing abdominal obesity, even among normal weight individuals, could potentially be an important intervention opportunity for liver cancer, and potentially other obesity-related cancers.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

For the Black Women's Health Study, pathology data were obtained from several of the following state cancer registries (AZ, CA, CO, CT, DE, DC, FL, GA, IL, IN, KY, LA, MD, MA, MI, NJ, NY, NC, OK, PA, SC, TN, TX, VA), and results reported do not necessarily represent their views.

For the Nurses' Health Study and the Health Professionals Follow-up Study, we would like to thank the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY. The authors assume full responsibility for analyses and interpretation of these data.

For NIH-AARP, the acknowledgement can be found at the following website: <https://dietandhealth.cancer.gov/acknowledgement.html>. For the Women's Health Initiative, the full list of investigators that have contributed can be found on the following website: <https://www.whi.org/researchers/Documents%20%20Write%20a%20Paper/WHI%20Investigator%20Long%20List.pdf>.

Funding: NIH Intramural Research Program, National Cancer Institute (JL Petrick, AA Florio, LE Beane-Freeman, JN Hofmann, CM Kitahara, ND Freedman, BI Graubard, J Koshiol, LM Liao, MS Linet, MP Purdue, C Schairer, R Sinha, KA McGlynn). National Institutes of Health grants CA047988 (I Lee, JE Buring), HL043851 (I Lee, JE Buring), HL080467 (I Lee, JE Buring), HL099355 (I Lee, JE Buring), K07 CA188126 (X Zhang), DK098311 (AT Chan), CA186107 (AT Chan), CA87969 (AT Chan), CA167552 (AT Chan), R01 CA39742 (K Robien, J Poynter), UM1 CA182934 (A Zeleniuch-Jacquotte), P30 CA016087 (A Zeleniuch-Jacquotte), P30 ES000260 (A Zeleniuch-Jacquotte), U01 CA164974 (L Rosenberg, JR Palmer), and R01 CA058420 (L Rosenberg, JR Palmer). American Cancer Society Research Scholar Grant RSG NEC-130476 (X Zhang). The WHI program (J Wactawski-Wende, TE Rohan) is funded by the National Institutes of Health contracts HHSN268201600018C, HHSN268201600001C, HHSN268201600002C, HHSN268201600003C, and HHSN268201600004C. The HPFS and NHS programs (X Zhang) were support by the National Cancer Institute, National Institutes of Health grant numbers UM1CA186107, P50CA127003, P01CA87969, and UM1CA167552.

Abbreviations

HCC	hepatocellular carcinoma
ICC	intrahepatic cholangiocarcinoma
HBV	hepatitis B virus
HCV	hepatitis C infection
BMI	body mass index
AARP	NIH-AARP Diet and Health Study
BCDDP	The Breast Cancer Detection Demonstration Project
WHS	Women's Health Study
PHS	Physicians' Health Study
HPFS	Health Professionals Follow-Up Study
NYUWHS	New York University Women's Health Study
CPS-II	Cancer Prevention Study—II Nutrition Cohort

IWHS	Iowa Women's Health Study
BWHS	Black Women's Health Study
WHI	Women's Health Initiative
NHS	Nurses' Health Study
CSDLH	Canadian Study of Diet, Lifestyle, and Health
WHR	waist-to-hip ratio
ICD-10	International Classification of Diseases, 10 th edition
ICD-O-3	International Classification of Diseases for Oncology, 3 rd edition
NAFLD	non-alcoholic fatty liver disease
HBsAg	hepatitis B surface antigen
anti-HCV	antibody to hepatitis C virus

References

1. Lauby-Secretan B, Scoccianti C, Loomis D, Grosse Y, Bianchini F, Straif K, Handbook IARC. Body Fatness and Cancer - Viewpoint of the IARC Working Group. *New Engl J Med* 2016;375: 794–8. [PubMed: 27557308]
2. Campbell PT, Newton CC, Freedman ND, Koshiol J, Alavanja MC, Beane Freeman LE, Buring JE, Chan AT, Chong DQ, Datta M, Gaudet MM, Gaziano JM, et al. Body Mass Index, Waist Circumference, Diabetes, and Risk of Liver Cancer for U.S. Adults. *Cancer research* 2016;76: 6076–83. [PubMed: 27742674]
3. Petrick JL, Thistle JE, Zeleniuch-Jacquotte A, Zhang X, Wactawski-Wende J, Van Dyke AL, Stampfer MJ, Sinha R, Sesso HD, Schairer C, Rosenberg L, Rohan TE, et al. Body Mass Index, Diabetes and Intrahepatic Cholangiocarcinoma Risk: The Liver Cancer Pooling Project and Meta-analysis. *The American journal of gastroenterology* 2018;113: 1494–505. [PubMed: 30177781]
4. McGlynn KA, Petrick JL, London WT. Global epidemiology of hepatocellular carcinoma: an emphasis on demographic and regional variability. *Clinics in liver disease* 2015;19: 223–38. [PubMed: 25921660]
5. Yang X, Smith U. Adipose tissue distribution and risk of metabolic disease: does thiazolidinedione-induced adipose tissue redistribution provide a clue to the answer? *Diabetologia* 2007;50: 1127–39. [PubMed: 17393135]
6. Sun B, Karin M. Obesity, inflammation, and liver cancer. *Journal of hepatology* 2012;56: 704–13. [PubMed: 22120206]
7. Ibrahim MM. Subcutaneous and visceral adipose tissue: structural and functional differences. *Obes Rev* 2010;11: 11–8. [PubMed: 19656312]
8. Manolopoulos KN, Karpe F, Frayn KN. Gluteofemoral body fat as a determinant of metabolic health. *International journal of obesity (2005)* 2010;34: 949–59. [PubMed: 20065965]
9. Snijder MB, Zimmet PZ, Visser M, Dekker JM, Seidell JC, Shaw JE. Independent and opposite associations of waist and hip circumferences with diabetes, hypertension and dyslipidemia: the AusDiab Study. *Int J Obes Relat Metab Disord* 2004;28: 402–9. [PubMed: 14724659]
10. Zhang C, Rexrode KM, van Dam RM, Li TY, Hu FB. Abdominal obesity and the risk of all-cause, cardiovascular, and cancer mortality: sixteen years of follow-up in US women. *Circulation* 2008;117: 1658–67. [PubMed: 18362231]

11. Parker ED, Pereira MA, Stevens J, Folsom AR. Association of hip circumference with incident diabetes and coronary heart disease: the Atherosclerosis Risk in Communities study. *American journal of epidemiology* 2009;169: 837–47. [PubMed: 19224980]
12. Lissner L, Bjorkelund C, Heitmann BL, Seidell JC, Bengtsson C. Larger hip circumference independently predicts health and longevity in a Swedish female cohort. *Obesity research* 2001;9: 644–6. [PubMed: 11595782]
13. Heitmann BL, Frederiksen P, Lissner L. Hip circumference and cardiovascular morbidity and mortality in men and women. *Obesity research* 2004;12: 482–7. [PubMed: 15044665]
14. Bigaard J, Frederiksen K, Tjonneland A, Thomsen BL, Overvad K, Heitmann BL, Sorensen TI. Waist and hip circumferences and all-cause mortality: usefulness of the waist-to-hip ratio? *Int J Obes Relat Metab Disord* 2004;28: 741–7. [PubMed: 15052280]
15. Freisling H, Arnold M, Soerjomataram I, O’Doherty MG, Ordonez-Mena JM, Bamia C, Kampman E, Leitzmann M, Romieu I, Kee F, Tsilidis K, Tjonneland A, et al. Comparison of general obesity and measures of body fat distribution in older adults in relation to cancer risk: meta-analysis of individual participant data of seven prospective cohorts in Europe. *British journal of cancer* 2017;116: 1486–97. [PubMed: 28441380]
16. McGlynn KA, Sahasrabudhe VV, Campbell PT, Graubard BI, Chen J, Schwartz LM, Petrick JL, Alavanja MC, Andreotti G, Boggs DA, Buring JE, Chan AT, et al. Reproductive factors, exogenous hormone use and risk of liver cancer among U.S. women: Results from the Liver Cancer Pooling Project. *British journal of cancer* 2015;(In Press).
17. Schatzkin A, Subar AF, Thompson FE, Harlan LC, Tangrea J, Hollenbeck AR, Hurwitz PE, Coyle L, Schussler N, Michaud DS, Freedman LS, Brown CC, et al. Design and serendipity in establishing a large cohort with wide dietary intake distributions : the National Institutes of Health-American Association of Retired Persons Diet and Health Study. *American journal of epidemiology* 2001;154: 1119–25. [PubMed: 11744517]
18. Flood A, Velie EM, Chatterjee N, Subar AF, Thompson FE, Lacey JV, Jr., Schairer C, Troisi R, Schatzkin A. sFruit and vegetable intakes and the risk of colorectal cancer in the Breast Cancer Detection Demonstration Project follow-up cohort. *The American journal of clinical nutrition* 2002;75: 936–43. [PubMed: 11976170]
19. Rexrode KM, Lee IM, Cook NR, Hennekens CH, Buring JE. Baseline characteristics of participants in the Women’s Health Study. *Journal of women’s health & gender-based medicine* 2000;9: 19–27.
20. Steering Committee of the PHS Research Group. Final report on the aspirin component of the ongoing Physicians’ Health Study. *The New England journal of medicine* 1989;321: 129–35. [PubMed: 2664509]
21. Grobbee DE, Rimm EB, Giovannucci E, Colditz G, Stampfer M, Willett W. Coffee, caffeine, and cardiovascular disease in men. *The New England journal of medicine* 1990;323: 1026–32. [PubMed: 2215561]
22. Toniolo PG, Pasternack BS, Shore RE, Sonnenschein E, Koenig KL, Rosenberg C, Strax P, Strax S. Endogenous hormones and breast cancer: a prospective cohort study. *Breast cancer research and treatment* 1991;18 Suppl 1: S23–6. [PubMed: 1873553]
23. Calle EE, Rodriguez C, Jacobs EJ, Almon ML, Chao A, McCullough ML, Feigelson HS, Thun MJ. The American Cancer Society Cancer Prevention Study II Nutrition Cohort: rationale, study design, and baseline characteristics. *Cancer* 2002;94: 2490–501. [PubMed: 12015775]
24. Munger RG, Folsom AR, Kushi LH, Kaye SA, Sellers TA. Dietary assessment of older Iowa women with a food frequency questionnaire: nutrient intake, reproducibility, and comparison with 24-hour dietary recall interviews. *American journal of epidemiology* 1992;136: 192–200. [PubMed: 1415141]
25. Rosenberg L, Adams-Campbell L, Palmer JR. The Black Women’s Health Study: a follow-up study for causes and preventions of illness. *Journal of the American Medical Women’s Association* 1995;50: 56–8.
26. Anderson GL, Manson J, Wallace R, Lund B, Hall D, Davis S, Shumaker S, Wang CY, Stein E, Prentice RL. Implementation of the Women’s Health Initiative study design. *Annals of epidemiology* 2003;13: S5–17. [PubMed: 14575938]

27. Belanger CF, Hennekens CH, Rosner B, Speizer FE. The nurses' health study. *The American journal of nursing* 1978;78: 1039–40. [PubMed: 248266]
28. Rohan TE, Soskolne CL, Carroll KK, Kreiger N. The Canadian Study of Diet, Lifestyle, and Health: design and characteristics of a new cohort study of cancer risk. *Cancer detection and prevention* 2007;31: 12–7. [PubMed: 17303348]
29. Spencer EA, Roddam AW, Key TJ. Accuracy of self-reported waist and hip measurements in 4492 EPIC-Oxford participants. *Public Health Nutrition* 2004;7: 723–7. [PubMed: 15369609]
30. Barrios P, Martin-Biggers J, Quick V, Byrd-Bredbenner C. Reliability and criterion validity of self-measured waist, hip, and neck circumferences. *BMC medical research methodology* 2016;16: 49. [PubMed: 27145829]
31. KUSHI LH, KAYE SA, FOLSOM AR, SOLER JT, PRINEAS RJ. ACCURACY AND RELIABILITY OF SELF-MEASUREMENT OF BODY GIRTHS. *American journal of epidemiology* 1988;128: 740–8. [PubMed: 3421240]
32. Freudenheim JL, Darrow SL. Accuracy of self-measurement of body fat distribution by waist, hip, and thigh circumferences. *Nutrition and Cancer* 1991;15: 179–86. [PubMed: 1866312]
33. World Health Organization. *Waist Circumference and Waist-Hip Ratio Report of a WHO Expert Consultation* 2008.
34. Dobbelsteyn CJ, Joffres MR, MacLean DR, Flowerdew G. A comparative evaluation of waist circumference, waist-to-hip ratio and body mass index as indicators of cardiovascular risk factors. *The Canadian Heart Health Surveys. Int J Obes Relat Metab Disord* 2001;25: 652–61. [PubMed: 11360147]
35. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;327: 557. [PubMed: 12958120]
36. Langholz B, Jiao J. Computational methods for case-cohort studies. *Comput Stat Data An* 2007;51: 3737–48.
37. Prentice RL, Kalbfleisch JD, Peterson AV Jr., Flournoy N, Farewell VT, Breslow NE. The analysis of failure times in the presence of competing risks. *Biometrics* 1978;34: 541–54. [PubMed: 373811]
38. Rothman KJ, Greenland S, Lash TL. *Modern epidemiology*, 3rd ed. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins, 2008 x, 758 p.
39. Kleinbaum DG, Klein M. *Survival analysis : a self-learning text*, 3rd ed. New York: Springer, 2012 xv, 700 p.
40. Janssen I, Katzmarzyk PT, Ross R. Waist circumference and not body mass index explains obesity-related health risk. *The American journal of clinical nutrition* 2004;79: 379–84. [PubMed: 14985210]
41. Grundy SM, Neeland IJ, Turer AT, Vega GL. Waist circumference as measure of abdominal fat compartments. *J Obes* 2013;2013: 454285. [PubMed: 23762536]
42. Swainson MG, Batterham AM, Tsakirides C, Rutherford ZH, Hind K. Prediction of whole-body fat percentage and visceral adipose tissue mass from five anthropometric variables. *PloS one* 2017;12: e0177175–e. [PubMed: 28493988]
43. Marin P, Andersson B, Ottosson M, Olbe L, Chowdhury B, Kvist H, Holm G, Sjostrom L, Bjorntorp P. The morphology and metabolism of intraabdominal adipose tissue in men. *Metabolism* 1992;41: 1242–8. [PubMed: 1435298]
44. Wajchenberg BL. Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev* 2000;21: 697–738. [PubMed: 11133069]
45. Renehan AG, Zwahlen M, Egger M. Adiposity and cancer risk: new mechanistic insights from epidemiology. *Nat Rev Cancer* 2015;15: 484–98. [PubMed: 26205341]
46. Karagozian R, Derdák Z, Baffy G. Obesity-associated mechanisms of hepatocarcinogenesis. *Metabolism* 2014;63: 607–17. [PubMed: 24629562]
47. Frayn K. Adipose tissue as a buffer for daily lipid flux. *Diabetologia* 2002;45: 1201–10. [PubMed: 12242452]
48. Yusuf S, Hawken S, Ôunpuu S, Bautista L, Franzosi MG, Commerford P, Lang CC, Rumboldt Z, Onen CL, Lisheng L, Tanomsup S, Wangai P, et al. Obesity and the risk of myocardial infarction in 27 000 participants from 52 countries: a case-control study. *The Lancet* 2005;366: 1640–9.

49. Snijder MB, Zimmet PZ, Visser M, Dekker JM, Seidell JC, Shaw JE. Independent and opposite associations of waist and hip circumferences with diabetes, hypertension and dyslipidemia: the AusDiab Study. *International Journal Of Obesity* 2004;28: 402. [PubMed: 14724659]
50. Canoy D Distribution of body fat and risk of coronary heart disease in men and women. *Current opinion in cardiology* 2008;23: 591–8. [PubMed: 18830075]
51. Bray F, Parkin DM. Evaluation of data quality in the cancer registry: Principles and methods. Part I: Comparability, validity and timeliness. *European Journal of Cancer* 2009;45: 747–55. [PubMed: 19117750]
52. Hetemaki N, Savolainen-Peltonen H, Tikkanen MJ, Wang F, Paatela H, Hamalainen E, Turpeinen U, Haanpaa M, Vihma V, Mikkola TS. Estrogen Metabolism in Abdominal Subcutaneous and Visceral Adipose Tissue in Postmenopausal Women. *J Clin Endocrinol Metab* 2017;102: 4588–95. [PubMed: 29029113]
53. Brown LM, Clegg DJ. Central effects of estradiol in the regulation of food intake, body weight, and adiposity. *J Steroid Biochem Mol Biol* 2010;122: 65–73. [PubMed: 20035866]
54. Ley CJ, Lees B, Stevenson JC. Sex- and menopause-associated changes in body-fat distribution. *The American journal of clinical nutrition* 1992;55: 950–4. [PubMed: 1570802]
55. Petrick JL, Braunlin M, Laversanne M, Valery PC, Bray F, McGlynn KA. International trends in liver cancer incidence, overall and by histologic subtype, 1978–2007. *Int J Cancer* 2016;139: 1534–45. [PubMed: 27244487]
56. Bigaard J, Spanggaard I, Thomsen BL, Overvad K, Tjonneland A. Self-reported and technician-measured waist circumferences differ in middle-aged men and women. *J Nutr* 2005;135: 2263–70. [PubMed: 16140909]
57. Connor Gorber S, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2007;8: 307–26. [PubMed: 17578381]

Novelty and Impact

Utilizing data from 12 North American-based, prospective studies with over 2,200 liver cancer cases within the Liver Cancer Pooling Project, we report that excess abdominal size is associated with an increased risk of liver cancer, even among individuals considered to have a normal body mass index. However, excess gluteofemoral size alone confers no increased risk.

Table 1.

Characteristics of participants in the Liver Cancer Pooling Project.

	Non-cases (N=1,165,036)	Primary Liver Cancer (N=2,208)	Hepatocellular Carcinoma (N=1,154)	Intrahepatic Cholangiocarcinoma (N=335)
Age at entry, N (%)				
<50	151,234 (13.0)	55 (2.5)	25 (2.2)	7 (2.1)
50–59	395,084 (33.9)	572 (25.9)	311 (26.9)	87 (26.0)
60–69	531,168 (45.6)	1404 (63.6)	735 (63.7)	210 (62.7)
70	87,550 (7.5)	177 (8.0)	83 (7.2)	31 (9.3)
Sex, N (%)				
Male	457,755 (39.3)	1495 (67.7)	835 (72.4)	180 (53.7)
Female	707,281 (60.7)	713 (32.3)	319 (27.6)	155 (46.3)
Race, N (%)				
White	1,010,667 (86.7)	1913 (86.6)	974 (84.4)	298 (89.0)
Black	97,143 (8.3)	99 (4.5)	54 (4.7)	12 (3.6)
Asian/Pacific Islander	9,888 (0.8)	25 (1.1)	15 (1.3)	4 (1.2)
American Indian/Alaskan Native	2,626 (0.2)	8 (0.4)	4 (0.3)	2 (0.6)
Other	31,104 (2.7)	126 (5.7)	84 (7.3)	18 (5.4)
Missing	13,608 (1.2)	37 (1.7)	23 (2.0)	1 (0.3)
Alcohol, ever, N (%)				
No	299,308 (25.7)	623 (28.2)	345 (29.9)	75 (22.4)
Yes	851,745 (73.1)	1562 (70.7)	797 (69.1)	256 (76.4)
Unknown	13,983 (1.2)	23 (1.0)	12 (1.0)	4 (1.2)
Alcohol, grams/day, N (%)				
Non-drinker	299,308 (25.7)	623 (28.2)	345 (29.9)	75 (22.4)
Quartile 1: 1.08	200,161 (17.2)	343 (15.5)	179 (15.5)	66 (19.7)
Quartile 2: 1.09–3.58	205,391 (17.6)	357 (16.2)	173 (15.0)	55 (16.4)
Quartile 3: 3.59–13.54	216,550 (18.6)	320 (14.5)	150 (13.0)	51 (15.2)
Quartile 4: >13.54	214,283 (18.4)	533 (24.1)	288 (25.0)	83 (24.8)
Missing	29,343 (2.5)	32 (1.4)	19 (1.6)	5 (1.5)
Cigarette smoking, N (%)				
Never smoker	499,215 (42.8)	670 (30.3)	335 (29.0)	114 (34.0)
Former smoker	493,902 (42.4)	1162 (52.6)	627 (54.3)	175 (52.2)
Current smoker	143,883 (12.4)	308 (13.9)	153 (13.3)	38 (11.3)
Ever smoker	1,445 (0.1)	0 (0.0)	0 (0.0)	0 (0.0)
Missing	26,591 (2.3)	68 (3.1)	39 (3.4)	8 (2.4)

Table 2.

Adjusted^d hazard ratios (HR) and 95% confidence intervals (CI) for associations between waist Circumference and hepatocellular carcinoma and intrahepatic cholangiocarcinoma risk, Liver Cancer Pooling Project.

	Primary Liver Cancer						Hepatocellular Carcinoma						Intrahepatic Cholangiocarcinoma												
	Waist Circumference		Non-case N		Hip Circumference Adjusted		HR (95% CI)		BMI Adjusted		Case N		HR (95% CI)		Hip Circumference Adjusted		BMI Adjusted		Case N		HR (95% CI)				
	Case N	Non-case N	HR (95% CI)	HR (95% CI)	Case N	Non-case N	HR (95% CI)	HR (95% CI)	Case N	Non-case N	HR (95% CI)	HR (95% CI)	Case N	Non-case N	HR (95% CI)	HR (95% CI)	Case N	Non-case N	HR (95% CI)	HR (95% CI)	Case N	Non-case N	HR (95% CI)	HR (95% CI)	
Centimeters																									
M: <90, F: <70	149	122,764	1.00	1.00	130	103,707	1.00	1.00	1.00	1.00	75	1.00	1.00	1.00	60	1.00	1.00	1.00	19	1.00	1.00	1.00	17	1.00	
M: 90-<100, F: 70-<80	337	257,505	1.16 (0.97-1.39)	1.10 (0.90-1.34)	286	212,187	1.10 (0.90-1.34)	1.15 (0.89-1.49)	1.03 (0.79-1.34)	1.12 (0.83-1.50)	174	1.15 (0.89-1.49)	1.03 (0.79-1.34)	1.11 (0.83-1.50)	139	1.03 (0.79-1.34)	1.11 (0.83-1.50)	1.11 (0.83-1.50)	58	1.11 (0.83-1.50)	1.11 (0.83-1.50)	1.11 (0.83-1.50)	50	1.37 (0.78-2.38)	
M: 100-<110, F: 80-<90	341	208,677	1.62 (1.35-1.95)	1.60 (1.28-1.99)	313	170,882	1.60 (1.28-1.99)	1.64 (1.26-2.12)	1.26 (0.95-1.68)	1.66 (1.20-2.30)	171	1.64 (1.26-2.12)	1.26 (0.95-1.68)	1.66 (1.20-2.30)	153	1.26 (0.95-1.68)	1.66 (1.20-2.30)	2.01 (1.20-3.38)	59	2.01 (1.20-3.38)	2.01 (1.20-3.38)	2.01 (1.20-3.38)	54	1.83 (1.03-3.25)	
M: 110, F: 90	320	192,688	2.06 (1.70-2.50)	1.88 (1.44-2.47)	284	161,050	1.88 (1.44-2.47)	2.25 (1.71-2.96)	1.35 (0.94-1.94)	2.04 (1.37-3.03)	177	2.25 (1.71-2.96)	1.35 (0.94-1.94)	2.04 (1.37-3.03)	151	1.35 (0.94-1.94)	2.04 (1.37-3.03)	2.58 (1.52-4.39)	61	2.58 (1.52-4.39)	2.02 (0.99-4.14)	2.02 (0.99-4.14)	55	2.08 (1.04-4.16)	
<i>P_{interaction}</i>			0.5	0.5			0.5	0.8				0.8						1.0							
WHO cutpoints, cm																									
M: <94, F: <80	365	324,570	1.00	1.00	321	275,332	1.00	1.00	1.12 (0.96-1.30)	1.12 (0.91-1.38)	177	1.00	1.00	1.00	146	1.00	1.00	1.00	60	1.00	1.00	1.00	54	1.00	
M: 94-<102, F: 80-<88	300	197,508	1.24 (1.08-1.43)	1.18 (1.00-1.38)	265	159,991	1.18 (1.00-1.38)	1.31 (1.07-1.60)	1.12 (0.91-1.38)	1.24 (0.97-1.57)	158	1.31 (1.07-1.60)	1.12 (0.91-1.38)	1.24 (0.97-1.57)	133	1.12 (0.91-1.38)	1.24 (0.97-1.57)	1.42 (0.98-2.05)	55	1.42 (0.98-2.05)	1.27 (0.86-1.87)	1.27 (0.86-1.87)	49	1.28 (0.86-1.91)	
M: 102, F: 88	482	259,556	1.66 (1.46-1.89)	1.47 (1.22-1.76)	427	212,503	1.47 (1.22-1.76)	1.84 (1.53-2.22)	1.22 (0.96-1.55)	1.61 (1.24-2.10)	262	1.84 (1.53-2.22)	1.22 (0.96-1.55)	1.61 (1.24-2.10)	224	1.22 (0.96-1.55)	1.61 (1.24-2.10)	1.72 (1.24-2.39)	82	1.72 (1.24-2.39)	1.29 (0.82-2.05)	1.29 (0.82-2.05)	73	1.32 (0.85-2.06)	
<i>P_{interaction}</i>			0.9	0.9			0.9	0.5				0.5						0.6							
Sex- and study-specific																									
Quartile 1	205	207,659	1.00	1.00	178	172,277	1.00	1.00	1.00	1.00	102	1.00	1.00	1.00	81	1.00	1.00	1.00	33	1.00	1.00	1.00	30	1.00	

Waist Circumference	Primary Liver Cancer										Hepatocellular Carcinoma										Intrahepatic Cholangiocarcinoma													
	Non-case		Case		HR (95% CI)		BMI Adjusted		Hip Circumference Adjusted		HR (95% CI)		Case N		HR (95% CI)		BMI Adjusted		Hip Circumference Adjusted		HR (95% CI)		Case N		HR (95% CI)		BMI Adjusted		Hip Circumference Adjusted		HR (95% CI)		Case N	
	N	N	N	N	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI	HR	95% CI		
Quartile 2	203,479	250	1.14 (0.95-1.36)	1.14 (0.95-1.36)	1.08 (0.89-1.31)	1.08 (0.89-1.31)	1.06 (0.89-1.27)	1.06 (0.89-1.27)	1.18 (0.92-1.51)	1.18 (0.92-1.51)	1.08 (0.89-1.31)	1.08 (0.89-1.31)	1.06 (0.82-1.36)	1.06 (0.82-1.36)	1.14 (0.86-1.51)	1.14 (0.86-1.51)	1.18 (0.92-1.51)	1.18 (0.92-1.51)	1.06 (0.82-1.36)	1.06 (0.82-1.36)	1.14 (0.86-1.51)	1.14 (0.86-1.51)	1.06 (0.82-1.36)	1.06 (0.82-1.36)	1.14 (0.86-1.51)	1.14 (0.86-1.51)	1.18 (0.92-1.51)	1.18 (0.92-1.51)	1.06 (0.82-1.36)	1.06 (0.82-1.36)	1.14 (0.86-1.51)	1.14 (0.86-1.51)		
Quartile 3	185,588	294	1.48 (1.25-1.75)	1.48 (1.25-1.75)	1.48 (1.21-1.80)	1.48 (1.21-1.80)	1.30 (1.08-1.56)	1.30 (1.08-1.56)	1.48 (1.16-1.89)	1.48 (1.16-1.89)	1.48 (1.21-1.80)	1.48 (1.21-1.80)	1.19 (0.92-1.54)	1.19 (0.92-1.54)	1.53 (1.14-2.05)	1.53 (1.14-2.05)	1.48 (1.16-1.89)	1.48 (1.16-1.89)	1.19 (0.92-1.54)	1.19 (0.92-1.54)	1.53 (1.14-2.05)	1.53 (1.14-2.05)	1.19 (0.92-1.54)	1.19 (0.92-1.54)	1.53 (1.14-2.05)	1.53 (1.14-2.05)	1.19 (0.92-1.54)	1.19 (0.92-1.54)	1.53 (1.14-2.05)	1.53 (1.14-2.05)				
Quartile 4	184,908	398	1.90 (1.61-2.23)	1.90 (1.61-2.23)	1.81 (1.45-2.26)	1.81 (1.45-2.26)	1.46 (1.18-1.80)	1.46 (1.18-1.80)	2.10 (1.67-2.64)	2.10 (1.67-2.64)	1.81 (1.45-2.26)	1.81 (1.45-2.26)	1.35 (1.00-1.82)	1.35 (1.00-1.82)	1.95 (1.41-2.71)	1.95 (1.41-2.71)	2.10 (1.67-2.64)	2.10 (1.67-2.64)	1.35 (1.00-1.82)	1.35 (1.00-1.82)	1.95 (1.41-2.71)	1.95 (1.41-2.71)	1.35 (1.00-1.82)	1.35 (1.00-1.82)	1.95 (1.41-2.71)	1.95 (1.41-2.71)	1.35 (1.00-1.82)	1.35 (1.00-1.82)	1.95 (1.41-2.71)	1.95 (1.41-2.71)				
$P_{interaction}^c$			0.3	0.3					0.3	0.3								0.3	0.3															
Per 5 cm increase			1.11 (1.09-1.14)	1.11 (1.09-1.14)	1.12 (1.08-1.17)	1.12 (1.08-1.17)	1.08 (1.04-1.12)	1.08 (1.04-1.12)	1.14 (1.10-1.17)	1.14 (1.10-1.17)	1.12 (1.08-1.17)	1.12 (1.08-1.17)	1.07 (1.02-1.13)	1.07 (1.02-1.13)	1.15 (1.09-1.21)	1.15 (1.09-1.21)	1.14 (1.10-1.17)	1.14 (1.10-1.17)	1.07 (1.02-1.13)	1.07 (1.02-1.13)	1.15 (1.09-1.21)	1.15 (1.09-1.21)	1.07 (1.02-1.13)	1.07 (1.02-1.13)	1.15 (1.09-1.21)	1.15 (1.09-1.21)	1.14 (1.10-1.17)	1.14 (1.10-1.17)	1.07 (1.02-1.13)	1.07 (1.02-1.13)				
P_{trend}^b			<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001			
$P_{interaction}^c$			0.6	0.6					0.5	0.5								0.5	0.5															

^a Adjusted for age (continuous), race (white, black, other), sex, alcohol consumption (nondrinker and AARP, BCDDP, WHS, PHS, HPFS, NYUWHS, CPS-II, IWHS, BWHS, WHI, NHS, and CSDLH).

^b P_{trend} from Wald tests.

^c $P_{interaction}$ with sex from log-likelihood ratio tests.

Table 3.

Adjusted^a hazard ratios (HR) and 95% confidence intervals (CI) for associations between waist circumference and hepatocellular carcinoma and intrahepatic cholangiocarcinoma risk, Liver Cancer Pooling Project.

Hip Circumference	Primary Liver Cancer						Hepatocellular Carcinoma						Intrahepatic Cholangiocarcinoma																		
	Non-case N		Case N		Waist Circumference Adjusted		HR (95% CI)		BMI Adjusted		Case N		HR (95% CI)		Waist Circumference Adjusted		Case N		HR (95% CI)		BMI Adjusted		Case N		HR (95% CI)		Waist Circumference Adjusted				
Sex- and study-specific																															
Quartile 1	183,297	238	1.00	1.00	182,631	236	1.00	1.00	1.00	1.00	110	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	
Quartile 2	155,733	198	1.00	1.00	155,346	198	0.88	0.88	1.14	1.14	107	1.03	1.03	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	
			(0.83-1.19)	(0.77-1.10)			(0.73-1.05)	(0.81-1.16)	(0.89-1.47)	(0.89-1.47)		(0.80-1.32)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	(0.76-1.27)	
Quartile 3	159,909	267	1.26	1.26	159,500	267	0.97	0.97	1.22	1.22	117	0.95	0.95	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89	0.89
			(1.07-1.48)	(0.89-1.26)			(0.81-1.16)	(0.81-1.16)	(0.95-1.56)	(0.95-1.56)		(0.73-1.23)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	(0.67-1.17)	
Quartile 4	150,757	313	1.53	1.53	150,349	312	0.92	0.92	1.76	1.76	169	1.03	1.03	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97	0.97
			(1.30-1.79)	(0.86-1.30)			(0.73-1.14)	(0.73-1.14)	(1.39-2.21)	(1.39-2.21)		(0.77-1.38)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	(0.70-1.33)	
<i>P</i> _{interaction} ^c			0.5	0.5			0.5	0.5	0.2	0.2		0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
Per 5 cm increase			1.09	1.01			0.99	0.99	1.12	1.12		0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99	0.99
			(1.06-1.12)	(0.97-1.05)			(0.94-1.03)	(0.94-1.03)	(1.08-1.16)	(1.08-1.16)		(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	(0.93-1.05)	
<i>P</i> _{trend} ^b			<0.001	0.7			0.5	0.5	<0.001	<0.001		0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
<i>P</i> _{interaction} ^c			0.8	0.8			0.8	0.8	0.5	0.5		0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7

^a Adjusted for age (continuous), race (white, black, other), sex, alcohol consumption (nondrinker and 1.08, 1.09–3.58, 3.59–13.54, >13.54 g/day), cigarette smoking (never, former, current), and study (AARP, BCDDP, WHS, PHS, HPFS, NYUWHS, CPS-II, IWHS, BWHS, WHI, and CSDLH).

^b *P*_{trend} from Wald tests.

^c *P*_{interaction} with sex from log-likelihood ratio tests.

Table 4.

Adjusted^a Hazard Ratios (HR) and 95% Confidence Intervals (CI) for Associations between Body Fat Distribution Metrics and Liver Cancer Risk, Liver Cancer Pooling Project.

	Non-case N	Primary Liver Cancer			Hepatocellular Carcinoma			Intrahepatic Cholangiocarcinoma		
		Case N	HR (95% CI)	BMI Adjusted HR (95% CI)	Case N	HR (95% CI)	BMI Adjusted HR (95% CI)	Case N	HR (95% CI)	BMI Adjusted HR (95% CI)
Waist-to-Hip Ratio										
WHR cutpoints										
M: <0.85, F: <0.80	230,387	164	1.00	1.00	74	1.00	1.00	40	1.00	1.00
M: 0.85-<0.90, F: 0.80-<0.85	136,303	181	1.31 (1.05-1.62)	1.19 (0.96-1.48)	88	1.36 (0.99-1.86)	1.20 (0.87-1.65)	36	1.23 (0.77-1.98)	1.13 (0.71-1.79)
M: 0.90-<0.95, F: 0.85-<0.90	135,443	275	1.42 (1.14-1.75)	1.24 (1.00-1.53)	134	1.46 (1.06-2.00)	1.20 (0.87-1.65)	46	1.41 (0.90-2.22)	1.23 (0.78-1.94)
M: 0.95, F: 0.90	145,693	393	1.61 (1.31-1.98)	1.29 (1.04-1.60)	207	1.73 (1.26-2.37)	1.26 (0.91-1.75)	54	1.47 (0.95-2.28)	1.19 (0.75-1.90)
WHO cutpoints										
M: <0.90, F: <0.85	366,690	345	1.00	1.00	162	1.00	1.00	76	1.00	1.00
M: 0.90, F: 0.85	281,136	668	1.29 (1.11-1.50)	1.13 (0.97-1.32)	341	1.33 (1.06-1.66)	1.10 (0.88-1.37)	100	1.29 (0.94-1.76)	1.13 (0.81-1.58)
Sex- and study-specific										
Quartile 1	163,980	190	1.00	1.00	87	1.00	1.00	30	1.00	1.00
Quartile 2	161,416	222	1.05 (0.87-1.27)	1.00 (0.83-1.21)	120	1.23 (0.95-1.61)	1.14 (0.88-1.50)	40	1.29 (0.81-2.08)	1.23 (0.76-2.00)
Quartile 3	162,073	245	1.17 (0.98-1.40)	1.06 (0.88-1.27)	110	1.14 (0.87-1.49)	0.99 (0.75-1.30)	53	1.70 (1.09-2.67)	1.55 (0.98-2.45)
Quartile 4	160,357	356	1.59 (1.34-1.89)	1.33 (1.12-1.60)	186	1.80 (1.40-2.32)	1.40 (1.08-1.82)	53	1.75 (1.11-2.73)	1.49 (0.90-2.45)
Per 0.05 unit increase										
			1.11 (1.08-1.14)	1.07 (1.04-1.11)		1.12 (1.08-1.16)	1.07 (1.02-1.13)		1.10 (1.04-1.16)	1.07 (0.99-1.15)
P_{trend}^b			<0.001	<0.001		<0.001	0.007		0.001	0.08
$P_{interaction}^c$			0.7			0.3			0.1	
Quartile Cross-Classification										
Sex- and study-specific										
Category 1: Hip and Waist <P75	448,999	595	1.00	1.00	281	1.00	1.00	107	1.00	1.00

Waist-to-Hip Ratio	Non-case N	Primary Liver Cancer			Hepatocellular Carcinoma			Intrahepatic Cholangiocarcinoma		
		Case N	HR (95% CI)	BMI Adjusted HR (95% CI)	Case N	HR (95% CI)	BMI Adjusted HR (95% CI)	Case N	HR (95% CI)	BMI Adjusted HR (95% CI)
Category 2: Hip >P75, Waist <P75	46,379	65	1.04 (0.82–1.32)	0.91 (0.71–1.17)	38	1.23 (0.90–1.68)	1.01 (0.74–1.40)	12	1.08 (0.59–1.98)	0.94 (0.50–1.77)
Category 3: Hip <P75, Waist >P75	48,478	106	1.46 (1.20–1.77)	1.25 (1.02–1.53)	53	1.57 (1.19–2.06)	1.22 (0.91–1.64)	17	1.47 (0.88–2.45)	1.27 (0.74–2.19)
Category 4: Hip and Waist >P75	103,970	247	1.67 (1.45–1.92)	1.24 (1.02–1.51)	131	1.87 (1.53–2.28)	1.18 (0.89–1.57)	40	1.60 (1.12–2.29)	1.19 (0.71–1.99)
$P_{interaction}^b$			0.9			0.3			0.8	

^a Adjusted for age (continuous), race (white, black, other), sex, alcohol consumption (nondrinker and 1.08, 1.09–3.58, 3.59–13.54, >13.54 g/day), cigarette smoking (never, former, current), and study (AARP, BCDDP, WHS, PHS, HPFS, NYUWHS, CPS-II, IWHS, BWHS, WHI, and CSDLH).

^b P_{trend} from Wald tests.

^c $P_{interaction}$ with sex from log-likelihood ratio tests.