

ARTICLE

Open Access

CRISPR/Cas9-mediated mutagenesis of *VvMLO3* results in enhanced resistance to powdery mildew in grapevine (*Vitis vinifera*)

Dong-Yan Wan^{1,2}, Ye Guo^{1,2}, Yuan Cheng^{1,2}, Yang Hu^{1,2}, Shunyuan Xiao³, Yuejin Wang^{1,2} and Ying-Qiang Wen^{1,2}

Abstract

Grapevine (*Vitis vinifera*), one of the most economically important fruit crops in the world, suffers significant yield losses from powdery mildew, a major fungal disease caused by *Erysiphe necator*. In addition to suppressing host immunity, phytopathogens modulate host proteins termed susceptibility (S) factors to promote their proliferation in plants. In this study, CRISPR/Cas9 (clustered regularly interspaced short palindromic repeats/CRISPR-associated 9) technology was used to enable the targeted mutagenesis of *MLO* (mildew resistance Locus O) family genes that are thought to serve as S factors for powdery mildew fungi. Small deletions or insertions were induced in one or both alleles of two grapevine *MLO* genes, *VvMLO3* and *VvMLO4*, in the transgenic plantlets of the powdery mildew-susceptible cultivar Thompson Seedless. The editing efficiency achieved with different CRISPR/Cas9 constructs varied from 0 to 38.5%. Among the 20 *VvMLO3/4*-edited lines obtained, one was homozygous for a single mutation, three harbored biallelic mutations, seven were heterozygous for the mutations, and nine were chimeric, as indicated by the presence of more than two mutated alleles in each line. Six of the 20 *VvMLO3/4*-edited grapevine lines showed normal growth, while the remaining lines exhibited senescence-like chlorosis and necrosis. Importantly, four *VvMLO3*-edited lines showed enhanced resistance to powdery mildew, which was associated with host cell death, cell wall apposition (CWA) and H₂O₂ accumulation. Taken together, our results demonstrate that CRISPR/Cas9 genome-editing technology can be successfully used to induce targeted mutations in genes of interest to improve traits of economic importance, such as disease resistance in grapevines.

Introduction

Grapevine is one of the most extensively cultivated and economically valuable horticultural crops in the world, with 7.6 million hectares in production and an annual value of ~\$3.6 billion in wine export markets alone (Organisation Internationale de la Vigne et du Vin, 2017). Unfortunately, grape production is limited by various diseases, including powdery mildew caused by the fungus *Erysiphe necator*. Powdery mildew can be controlled to a

certain extent by fungicides, but this increases production costs and can negatively affect the environment and human health. In addition, the indiscriminate use of pesticides increases carbon emissions¹.

A more cost-effective and environmentally friendly strategy for limiting losses due to diseases is the development of disease-resistant cultivars through traditional breeding and/or precision breeding with the aid of biotechnology. The identification and characterization of the genes and genetic pathways that confer resistance can greatly facilitate the latter approach. Advances in functional genomics are changing the preferred experimental strategy from using traditional forward genetics (i.e., from phenotype to genotype) to employing both forward genetics and reverse genetics (i.e., inferring gene function

Correspondence: Ying-Qiang Wen (wenyq@nwsuaf.edu.cn)

¹State Key Laboratory of Crop Stress Biology for Arid Areas, College of Horticulture, Northwest A&F University, Yangling 712100 Shaanxi, China

²Key Laboratory of Horticultural Plant Biology and Germplasm Innovation in Northwest China, Ministry of Agriculture, Yangling 712100 Shaanxi, China
Full list of author information is available at the end of the article

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

based on the phenotypes caused by mutations of a target gene)². The introduction of functional mutations through targeted genome editing is a powerful reverse genetics approach for understanding gene function. CRISPR-Cas9 technology is a relatively simple yet effective technique that can be exploited for gene mutation, the repression/activation of gene expression, subtle modification of gene function and epigenome editing³. In 2013, three research groups simultaneously reported the use of the CRISPR-Cas9 system for targeted genome modification in *Arabidopsis thaliana* (Arabidopsis), tobacco, rice and wheat for the first time^{4–6}. This system has also been successfully applied in maize⁷, cotton⁸, and other plants. Grapevine is a woody perennial plant with a highly heterozygous genome that can be genetically transformed only through lengthy and difficult procedures. To date, there have been only a few reports on the use of CRISPR-Cas9 for genome editing in grapevine^{5,9–11}.

Over the past 25 years, the plant immune system has been extensively studied, and fundamental discoveries have led to a better understanding of the molecular basis of plant disease resistance. While the development of disease-resistant crop varieties has typically relied on the introduction of dominant resistance (*R*) genes into elite cultivars via classical breeding, recessive mutations in host susceptibility (*S*) genes have also been utilized to confer resistance. *R*-gene-induced defense responses often result in localized cell death or a hypersensitive response (HR)¹²; by contrast, the loss of *S*-genes required for the successful invasion of a particular pathogen may not display strong defense activation. For example, the disruption of *S*-genes such as *PMR5*¹³, *DMR6*^{14,15}, *LOB1*¹⁶, and *SWEET14*^{17,18} can confer broad-spectrum disease resistance with no or minimal activation of defenses in several economically important plant species. *S*-gene-mediated resistance is generally more durable than *R*-gene-based resistance because the pathogen must overcome dependence on a host *S* factor¹⁵. The disruption of the *MLO* gene in barley confers durable powdery mildew resistance, as demonstrated by the fact that no powdery mildew strains capable of overcoming *mlo*-mediated resistance have been found in the field for many decades¹⁹.

Studies suggest that the pathogenicity of adapted powdery mildew species relies on a functional host *MLO* gene^{20,21}. *MLO* proteins constitute a family of highly conserved²², plant-specific, seven-transmembrane domain proteins that are structurally similar to G protein-coupled receptors (GPCRs) with a calmodulin-binding domain²³. The mechanism by which *MLO* proteins act as powdery mildew susceptibility factors is unknown, and the biochemical function of *MLO* proteins in the host remains largely unexplored²⁴. In barley, an *mlo* mutation was found to confer broad-spectrum and durable resistance to barley powdery mildew isolates of *Blumeria graminis* f. sp. *hordei*²⁵, but a range of developmentally controlled

pleiotropic effects occurred in the host, such as the formation of callose-containing cell wall appositions (papillae) in the absence of any pathogen and the premature onset of leaf senescence. These effects reduced the grain yield of the *mlo* mutants; thus, the dysfunctional *MLO* proteins may indeed result in a competitive disadvantage in natural populations²⁶. In bread wheat, TALEN-induced mutations in all three *TaMLO* homologs confer heritable broad-spectrum resistance to powdery mildew²⁷. In apple, the suppression of *MdMLO11* and *MdMLO19* through RNA interference (RNAi) reduces susceptibility to powdery mildew (*Podosphaera leucotricha*)²⁸. In *Arabidopsis*, the *Atmlo2 Atmlo6 Atmlo12* (*Atmlo2/6/12*) triple mutant is completely resistant to powdery mildew²¹. The grapevine genes *VvMLO3*, *VvMLO4*, and *VvMLO17*, orthologous to the *AtMLO2*, *AtMLO6*, and *AtMLO12* genes, respectively²⁹, are reported to exhibit rapid induction upon powdery mildew infection, suggesting that these *VvMLO* proteins may play a role in modulating antifungal defense responses in grapevine. One previous study used RNAi to suppress specific *VvMLO* genes and found that the simultaneous knockdown of *VvMLO6* and *VvMLO7* in grapevine reduced susceptibility to powdery mildew³⁰.

The grapevine cultivar Thompson Seedless is an ideal material for studying powdery mildew resistance in grapevine because it is amenable to transformation and susceptible to powdery mildew infection. There are 17 *VvMLOs* in Thompson Seedless. In this study, we used CRISPR/Cas9 technology to target two grapevine *MLO* genes, *VvMLO3* and *VvMLO4*, and obtained *VvMLO3*-edited grapevine lines with enhanced resistance to powdery mildew. Our work suggests that CRISPR/Cas9-targeted mutagenesis is a valuable tool for generating disease-resistant grapevines.

Results

Target selection within *MLOs* and vector construction

For each *VvMLO* gene, two target loci were selected (Fig. 1a). Schematic representations of the target sites of two small guide (sg) RNAs and their gene expression cassettes used for the *Agrobacterium*-mediated transformation of grapevine embryogenic calli is shown in Fig. 1a, b. The sgRNA intermediate vectors are based on the pUC18 backbone described previously³¹. The binary vector for plant transformation contains a plant codon-optimized *Cas9* gene. In total, four DNA constructs, designated CM3G1 (*Cas9/sgRNA1-MLO3-Grapevine*), CM3G2, and CM4G3 (*Cas9/sgRNA3-MLO4-Grapevine*) and CM4G4, were produced and used for transformation (Fig. 1b, c).

Targeted mutagenesis in regenerated transgenic grapevines

In total, 210 plantlets were regenerated from the grapevine anther filament-induced embryogenic calli




Fig. 1 Target selection and expression constructs for Cas9 and sgRNA. **a** Target selection of *VvMLO3* and *VvMLO4*. CRISPR/Cas9-targeted sites were selected within exons 1 (sgRNA1) and 13 (sgRNA2) of *VvMLO3* and exons 1 (sgRNA3) and 10 (sgRNA4) of *VvMLO4*. The protospacer-adjacent motif (PAM) sequence is highlighted in red. **b**, **c** Illustration of the assembly of the pYLCRISPR/Cas9 binary vector combined with the sgRNA by using Golden Gate cloning.

subjected to coinubation with *Agrobacterium* cells (Fig. 2). To verify the presence of the CRISPR-Cas9 sequence, genomic DNA from each regenerated plantlet was analyzed by PCR. As shown in Fig. 3a, a fragment of ~508 bp was amplified from *Cas9*-positive plantlets, whereas *Cas9*-negative and nontransgenic control plantlets showed no amplification. A total of 125 plantlets were identified as *Cas9* positive for any of the four targets of the two *VvMLO* genes, and these plantlets were further analyzed for the presence of the intended sequence variation. A total of 68 independent PCR-positive plantlets for *VvMLO3*-sgRNA1, 39 for *VvMLO3*-sgRNA2, 13 for *VvMLO4*-sgRNA3, and 5 for *VvMLO4*-sgRNA4 were analyzed by Sanger sequencing (Table 1). The corresponding DNA target sites of sgRNA1, sgRNA2, or sgRNA3 contained mutations (Table 1), indicating that three of the four CRISPR/Cas9-DNA constructs (i.e., CM3G1, CM3G2, CM4G3) enabled successful gene editing in the regenerated grapevine plantlets. Different target sites were mutated with different editing efficiencies, varying from 0 to 38.5% (Table 1). Single-nucleotide insertions and nucleotide deletions of various lengths were identified among these edited lines (Fig. 3b). The mutations detected in the sgRNA1 lines were mainly small deletions and insertions, while those of the sgRNA2

lines were mainly small insertions, and those of the sgRNA3 lines were mainly small insertions (Fig. 3b and Table S1). Most of the insertions consisted of only 1 bp, and the majority of deletions were short in the sgRNA1 and sgRNA2 lines. Double-strand breaks (DSBs) usually occur at a position three base pairs upstream of the PAM sequence³². Among the 20 *VvMLO*-edited lines, 16 showed mutation sites exactly at the 4th base from the PAM site (Fig. S1 and Table S1).

Four different types of mutation sequence chromatograms are shown in Fig. 3c. These 20 edited lines were detected among the 125 transgenic plantlets that were obtained and analyzed (Table 1); thus, the overall gene editing efficiency was 16.5%. It was interesting that no edited lines were detected among any of the five transgenic lines containing the CM4G4 construct. Among the 20 edited lines, one (5%) was homozygous for a 1-bp insertion, three (15%) were biallelic for small deletions and insertions (Indels), seven (35%) were heterozygous (i.e., only one allele contained a mutation), and nine (45%) were chimeric (i.e., more than two mutated alleles were detected in one line) (Fig. 3d). Detailed information about the mutation found in one edited line, CM3G2-12, is provided as an example in Fig. 3e; both alleles of *VvMLO3* in this line contained a single-nucleotide (A) insertion in




Fig. 2 Genetic transformation of *Vitis vinifera* cv. Thompson Seedless by *Agrobacterium tumefaciens* and the regeneration of transgenic lines. **a** Coculture of proembryonic masses (PEMs) and *Agrobacterium tumefaciens* on two layers of filter paper soaked in 2 ml suspension liquid (MS + 200 μmol AS). **b** The transformed PEMs cultured on X3CC (X3 + 200 $\text{mg}\cdot\text{L}^{-1}$ carb + 200 $\text{mg}\cdot\text{L}^{-1}$ Cef) medium after the elimination of the bacteria. **c** The transformed PEMs cultured on X3CCK75 (X3 + 200 $\text{mg}\cdot\text{L}^{-1}$ carb + 200 $\text{mg}\cdot\text{L}^{-1}$ Cef + 75 $\text{mg}\cdot\text{L}^{-1}$ kanamycin) medium. **d-f** Appearance of kanamycin-resistant somatic embryos. **g** Kanamycin-resistant somatic embryos transferred to germination medium and cultured under light. **h** Germination of resistant plantlets. **i** *MLO*-edited lines domesticated in the phytotron.

the target site, resulting in the formation of a premature stop codon, which in turn was predicted to produce a *VvMLO3* mutant with a C-terminal truncation.

Phenotypic characterization of genome-edited lines

The vast majority (93%) of the regenerants obtained after *Agrobacterium* infection in this study showed abnormal growth and development (not shown). When grown under axenic conditions, 14 of 20 3-month-old *VvMLO3*- or *VvMLO4*-edited plantlets (e.g., CM3G1-65 and CM3G2-25; Fig. 4a) showed similar phenotypes, which were characterized by senescence-like chlorosis and necrosis. Among these 14 plantlets, CM3G2-12, homozygous for a 1-bp insertion, died in the medium when it was two months old. However, the nonedited plants displayed normal growth and development during the same period.

We found that when the *VvMLO*-edited plantlets were subcultured to induce proliferation, their rooting

potential was very low, even in the presence of IBA at a concentration as high as 1.0 $\text{mg}\cdot\text{L}^{-1}$. Only a few lines (i.e., CM3G2-30 and CM3G1-51) showed normal rooting (Fig. 4b). Compared with the nontransgenic control, the growth of the *VvMLO*-edited plantlets was poor, and the lower part of the plantlets showed accelerated defoliation.

Powdery mildew-triggered mesophyll cell death, cell wall apposition and H_2O_2 production are associated with *mlo*-mediated resistance

To evaluate the effect of targeted mutations in the two *VvMLO* genes on resistance to powdery mildew, two *VvMLO3*-edited lines, CM3G2-30 and CM3G1-51 (sequencing chromatograms are shown in Fig. 5a), and one nontransgenic control plant were transplanted to pots (Fig. 5b), and detached leaves of these plants were inoculated with the powdery mildew isolate *En. NAFU1*³³ 15 days after transplantation. Leaves inoculated with *En. NAFU1* were collected at 24 and 72 hpi and stained




Fig. 3 Targeted mutagenesis of *VvMLO3* and *VvMLO4* using the CRISPR-Cas9 system. **a** PCR amplification of a 507-bp DNA fragment of genomic DNA from regenerated plantlets using gene-specific primers. M: marker, P: plasmid, C: control, CM3G1-53 to CM3G1-62: different regenerated lines expressing *VvMLO3*-sgRNA1. **b** The numbers on the right indicate the type of mutation and the number of nucleotides involved, “-” and “+” indicate deletions and insertions, respectively. The insertions are highlighted in red letters, and the protospacer-adjacent motif (PAM) is indicated in blue letters. **c** Four different types of mutation sequencing chromatograms: biallelic, homozygous, heterozygous, and chimeric mutations. The first substitution or indel sites are indicated with red arrowheads. The protospacer-adjacent motif (PAM) sequences are highlighted in blue, the indels or substituted bases are highlighted in red, and “-” indicates deletions. **d** The frequency of different mutation types found in the edited lines. **e** Amino acid sequence alignment between one representative mutant and the wild-type for the target region of *VvMLO3*.

Table 1 Summary of genome editing results in regenerated plantlets.

	CM3G1	CM3G2	CM4G3	CM4G4
Regenerated lines	117	63	22	8
Cas9 positive lines	68	39	13	5
Homozygous ^a lines	0	1	0	0
Heterozygous ^b lines	1	3	3	0
Biallelic ^c lines	2	0	1	0
Chimeric ^d lines	7	1	1	0
Nonedited lines	107	58	17	8
Editing efficiency	14.7%	12.8%	38.5%	0

^aHomozygous = both alleles contain the same mutation.

^bHeterozygous = only one allele is mutated.

^cBiallelic = both alleles are mutated, but the mutations are not identical.

^dChimeric = presence of more than two mutant alleles, indicative of the presence of different edited cell lines.

with trypan blue for the visualization of fungal structures by microscopy. At 24 hpi, both the *VvMLO3*-edited and nontransgenic lines produced appressoria with no apparent differences. However, at 72 hpi, the edited line showed much less fungal growth compared with the control line (Fig. 5c), with only ~1/4 of the total hyphal length per colony being observed on CM3G2-30 and ~1/10 on CM3G1-51 compared with that in the control line ($P < 0.01$) (Fig. 5d). In addition, there were fewer secondary hyphae on the two *VvMLO3*-edited lines compared with the nontransgenic control (not shown), indicating that both edited lines exhibited enhanced resistance to powdery mildew. Unfortunately, all five *VvMLO4*-edited lines produced deformed leaves and died. Thus, it remains to be determined whether *VvMLO4* plays a role in powdery mildew pathogenesis. The potted *VvMLO3*-edited plants were transferred to a greenhouse for further growth and evaluation (Fig. 6a). In the edited




Fig. 4 *VvMLO* mutant grapevine plantlets showed early senescence-like phenotypes and difficulty in rooting. **a** Targeted *VvMLO*-edited plantlets showed early senescence-like phenotypes. **A**: 1-month-old *VvMLO*-edited plantlets grown in the rooting medium, **B**: 3-month-old *VvMLO*-edited plantlets grown in the rooting medium. Red arrows indicate chlorosis, and red box frames indicate the regenerated nonedited plantlets. Bars = 1 cm. **b** Rooting of the indicated *VvMLO*-edited and nonedited wild-type (WT) plantlets in rooting medium. Bars = 1 cm.




Fig. 5 (See legend on next page.)

(see figure on previous page)

Fig. 5 Different types of mutations confer different levels of resistance. **a** Sequence chromatograms of mutation lines CM3G2-30 and CM3G1-51. The first substitution or indel sites are indicated with red arrowheads. The protospacer-adjacent motif (PAM) sequences are highlighted in blue, and the indels or substituted bases are highlighted in red. ‘—’ indicates deletions. **b** Comparison of the wild-type (WT) and different types of *VvMLO3*-edited lines 15 days after transplantation from subculture medium. **c** Representative micrographs showing the growth of powdery mildew in the WT and *VvMLO3*-edited lines at 24 and 72 hpi. Bars = 100 μ m except for that (20 μ m) in the inset. **d** Total hyphal length per powdery mildew colony on the leaves of the *VvMLO3*-edited lines and WT inoculated with *En* NAFU1 at 72 hpi. Data are the means \pm SE calculated from three replicate experiments. Asterisks indicate significant differences between the edited lines and the Thompson Seedless WT control ($P < 0.01$; $n = 3$, Student's t test).

lines harboring heterozygous mutations in *VvMLO3*, clear powdery mildew-induced cell death was observed (Fig. 6b). Careful examination of the trypan blue-stained leaf sections revealed that in many cases, the mesophyll cells underneath the infected epidermal cells were stained blue (Fig. 6c), suggesting that signals from mildew-invaded epidermal cells trigger the collapse of neighboring mesophyll cells as a result of the mutation of *VvMLO3* in these heterozygous lines.

In a previous report, increased cell wall apposition (CWA) or papilla formation was observed beneath the site of attempted penetration in *mlo* mutant plants²¹. We used aniline blue staining to check for the presence of callose in the CWAs after the inoculation of powdery mildew onto the detached leaves of *VvMLO3*-edited lines. At 7 dpi, clear CWAs were observed in the heterozygous mutant line CM3G2-40 but not in nontransgenic control plants (Fig. 6d). In addition, H₂O₂ production and accumulation visualized as brownish precipitates upon 3,3'-diaminobenzidine (DAB) staining were more pronounced in mildew-infected cells of the heterozygous mutant lines CM3G1-91 and CM3G2-40 at 7 dpi (Fig. 6b and Fig. S3a). The quantification of H₂O₂ accumulation using ZEN 2012 software showed that ~22% and ~42% of the infected leaf epidermal cells of the two *VvMLO*-edited plants exhibited H₂O₂ accumulation, which was three to six times higher than that (6.5%) in the infected leaf epidermal cells of the wild-type plants (Fig. S2a, b). Taken together, our data suggest that *Vvmlo3*-mediated resistance to powdery mildew is associated with callose deposition and H₂O₂ accumulation in powdery mildew-infected epidermal cells and possibly the demise of mesophyll cells underneath.

To assess the impact of the *VvMLO3* gene mutation on whole-plant resistance to powdery mildew, two independent experiments were carried out. Potted transgenic or nontransgenic control plants were inoculated with fresh conidia of powdery mildew *En*. NAFU1. This strain was able to infect both the WT control and heterozygous mutant lines CM3G1-91 and CM3G2-40 (Fig. 7a). However, it was noted that the leaves of the edited lines, but not those of the control line, were covered with abundant white velvety fungus at 15 dpi, which seemed to correlate with the presence of necrotic lesions in the infected leaf

tissue in the edited lines but not in the control line at 15 dpi (Fig. 7b). The quantification of the number of spores per mg of fresh infected leaf tissue at 20 dpi revealed that there was an ~2-fold reduction in fungal sporulation in the edited line CM3G1-91 in comparison with the WT line, while a small, yet still statistically significant reduction was detected in the other edited line, CM3G2-40 (Fig. 7c).

Discussion

Since the first report on the application of CRISPR/Cas9 technology for the editing of grapevine genes in 2016³⁴, two more recent studies have explored the use of this new genome-editing technology in grapevine using protoplasts¹¹ or embryogenic callus cells⁹. Two other examples of successful targeted mutagenesis using the CRISPR/Cas9 system in grapevine resulted from studies conducted for the functional characterization of the *WRKY52* gene¹⁰ and the *VvCCD8* gene³⁵. In this study, two grapevine *MLO* genes, *VvMLO3* and *VvMLO4*, were edited by using CRISPR/Cas9 technology, and the reduction of *VvMLO3* expression by mutating one allele significantly improved the resistance of a susceptible grapevine cultivar against the powdery mildew isolate *En*. NAFU1.

Charrier et al. performed targeted mutagenesis in apple and pear using the CRISPR-Cas9 system, and they classified the obtained mutants as homozygous, heterozygous, biallelic, and chimeric³⁶. According to their classification, in a “homozygous” line, both alleles of the target gene contain the same mutation; in a “heterozygous” line, only one allele is mutated; in a “biallelic” line, both alleles are mutated, but the mutations are not identical; and finally, in a “chimeric” line, more than two alleles are detected, which is indicative of the presence of cell lineages derived from different edited cells during somatic embryogenesis. In this study, all four abovementioned types of grapevine mutants were detected, and the “chimeric” type accounted for 45% of the mutants obtained, suggesting that CRISPR/Cas9 is enzymatically active during the early stage of embryogenesis initiated from a single transformed somatic cell.

As a woody perennial fruit crop, grapevine is known to be recalcitrant to genetic transformation. To increase the chance of success, we designed two sgRNAs for each of




Fig. 6 (See legend on next page.)

(see figure on previous page)

Fig. 6 *VvMLO3*-edited grapevine plants show infection-triggered cell death, H₂O₂ accumulation and CWAs. **a** The wild-type (WT) and two *VvMLO3*-edited heterozygous mutant grapevine lines grown under phytotron conditions for 6 months (bar = 10 cm). **b** Representative micrographs showing DAB- and trypan blue-stained epidermal cells of the WT and *VvMLO3*-edited lines at 5 or 7 dpi. Red arrowheads indicate trypan blue retention, and black arrowheads indicate H₂O₂ accumulation (bar = 50 μm). **c** Representative images showing a trypan blue-stained leaf section of WT or CM3G2-40 with a focus on either the epidermal layer or the mesophyll cell layer at 7 dpi (bars = 50 μm). **d** Histochemical analysis of infection-triggered CWAs of epidermal cells of the WT and the heterozygous CM3G2-40 mutant line at 7 dpi. Red arrowheads indicate haustoria (H), and black arrowheads indicate infection-triggered CWAs (bars = 50 μm).




Fig. 7 Targeted editing of *VvMLO3* improved the resistance of *V. vinifera* cv. Thompson Seedless to powdery mildew. **a** Representative *En. NAFU1*-infected leaves of the *MLO*-edited lines in comparison with those of the WT control line at 15 dpi (bar = 5 cm). **b** Detection of cell death by trypan blue staining in leaves of *VvMLO3*-edited grapevine at 15 dpi (bars = 50 μm). Black arrows indicate leaf cell necrotic lesions, while red arrows denote clustered dead or dying mesophyll cells stained by trypan blue. **c** Quantification of the number of spores per mg of fresh leaf tissue in WT and *VvMLO3*-edited grapevines at 20 dpi. Each data point is the mean of three independent experiments (±standard deviation); asterisks indicate significant differences in comparison with the WT as determined by Student's *t* test (**P* < 0.05, ***P* < 0.01).

the two *MLO* target genes, one of which targeted a site near the initiation codon, while the other targeted a site in the middle of the coding sequence for each gene (Fig. 1a).

This strategy was found to be effective for *VvMLO3*, as mutations were identified in two target sites. For *VvMLO4*, sgRNA3 (CM4G3) achieved an ~23% editing

efficiency, but sgRNA4 (CM4G4) failed to produce a single mutant line (Table 1). The latter situation may simply be due to the insufficient number (eight) of transgenic lines generated or to the presence of unusual local chromatin structures that are less accessible by CRISPR-Cas9³⁷. The DNA mutations induced by CRISPR/Cas9 in plants mainly consist of short indels and replacements that occur during the process of DSB repair via nonhomologous end joining (NHEJ)³⁸. Such diverse mutations are expected to occur during NHEJ-mediated DNA repair. This mutation pattern was also observed in this study, as most of the mutations induced in *VvMLO3* were indels. Among the 20 edited lines obtained, only one was homozygous for a single 1-bp insertion, while all other lines showed a combination of different types of mutations (Fig. 3c).

MLO homologs have been subjected to targeted mutagenesis in many plants for the engineering of powdery mildew resistance^{20,21,27,39,40}. The loss of *AtMLO2*, but not *AtMLO6* or *AtMLO12*, in *Arabidopsis thaliana* enhances the resistance of the mutant plants to powdery mildew, while the loss of all three genes results in complete resistance²¹. These findings indicate that while *AtMLO2* is the major host S protein required for powdery mildew infection, *AtMLO6* and *AtMLO12* possess functions that overlap with those of *AtMLO2*. In this study, *VvMLO3* and *VvMLO4*, from the clade corresponding to *AtMLO2*, 6, and 12 in the phylogenetic tree²⁹ (Fig. S3), were selected for CRISPR/Cas9-mediated mutagenesis. Unfortunately, the only *Vvml03* homozygous mutant plantlet in which *VvMLO3* was knocked out died of massive leaf necrosis in the tissue culture medium (data not shown). This implies that either the loss of *VvMLO3* or a genetic mutation due to the T-DNA insertion might have triggered lethal cell death in this *Vvml03* mutant. Nevertheless, although multiple *VvMLO3*-edited lines were heterozygous lines, qRT-PCR revealed that there was an ~4-8-fold reduction in the *VvMLO3* expression level in the edited lines CM3G1-91 and CM3G2-40 in comparison with the WT line (Fig. S4), which indicated that the *VvMLO3* expression level was indeed negatively impacted by such editing. Significantly enhanced mildew resistance was observed in multiple *VvMLO3* heterozygous lines in comparison with the WT plants from 2 to 7 dpi (Figs. 5 and 6), suggesting that *VvMLO3* is one of the functional homologs of *AtMLO2* in grapevine. Taken together, our results suggest that *VvMLO3* is an important (though not the only) host susceptibility gene required for powdery mildew infection. It should note that the nomenclature for grapevine *MLO* genes differed in two previous studies by Feechan²⁹ and Winterhagen⁴¹ that were published around the same time. In the present study, the naming of *VvMLO3* and *VvMLO4* followed the Feechan nomenclature²⁹, whereas these genes were named *VvMLO11(W)*

and *VvMLO13(W)*, respectively, according to Winterhagen⁴¹ (for clarity, when we describe *MLO* genes using the Winterhagen nomenclature, we use *VvMLO6 (W)* and *VvMLO7 (W)*) (Fig. S3). Previous studies showed that *VvMLO6 (W)* and *VvMLO7 (W)*, rather than *VvMLO11 (W)* or *VvMLO13 (W)* (i.e., *VvMLO3* and *VvMLO4* according to the Feechan nomenclature), may negatively regulate disease resistance^{29,42}, and the silencing of the *VvMLO6 (W)* and *VvMLO7 (W)* genes resulted in enhanced resistance to powdery mildew³⁰. Hence, it is likely that *VvMLO6/7 (W)* may also be a functional homolog of *AtMLO2*. Simultaneous targeted mutagenesis of all three grapevine *MLO* genes may therefore be necessary to develop complete mildew resistance in grapevine in the future.

In addition to powdery mildew resistance, *mlo* mutant plants display premature leaf chlorosis and a reduced grain yield in barley¹⁹, increased callose deposition in the cell wall of leaves in pathogen-free plants and slower growth and development in *A. thaliana*^{21,43,44}, and reduced plant size in pepper⁴⁵. Not surprisingly, similar growth phenotypes, such as early leaf senescence, were observed in our *VvMLO3* heterozygous mutant grapevine plantlets (Fig. 4a). Interestingly, mildew-induced mesophyll cell death was observed in the *VvMLO3* heterozygous mutant lines (Fig. 6), which, along with H₂O₂ accumulation and callose deposition in mildew-infected epidermal cells, seemed to be associated with the increased resistance of these mutant lines to powdery mildew (Fig. 6). In this regard, it is worth pointing out that no obvious pleiotropic phenotypes were observed when *VvMLO6*, 7, 11, or 13 (W) was knocked down by RNAi³⁰. This implies that *VvMLO6*, 7, and 13 (W) may be functionally distinct from *VvMLO3* and/or that these four genes are functionally redundant or were not adequately silenced.

Conclusion

CRISPR/Cas9-enabled targeted mutagenesis suggested that *VvMLO3* is an S-gene required for the powdery mildew infection of grapevine and that the targeted mutation of *VvMLO3* in a grapevine cultivar results in enhanced resistance to powdery mildew. This demonstrates that CRISPR/Cas9-targeted mutagenesis can be used to develop disease-resistant cultivars and facilitate the functional characterization of genes of interest in grapevine.

Materials and methods

Plant materials

V. vinifera cv. Thompson Seedless was used for gene cloning and transformation and as a control in gene functional analysis and phenotypic observations of powdery mildew infections. Plants were grown in a phytotron at Northwest A&F University, Yangling, Shaanxi, People's

Republic of China. The temperature was controlled between 22 and 26 °C under a 14 h/10 h (day/night) light cycle. Embryogenic calli were induced from anther filaments of Thompson Seedless. Proembryogenic masses (PEMs) and somatic embryos of Thompson Seedless at the mid-cotyledonary stage of development were used for transformation. PEM cultures were maintained on MS basal medium supplemented with 60 g·L⁻¹ sucrose, 3 g·L⁻¹ Phytigel, and 1 g·L⁻¹ activated charcoal with a pH of 5.8–6.0. PEM cultures were reinduced by placing chopped torpedo-shaped-stage or mid-cotyledonary-stage embryos on KBN medium (MS, sucrose, 1 g·L⁻¹ myo-inositol, 0.3 g·L⁻¹ KNO₃, 1.126 mg·L⁻¹ 6-BA, 0.552 mg·L⁻¹ 2,4-D and 0.505 mg·L⁻¹ NOA). The secondary regeneration of PEMs was typically observed within 3–7 months. All cultures were maintained in the dark at 25 °C and transferred to fresh media monthly⁴⁶.

Cloning of grapevine *MLO* genes

Total RNA was extracted from Thompson Seedless leaves using the E.Z.N.A. Plant RNA Kit (Omega, Guangzhou, China) according to the manufacturer's instructions. First-strand cDNA was generated from 1.5 µg of total RNA using PrimeScript RTase (Takara Bio Inc., Dalian, China). The synthesized cDNA product was diluted 10-fold and used as a template for the cloning of *MLO* genes. Two *VvMLO* genes, *VvMLO3* and *VvMLO4*, identified in a previous study²⁹ were selectively amplified using oligonucleotide primers derived from *VvMLO* gene sequences available in the National Center for Biotechnology Information (NCBI) database (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>). PrimeSTAR HS DNA Polymerase was used for amplification. The PCR products were cloned into the pMD18-T vector (Takara Bio Inc., Dalian, China) and sequenced at the Beijing AuGCT Biotech, Yangling, Sequencing Department. The sequenced DNAs were aligned with the genome of *Vitis vinifera* PN40024.

CRISPR-Cas9 binary construct design

Target sequences were selected within two *VvMLO* exons, and four sgRNAs, designated sgRNA1 to sgRNA4, were designed within the two coding regions (Table S3). To increase the editing success rate for each gene, we selected two target loci in each *MLO* gene, seeking loci with a moderate GC content. We screened the target sequences to minimize potential off-target alignments using the online software Grape-CRISPR (<http://biodb.sdau.edu.cn/gc/>) and CRISPR-P (<http://crispr.hzau.edu.cn/CRISPR2/>)⁴⁷. For each gene, we located one target site near the start codon and another in a region corresponding to the functional domain. Next, we performed a BLAST-P search based on the National Center for Biotechnology Information (NCBI) database ([\[ncbi.nlm.nih.gov/Blast.cgi\]\(https://blast.ncbi.nlm.nih.gov/Blast.cgi\)\) of the target sequences \(including PAM\) against the grapevine genome sequence to ensure that the target sequence was as specific as possible. We avoided sequences containing more than four consecutive Ts to prevent RNA Pol III from using such sites as a transcription termination signal. \(The selected target sequences are shown in Fig. 1a.\) Target adapter oligos were synthesized for each target locus using Ao ke ding sheng \(<http://www.augct.com/?ClassID=2>\) in the Yangling department \(Table S2\).](https://blast.</p></div><div data-bbox=)

The binary pYLCRISPR/Cas9-N⁴⁸ vector and two sgRNA cassettes driven by the *AtU3b* and *AtU6-1* promoters were used to generate vectors expressing CRISPR/Cas9-*MLO*s. For each sgRNA, a pair of DNA oligonucleotides was synthesized by Beijing AuGCT Biotech and annealed to generate dimers, which were subsequently integrated upstream of the sgRNA scaffolds in the plasmid vector. Two sequential rounds of overlapping PCR were carried out to insert double-stranded, gRNA spacer DNA sequences between double *Bsa* I restriction enzyme sites to generate intermediate vectors with one or two target sites. Intermediate sgRNA expression cassettes with target sequences were generated by overlapping PCR (Figs. S5 and S6). We used a binary CRISPR/Cas9 vector, pYL-CRISPR/Cas9P35S-N containing the CaMV 35S promoter and the NPTII gene as a selectable marker. The Cas9 vector was built in the backbone of pCAMBIA1300. The final purified PCR products were assembled into the binary pYLCRISPR/Cas9 vector containing the NPTII gene and 2 × 35 s promoter by Golden Gate ligation⁴⁹. The ligated products with one or two sgRNA expression cassettes were directly used to transform DH5α *E. coli* competent cells.

Agrobacterium-mediated transformation of grapevine

Transformation was carried out via the *Agrobacterium*-mediated transformation system as previously described⁵⁰ with minor modifications. *Agrobacterium tumefaciens* strain GV 3101 was used for transformation. Transformation was carried out in several batches to avoid failure. The bacterial solution (OD600, 0.6–0.8) was poured into a sterile flask, and the grapevine tissue was soaked in the flask for 15–20 min with gentle shaking. The tissue was then placed on two layers of sterile filter paper to remove excess bacterial culture and transferred to 2 ml of liquid MS medium (containing 20 mg·L⁻¹ AS and 3% sucrose) for cocultivation. After 2 days of cocultivation at 25 °C in the dark, the material was further maintained for 1 month without any antibiotics before screening to allow the transformed material to recover. The tissue was then washed three times in sterile water and transferred to KBN delayed-screening medium (30 g·L⁻¹ sucrose, MS, 0.3 g·L⁻¹ KNO₃, 1.126 mg·L⁻¹ 6-BA, 1 g·L⁻¹ myo-inositol, 0.505 mg·L⁻¹ NOA, Phytigel, 200 mg·L⁻¹ Cef) for

3 weeks, followed by incubation for 1 week on X3 delayed-screening medium (30 g·L⁻¹ sucrose, MS, 1 g·L⁻¹ activated charcoal, Phytigel, 200 mg·L⁻¹ Cef). After this delayed screening, the transformed PEM cultures were transferred to resistance screening medium containing Cef and Kan (30 g·L⁻¹ sucrose, MS, 1 g·L⁻¹ activated charcoal, Phytigel, 200 mg·L⁻¹ Cef, 75 mg·L⁻¹ Kan). Screening was continued until germinated embryos developed. Germinated resistant embryos were then transferred to germination medium (15 g·L⁻¹ sucrose, MS, Phytigel) and placed under light for further development. After true leaves appeared, the cultured plantlets were transferred to 30 mL rooting medium (30 g·L⁻¹ sucrose, MS, 1 g·L⁻¹ activated carbon, Phytigel, 1 mg·L⁻¹ IBA) for further development. Rooted plantlets were transplanted into pots (diameter: 14 cm, height: 10 cm) containing potting mix (perlite, vermiculite, and peat) and covered with a transparent plastic cup, then placed in a phytotron.

Genotyping and mutant verification

The genotyping of transgenic lines was carried out essentially as described previously³¹. The genomic DNA of each line was isolated from 0.5 to 1.0 g of regenerated plantlet tissue using a CTAB-based method⁵¹. To verify the presence of the engineered sequences, specific oligonucleotide primers (SP-L-35S and SP-R) were used to amplify the CRISPR/Cas9 sequence (Table S2). The PCR products (ca. 400–600 bp) were sequenced directly using internal specific primers, for which the ideal binding positions are ~150–250 bp upstream of the target sites (Beijing AuGCT Biotech Yangling Sequencing Department). The detection of target gene editing requires the sequencing of multiple clones of the PCR amplicons derived from the same genomic DNA sample. The PCR amplicons were cloned into a pMD18-T plasmid vector, and five to ten randomly selected *E. coli* colonies were used for plasmid preparation and insert sequencing. SnapGene Viewer software was used for mutation analysis. The sequence results are shown in Table S1.

Pathogen inoculation and evaluation of resistance to powdery mildew

Four *MLO*-edited lines, CM3G1-51 and -91 and CM3G2-30 and -40, were assessed for resistance to powdery mildew. The transgenic mutant and non-transgenic control lines were inoculated with the powdery mildew isolate *En* NAFU1 as described previously³³. For detached leaf inoculation, CM3G2-30, CM3G1-51, and wild-type (WT) plants were used at 15 days after transplantation from the subculture medium. Fully expanded leaves at the third position from the shoot tip were chosen for inoculation. Representative micrographs and hyphal development were observed at 24 and 72 hpi, respectively.

For plant leaf inoculation, CM3G2-40, CM3G1-91, and wild-type (WT) plants were grown under phytotron conditions for 6 months for propagation and then subjected to inoculation. DAB and trypan blue-stained epidermal cells of WT, CM3G2-40 and CM3G1-91 plants were observed at 5 or 7 dpi. To assess the impact of the *VvMLO3* gene mutation on whole-plant resistance to powdery mildew, plants of the transgenic lines CM3G2-40 and CM3G1-91 or nontransgenic control plants that had been grown in pots for six months were inoculated with fresh conidia of *En*. NAFU1 powdery mildew, and the detection of cell death was performed by trypan blue staining in the leaves of CM3G2-40 or CM3G1-91 plants at 15 dpi. Three individuals from each CM3G2-40 and CM3G1-91 line were evaluated. Symptoms of powdery mildew infection were evaluated after inoculation using an Olympus BX51 microscope under visible light. DAB staining, trypan blue staining and aniline blue staining were carried out as previously described to detect the accumulation of H₂O₂, fungal structures, cell death and CWAs⁵².

Statistical analysis

Disease severity data were analyzed with ZEN 2012 and SigmaPlot 10.0 software. All data were analyzed using paired Student's *t*-tests (<http://www.physics.csbsju.edu/stats/>). The mean values ± standard deviation of the mean (SD) were calculated based on the results of at least three replicates, and significant differences compared with controls are represented by **p* < 0.05 and ***p* < 0.01.

Acknowledgements

We thank Dr. Yaoguang Liu of South China Agricultural University for providing the pYL-CRISPR/Cas9 system. We also thank Dr. Steve van Nocker of Michigan State University and Bruce Levine of University of Maryland for careful editing of this paper. This work was supported by the National Key Research and Development Program of China (2018YFD1000300) and the National Natural Science Foundation of China (Grant No. 31772264) to Y.-Q.W., and NSF support (IOS-1901566) to S.X.

Author details

¹State Key Laboratory of Crop Stress Biology for Arid Areas, College of Horticulture, Northwest A&F University, Yangling 712100 Shaanxi, China. ²Key Laboratory of Horticultural Plant Biology and Germplasm Innovation in Northwest China, Ministry of Agriculture, Yangling 712100 Shaanxi, China. ³Institute for Bioscience and Biotechnology Research & Department of Plant Sciences and Landscape Architecture, University of Maryland College Park, Rockville, MD 20850, USA

Author contributions

Y.-Q.W. conceived the project. D.Y.W. and Y.Q.W. designed experiments and wrote the paper. D.Y.W. and Y.G. performed most of the experiments. Y.C. performed the *Agrobacterium*-mediated transformation of grapevine. Y.H. assisted with the data analysis. S.X. assisted with the data analysis and wrote the paper. Y.W. contributed to the study via consultation. All of the authors approved the final paper.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-0339-8>).

Received: 24 April 2019 Revised: 8 May 2020 Accepted: 12 May 2020

Published online: 01 August 2020

References

- Qiu, W., Feechan, A. & Dry, I. Current understanding of grapevine defense mechanisms against the biotrophic fungus (*Erysiphe necator*), the causal agent of powdery mildew disease. *Hortic. Res.* **2**, 15020 (2015).
- Bouchez, D. & Höfte, H. Update on genomics functional genomics in plants. *Plant Physiol.* **118**, 725 (1998).
- Kumar, V. & Jain, M. The CRISPR–Cas system for plant genome editing: advances and opportunities. *J. Exp. Bot.* **66**, 47 (2015).
- Li, J. F. et al. Multiplex and homologous recombination-mediated genome editing in *Arabidopsis* and *Nicotiana benthamiana* using guide RNA and Cas9. *Nat. Biotechnol.* **31**, 688 (2013).
- Nakajima, I. et al. CRISPR/Cas9-mediated targeted mutagenesis in grape. *PLoS ONE* **12**, e177966 (2017).
- Shan, Q. et al. Targeted genome modification of crop plants using a CRISPR–Cas system. *Nat. Biotechnol.* **31**, 686 (2013).
- Liang, Z., Zhang, K., Chen, K. & Gao, C. Targeted mutagenesis in *Zea mays* using TALENs and the CRISPR/Cas system. *J. Genet. Genomics* **41**, 63–68 (2014).
- Gao, W. et al. Genome editing in cotton with the CRISPR/Cas9 system. *Front. Plant Sci.* **8**, 1364 (2017).
- Ren, C. et al. CRISPR/Cas9-mediated efficient targeted mutagenesis in Chardonnay (*Vitis vinifera* L.). *Sci. Rep.* **6**, 32289 (2016).
- Wang, X. et al. CRISPR/Cas9-mediated efficient targeted mutagenesis in grape in the first generation. *Plant Biotechnol. J.* **16**, 844 (2018).
- Malnoy, M. et al. DNA-free genetically edited grapevine and apple protoplast using CRISPR/Cas9 ribonucleoproteins. *Front. Plant Sci.* **7**, 1904 (2016).
- Dangl, J. L. & Jones, J. D. Plant pathogens and integrated defence responses to infection. *Nature* **411**, 826 (2001).
- Vogel, J. P., Raab, T. K., Somerville, C. R. & Somerville, S. C. Mutations in *PMR5* result in powdery mildew resistance and altered cell wall composition. *Plant J.* **40**, 968 (2004).
- de Toledo Thomazella, D. P., Brail, Q., Dahlbeck, D. & Staskawicz, B. CRISPR–Cas9 mediated mutagenesis of a *DMR6* ortholog in tomato confers broad-spectrum disease resistance. Preprint at <http://biorxiv.org/content/early/2016/07/20/064824> (2016).
- van Schie, C. C. N. & Takken, F. L. W. Susceptibility genes 101: how to be a good host. *Annu. Rev. Phytopathol.* **52**, 551 (2014).
- Peng, A. et al. Engineering canker-resistant plants through CRISPR/Cas9-targeted editing of the susceptibility gene *CsLOB1* promoter in citrus. *Plant Biotechnol. J.* **15**, 1509 (2017).
- Char, S. N. et al. An *Agrobacterium*-delivered CRISPR/Cas9 system for high-frequency targeted mutagenesis in maize. *Plant Biotechnol. J.* **15**, 257 (2017).
- Li, T., Liu, B., Spalding, M. H., Weeks, D. P. & Yang, B. High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nat. Biotechnol.* **30**, 390 (2012).
- Jørgensen, I. H. Discovery, characterization and exploitation of *Mlo* powdery mildew resistance in barley. *Euphytica* **63**, 141 (1992).
- Buschges, R. et al. The barley *Mlo* gene: a novel control element of plant pathogen resistance. *Cell* **88**, 695 (1997).
- Consonni, C. et al. Conserved requirement for a plant host cell protein in powdery mildew pathogenesis. *Nat. Genet.* **38**, 716 (2006).
- Elliott, C. et al. Conserved extracellular cysteine residues and cytoplasmic loop-loop interplay are required for functionality of the heptahelical MLO protein. *Biochem. J.* **385**, 243 (2005).
- Devoto, A. et al. Molecular phylogeny and evolution of the plant-specific seven-transmembrane MLO family. *J. Mol. Evol.* **56**, 77 (2003).
- Kusch, S., Pesch, L. & Panstruga, R. Comprehensive phylogenetic analysis sheds light on the diversity and origin of the MLO family of integral membrane proteins. *Genome. Biol. Evol.* **8**, 878 (2016).
- Wolter, M., Hollricher, K., Salamini, F. & Schulze-Lefert, P. The *mlo* resistance alleles to powdery mildew infection in barley trigger a developmentally controlled defence mimic phenotype. *Mol. Gen. Genet.* **239**, 122 (1993).
- Panstruga, R. Serpentine plant MLO proteins as entry portals for powdery mildew fungi. *Biochem. Soc. Trans.* **33**, 389–392 (2005).
- Wang, Y. et al. Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* **32**, 947 (2014).
- Pessina, S. et al. The knock-down of the expression of *MdMLO19* reduces susceptibility to powdery mildew (*Podosphaera leucotricha*) in apple (*Malus domestica*). *Plant Biotechnol. J.* **14**, 2033 (2016).
- Feechan, A., Jermakow, A. M., Torregrosa, L., Panstruga, R. & Dry, I. B. Identification of grapevine *MLO* gene candidates involved in susceptibility to powdery mildew. *Funct. PLANT Biol.* **35**, 1255 (2008).
- Pessina, S. et al. Knockdown of *MLO* genes reduces susceptibility to powdery mildew in grapevine. *Hortic. Res.* **3**, 16016 (2016).
- Ma, X. et al. A robust CRISPR/Cas9 system for convenient, high-efficiency multiplex genome editing in monocot and dicot plants. *Mol. Plant* **8**, 1274 (2015).
- Jinek, M. et al. A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816 (2012).
- Gao, Y. et al. Identification and utilization of a new *Erysiphe necator* isolate NAFU1 to quickly evaluate powdery mildew resistance in wild Chinese grapevine species using detached leaves. *Plant Physiol. Bioch.* **98**, 12 (2016).
- Wang, Y. et al. Identification of genomic sites for CRISPR/Cas9-based genome editing in the *Vitis vinifera* genome. *BMC Plant Biol.* **16**, 96 (2016).
- Ren, C. et al. Knockout of *WCCD8* gene in grapevine affects shoot branching. *BMC Plant Biol.* **20**, 47 (2020).
- Charrier, A. et al. Efficient targeted mutagenesis in apple and first time edition of pear using the CRISPR–Cas9 system. *Front. Plant Sci.* **10**, 40 (2019).
- Kusch, C., Arslan, S., Singh, R., Thorpe, J. & Adli, M. Genome-wide analysis reveals characteristics of off-target sites bound by the Cas9 endonuclease. *Nat. Biotechnol.* **32**, 677 (2014).
- Feng, Z. et al. Multigeneration analysis reveals the inheritance, specificity, and patterns of CRISPR/Cas-induced gene modifications in *Arabidopsis*. *Proc. Natl Acad. Sci. USA* **111**, 4632 (2014).
- Kusch, S. & Panstruga, R. *mlo*-Based resistance: an apparently universal “weapon” to defeat powdery mildew disease. *Mol. Plant-Microbe Interact.* **30**, 179 (2017).
- Nekrasov, V. et al. Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. *Sci. Rep.* **7**, 482 (2017).
- Winterhagen, P., Howard, S. F., Qiu, W. & Kovács, L. G. Transcriptional up-regulation of grapevine *MLO* genes in response to powdery mildew infection. *Am. J. Enol. Viticult.* **59**, 159 (2008).
- Feechan, A. et al. Host cell entry of powdery mildew is correlated with endosomal transport of antagonistically acting *VvPEN1* and *VvMLO* to the papilla. *Mol. Plant Microbe.* **26**, 1138 (2013).
- Nsel, C. P., Freialdenhoven, A., Kurth, J., Kolsch, R. & Schulze-Lefert, P. Interaction analyses of genes required for resistance responses to powdery mildew in barley reveal distinct pathways leading to leaf cell death. *Plant Cell* **9**, 1397–1409 (1997).
- Piffanelli, P. et al. The barley *MLO* modulator of defense and cell death is responsive to biotic and abiotic stress stimuli. *Plant Physiol.* **129**, 1076 (2002).
- Zheng, Z. et al. Loss of function in *Mlo* orthologs reduces susceptibility of pepper and tomato to powdery mildew disease caused by *Leveillula taurica*. *PLoS ONE* **8**, e70723 (2013).
- Cheng, Y., Wan, D., Zhang, B., Wang, Y. & Wen, Y. Induction, preservation, propagation and cytological observation of proembryonic masses of grapevine. *J. Fruit. Sci.* **8**, 968 (2017).
- Lei, Y. et al. CRISPR-P: a web tool for synthetic single-guide ma design of CRISPR-system in plants. *Mol. Plant* **7**, 1494 (2014).
- Ma, X. & Liu, Y. G. CRISPR/Cas9-based multiplex genome editing in monocot and dicot plants. *Curr. Protoc. Mol. Biol.* **115**, 31 (2016).
- Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision cloning method with high throughput capability. *PLoS ONE* **3**, e3647 (2008).
- Hu, Y. et al. Ectopic expression of *Arabidopsis* broad-spectrum resistance gene *RPW8.2* improves the resistance to powdery mildew in grapevine (*Vitis vinifera*). *Plant Sci.* **267**, 20 (2018).
- Doyle, J. J. & Doyle, J. L. Isolation of plant DNA from fresh tissue. *Focus* **12**, 12 (1990).
- Xiao, S., Brown, S., Patrick, E., Brearley, C. & Turner, J. G. Enhanced transcription of the *Arabidopsis* disease resistance genes *RPW8.1* and *RPW8.2* via a salicylic acid-dependent amplification circuit is required for hypersensitive cell death. *Plant Cell* **15**, 33 (2003).