

RESEARCH

Open Access



# Variable phenotypes and outcomes associated with the *MMACHC* c.609G>A homologous mutation: long term follow-up in a large cohort of cases

Ruxuan He<sup>1†</sup>, Ruo Mo<sup>1†</sup>, Ming Shen<sup>2†</sup>, Lulu Kang<sup>1</sup>, Jinqing Song<sup>1</sup>, Yi Liu<sup>1</sup>, Zhehui Chen<sup>1</sup>, Hongwu Zhang<sup>3</sup>, Hongxin Yao<sup>3</sup>, Yupeng Liu<sup>4</sup>, Yao Zhang<sup>1</sup>, Hui Dong<sup>1</sup>, Ying Jin<sup>1</sup>, Mengqiu Li<sup>1</sup>, Jiong Qin<sup>4</sup>, Hong Zheng<sup>5</sup>, Yongxing Chen<sup>6</sup>, Dongxiao Li<sup>6</sup>, Haiyan Wei<sup>6</sup>, Xiyuan Li<sup>7</sup>, Huifeng Zhang<sup>8</sup>, Min Huang<sup>9</sup>, Chunyan Zhang<sup>2</sup>, Yuwu Jiang<sup>1</sup>, Desheng Liang<sup>10\*</sup>, Yaping Tian<sup>2\*</sup> and Yanling Yang<sup>1\*</sup>

## Abstract

**Background:** Cobalamin C deficiency (*cbfC*) caused by the *MMACHC* mutations is the most common type of the disorders of intracellular cobalamin metabolism. While the c.609G > A mutation is most frequent in Chinese *cbfC* patients, its correlation with phenotype has not been delineated. Here we aim to investigate the factors affecting variable phenotypes and outcomes associated with the *MMACHC* c.609G > A homologous mutation in 149 Chinese cases to have implications for treatment and prevention.

**Methods:** We assessed 149 *cbfC* patients caused by *MMACHC* c.609G > A homozygous mutation. The clinical manifestations, complications, treatment, and outcomes were evaluated; 120 patients were followed-up till December 2019.

(Continued on next page)

\* Correspondence: [liangdesheng@sklmg.edu.cn](mailto:liangdesheng@sklmg.edu.cn); [tianyp@301hospital.com.cn](mailto:tianyp@301hospital.com.cn); [organic.acid@vip.126.com](mailto:organic.acid@vip.126.com)

<sup>†</sup>Ruxuan He, Ruo Mo and Ming Shen contributed equally to this work as co-first authors

<sup>10</sup>School of Life Sciences, Central South University, Changsha 410013, China

<sup>2</sup>Research Center for Translational Medicine, Medical Innovation Research Division of Chinese PLA General Hospital, Beijing 100853, China

<sup>1</sup>Department of Pediatrics, Peking University First Hospital, Beijing 100034, China

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

**Results:** Two patients (1.3%) were prenatally diagnosed, treated after birth and consequently showed normal development. In 15 patients (10.1%) detected by newborn screening, 10 were treated at the age of 2 weeks and showed normal development, while the other 5 were treated after onset and showed neurologic disorders. All 132 clinically diagnosed patients (88.6%) developed symptoms at age from few minutes after birth to 72 months. Among them, 101 (76.5%) had early-onset (before the age of 12 months) and 31 (23.5%) had late-onset (after the age of 12 months). Totally 5 patients died and 24 were lost to follow-up. Of the 132 clinical diagnosed patients, 92 (69.7%) presented with developmental delay, 65 (49.2%) had seizures, 37 (28.0%) had anemia, 24 (18.2%) had feeding difficulty, 23 (17.4%) had ocular problems, and 22 (16.7%) had hydrocephalus. Compared with the non-developmental delay group, the onset age, the age at treatment initiation and the time from onset to treatment initiation were later in the developmental delay group. Seizure group showed significantly higher urinary methylmalonic acid concentration. During long-term follow-up, plasma total homocysteine (tHcy) levels were significantly higher in patients in the uncontrolled group than those in the seizure-free group.

**Conclusions:** Most *cbIC* patients caused by *MMACHC* c.609G > A homozygous mutation showed early-onset. The clinically diagnosed patients usually showed the presence of irreversible brain disorders. Patients treated from the pre-symptomatic stage showed favorable outcomes. Therefore, newborn screening, prenatal diagnosis and early treatment are crucial and the c.609G > A mutant allele should be listed in the pre-pregnancy carrier screening panel in China.

**Keywords:** Cobalamin C deficiency (*cbIC*), The *MMACHC* gene, C.609G > A, Methylmalonic acidemia and homocysteinemia

## Background

The *cbIC* type (*cbIC*) of combined methylmalonic acidemia and homocystinuria is the most common defect in the intracellular cobalamin metabolism pathway, characterized by variable and non-specific symptoms especially in childhood [1–3]. An increasing hospitalization with various presentations and a heavy financial burden per hospitalization were observed in Mainland China, while the medical resources were still relatively centralized in some districts, such as Beijing, Shanghai and Guangzhou [4]. All the disorders of intracellular cobalamin metabolism are inherited in an autosomal recessive manner except for *cbIX*, which is inherited in an X-linked manner [5]. In the patients with *cbIC*, age of onset ranges from prenatal to adult and clinical manifestations vary from mild to life-threatening [6]. The most common *MMACHC* pathogenic variant previously reported, c.609G > A (p.Trp203X), results in a premature termination codon that is predicted to cause a truncated or absent *MMACHC* protein due to nonsense mediated decay and accounts for 48.1% of mutant alleles in 79 Chinese patients with *cbIC* [3]. This variant is listed in the Genome Aggregation Database (gnomAD) with an overall population frequency of 0.004% and has been reported in numerous affected individuals in the homozygous and compound heterozygous state, and there is evidence that the variant may be a founder mutation in the Chinese population [3, 7].

The *MMACHC* c.609G > A is the hot spot mutation in Chinese patients with early-onset *cbIC* [2]. Neurological disorders are common complications of *cbIC*, which may

result in varying degrees of sequelae [2, 3]. Epilepsy is a frequent symptom in children with early-onset *cbIC* [8]. With an improvement in clinical diagnosis, there has been an increase in the number of cases being diagnosed with *cbIC*-related epilepsy or other neurological diseases. Patients with *cbIC* caused by the *MMACHC* c.609G > A homozygous mutation have been found to show significant differences in phenotypes and outcomes.

In the present study, we carried out a retrospective analysis of 149 Chinese *cbIC* patients with the *MMACHC* c.609G > A homozygous mutation who were followed up in Department of Pediatrics, Peking University First Hospital, Beijing. Clinical features, metabolites, diagnosis, process of management and outcomes were reviewed and factors affecting variable phenotypes and outcomes were investigated.

## Methods

### Patients and data collection (Table 1)

From January 1998 to December 2019, 149 patients with *cbIC* attributable to *MMACHC* c.609G > A homozygous mutation were diagnosed by gene sequencing at the Peking University First Hospital in China. These patients came from 11 provinces or cities of Mainland China. Five cases died and 24 cases were lost to follow-up. The remaining 120 cases received personalized treatment and regular follow-up. Patients with definite pathogenic variants in genes besides *MMACHC* were excluded.

General data were collected, including the age at onset and diagnosis, clinical manifestations, family history, treatment and outcomes. Electroencephalography and

**Table 1** General information pertaining to 149 patients caused by the *MMACHC* c.609G > A homozygous mutation

Patients	n	Male		Female	
		n	%	n	%
Diagnosed prenatally and treated at birth	2	1	0.7	1	0.7
Detected by newborn screening	15	7	4.7	8	5.4
Treated at birth	10	6	4.0	4	2.7
Treated after onset	5	1	0.7	4	2.7
Clinically diagnosed	132	73	49.0	59	39.6
Death	5	2	1.3	3	2.0
Regular follow-up	103	60	40.3	43	28.9
Lost to follow-up	24	11	7.4	13	8.7
Total	149	81	54.4	68	45.6

Note: n number

brain imaging by MRI or CT were performed. Elevated blood propionylcarnitine and urinary methylmalonic acid levels confirmed the diagnosis of MMA. All patients had significantly elevated plasma total homocysteine (tHcy) (34.3–278  $\mu\text{mol/L}$ , normal control  $\leq 15 \mu\text{mol/L}$ ). Moreover, routine blood and urine examination were performed to evaluate liver, renal, and heart function.

#### Biochemical examination

Dried blood spots were collected. Blood amino acids, free carnitine, and acylcarnitines were analyzed using liquid chromatography–tandem mass spectrometry (Waters MS/MS system A, 1445–002; API3200, Applied Biosystems, CA, USA), as previously described [9, 10]. Metabolite concentrations were automatically calculated using the Chemoview software [2, 9]. The reference range for propionylcarnitine was 1.00–5.00  $\mu\text{mol/L}$ .

Gas chromatography–mass spectrometry (GC–MS), which was performed on a Shimadzu GCMS-QP2010 system (Kyoto, Japan), was used to analyze urinary organic acids, as previously reported. Mass spectra were obtained by standard electron impact ionization scanning from 50  $m/z$  to 500  $m/z$ . Data were collected using a GC–MS solution software [11, 12].

Plasma total homocysteine (tHcy) were assessed using fluorescence polarization immunoassay [2].

#### Treatment

For patients during acute decompensation, the initial therapy involved intramuscular or intravenous injections of cobalamin (hydroxocobalamin was the first choice at a dose of 1 mg per day), L-carnitine (50–200 mg/kg per day), intravenous fluid therapy with glucose and electrolytes, oral betaine (100–500 mg/kg per day), a high-calorie diet with symptomatic treatment. If hydroxocobalamin was not available, methylcobalamin was used

for injection. All patients followed a normal diet. For the patients with methionine deficiency, methionine was supplemented by oral. The long-term treatment was then adjusted depending on the condition of individual patients [2, 13].

#### Genomic analyses

Genomic DNA was extracted from peripheral blood of patients and their parents using a TIANamp® Blood DNA Kit (Tiagen Biotech Co. Ltd., Beijing, China), as per manufacturer instructions. Next-generation sequencing (NGS) was performed to screen for mutations (Running Gene Inc., Beijing, China; Berry Genomics Corporation, Beijing, China; Translational Medicine Laboratory, Chinese People's Liberation Army General Hospital, Beijing, China). The *MMACHC* c.609G > A homozygous mutation was found in 149 patients. Patients with other gene mutations associated with MMA and homocystinemia were excluded.

#### Statistical analysis

SPSS 24.0 (SPSS Inc., Chicago, IL, USA) was used for statistical analyses. Analyses using the Shapiro–Wilk method showed that the data did not conform to normal distribution, so the values are expressed as median (quartile), and nonparametric unpaired Mann–Whitney U test was also performed. All tests were set as two-tailed, and  $P < 0.05$  indicated statistical significance.

## Results

#### Patients (Table 1)

In total, 149 patients with *cbIC* attributable to the *MMACHC* c.609G > A homozygous mutation were analyzed. Among them, 81 (54.4%) were males and 68 (45.6%) were females. Two unrelated cases (one male and one female) were prenatally diagnosed based on the family history that their elder siblings had died of *cbIC* [14, 15]. *CbIC* was diagnosed by the increasing of methylmalonic acid and total homocysteine in amniotic fluid and *MMACHC* c.609G > A homozygous mutation in amniocytes at the gestational age of 20 weeks. No abnormal echo was found in the two fetuses by ultrasound examination. Mild carnitine deficiency, slight methylmalonic aciduria and normal plasma total homocysteine were observed in their mothers. The mothers chose to continue their pregnancy. Oral methylcobalamin (1 mg per day), L-carnitine (1 g per day) and multivitamins were supplemented during their pregnancy to support the growth and development of fetuses. Their blood free carnitine levels were maintained at the normal range. Two patients were treated from the first day of birth. At present, they are aged 3 and 11 years respectively, with normal physical and neurocognitive development. No abnormal finding was found by eye test.

Fifteen cases (10.1%) were detected by newborn screening. Among them, treatment was administered to ten cases (six males and four females; intramuscular cobalamin injection, oral administration of L-carnitine, and betaine) at the age of 15 days. They are currently 2 to 8 years old, with normal psychomotor and physical development. Unfortunately, parents of the other five cases (one male and four females) refused the confirmative testing and presymptomatic treatment. These cases developed symptoms at the age of 1–4 months, including developmental delay in all five cases, hydrocephalus in two cases, and seizures in one case.

### Phenotypic features

One hundred and thirty-two cases (88.6%) were clinically diagnosed. The onset age ranged from a few minutes after birth to age 72 months (median age 3 months). The average time from onset to confirmative diagnosis was 10 months. Early onset (before the age of 12 months) was observed in 101 cases (76.5%), of whom 36 (27.3%) exhibited symptoms during the neonatal period. Late onset (after the age of 12 months) was observed in 31 cases (23.5%). The most common clinical manifestations were developmental delay (69.7%), followed by seizures (49.2%), anemia (28.0%), feeding difficulty (18.2%), ocular problems (17.4%), and hydrocephalus (16.7%) (Table 2). Of the 132 clinically diagnosed patients, strabismus, nystagmus, visual impairment, and maculopathy were observed in 23 patients (17.4%) at their first visit by eye tests. Frequencies were compared between early-onset ( $n = 101$ ) and late-onset ( $n = 31$ ) groups. As shown

in Table 2, the frequency of developmental delay was significantly higher in the late-onset group, and the frequency of hydrocephalus was significantly higher in the early-onset group.

### Follow-up and outcomes

The two cases diagnosed prenatally and 15 cases detected by newborn screening were followed up regularly.

Of the 132 clinical diagnosed cases (Tables 1, 2), 37 were misdiagnosed as nutritional anemia ( $n = 9$ ), cerebral palsy ( $n = 9$ ), hypoxic ischemic encephalopathy ( $n = 8$ ), epilepsy ( $n = 5$ ), pneumonia ( $n = 4$ ), and autism ( $n = 2$ ). The age at diagnosis ranged from 3 days to 101 months (median 6 months). Time delays from the onset to treatment initiation was 7 days to 96 months (median 2 months, average = 12 months). Early onset was observed in 101 cases (76.5%), with symptoms appearing before the age of 1 year. Late onset was observed in 31 cases (23.5%), with symptoms appearing from the age of 14 months to 72 months (median 24 months), and median age of diagnosis is 39 months. Five (3.8%) patients died. Two of them died of intractable epilepsy at the age of 9 months and 10 years, respectively. The three others died of multiple organ failure induced by infection at the age of 3 months, 2 years and 4 years, respectively. Twenty-four cases (16.1%) were lost to follow-up.

### Factors affecting variable phenotypes and outcomes

One hundred and thirty-two clinical diagnosed patients were divided into two groups depending on whether they showed a developmental delay, epilepsy, hydrocephalus,

**Table 2** Clinical manifestations of 132 clinical diagnosed *cb1C* patients caused by the *MMACHC* c.609G > A homozygous mutation

Clinical manifestations	Cases		Early-onset ( $n = 101$ )		Late-onset ( $n = 31$ )		<b>P</b> value
	n	%	n	%	n	%	
Developmental delay	92	69.7	63	62.4	29	93.5	<b>0.003</b>
Epilepsy	65	49.2	54	53.5	11	35.5	0.053
Anemia	37	28.0	32	31.7	5	16.1	0.073
Feeding difficulty	24	18.2	21	20.8	3	9.7	0.138
Hydrocephalus	22	16.7	21	20.8	1	3.2	<b>0.018</b>
Ocular problems	23	17.4	19	18.8	4	12.9	0.399
Vomiting	19	14.4	15	14.9	4	12.9	0.951
Hypotonia/abnormal muscle tone	15	11.4	10	9.9	4	12.9	0.944
Skin lesions	12	9.1	11	10.9	1	3.2	0.319
Lactic acidosis	6	4.5	6	5.9	0	0.0	0.335
Liver damage	5	3.8	5	5.0	0	0.0	0.335
Renal damage	2	1.5	1	1.0	1	3.2	0.427
Cardiomyopathy	0	0.0	0	0.0	0	0.0	
Pulmonary hypertension	0	0.0	0	0.0	0	0.0	

Notes:  $n$  number;  $P$  value was calculated by using a chi-square test.  $P < 0.05$  was considered as statistically significant

and anemia. Factors including age at onset, age at treatment initiation, time from onset to treatment initiation, and biochemical metabolic markers at first visit were analyzed, except for two cases whose data was not available (Table 3). There was significant difference in the onset time, the age at treatment initiation and the time from onset to treatment initiation between developmental delay and non-developmental delay groups.

In comparison with patients in the non-epileptic group, those in the epileptic group showed significantly higher levels of urinary methylmalonic acid. There were no significant differences in age at onset, age at treatment initiation, time from onset to treatment initiation, and biochemical metabolic markers (tHcy, methionine, free carnitine, acetylcarnitine, propionylcarnitine, propionylcarnitine/free carnitine ratio, and propionylcarnitine/acetylcarnitine ratio) between the groups (Table 3).

Forty-one patients with seizures were divided into two groups, seizure-free ( $n = 21$ ) and uncontrolled ( $n = 20$ ) groups, depending on whether they still had clinical

seizures in the last 6 months (up to December 2019). As of December 2019, the age of the seizure-free group ranged from 1 year 5 months to 26 years 7 months (median 7 years 7 months). The age of the uncontrolled group ranged from 6 years to 16 years 3 months (median 10 years 8 months). The treatment recommendations for the two groups were the same. However, it is difficult to confirm whether the injection is missed because all the patients were given parenteral route of administration IM at home for long-term treatment. Some of the parents had poor compliance to frequent injection for their children. To reduce the plasma total homocysteine, the dose of cobalamin should be increased for the patients of uncontrolled group. General information pertaining to patients and recent condition were compared between the groups depending on whether clinical seizures had been controlled (Table 4).

During long-term follow-up, based on recent metabolic studies, we noted a significant difference in plasma total homocysteine (tHcy) levels among patients in the

**Table 3** Variable phenotypes and factors in 130 patients with *MMACHC* c. 609G > A homozygous mutation

Symptoms and signs	Cases	Age at onset	Age at treatment initiation	Time delays from onset to treatment initiation	Urinary methylmalonic acid pre-treatment	Blood tHcy pre-treatment	Blood methionine pre-treatment	
	n	(months) Median (P25, P75)	(months) Median (P25, P75)	(months) Median (P25, P75)	(mmol/mol creatinine) Median (P25, P75)	( $\mu$ mol/L) Median (P25, P75)	( $\mu$ mol/L) Median (P25, P75)	
<b>Seizures</b>	N	65	3.0 (1.0, 15.0)	6.5 (2.0, 34.0)	1.5 (0.0, 18.3)	184.7 (43.5, 560.1)	99.8 (73.7, 129.6)	
	Y	65	2.0 (0.9, 5.0)	5.5 (3.0, 27.0)	2.3 (1.0, 12.0)	493.4 (259.9, 906.8)	96.0 (58.8, 151.7)	
	Z		-1.407	-0.031	-0.893	-2.239	-0.304	-0.87
	P		0.159	0.975	0.372	<b>0.025</b>	0.761	0.406
<b>Developmental delay</b>	N	38	1.0 (0.7, 3.0)	3.0 (1.5, 5.5)	1.5 (0.9, 2.3)	164.6 (35.0, 664.9)	102.9 (65.3, 142.6)	
	Y	92	3.0 (1.0, 12)	12.0 (3.3, 36)	6.0 (0.5, 25.0)	443.6 (199.2, 722.1)	88.0 (59.1, 147.7)	
	Z		-2.892	-3.746	-2.017	-1.481	-0.313	-0.707
	P		<b>0.004</b>	<b>&lt; 0.001</b>	<b>0.044</b>	0.14	0.754	0.503
<b>Hydrocephalus</b>	N	108	2.0 (1.0, 10.0)	8.0 (3.0, 34.0)	2.0 (0.5, 18.0)	282.1 (76.7, 768.1)	93.0 (63.0, 143.7)	
	Y	22	2.0 (1.0-4.0)	4.0 (2.8-0.5)	1.8 (0.9-6.8)	500.0 (257.3-560.1)	101.0 (59.1-161.9)	14.0 (5.7-21.5)
	Z		-0.487	-0.92	-0.531	-0.608	-0.546	-0.445
	P		0.626	0.357	0.595	0.552	0.593	0.695
<b>Anemia</b>	N	93	3.0 (1.0, 7.8)	8.0 (3.0, 34)	3.0 (0.0, 24)	292.6 (92.3, 676.1)	87.2 (58.8, 127.1)	
	Y	37	2.0 (1.0, 5.0)	3.0 (2.0, 18.0)	1.5 (1.0, 9.0)	442.3 (97.0, 672.9)	137.3 (74.3, 164.9)	
	Z		-0.622	-2.185	-0.777	-0.014	-2.172	-0.64
	P		0.534	0.029	0.437	0.989	<b>0.03</b>	0.56

Notes: n number, N No, Y Yes, Z Mann-Whitney U Z-score; P value was calculated by using a nonparametric unpaired Mann-Whitney U test.  $P < 0.05$  was considered as statistically significant

**Table 4** Data of 41 long-term follow-up patients caused by the *MMACHC* c.609G > A homozygous mutation with seizures

Group of the patients	Uncontrolled			Seizure-free			Z	P
	(n = 20)			(n = 21)				
	Median	P25	P75	Median	P25	P75		
Age at onset (months)	1	0.3	10.0	2	1.0	5.0	-0.756	0.472
Age at treatment initiation (months)	13	1.3	59.0	6.5	2.3	28.5	-0.047	0.982
Time from onset to treatment initiation (months)	8	1.3	52.0	5.5	0.7	23.3	-0.875	0.39
Latest urinary methylmalonic acid	20.5	6.1	75.0	16.5	12.4	47.9	-0.279	0.802
Latest blood metabolic markers								
Methionine (μmol/L)	17.1	14.9	23.5	15.7	12.8	27.1	-0.537	0.603
tHcy (μmol/L)	82.9	58.2	96.6	51.4	36.2	66.0	-2.413	<b>0.015</b>
Free carnitine (μmol/L)	56.8	40.0	77.7	56.6	27.6	69.2	-0.67	0.535
Acetylcarnitine (μmol/L)	26.1	14.8	33.2	23.5	9.9	28.3	-0.952	0.368
Propionylcarnitine (μmol/L)	5.4	3.7	7.7	3.7	0.8	6.5	-1.093	0.298
Propionylcarnitine/free carnitine ratio	0.09	0.08	0.12	0.08	0.06	0.18	-0.462	0.68
Propionylcarnitine/ acetylcarnitine ratio	0.27	0.18	0.31	0.2	0.19	0.25	-1.029	0.332

Notes: n number, Z Mann-Whitney U Z-score; P value was calculated by using a nonparametric unpaired Mann-Whitney U test.  $P < 0.05$  was considered as statistically significant

seizure-free and uncontrolled groups. Plasma tHcy in patients in the uncontrolled group were significantly higher than those in the seizure-free group. There were no significant differences in the age at onset, age at diagnosis, time from onset to diagnosis, latest urinary methylmalonic acid levels, blood free carnitine, acetylcarnitine, propionylcarnitine, propionylcarnitine/free carnitine ratio, and propionyl carnitine/acetylcarnitine ratio between the groups.

Skin lesions were present in 12 patients. Their blood methionine level was significantly low (5.6–10 μmol/L, normal control 12–50 μmol/L), and they presented with feeding difficulties, malnutrition, and failure to thrive. Five patients showed leucine, valine, and threonine deficiencies. Skin erosions occurred in three infants 3 to 5 days after feeding by medical formula for cobalamin non-responsive isolated methylmalonic aciduria. The condition of all patients improved soon after metabolic treatment with normal diet.

## Discussion

*CblC* deficiency caused by *MMACHC* mutation is the most common type of MMA combined with homocysteinemia. Although the genetic epidemiological data on *CblC* deficiency are limited, the carrier prevalence of the *MMACHC* mutation has been deduced to be 1/28–31 from the reported *cblC* incidences in the population in Shandong province, China [16, 17]. The c.609G > A pathogenic variant has been reported the hot spot mutation in Chinese patients with early onset *cblC* deficiency [18–20], however, the correlation of phenotype with the c.609G > A mutant allele has not been described.

In the present study, we investigated the clinical and biochemical characteristics of the patients with *cblC* deficiency caused by *MMACHC* c.609G > A homozygous mutation to discuss the optimal strategies of treatment and prevention. All of the 149 patients with the same genotype of c.609G > A homozygous alleles presented with significant differences in clinical phenotypes, ranging from asymptomatic condition to severe brain damage to death.

With early treatment, developmental milestones of the two prenatally diagnosed patients and the 10 patients detected by newborn screening were normal. No organ damage occurred. These cases benefited from early diagnosis, treatment, and prevention of severe complications. Fortunately, newborn screening for MMA is widely promoted around China in recent years, which has proven to be helpful to improve the outcome of patients with MMA [21, 22]. The five patients detected by newborn screening but treated after disease onset because of poor treatment compliance showed severe mental and motor retardation. Two of them had hydrocephalus, and one had seizures. A delay in treatment leads to irreversible brain damage, pointing out emphatically the pre-symptomatic treatment and education to improve parents' understanding and medication compliance.

One hundred and thirty-two patients who were diagnosed after onset with the *MMACHC* c.609G > A homozygous mutation showed various symptoms. The most common symptoms were developmental delay, epilepsy, anemia, feeding difficulty and hydrocephalus. Brain damage is also commonly observed in patients with the *MMACHC* c.482G > A mutation [23]. Although most

patients showed an improvement after metabolic treatment, there are varied neurological sequelae. The level of urinary methylmalonic acid was higher in patients with seizures pre-treatment, indicating an increase in toxic metabolites. Seizures have been reported to occur after injecting methylmalonic acid into the striatum of rats [24, 25]. Patients with no seizures show lower urinary methylmalonic acid levels, which is considered to be associated with severe brain damage in patients with higher levels of methylmalonic acid [26, 27]. It is more difficult to control seizures in patients with persistently high plasma tHcy levels. Excluding poor medication compliance, it is suggested that homocysteine may play a role in the pathogenesis. In vitro studies have suggested that homocysteine-induced phosphorylation disorder, overactivation of N-methyl-D-aspartate receptor [28], vascular oxidative stress, and inflammation [29] are related to neurological dysfunction [30]. Moreover, neurologic involvement in *CblC* patients was reported due to brain choline deficiency caused by the transmethylation defect [31]. There was also a small cohort study that found no significant correlation between plasma homocysteine or methylmalonic acid and the long-term prognosis of neurodevelopment [32]. Therefore, the relevant metabolic indicators still need to be further explored.

Methylcobalamin deficiency in patients with *cblC* deficiency is reportedly caused by elevated tHcy and reduced tetrahydrofolic acid levels. Impaired folic acid metabolism affects the synthesis of nucleotides [33]. In the 132 patients of this study, the main hematological abnormality was megaloblastic anemia, which is consistent with the results of a previous study [26]. At the first visit, blood tHcy levels in patients with anemia were higher than in those without anemia.

Eye diseases are common in patients with the *MMAC HC* gene *c.271dupA* homozygous mutation and are characterized by early maculopathy. However, in this study, the incidence of ocular problems in patients with the *c.609G > A* homozygous mutation was 17.4%, which is much lower [34, 35]. It is considered that some patients didn't undergo a detailed examination. An ophthalmological examination should be paid attention to in future diagnosis and treatment.

The *MMACHC c.80A > G* mutation is associated with pulmonary hypertension in patients with *cblC* deficiency [36]. It has been reported that *c.271dupA*, *c.276G > T*, and *c.565C > A* are more common in patients with renal thrombotic microangiopathy caused by MMA [37]. However, only two cases with proteinuria were observed in this study. After metabolic treatment, their proteinuria disappeared. Cardiovascular diseases have not been observed in patients with the *c.609G > A* homozygous mutation.

Skin lesion was found in 12 patients in this study. They are considered to be associated with essential amino acid deficiency due to malnutrition. In three cases, eczema and perineal erosion occurred after a low protein diet and taking the special medical formula for cobalamin non-responsive isolated methylmalonic aciduria. Nine patients had skin lesions and feeding difficulties at the onset of the disease. The condition of all patients improved soon after metabolic treatment with a normal diet.

In the present study, most (76.5%) patients with the *MMACHC c.609G > A* homozygous mutation showed early onset. The *c.609G > A* variant is nonsense and results in pre-mature termination codon, which is predicted to cause a truncated or absent *MMACHC* protein. Similarly, two nonsense mutations, *c.271dupA* and *c.331C > T*, either homozygous or compound heterozygous, have been mainly reported in early onset patients; while *c.394C > T* and *c.482G > A* were reported to be related with late-onset patients [38]. This could be related to the low mRNA transcription level of the *MMACHC* gene and residual function of the enzyme [23, 39].

## Conclusion

We assessed a cohort of 149 patients with *cblC* deficiency caused by the *MMACHC c.609G > A* homozygous mutation to determine phenotypic differences in patients with the same genetic defect. Most patients showed early-onset, culminating in irreversible brain damage. Delayed treatment resulted in developmental delay. Newborn screening and early treatment are pivotal to prevent disabilities, however, some severe cases showed symptoms as early as a few minutes after birth. Newborn screening seemed to be late for very early onset patients. Moreover, given that a much higher carrier prevalence of the *MMACHC c.609G > A* mutation has been deduced in Chinese population, we propose that the *c.609G > A* mutant allele should thus be listed in the pre-pregnancy carrier screening panel in China.

## Acknowledgments

We would like to thank all patients and their families for participating in this study. We are grateful to Berry Genomics Corporation (Beijing, China), Running Gene Inc. (Beijing, China) and Chigene (Beijing) Translational Medical Research Center Co. Ltd. for performing sequencing and analysis. We are also thankful to Similan Clinic (Beijing, China) for carrying out biochemical examinations. We are greatly indebted to Professor Seiji Yamaguchi and his team of Department of Pediatrics, Shimane Medical University, Japan, for their expert technical assistance in diagnosis and treatment of methylmalonic acidemia.

## Authors' contributions

YL, DS L and YP T conceived the study. RX H, R M, and M S are joint first authors, drafted the manuscript. LL K, JQ S, Y L, ZH C, HW Z, HX Y, YP L, Y Z, H D, Y J, MQ L, J Q, H Z, YX C, DX L, HY W, HF Z M H, CY Z and YW J participated in clinical management and patient data collection. XY L and M S interpreted the function of some mutations. DS L verified the genetic data. All authors read and approved the final manuscript.

### Funding

This study is supported by grants from the National Key Research and Development Program of China (No. 2016YFC0905102, 2017YFC1001700, 2019YFC1005100), the National Natural Science Foundation of China (No. 81471097), and Beijing Municipal Science and Technology Commission (No. Z141107004414036, Z151100003915126).

### Availability of data and materials

The datasets used and/or analyzed during the current study is available from the corresponding author on request.

### Ethics approval and consent to participate

This study was approved by the Hospital Institutional Ethics Committee and was conducted in accordance with the Declaration of Helsinki. Written informed consent was obtained from the parents of all patients for sample collection and publication of medical data.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>Department of Pediatrics, Peking University First Hospital, Beijing 100034, China. <sup>2</sup>Research Center for Translational Medicine, Medical Innovation Research Division of Chinese PLA General Hospital, Beijing 100853, China. <sup>3</sup>Department of Pediatric Surgery, Peking University First Hospital, Beijing, China. <sup>4</sup>Department of Pediatrics, People's Hospital of Peking University, Beijing, China. <sup>5</sup>Department of Pediatrics, First Affiliated Hospital of Henan University of Traditional Chinese Medicine, Zhengzhou, China. <sup>6</sup>Department of Endocrinology and Inherited Metabolic, Henan Children's Hospital, Zhengzhou, China. <sup>7</sup>Precision Medicine Center, General Hospital of Tianjin Medical University, Tianjin, China. <sup>8</sup>Department of Pediatrics, Hebei Medical University Second Hospital, Shijiazhuang, China. <sup>9</sup>Similan Clinic, Beijing, China. <sup>10</sup>School of Life Sciences, Central South University, Changsha 410013, China.

Received: 16 March 2020 Accepted: 26 July 2020

Published online: 03 August 2020

### References

- Shibata N, Hasegawa Y, Yamada K, Kobayashi H, Purevsuren J, Yang Y, et al. Diversity in the incidence and spectrum of organic acidemias, fatty acid oxidation disorders, and amino acid disorders in Asian countries: selective screening vs. expanded newborn screening. *Mol Genet Metab Rep*. 2018;16:5–10.
- Liu Y, Liu YP, Zhang Y, Song JQ, Yang YL. Heterogeneous phenotypes, genotypes, treatment and prevention of 1 003 patients with methylmalonic acidemia in the mainland of China. *Zhonghua Er Ke Za Zhi*. 2018;56(6):414–20.
- Liu M-Y, Yang Y-L, Chang Y-C, Chiang S-H, Lin S-P, Han L-S, et al. Mutation spectrum of MMACHC in Chinese patients with combined methylmalonic aciduria and homocystinuria. *J Hum Genet*. 2010;55(9):621–6.
- Jiang Y-Z, Shi Y, Shi Y, Gan L-X, Kong Y-Y, Zhu Z-J, et al. Methylmalonic and propionic acidemia among hospitalized pediatric patients: a nationwide report. *Orphanet J Rare Dis*. 2019;14(1):292.
- Yu H-C, Sloan JL, Scharer G, Brebner A, Quintana AM, Achilly NP, et al. An X-linked cobalamin disorder caused by mutations in transcriptional coregulator HCF1. *Am J Hum Genet*. 2013;93(3):506–14.
- Lin HJ, Neidich JA, Salazar D, Thomas-Johnson E, Ferreira BF, Kwong AM, et al. Asymptomatic maternal combined Homocystinuria and Methylmalonic Aciduria (cblC) detected through low Carnitine levels on newborn screening. *J Pediatr*. 2009;155(6):924–7.
- Lerner-Ellis JP, Anastasio N, Liu J, Coelho D, Suormala T, Stucki M, et al. Spectrum of mutations in MMACHC, allelic expression, and evidence for genotype–phenotype correlations. *Hum Mutat*. 2009;30(7):1072–81.
- Wolf NI, Bast T, Surtees R. Epilepsy in inborn errors of metabolism. *Epileptic Disord*. 2005;7(2):67–81.
- Chace DH, Kalas TA, Naylor EW. Use of tandem mass spectrometry for multianalyte screening of dried blood specimens from newborns. *Clin Chem*. 2003;49(11):1797–817.
- la Marca G, Malvagia S, Casetta B, Pasquini E, Donati MA, Zammarchi E. Progress in expanded newborn screening for metabolic conditions by LC-MS/MS in Tuscany: update on methods to reduce false tests. *J Inherit Metab Dis*. 2008;31:S395–404.
- Kimura M, Yamamoto T, Yamaguchi S. Automated metabolic profiling and interpretation of GC/MS data for organic acidemia screening: a personal computer-based system. *Tohoku J Exp Med*. 1999;188(4):317–34.
- X-w F, Iga M, Kimura M, Yamaguchi S. Simplified screening for organic acidemia using GC/MS and dried urine filter paper: a study on neonatal mass screening. *Early Hum Dev*. 2000;58(1):41–55.
- Baumgartner MR, Hörster F, Dionisi-Vici C, Halliloglu G, Karall D, Chapman KA, et al. Proposed guidelines for the diagnosis and management of methylmalonic and propionic acidemia. *Orphanet J Rare Dis*. 2014;9(1):130.
- Shigematsu Y, Hata I, Nakai A, Kikawa Y, Sudo M, Tanaka Y, et al. Prenatal diagnosis of organic acidemias based on amniotic fluid levels of acylcarnitines. *Pediatr Res*. 1996;39(4):680–4.
- Hasegawa Y, Iga M, Kimura M, Shigematsu Y, Yamaguchi S. Prenatal diagnosis for organic acid disorders using two mass spectrometric methods, gas chromatography mass spectrometry and tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2005;823(1):13–7.
- Guo KJ, Zhou X, Chen XG, Wu YL, Liu CX, Kong QS. Expanded newborn screening for inborn errors of metabolism and genetic characteristics in a Chinese population. *Front Genet*. 2018;9:7.
- Han B, Cao Z, Tian L, Zou H, Yang L, Zhu W, et al. Clinical presentation, gene analysis and outcomes in young patients with early-treated combined methylmalonic acidemia and homocystinemia (cblC type) in Shandong province, China. *Brain Dev*. 2016;38(5):491–7.
- Wang F, Han L, Yang Y, Gu X, Ye J, Qiu W, et al. Clinical, biochemical, and molecular analysis of combined methylmalonic acidemia and hyperhomocystinemia (cblC type) in China. *J Inherit Metab Dis*. 2010;33(Suppl 3):S435–42.
- Hu S, Mei S, Liu N, Kong X. Molecular genetic characterization of cblC defects in 126 pedigrees and prenatal genetic diagnosis of pedigrees with combined methylmalonic aciduria and homocystinuria. *BMC Med Genet*. 2018;19(1):154.
- Wang C, Li D, Cai F, Zhang X, Xu X, Liu X, et al. Mutation spectrum of MMACHC in Chinese pediatric patients with cobalamin C disease: a case series and literature review. *Eur J Med Genet*. 2019;62(10):103713.
- Worgan LC, Niles K, Tirone JC, Hofmann A, Verner A, Sammak A, et al. Spectrum of mutations in Mut methylmalonic acidemia and identification of a common hispanic mutation and haplotype. *Hum Mutat*. 2006;27(1):31–43.
- Han L, Wu S, Ye J, Qiu W, Zhang H, Gao X, et al. Biochemical, molecular and outcome analysis of eight chinese asymptomatic individuals with methyl malonic acidemia detected through newborn screening. *Am J Med Genet A*. 2015;167(10):2300–5.
- Almannai M, Marom R, Divin K, Scaglia F, Sutton VR, Craigen WJ, et al. Milder clinical and biochemical phenotypes associated with the c. 482G>A (p. Arg161Gln) pathogenic variant in cobalamin C disease: implications for management and screening. *Mol Genet Metab*. 2017;122(1–2):60–6.
- Figuera MR, Queiroz CM, Stracke MP, Brauer MCN, González-Rodríguez LL, Frussa-Filho R, et al. Ascorbic acid and  $\alpha$ -tocopherol attenuate methylmalonic acid-induced convulsions. *Neuroreport*. 1999;10(10):2039–43.
- deMello CF, Begnini J, JimenezBernal RE, Rubin MA, deBastiani J, daCosta EM, et al. Intrastriatal methylmalonic acid administration induces rotational behavior and convulsions through glutamatergic mechanisms. *Brain Res*. 1996;721(1–2):120–5.
- Kruman II, Culmsee C, Chan SL, Kruman Y, Guo Z, Penix L, et al. Homocysteine elicits a DNA damage response in neurons that promotes apoptosis and hypersensitivity to excitotoxicity. *J Neurosci*. 2000;20(18):6920–6.
- Huemer M, Diodato D, Schwahn B, Schiff M, Bandeira A, Benoist J-F, et al. Guidelines for diagnosis and management of the cobalamin-related remethylation disorders cblC, cblD, cblE, cblF, cblG, cblJ and MTHFR deficiency. *J Inherit Metab Dis*. 2017;40(1):21–48.
- Lipton SA, Kim WK, Choi YB, Kumar S, D'Emilia DM, Rayudu PV, et al. Neurotoxicity associated with dual actions of homocysteine at the N-methyl-D-aspartate receptor. *Proc Natl Acad Sci U S A*. 1997;94(11):5923–8.
- Papatheodorou L, Weiss N. Vascular oxidant stress and inflammation in hyperhomocystinemia. *Antioxid Redox Signal*. 2007;9(11):1941–58.
- De Almeida LMV, Funchal C, de Lima PP, Pessutto FDB, Loureiro SO, Vivian L, et al. Effect of propionic and methylmalonic acids on the in vitro



- phosphorylation of intermediate filaments from cerebral cortex of rats during development. *Metab Brain Dis.* 2003;18(3):207–19.
31. Debray FG, Boulanger Y, Khiat A, Decarie JC, Orquin J, Roy MS, et al. Reduced brain choline in homocystinuria due to remethylation defects. *Neurology.* 2008;71(1):44–9.
  32. Weisfeld-Adams JD, Bender HA, Miley-Akerstedt A, Frempong T, Schragr NL, Patel K, et al. Neurologic and neurodevelopmental phenotypes in young children with early-treated combined methylmalonic acidemia and homocystinuria, cobalamin C type. *Mol Genet Metab.* 2013;110(3):241–7.
  33. Greibe E, Nexø E. Forms and amounts of vitamin B12 in infant formula: a pilot study. *PLoS One.* 2016;11(11):e0165458.
  34. Gizicki R, Robert M-C, Gomez-Lopez L, Orquin J, Decarie J-C, Mitchell GA, et al. Long-term visual outcome of Methylmalonic Aciduria and Homocystinuria, Cobalamin C Type. *Ophthalmology.* 2014;121(1):381–6.
  35. Brooks BP, Thompson AH, Sloan JL, Manoli I, Carrillo-Carrasco N, Zein WM, et al. Ophthalmic manifestations and long-term visual outcomes in patients with Cobalamin C deficiency. *Ophthalmology.* 2016;123(3):571–82.
  36. Liu XQ, Yan H, Qiu JX, Zhang CY, Qi JG, Zhang X, et al. Pulmonary arterial hypertension as leading manifestation of methylmalonic aciduria: clinical characteristics and gene testing in 15 cases. *Beijing Da Xue Xue Bao.* 2017;49(5):768.
  37. Beck BB, van Spronsen F, Diepstra A, Berger RMF, Komhoff M. Renal thrombotic microangiopathy in patients with cblC defect: review of an under-recognized entity. *Pediatr Nephrol.* 2017;32(5):733–41.
  38. Huemer M, Scholl-Bürgi S, Hadaya K, Kern I, Beer R, Seppi K, et al. Three new cases of late-onset cblC defect and review of the literature illustrating when to consider inborn errors of metabolism beyond infancy. *Orphanet J Rare Dis.* 2014;9:161.
  39. Morel CF, Lerner-Ellis JP, Rosenblatt DS. Combined methylmalonic aciduria and homocystinuria (cblC): phenotype–genotype correlations and ethnic-specific observations. *Mol Genet Metab.* 2006;88(4):315–21.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

