

CORRECTION



Correction: The necroptosis machinery mediates axonal degeneration in a model of Parkinson disease

Maritza Oñate · Alejandra Catenaccio · Natalia Salvadores · Cristian Saquel · Alexis Martinez · Ines Moreno-Gonzalez · Nazaret Gamez · Paulina Soto · Claudio Soto · Claudio Hetz · Felipe A. Court

Published online: 5 February 2020

© The Author(s), under exclusive licence to ADMC Associazione Differenziamento e Morte Cellulare 2020

Correction to: Cell Death & Differentiation

<https://doi.org/10.1038/s41418-019-0408-4>

Since online publication of this article, the authors noticed that reference details provided for references 65 and 66 were incorrect. In addition, four references were erroneously missed out. The correct references are provided below. The references have been updated and renumbered in the PDF and HTML versions of the manuscript.

- (65) Dillon CP, Weinlich R, Rodriguez DA, Cripps JG, Quarato G, Gurung P, et al. RIPK1 blocks early postnatal lethality mediated by caspase-8 and RIPK3. *Cell*. 2014;5:1189–1202.
(66) Oberst A, Dillon CP, Weinlich R, McCormick LL, Fitzgerald P, Pop C, et al. Catalytic activity of the caspase-

8–FLIPL complex inhibits RIPK3-dependent necrosis. *Nature*. 2011;471:363–7.

- (67) Murphy JM, Czabotar PE, Hildebrand JM, Lucet IS, Zhang JG, Alvarez-Diaz S, et al. The pseudokinase MLKL mediates necroptosis via a molecular switch mechanism. *Immunity*. 2013;39:443–53.
(68) Newton K, Sun X, Dixit VM. Kinase RIP3 is dispensable for normal NF- κ Bs, signaling by the B-cell and T-cell receptors, tumor necrosis factor receptor 1, and Toll-like receptors 2 and 4. *Mol Cell Biol*. 2004;24:1464–9.
(69) Paxinos G, Franklin K. The mouse brain in stereotaxic coordinates. 3rd ed. Amsterdam: Academic Press. 2008. p. 256.
(70) Castillo V, Oñate M, Woehlbier U, Rozas P, Andreu C, Medinas D, et al. Functional role of the disulfide isomerase ERp57 in axonal regeneration. *PLoS ONE*.