



# HHS Public Access

Author manuscript

*Am J Med.* Author manuscript; available in PMC 2020 August 10.

Published in final edited form as:

*Am J Med.* 2020 August ; 133(8): 901–907. doi:10.1016/j.amjmed.2020.03.030.

## Intermittent Fasting: A Heart Healthy Dietary Pattern?

Tiffany A. DONG, MD<sup>1</sup>, Pratik B. SANDESARA, MD<sup>2</sup>, Devinder S. DHINDSA, MD<sup>2</sup>, Anurag MEHTA, MD<sup>2</sup>, Laura C. ARNESON, MD<sup>3</sup>, Allen L. DOLLAR, MD<sup>4</sup>, Pam R. TAUB, MD<sup>5</sup>, Laurence S. SPERLING, MD<sup>6</sup>

<sup>1</sup>Department of Medicine, Emory University School of Medicine Atlanta, Georgia, USA; Emory Clinical Cardiovascular Research Institute, Emory University School of Medicine Atlanta, Georgia, USA

<sup>2</sup>Emory Clinical Cardiovascular Research Institute, Emory University School of Medicine Atlanta, Georgia, USA; Division of Cardiology, Emory University, Atlanta, GA, USA

<sup>3</sup>Department of Medicine at Northwestern University Feinberg School of Medicine, Chicago, IL, USA

<sup>4</sup>Division of Cardiology, Emory University, Atlanta, GA, USA

<sup>5</sup>Division of Cardiovascular Medicine, UC San Diego School of Medicine, La Jolla, CA USA

<sup>6</sup>Emory Clinical Cardiovascular Research Institute, Emory University School of Medicine Atlanta, Georgia, USA; Division of Cardiology, Emory University, Atlanta, GA, USA; Emory School of Medicine, Atlanta, GA, USA

### Abstract

Dietary patterns, such as the Dietary Approaches to Stop Hypertension and the Mediterranean diet, have been shown to improve cardiac health. Intermittent fasting is another type of popular dietary pattern that is based upon timed periods of fasting. Two different regimens are Alternative Day Fasting and Time-Restricted Eating. Although there are no large, randomized control trials examining the relationship between intermittent fasting and cardiovascular outcomes, current human studies that suggest this diet could reduce the risk for cardiovascular disease with improvement in weight control, hypertension, dyslipidemia and diabetes. Intermittent fasting may exert its effects through multiple pathways including reducing oxidative stress, optimization of circadian rhythms and ketogenesis. This review evaluates current literature regarding the potential cardiovascular benefits of intermittent fasting and proposes directions for future research.

### Keywords

Alternative Day Fast; Time Restricted Feeding; Circadian; Diabetes; Hypertension; Dyslipidemia

---

Electronic address: lsperli@emory.edu.

All authors were involved in the writing of the manuscript.

**Disclosures:** Pam Taub: No relevant financial disclosures related to this topic. General disclosures: Consultant/Speaker for Sanofi/Regeneron, Novo-Nordisk, Boehringer-Ingelheim, Janssen, Pfizer, Amarin and Amgen. She is a shareholder of Cardero Therapeutics. The remaining authors have nothing to disclose.

## Introduction

Although cardiovascular mortality rates have improved, the decline in mortality has recently ceased and there has been an increase in mortality in 35–64 year old males and females in the United States.<sup>1</sup> Obesity along with poor diet are important, modifiable contributors to the rise of cardiovascular disease with an estimated attributable risk of 13% to cardiovascular mortality.<sup>2</sup> There are several dietary interventions that have been shown to improve cardiovascular risk including caloric restriction, which involves limiting calories consumed during a given period. Caloric restriction is linked to improvement in weight, blood pressure and insulin sensitivity in humans.<sup>3</sup>

Intermittent fasting is a dietary intervention similar to caloric restriction, as it utilizes the principle of restricting food intake. However, intermittent fasting focuses on the timing of when one can consume meals either within a day or a week. Two overarching types of intermittent fasting are alternative day fasting and time-restricted fasting. In alternative day fasting, subset may consist of 24-hour fasts followed by a 24-hour eating period that can be done several times a week such as a 5:2 strategy when there are 2 fast days mixed into 5 nonrestrictive days. For time restricted fast programs, variations include 16-hour fasts with 8hour feeding times, 20-hour fasts with 4-hour feed times or other similar versions. While both caloric restriction and intermittent fasting may result in overall decreased caloric intake, this is not integral to intermittent fasting. Intermittent fasting has been linked to better glucose control in both humans and animals.<sup>4,5</sup> However, long-term adherence to caloric restriction is low while adherence intermittent fasting may be more promising.

Given the similarity between these two diets, it is plausible that intermittent fasting could also confer cardiovascular benefits. This dietary pattern has also shown potential benefit in slowing the progression of neurodegenerative diseases like Alzheimer's and Parkinson's.<sup>6</sup> In this review, we explore the potential benefits of intermittent fasting for improving cardiovascular health.

## Mechanisms

There are several proposed mechanisms for how intermittent fasting could lead to better cardiovascular outcomes (Figure 1). The Oxidative Stress Hypothesis supports decreased oxidative insult.<sup>7</sup> A second theory, the circadian rhythm hypothesis, is associated more with intermittent fasting than caloric restriction, indicating a mechanism unique to intermittent fasting. A third theory involves intermittent fasting inducing a ketogenic state, which has been linked to decreases in cardiovascular risk factors.

## Oxidative Stress Hypothesis

The Oxidative Stress Hypothesis states that decrease energy intake cause mitochondria to produce fewer free radicals.<sup>7</sup> After 8 weeks of alternative day fasting, obese patients with asthma showed lower levels of inflammation such as tumor necrosis factor-alpha and brain-derived neutrophilic factors as well as oxidative stress including nitrotyrosine, 8-isoprostane, protein carbonyls and 4-hydroxynoneal adducts. Moreover, they had higher levels of the antioxidant uric acid.<sup>8</sup>

## Circadian Rhythm Theory

The circadian rhythm theory assumes that physiologic processes occur at the most advantageous time as dictated by evolution.<sup>9</sup> Fasting properly may allow optimization with our organs' peripheral clocks such as those in the liver, adipose and skeletal tissues. Dysregulation of this system increases the risk for chronic diseases, as evidenced by higher rate of cardiometabolic diseases in shift workers.<sup>10</sup> One circadian example relevant to intermittent fasting is decreasing insulin levels later in the day.<sup>11</sup> Late dinners are associated with higher postprandial glucose levels than daytime meals, increasing the risk of diabetes. In humans, circadian misalignment increases insulin resistance after only 3 days.<sup>10</sup> Nighttime eating decrease both quality and quantity of sleep, which also leads to increased insulin resistance, obesity and cardiovascular disease.<sup>12,13</sup> Different time restricted fast regimens have demonstrated variable results based on the timing of the fast, which emphasizes the role of circadian system in this dietary pattern. Subjects who were allowed to eat during the middle of the day had better weight loss with less adipose, glucose control, lipid levels and inflammation.<sup>14</sup> In contrast, those on a time restricted fast regimen that allowed late afternoon or evening intake defined as beyond 16:00 had no improvement and even worsening of glucose control, blood pressure and lipid levels.<sup>15,16</sup> Thus, intermittent fasting when timed properly, may sync with one's circadian rhythm and thus improve cardiac health.

## Ketogenic State

Intermittent fasting induces a ketogenic state, as evidenced by the rise in  $\beta$ -hydroxybutyrate levels in overweight individual who fast.<sup>8</sup> After 6–8 hours of fasting, ketone levels become detectable, which signals a switch from fat storage to fat utilization with decrease in low-density lipoproteins (LDL) and increase in high-density lipoproteins (HDL) levels.<sup>17,18</sup> This change from using glucose as energy to using fatty acids and ketones for energy is called intermittent metabolic switching. Furthermore, the ketogenic diet promotes weight loss, as processing ketones requires greater energy.<sup>19</sup> Intermittent fasting contains elements of the ketogenic diet, benefitting from increased adipose metabolism leading to improvement in weight and lipids. Importantly, intermittent fasting may be more beneficial than the ketogenic diet, as the latter involves high consumption of animal fats. Excessive fat intake can be detrimental since it is associated with higher levels of trimethylamine N-oxide, a metabolite associated with increased cardiovascular risk that has been found to be higher in a ketogenic diet.<sup>20</sup>

## The Effect of Intermittent Fasting on Cardiovascular Risk Factors (Table 1)

### Obesity

In a study of overweight men with type II diabetes, subjects in both caloric restriction and intermittent fasting regimens experienced weight loss with intermittent fasting subjects losing 1.1% of body fat with a mean 6.5% weight loss after 12 weeks.<sup>21</sup> Similar findings were observed in both overweight and obese premenopausal females who were randomized to intermittent fasting and caloric restriction for 6 months. The intermittent fasting and caloric restriction groups had comparable results with the intermittent fasting group losing

6.4 kilograms (95% CI 4.8 to 7.9 kg) and the caloric restriction group losing 5.6 kilograms (95% CI 4.4 to 6.9 kg).<sup>22</sup> A study with 16 nonobese men and women who underwent alternative day fasting for 22 days did lose  $2.5 \pm 0.5\%$  of initial body weight ( $p < 0.001$ ) and  $4 \pm 1\%$  of their fat mass ( $p < 0.001$ ).<sup>23</sup> Interestingly, obese subjects who were randomized to fasting except for lunch or dinner for 8 weeks had similar weight loss with the lunch group losing  $3.5 \pm 0.4$  kg ( $p < 0.001$ ) and the dinner group  $4.1 \pm 0.5$  kg ( $p < 0.001$ ).<sup>24</sup> Despite not adhering to the circadian rhythm, weight loss likely occurred due to limited calorie consumption. The change in weight may also be related to the utilization of fatty acids for energy consistent with a ketogenic state.

### Blood Pressure

Human studies have shown reductions in both systolic and diastolic blood pressure with intermittent fasting. A small study of men with prediabetes had an average reduction of systolic blood pressure of  $11 \pm 4$  mmHg and a diastolic blood pressure reduction of  $10 \pm 4$  mmHg after 5-weeks of fasting for 18-hour periods.<sup>25</sup> Similarly, a prospective observational study of 82 Muslims who celebrated Ramadan, a month-long religious holiday involving daytime fasting, showed a 3-point reduction in systolic blood pressure although diastolic change was not significant.<sup>26</sup> One potential explanation for this is a decrease in sympathetic tone and increase in parasympathetic tone. Using power spectral analysis of heart rate and arterial pressure, rats placed on intermittent fasting have a lower frequency component in diastolic blood pressure variability, a marker for sympathetic tone. Additionally, these rats have a the higher frequency component of the heart rate variability spectra, a marker for parasympathetic tone.<sup>4</sup> Higher vagal activity has been associated with decreased levels of inflammatory cytokines including tumor necrosis factor alpha, interleukin-1b, interleukin-6 and interleukin-8, which are implicated in the pathogenesis of atherosclerosis.<sup>27</sup> Thus, intermittent fasting's appears to have the ability to lower blood pressure, which thus could improve CVD mortality.

### Dyslipidemia

In addition to blood pressure, intermittent fasting seems to have a positive impact on lipid values. In a study with 60 overweight and obese adults, the alternative day fasting group who underwent a 75% caloric restriction every other day had a reduction in LDL by  $10 \pm 4\%$  and reduction in triglyceride  $17 \pm 5\%$  after 12-weeks.<sup>28</sup> However, these changes could be explained by weight loss observed. Muslims celebrating Ramadan had better HDL, LDL, triglycerides and very low-density lipoprotein levels resulting in an decrease in average Framingham risk score 13.8 to 10.8.<sup>26</sup> Similar to this study, another study with 83 obese participants also showed improvement in HDL and LDL after 12 weeks of alternative day fasting combined with exercise.<sup>29</sup> It is unclear why discrepancies among these studies regarding HDL exist. One explanation could be the difference in timing in relation to circadian rhythms as the first study involved fasting every other day while Ramadan involves fasting from sunrise to sundown. In mice models, intermittent fasting does appear to be more beneficial when food intake occurs during times of activity as compared to more dormant times, as measured by hepatic production of circadian genes such as mPER and mClock.<sup>30</sup>

## Diabetes

In contrast to blood pressure and dyslipidemia, the relationship between intermittent fasting and diabetes is not as straightforward. Nonobese males who fasted for 20-hour intervals then ate without restriction on alternative days showed increased insulin-mediated glucose uptake after two weeks.<sup>31</sup> Another small study of men with prediabetes had better insulin sensitivity and increased beta cell responsiveness, as measured by a higher insulinogenic index calculated by the change in insulin divided by change in glucose within the first thirty minutes of an oral glucose tolerance test.<sup>25</sup> However, the data does not consistently support improvements in fasting glucose levels, which may be due to a latency period or missed glucose fluctuations during the day.<sup>25,31</sup> In a study with obese individuals on alternative day fasting, insulin sensitivity did not appear to change after the 8-week intervention.<sup>5</sup> The difference in the outcomes of these studies may be explained by the different populations studied (obese versus non-obese), suggesting distinct impact based on subgroups.

## The Effect of Intermittent Fasting after a Cardiovascular Event

Even after a cardiovascular event, intermittent fasting may confer cardiac protection. In observational studies, Muslims with a history of ischemic cardiomyopathy have a decreased incidence of acute decompensated heart failure during Ramadan compared to other parts of the year.<sup>32</sup> The Intermountain Heart Collaborative Study Group performed a meta-analysis of two studies involving Latter Day Saints that combined about 648 patients. They compared the incidence of coronary heart disease defined as at least one coronary artery with 70% stenosis in those who underwent a monthly 1-day religious fast to those who did not. The subjects who followed the fast had a lower risk for coronary heart disease with odds ratio 0.65 (CI 0.460.94).<sup>33</sup> Although human data is sparse regarding intermittent fasting after a cardiovascular event, these observational studies suggest a positive impact.

## Intermittent Fasting and Longevity

Currently, there are no randomized controlled trials with humans regarding longevity and intermittent fasting. On the cellular level, human skin fibroblasts in vitro conditions simulating intermittent fasting had longer lifespans than controls. In addition, this group retained their youthful morphology while the controls developed a senescent morphology, which is associated with a smaller, thinner appearance. Thus, this ex vivo study in human fibroblasts suggests that intermittent fasting could delay aging at the cellular level.<sup>34</sup>

## Intermittent Fasting versus Caloric Restriction (Figure 2)

While intermittent fasting and caloric restriction are similar, it is important to make the distinction between these two dietary patterns, as they may lead to different biologic outcomes. One important distinction is that intermittent fasting does not necessarily involve limiting calories as caloric restriction does. In humans, the impact of intermittent fasting on cardiovascular risk factors (i.e. blood pressure, fasting glucose, and lipid profile) can be still seen during Ramadan without decreasing caloric intake. On average, they have lower blood pressures during this month.<sup>26</sup> In obese adults, intermittent fasting and caloric restriction

appear to have similar effects on improving lipid panels while alternative day fasting groups had a significantly better impact on fasting glucose.<sup>5</sup>

In terms of practicality, it may be easier for individual to adhere to intermittent fasting rather than caloric restriction, as caloric restriction has poor long-term compliance rates with one study citing a dropout rate of 21% after 2 months and a 42% dropout after 1-year.<sup>35, 36, 37</sup> Thus far, long-term trials of 1 year show either similar or worse compliance rates with intermittent fasting compared to CF.<sup>38,39</sup> However, both these trials involved alternative day fasting regimens with 2 days of fasting interspersed within 1 week. Time restricted fast regimens such as 16-hours of fasting and 8-hours of eating may have better adherence rates when compared to caloric restriction.

Although intermittent fasting is distinct from caloric restriction, this type of dietary regimen may also lead to better cardiovascular outcomes since the literature demonstrates fasting improves various cardiovascular risk factors such as diabetes, hypertension and cholesterol. Despite the limited number of studies, the ability of intermittent fasting to improve cardiac health appears promising.

## Future Directions and Limitations

More studies are needed to evaluate mechanisms, efficacy in humans, target populations and safety of intermittent fasting. There are numerous intermittent fasting regimens ranging from 12 to 16-hour daily fasts to 5:2 strategy and it remains uncertain which strategy is the best for cardiovascular health especially with evidence suggesting that intermittent fasting regimens should follow circadian rhythms.<sup>14</sup> Certain regimens may be easier to adhere to than others. Future studies should also investigate the safety of each intermittent fasting strategy as well.

In addition to finding effective intermittent fasting regimens tailored to different patient populations, future studies should establish the duration of intermittent fasting needed before cardiovascular benefits occurred. For humans, benefits appeared within a month during observational Ramadan studies. Extending intermittent fasting for longer may lead to continued improvement in both cardiovascular risk factors and events. Furthermore, it should be established if these benefits extend beyond the duration of intermittent fasting. Rats that were placed initially on intermittent fasting with subsequent improvement of blood pressure had reversal of this improvement 3–4 weeks after returning to an ad libitum diet.<sup>4</sup> However, for obese adults who were on intermittent fasting for 8 weeks then transitioned back to their regular diet for 24 weeks still maintained their lower cholesterol and glucose levels.<sup>5</sup> While intermittent fasting was effective for this population of obese patients, it is unclear if intermittent fasting affects all populations similarly.

## Conclusions

Human studies show promise for cardiovascular benefit in intermittent fasting. Although the exact mechanisms remain to be elucidated, intermittent fasting appears to positively impact multiple cardiovascular risk factors including obesity, hypertension, dyslipidemia, and diabetes. Furthermore, intermittent fasting has been associated with improved outcome after

a cardiac event. These results should encourage future studies to optimize intermittent fasting's potential to improve cardiovascular outcomes.

## References

1. Ritchey MD, et al. Million Hearts: Description of the National Surveillance and Modeling Methodology Used to Monitor the Number of Cardiovascular Events Prevented During 2012–2016 *J Am Heart Assoc*, 6 (2017) Google Scholar
2. Go AS, et al. Executive summary: heart disease and stroke statistics-2014 update: a report from the American Heart Association *Circulation*, 129 (2014), pp. 399–410 CrossRef View Record in Scopus Google Scholar [PubMed: 24446411]
3. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults–The Evidence Report National Institutes of Health *Obes Res*, 6 (Suppl 2) (1998), pp. 51s–209s Google Scholar [PubMed: 9813653]
4. Mager DE, et al. Caloric restriction and intermittent fasting alter spectral measures of heart rate and blood pressure variability in rats *Faseb j*, 20 (2006), pp. 631–637 CrossRef View Record in Scopus Google Scholar [PubMed: 16581971]
5. Catenacci VA, et al. A randomized pilot study comparing zero-calorie alternate-day fasting to daily caloric restriction in adults with obesity *Obesity (Silver Spring)*, 24 (2016), pp. 1874–1883 CrossRef View Record in Scopus Google Scholar [PubMed: 27569118]
6. Martin B, Mattson MP, Maudsley S Caloric restriction and intermittent fasting: two potential diets for successful brain aging *Ageing Res Rev*, 5 (2006), pp. 332–353 Article Download PDF View Record in Scopus Google Scholar [PubMed: 16899414]
7. Merry BJ Oxidative stress and mitochondrial function with aging—the effects of calorie restriction *Ageing Cell*, 3 (2004), pp. 7–12 CrossRef View Record in Scopus Google Scholar [PubMed: 14965349]
8. Johnson JB, et al. Alternate day calorie restriction improves clinical findings and reduces markers of oxidative stress and inflammation in overweight adults with moderate asthma *Free Radic Biol Med*, 42 (2007), pp. 665–674 Article Download PDF View Record in Scopus Google Scholar [PubMed: 17291990]
9. Panda S, Hogenesch JB, Kay SA Circadian rhythms from flies to human *Nature*, 417 (2002), pp. 329–335 View Record in Scopus Google Scholar [PubMed: 12015613]
10. Scheer FA, Hilton MF, Mantzoros CS, Shea SA Adverse metabolic and cardiovascular consequences of circadian misalignment *Proc Natl Acad Sci U S A*, 106 (2009), pp. 4453–4458 CrossRef View Record in Scopus Google Scholar [PubMed: 19255424]
11. Morgan L, Hampton S, Gibbs M, Arendt J Circadian aspects of postprandial metabolism *Chronobiol Int*, 20 (2003), pp. 795–808 View Record in Scopus Google Scholar [PubMed: 14535354]
12. Buxton OM, Marcelli E Short and long sleep are positively associated with obesity, diabetes, hypertension, and cardiovascular disease among adults in the United States *Soc Sci Med*, 71 (2010), pp. 1027–1036 Article Download PDF View Record in Scopus Google Scholar [PubMed: 20621406]
13. Spiegel K, Knutson K, Leproult R, Tasali E, VanCauter E Sleep loss: a novel risk factor for insulin resistance and Type 2 diabetes *J Appl Physiol* (1985), 99 (2005), pp. 2008–2019 CrossRef View Record in Scopus Google Scholar
14. Moro T, et al. Effects of eight weeks of time-restricted feeding (16/8) on basal metabolism, maximal strength, body composition, inflammation, and cardiovascular risk factors in resistance-trained males *J Transl Med*, 14 (2016), p. 290 Google Scholar [PubMed: 27737674]
15. Carlson O, et al. Impact of reduced meal frequency without caloric restriction on glucose regulation in healthy, normal-weight middle-aged men and women *Metabolism*, 56 (2007), pp. 1729–1734 Article Download PDF View Record in Scopus Google Scholar [PubMed: 17998028]
16. Stole KS, et al. A controlled trial of reduced meal frequency without caloric restriction in healthy, normal-weight, middle-aged adults *Am J Clin Nutr*, 85 (2007), pp. 981–988 View Record in Scopus Google Scholar [PubMed: 17413096]


17. Anton SD, et al. Flipping the Metabolic Switch: Understanding and Applying the Health Benefits of Fasting Obesity (Silver Spring), 26 (2018), pp. 254–268 CrossRef View Record:n Scopus Google Scholar [PubMed: 29086496]
18. Dashti HM, et al. Long term effects of ketogenic diet in obese subjects with high cholesterol level Mol Cell Biochem, 286 (2006), pp. 1–9 CrossRef View Record n Scopus Google Scholar [PubMed: 16652223]
19. Jornayvaz FR, et al. A high-fat, ketogenic diet causes hepatic insulin resistance in mice, despite increasing energy expenditure and preventing weight gain Am J Physiol Endocrinol Metab, 299 (2010), pp. E808–E815 CrossRef View Record in Scopus Google Scholar [PubMed: 20807839]
20. Park JE, Miller M, Rhyne J, Wang Z, Hazen SL Differential effect of short-term popular diets on TMAO and other cardio-metabolic risk markers Nutr Metab Cardiovasc Dis (2019) Google Scholar
21. Ash S, et al. Effect of intensive dietetic interventions on weight and glycaemic control in overweight men with Type II diabetes: a randomised trial Int J Obes Relat Metab Disord, 27 (2003), pp. 797–802 CrossRef View Record n Scopus Google Scholar [PubMed: 12821964]
22. Harvie MN, et al. The effects of intermittent or continuous energy restriction on weight loss and metabolic disease risk markers: a randomized trial in young overweight women Int J Obes (Lond), 35 (2011), pp. 714–727 CrossRef View Record in Scopus Google Scholar [PubMed: 20921964]
23. Heilbronn LK,Smith SR,Martin CK,Anton SD,Ravussin E Alternate-day fasting in nonobese subjects: effects on body weight, body composition, and energy metabolism Am J Clin Nutr, 81 (2005), pp. 69–73 View Record in Scopus Google Scholar [PubMed: 15640462]
24. Hoddy KK,et al. Meal timing during alternate day fasting: Impact on body weight and cardiovascular disease risk in obese adults Obesity (Silver Spring), 22 (2014), pp. 2524–2531 View Record in Scopus Google Scholar [PubMed: 25251676]
25. Sutton EF,et al. Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes Cell Metab, 27 (2018), pp. 1212–1221 View Record in Scopus Google Scholar [PubMed: 29754952]
26. Nematy M,et al. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study Nutr J, 11 (2012), p. 69 Google Scholar [PubMed: 22963582]
27. Chandrasekar B,Nelson JF,Colston JT,Freeman GL Calorie restriction attenuates inflammatory responses to myocardial ischemiareperfusion injury Am J Physiol Heart Circ Physiol, 280 (2001), pp. H2094–H2102 CrossRef View Record in Scopus Google Scholar [PubMed: 11299211]
28. Varady KA,Bhutani S,Klempel MC,Kroeger CM Comparison of effects of diet versus exercise weight loss regimens on LDL and HDL particle size in obese adults Lipids Health Dis, 10 (2011), p. 119 CrossRef Google Scholar [PubMed: 21767400]
29. Bhutani S,Klempel MC,Kroeger CM,Trepanowski JF,Varady KA Alternate day fasting and endurance exercise combine to reduce body weight and favorably alter plasma lipids in obese humans Obesity (Silver Spring), 21 (2013), pp. 1370–1379 CrossRef View Record in Scopus Google Scholar [PubMed: 23408502]
30. Froy O,Chapnik N,Miskin R Effect of intermittent fasting on circadian rhythms in mice depends on feeding time Mech Ageing Dev, 130 (2009), pp. 154–160 Article DownloadPDF View RecordinScopus Google Scholar [PubMed: 19041664]
31. Halberg N,et al. Effect of intermittent fasting and refeeding on insulin action in healthy men J Appl Physiol (1985), 99 (2005), pp. 2128–2136 CrossRef View Record in Scopus Google Scholar
32. Salim I,Al Suwaidi J,Ghadban W,Alkilani H,Salam AM Impact of religious Ramadan fasting on cardiovascular disease: a systematic review of the literature Curr Med Res Opin, 29 (2013), pp. 343–354 CrossRef View Record in Scopus Google Scholar [PubMed: 23391328]
33. Horne BD,et al. Relation of routine, periodic fasting to risk of diabetes mellitus, and coronary artery disease in patients undergoing coronary angiography Am J Cardiol, 109 (2012), pp. 1558–1562 Article DownloadPDF View RecordinScopus Google Scholar [PubMed: 22425331]
34. Sodagam L,Lewinska A,Wnuk M,Rattan SIS Chronic exposure to rapamycin and episodic serum starvation modulate ageing of human fibroblasts in vitro Biogerontology, 18 (2017), pp. 841–854 CrossRef View Record in Scopus Google Scholar [PubMed: 28884409]



35. Blomain ES,Dirhan DA,Valentino MA,Kim GW,Waldman SA Mechanisms of Weight Regain following Weight Loss *ISRN Obes*, 2013 (2013), Article 210524 Google Scholar
36. Dansinger ML,Gleason JA,Griffith JL,Selker HP,Schaefer EJ Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial *Jama*, 293 (2005), pp. 43–53 CrossRef View Record in Scopus Google Scholar [PubMed: 15632335]
37. Sandesara PB,Sperling LS Caloric Restriction as a Therapeutic Approach to Heart Failure: Can Less Be More in (Mice) and Men? *Circ Heart Fail*, 11 (2018), Article e004930 View Record in Scopus Google Scholar
38. Carter S,Clifton PM,Keogh JB Effect of Intermittent Compared With Continuous Energy Restricted Diet on Glycemic Control in Patients With Type 2 Diabetes: A Randomized Noninferiority Trial *JAMA Netw Open*, 1 (2018), Article e180756 CrossRef View Record in Scopus Google Scholar
39. Trepanowski JF,et al. Effect of Alternate-Day Fasting on Weight Loss, Weight Maintenance, and Cardioprotection Among Metabolically Healthy Obese Adults: A Randomized Clinical Trial *JAMA Intern Med*, 177 (2017), pp. 930–938 CrossRef View Record in Scopus Google Scholar [PubMed: 28459931]

### Clinical Significance


- Modifying one's dietary pattern can lead to better cardiovascular outcomes
- Intermittent Fasting may benefit cardiovascular health by improving obesity, hypertension, dyslipidemia and diabetes.
- Potential mechanisms of this diet involve reducing oxidative stress, syncing with the circadian system and inducing ketogenesis.



HDL = high-density lipoprotein, LDL = low-density lipoprotein

#### Figure 1. Mechanisms of Intermittent Fasting

Proposed mechanisms of how intermittent fasting reduces cardiovascular risk factors. There are three main theories: Oxidative Stress Hypothesis, Circadian Rhythm and Ketogenic State. The Oxidative Stress Hypothesis postulates that fasting reduces stress leading to fewer free radical with less mitochondrial energy production ultimately lowering the body's oxidative stress. The Circadian Rhythm component focuses on syncing eating periods to the organ's circadian rhythm, optimizing glucose and fat utilization. The third mechanism, Ketogenic State, recognizes that Intermittent Fasting induces ketogenesis, which decreases blood pressure and adipose tissue.



**Figure 2. Similarities and differences between intermittent fasting and caloric restriction**  
Intermittent Fasting shares commonalities as well as differences with Caloric Restriction. Both have been shown to reduce cardiovascular risk factors including improving blood pressure, insulin sensitivity and dyslipidemia. In addition, it shares the common pathway of reducing stress response. Intermittent fasting revolves around defined periods of fasting syncing with the circadian rhythm while caloric restriction focuses on restricting overall calories. Thus far, intermittent fasting appears promising for overweight and obese individual and it remains to be seen if adherence is easier with intermittent fasting regimens.

**Table 1.**

## Effects of Intermittent Fasting on Cardiovascular Risk Factors

Study Title	Duration (weeks)	Subjects	Intervention	Results
<i>Obesity</i>				
Ash (2003)	12	n=51, M only Mean age 54yo Overweight DM2	ADF	Mean weight loss of 6.4±4.6kg Reduction in waist circumference 8.1±4.6cm Loss of body fat 1.9±1.5%
Harvie (2011)	24	n=107, F only Mean age 40yo overweight and obese premenopausal	2 day a week fast (75% caloric restriction)	Lost 6.4 kg (CI 4.8 to 7.8)
Heilbronn (2005)	3	n=16, M % F Male mean age 34yo; Female Mean age 30yo nonobese	ADF	Decrease in body fat 2.5±0.5% of initial body weight 4±1% of their fat mass (p<0.001)
Hoddy (2014)	8	n=74, M and F Mean age 45yo obese	ADF for either lunch, dinner or small meals	ADF-lunch 3.5±0.4 kg ADF-dinner for 4.1 ± 0.5 kg ADF-small meals 4.0 ± 0.5 kg
Wilkinson (2019)	12	n=19, M and F Mean age 59yo Metabolic Syndrome	10-hour fast	Weight reduction 3.3±3.2kg BMI reduction 1.1±0.97kg/m <sup>2</sup> Waist Circumference 4.5±6.7cm
<i>Hypertension</i>				
Eshghina (2013)	6	n=15, F only Mean age 34 Overweight and obese	3 days of a week fast (75% caloric restriction)	SBP ↓ 115±9mmHg to 105±10mmHg DBP ↓ 83±11mmHg to 75±11mmHg)
Nematy (2012)	4	n=82, M and F Mean age 54yo 1 Cardiovascular Risk Factor	Ramadan	SBP ↓ 133±6mmHg to 130±7mmHg NS DBP
Sutton (2018)	5	n=8, M only Mean age 56yo Prediabetic	18-hour daily fasts	SBP ↓ 11 ± 4mmHg DBP ↓ 10 ± 4mmHg
Wilkinson (2019)	12	n=19, M and F Mean age 59yo Metabolic Syndrome	10-hour fast	SBP ↓ 5±10mmHg DBP ↓ 7±8 mmHg
<i>Dyslipidemia</i>				
Bhutani (2013)	12	n=83, M and F Mean age 42yo obese	ADF (75% caloric reduction) ADF combined with exercise	LDL ↓ 12±5% NS TG HDL ↑ 18±9%
Nematy (2012)	4	n=82, M and F Mean age 54yo 1 Cardiovascular Risk Factor	Ramadan	LDL ↓ 13 (110±46 to 97±35) TG ↓ 41 (225±129 to 183±112) HDL ↑ 4 (43±9 to 48±8)
Varady (2011)	12	n=60, M and F Mean age 47yo overweight and obese	ADF (75% caloric reduction)	LDL ↓ 10± 4% TG ↓ 17± 5% HDL ↑ 16±5%
<i>Diabetes Mellitus</i>				
Bhutani (2013)	12	n=83, M and F Mean age 42yo obese	ADF (75% caloric reduction) ADF combined with exercise	NS fasting glucose NS insulin
Catenacci (2016)	8	n=14, M and F Mean age 40yo obese	ADF (100% calorie reduction)	Fasting glucose ↓6.0±2mg/dL NS insulin
Klempel (2012)	8	n=54, F only Mean age 48yo Prediabetic, obese	Total Fast 24-hour then 6 days of 70% liquid intake Total Fast 24-hour then 6 day of 70% food intake	Insulin ↓ 3.0±3.0uIU/ml Glucose ↓ 4.0±3.0mg/dL
Sutton (2018)	5	n=8, M only Mean age 56yo Prediabetic	18-hour daily fasts	Fasting insulin ↓3.4±1.6mU/L Insulinogenic Index ↑14±7U/mg

ADF=Alternative Day Fast, DM 2=Diabetes Mellitus type II, DBP=Diastolic Blood Pressure, HDL=High Density Lipoprotein, F=Female, LDL=Low Density Lipoprotein, M=Male, NS=Not Significant, SBP=Systolic Blood Pressure, TG=Triglyceride

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript