



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Perspective

Approaches and Challenges in SARS-CoV-2 Vaccine Development

Gabriel Dagotto,^{1,2,5} Jingyou Yu,^{1,2,5} and Dan H. Barouch^{1,2,3,4,*}¹Center for Virology and Vaccine Research, Beth Israel Deaconess Medical Center, Boston, MA 02215, USA²Harvard Medical School, Boston, MA 02115, USA³Ragon Institute of MGH, MIT, and Harvard, Cambridge, MA 02139, USA⁴Massachusetts Consortium on Pathogen Readiness, Boston, MA 02215, USA⁵These authors contributed equally to this work*Correspondence: dbarouch@bidmc.harvard.edu<https://doi.org/10.1016/j.chom.2020.08.002>

SUMMARY

The explosive spread of SARS-CoV-2 suggests that a vaccine will be required to end this global pandemic. Progress in SARS-CoV-2 vaccine development to date has been faster than for any other pathogen in history. Multiple SARS-CoV-2 vaccine candidates have been evaluated in preclinical models and are currently in clinical trials. In this Perspective, we discuss three topics that are critical for SARS-CoV-2 vaccine development: antigen selection and engineering, preclinical challenge studies in non-human primate models, and immune correlates of protection.

INTRODUCTION

SARS-CoV-2 was initially identified as the etiologic agent responsible for a cluster of severe pneumonia cases in Wuhan, China in December 2019 (Chan et al., 2020; Huang et al., 2020; Li et al., 2020; Wu et al., 2020; Zhu et al., 2020c). On March 11, 2020, the World Health Organization (WHO) declared SARS-CoV-2 a global pandemic, and the disease was named COVID-19. In response to this pandemic, industry, government, philanthropy, non-governmental organizations, and academia are collaborating to develop therapeutic and prophylactic countermeasures, including vaccines.

SARS-CoV-2 is a member of the *Coronaviridae* family, joining severe acute respiratory syndrome-associated coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus (MERS-CoV) as coronaviruses causing severe disease in humans. These pathogens are single-stranded, positive-sense RNA viruses and encode four main structural proteins, spike (S), envelope (E), membrane (M), and nucleocapsid (N), as well as multiple non-structural proteins (Srinivasan et al., 2020). Similar to SARS-CoV, the SARS-CoV-2 S protein binds angiotensin-converting enzyme 2 (ACE2) as its primary host receptor to mediate viral entry (Hoffmann et al., 2020), and S is the main target for neutralizing antibodies.

SARS-CoV-2 has proven to be highly transmissible, including from asymptomatic and presymptomatic individuals (Arons et al., 2020; McMichael et al., 2020). Currently, at least six vaccine candidates have been tested in non-human primates (NHPs) and have reported either partial or complete protection (Corbett et al., 2020; Gao et al., 2020; Mercado et al., 2020; van Doremalen et al., 2020; Wang et al., 2020; Yu et al., 2020). Moreover, multiple promising vaccine candidates have entered clinical trials, and early phase clinical trial data have also been reported (Folegatti et al., 2020; Jackson et al., 2020; Mulligan et al.,

2020; Zhu et al., 2020a). In this Perspective, we focus on three core issues related to SARS-CoV-2 vaccine development: antigen selection and engineering, preclinical challenge studies, and immune correlates of protection. A discussion of clinical development and deployment strategies for SARS-CoV-2 vaccines, while also important, is beyond the scope of this manuscript.

ANTIGEN SELECTION AND ENGINEERING

Coronaviruses Encode Multiple Structural and Non-structural Proteins that Could Potentially Serve as Immunogens for a SARS-CoV-2 Vaccine

The best characterized proteins are S, N, M, and E. S has most commonly been utilized in coronavirus vaccine studies, due to its pivotal role in mediating viral entry into cells (Song et al., 2019). Mature S is a trimeric class I fusion protein located on the surface of the virion. Many coronaviruses proteolytically process S into the S1 and S2 domains. The S1 fragment contains the receptor binding domain (RBD) and the S2 fragment contains the fusion peptide, which are responsible for receptor binding and cell fusion, respectively. For SARS-CoV, S has been demonstrated to be the primary target of neutralizing antibodies. In mouse models of SARS-CoV, passive transfer and vaccine studies have shown that S-specific antibodies confer protective immunity (Enjuanes et al., 2008; Yang et al., 2004). For SARS-CoV-2, studies with monoclonal antibodies have shown that SARS-CoV-2-infected humans develop robust neutralizing antibody responses against S and in particular the RBD (Baum et al., 2020; Hansen et al., 2020; Ju et al., 2020; Rogers et al., 2020; Shi et al., 2020; Zost et al., 2020).

In addition to S, early studies with SARS-CoV suggested that most infected individuals developed an antibody response to N (Pei et al., 2005). N-protein-immunized BALB/c mice also






Figure 1. SARS-CoV-2 Spike

Graphical representation of the SARS-CoV-2 S protein sequence and crystal structure of SARS-CoV-2 S protein ectodomain (PDB: 6VSB) (Wrapp et al., 2020) created by using PyMol software. The transmembrane domain and cytoplasmic tail were not crystallized. TM, transmembrane domain; CT, cytoplasmic tail.

induced CD4⁺ and CD8⁺ T cells (Liu et al., 2006). However, vaccination with vaccines expressing N resulted in no protection against SARS-CoV challenge as well as enhanced infection, which was characterized by increased pulmonary eosinophil infiltration (Deming et al., 2006). Passive transfer of anti-N antibodies did not generate an enhanced response, leading the authors to believe that the increased severity was linked to a T cell response (Deming et al., 2006). This was shown upon vaccination with VV expressing S, M, N, and E, as well as VV expressing N, causing the authors to trace back the response to nucleocapsid as an immunogen. Similarly, BALB/c mice vaccinated with a vaccinia virus (VV) expressing N showed enhanced viral infection upon SARS-CoV challenge, which was associated with a T_H2 response with pulmonary infiltration of neutrophils, eosinophils, and lymphocytes (Yasui et al., 2008).

The M and E proteins have garnered less interest as vaccine targets due to lower immunogenicity (Du et al., 2008a), although SARS-CoV patient sera has been shown to be reactive to M peptides (Wang et al., 2003). Of the less-studied proteins, Orf3a has been shown to be capable of raising a neutralizing polyclonal antibody response in rabbits (Akerström et al., 2006).

Currently, Most Vaccines for SARS-CoV-2 Are Focused on S

To target the SARS-CoV-2 S protein in its native prefusion form, antigen stabilization strategies have been used (Figure 1). For MERS-CoV and SARS-CoV, introduction of two prolines in the S2 subunit effectively stabilized the S in its prefusion conforma-

tion (Pallesen et al., 2017). The prefusion-stabilized MERS-CoV S generated higher neutralizing antibody titers when compared to native S trimer (Pallesen et al., 2017). This mutation prevented conformational changes of S in the presence of its receptor, ACE2, or trypsin (Kirchdoerfer et al., 2018). This stabilization method has also been demonstrated to stabilize the SARS-CoV-2 S (Wrapp et al., 2020) and has been applied to multiple SARS-CoV-2 vaccine candidates (Corbett et al., 2020; Jackson et al., 2020; Mercado et al., 2020). Unlike SARS-CoV, but similar to MERS-CoV, SARS-CoV-2 includes a furin cleavage site between the S1 and S2 domains. For MERS-CoV, researchers mutated the furin cleavage site to enhance homogeneity and stability (Walls et al., 2019). Some investigators have also added a foldon trimerization tag to the C terminus (Walls et al., 2020), and deletion of the cytoplasmic tail of the SARS-CoV S has been shown to increase neutralizing antibody titers (Yang et al., 2004). Current vaccine candidates in clinical trials are also exploring the inclusion of a tissue plasminogen activator leader sequence (tPA) with S (Folegatti et al., 2020; Mercado et al., 2020; Zhu et al., 2020a; Zhu et al., 2020). The tPA leader can increase immunogen secretion and elicit increased humoral immune responses in influenza (Luo et al., 2008) and HIV vaccines (Wallace et al., 2013), although cellular immunogenicity was only increased in the influenza vaccine.

The RBD alone has also been explored as an immunogen. In preclinical studies in mice, rabbits, and monkeys, vaccination with modified vaccinia Ankara (MVA) expressing the full-length SARS-CoV S induced neutralizing antibodies that targeted RBD (Chen et al., 2005). Moreover, depletion of RBD-specific antibodies significantly reduced convalescent plasma neutralizing capabilities (He et al., 2005), consistent with the concept that RBD is the primary target of S-specific neutralizing antibodies. When rabbits were immunized with a variety of SARS-CoV-2 S immunogens, including RBD, S1, S2, and modified variants, RBD was found to elicit five-fold higher affinity antibodies than the other immunogens (Ravichandran et al., 2020). Sprague-Dawley rats immunized with SARS-CoV-2 RBD also elicited a strong neutralizing antibody response (Quinlan et al., 2020). RBD immunogens have been combined with a foldon trimerization tag in the BNT162b1 COVID-19 RNA vaccine candidate (Mulligan et al., 2020).

We recently compared a series of SARS-CoV-2 S immunogens in the context of both DNA vaccines and Ad26 vectors (Mercado et al., 2020; Yu et al., 2020). DNA vaccines encoding the full-length S elicited higher neutralizing antibody titers than did DNA vaccines encoding several S deletion mutants and also afforded improved protection against SARS-CoV-2 challenge in rhesus macaques (Yu et al., 2020). Ad26 vectors were also generated encoding a series of S variants, and the full-length S with the PP-stabilizing mutations proved the most immunogenic and afforded the best protection against SARS-CoV-2 challenge in rhesus macaques (Mercado et al., 2020).

PRECLINICAL CHALLENGE STUDIES IN NHPs

An optimal model for SARS-CoV-2 infection studies would involve an animal species permissive to viral replication and that develops pathologic and clinical features consistent with the human disease. Clinical manifestations of COVID-19 in

Table 1. NHP Challenge Studies of SARS-CoV-2 Vaccine Candidates

Vaccine Name	Vaccine Type	Vaccine Immunogen	Vaccine Dose	Number of Injections	Challenge Route	Reference
PiCoVacc	inactivated	whole virus	3 or 6 μ g	3	intratracheal	Gao et al., 2020
BBIBP-CorV	inactivated	whole virus	2 or 8 μ g	2	intratracheal	Wang et al., 2020
DNA-S	DNA	engineered S	5 mg	2	intranasal and intratracheal	Yu et al., 2020
mRNA-1273	RNA	engineered S	10 or 100 μ g	2	intranasal and intratracheal	Corbett et al., 2020
ChAdOx1 nCoV-19	adenoviral vector	tPA-S	2.5×10^{10} VP	1 or 2	intranasal, intratracheal, ocular, and oral	van Doremalen et al., 2020
Ad26.COV2.S (Ad26-S.PP)	adenoviral vector	engineered S	1×10^{11} VP	1	intranasal and intratracheal	Mercado et al., 2020

humans are usually mild but can include cough, fever, pneumonia, and occasionally respiratory failure and death (Yang et al., 2020). NHPs have significant genetic homology to humans and are often useful models for infectious diseases, although cost and availability can be limiting. In both rhesus and cynomolgus macaques, virus shedding could be detected in nasal, throat, and rectal swabs and in bronchoalveolar lavage for approximately 2 weeks after infection with SARS-CoV-2 using either intratracheal and intranasal infection or intratracheal, intranasal, ocular, and oral infection (Chandrashekar et al., 2020; Munster et al., 2020; Rockx et al., 2020). Early data suggest the utility of both macaque species as animal models for SARS-CoV-2 infection, although respiratory disease has been reported to be mild. The potential utility of African green monkeys as a model is also being explored. We recently demonstrated that SARS-CoV-2 infected macaques were also robustly protected against re-challenge, demonstrating natural protective immunity (Chandrashekar et al., 2020).

At least six SARS-CoV-2 vaccine challenge studies in macaques have been published at the time of this writing (Corbett et al., 2020; Gao et al., 2020; Mercado et al., 2020; van Doremalen et al., 2020; Wang et al., 2020; Yu et al., 2020) (Table 1). These vaccine studies in NHPs have included inactivated vaccines (PiCoVacc [Gao et al., 2020], BBIBP-CorV [Wang et al., 2020]), DNA vaccines (Yu et al., 2020), RNA vaccines (mRNA-1273 [Corbett et al., 2020]), and adenovirus-based vaccines (ChAdOx1 [van Doremalen et al., 2020], Ad26 [Mercado et al., 2020]). PiCoVacc and BBIBP-CorV are based on SARS-CoV-2 CN2 and SARS-CoV-2 HB02 strains, respectively. The viruses were grown in Vero cells and inactivated by using β -propiolactone and were evaluated as two- or three-shot immunization regimens (Gao et al., 2020; Wang et al., 2020). The DNA and mRNA-1273 vaccines encoded stabilized S immunogens and were tested as two-shot immunization regimens (Yu et al., 2020; Corbett et al., 2020). The ChAdOx1 vaccine expressed a codon-optimized full-length S with a human tPA leader sequence and was tested as a single-shot and a two-shot vaccine regimen (van Doremalen et al., 2020). The optimal Ad26 vaccine expressed a prefusion-stabilized S immunogen and was tested as a single-shot vaccine (Mercado et al., 2020).

After vaccination, macaques were challenged by SARS-CoV-2 by the intratracheal, intranasal, oral, and/or ocular routes. Efficacy was determined in these studies by assessment of viral loads in the upper and lower respiratory tracts. Most studies are now focusing on analysis of subgenomic RNA rather than genomic RNA (Wölfel et al., 2020), because subgenomic RNA

is believed to be more reflective of replicating virus, rather than input challenge virus. The inactivated vaccine PiCoVacc resulted in reduced viral loads in throat swabs (Gao et al., 2020). The inactivated vaccine BBIBP-CorV also decreased viral loads in throat swabs (Wang et al., 2020). The DNA vaccine encoding the full-length S resulted in >3.1 and >3.7 \log_{10} reductions in median subgenomic RNA levels in BAL and nasal swabs, respectively (Yu et al., 2020). The mRNA-1273 vaccine resulted in undetectable subgenomic RNA in lungs in all but one macaque, and more rapid clearance of virus from nasal swabs (Corbett et al., 2020). The optimal Ad26-S.PP vaccine resulted in undetectable subgenomic RNA in lungs and only one breakthrough in nasal swabs (Mercado et al., 2020). The ChAdOx1 vaccine resulted in reduced subgenomic RNA in BAL but no decrease in nasal swabs (van Doremalen et al., 2020).

The different challenge doses, strains, routes, and assays that were utilized make direct comparisons among these studies difficult. Nevertheless, these studies provide a substantial amount of preclinical data demonstrating protective efficacy of multiple vaccine candidates in NHPs. These data help inform the ongoing clinical development programs for these vaccines. A limitation of all these studies is that macaques do not develop severe disease, respiratory failure, or death. Small animal models are currently being developed that could model severe disease, including hamsters and transgenic mice.

IMMUNE CORRELATES OF PROTECTION

Determining immune correlates of protection (CoP) for SARS-CoV-2 will be critical for guiding vaccine development. Currently, mechanistic CoP for SARS-CoV-2 have not yet been fully determined, although several studies point to the importance of both humoral and cellular immunity.

Humoral Immunity

Neutralizing antibodies (nAbs) represent a commonly studied immune correlate of protection. Early human challenge studies reported that volunteers with higher pre-existing anti-CoV 229E nAb titers (> 5) demonstrated lower proportions of virus isolation and upper respiratory infection than those with low neutralizing titers (≤ 5) (Bradburne et al., 1967). Pre-challenge serum antibody titers were also negatively associated with upper respiratory infections, nasal virus shedding, and disease severity (Barrow et al., 1990; Callow, 1985). Passive administration of convalescent sera, purified IgG, or monoclonal antibodies have also been shown to suppress SARS-CoV challenge/infection

and associated disease progression in mice, hamsters, ferrets, and humans (Cheng et al., 2005; Roberts et al., 2006; Subbarao et al., 2004; Sui et al., 2005; Yuan et al., 2015).

For SARS-CoV-2, neutralizing monoclonal antibodies isolated from convalescent COVID-19 patients have been shown to inhibit SARS-CoV-2 infection in both prophylactic and therapeutic settings in rhesus macaques and hamsters (Rogers et al., 2020; Shi et al., 2020; Zost et al., 2020). We also reported that DNA vaccines and Ad26-based vaccines induced nAbs that strongly correlated with a reduction of viral loads in rhesus macaques (Mercado et al., 2020; Yu et al., 2020). SARS-CoV-2 inactivated virus vaccines and mRNA vaccines also induced nAbs and conferred protection in macaques (Corbett et al., 2020; Gao et al., 2020; Wang et al., 2020). Collectively, these studies suggest that nAb titers could serve as a useful biomarker for evaluating SARS-CoV-2 vaccines in both preclinical and clinical studies, although these correlates need to be confirmed in humans.

A question that has been raised is whether sub-neutralizing levels of antibodies could have detrimental effects. A prior study revealed that a vaccinia vector-based vaccine expressing feline coronavirus S induced low titers of neutralizing antibodies, which led to early cat mortality upon challenge (Vennema et al., 1990). Consistent with this observation, several studies observed enhanced respiratory disease (ERD) in response to SARS vaccines when antibodies had suboptimal potency or low binding affinity (Jaume et al., 2011; Wan et al., 2020; Wang et al., 2014). A major effort in the field will therefore be the development of animal models for ERD for SARS-CoV-2.

In addition to neutralization, emerging evidence suggests that certain antibody Fc-mediated functions could also contribute to protective efficacy (Yu et al., 2020), including antibody-dependent complement deposition (ADCD), antibody-dependent cellular phagocytosis (ADCP), and antibody-dependent NK cell activation (ADNKA) (Zohar and Alter, 2020). Mucosal immunity is also likely important for protection, because coronavirus infection occurs in the respiratory tract and potentially in the gastrointestinal tract (Xiao et al., 2020). Pulmonary immunoglobulin (Ig)A was observed to be inversely associated with MERS-CoV infectivity in humans (Muth et al., 2015), and animal studies have highlighted the potential role of mucosal immunity in defending SARS-CoV infection (Du et al., 2008b; Huang et al., 2009). Given that intranasal administration could induce stronger mucosal immunity in the respiratory tract than parenteral routes, it could prove useful to evaluate intranasal routes for vaccines (Roper and Rehm, 2009).

Cellular Immunity

Cellular immunity also appears important in the control of coronavirus infections. Current evidence suggests that not all patients develop protective humoral immune responses, and asymptomatic patients or individuals with mild disease typically develop robust T cell responses (Mathew et al., 2020; Sekine et al., 2020). In a mouse model, SARS-CoV-specific CD8⁺ T cell numbers correlated with virus clearance and increased survival (Zhao et al., 2010). In addition, memory CD4⁺ T cells were associated with protective immunity against MERS-CoV (Zhao et al., 2016). Recent studies have reported that neutralizing antibody titers correlated with SARS-CoV-2-specific

T cell responses (Ni et al., 2020) and that IgG and IgA antibody titers correlated with S-specific CD4⁺ T cell responses (Grifoni et al., 2020), suggesting that T cells could indirectly modulate the virus infection by orchestrating antibody production. However, in DNA vaccine studies in mice and NHPs, T cell responses did not correlate with protection (Yang et al., 2004; Yu et al., 2020). Nevertheless, severe patients tend to have a higher frequency of polyfunctional CD4 cells expressing interferon (IFN) γ , interleukin (IL)-2, and tumor necrosis factor alpha (TNF- α) (Sekine et al., 2020; Thieme et al., 2020), although it is possible that these high-frequency T cells could simply represent long-term viral exposure (Altmann and Boyton, 2020).

Innate Immunity

There is limited data on innate immune correlates of protection. Existing pathogenesis studies suggest that the excessive production of proinflammatory cytokines and chemokines (cytokine storm) is associated with poor clinical outcome (Channappanavar and Perlman, 2017). IFN-I and IFN-III have shown to be able to suppress SARS-CoV-2 infection *in vitro* (Sallard et al., 2020; Stanifer et al., 2020) and could represent innate immune responses that assist in controlling SARS-CoV-2 infection. Blocking IFN-I signaling in mice has been shown to increase viral load and mortality for SARS-CoV and MERS-CoV (Channappanavar et al., 2019; Frieman et al., 2010). Retrospective cohort studies have also suggested that induction of IFNs correlated with disease severity and viral load in both SARS and MERS patients (Cameron et al., 2007; Kim et al., 2016). It is possible that early induction promotes viral clearance, whereas delayed IFN responses can cause viral persistence, inflammation, and immunopathology (Park and Iwasaki, 2020).

The potential off-target effects of BCG, historically used as a vaccine for tuberculosis, and other live attenuated organisms are also being explored for SARS-CoV-2 (Curtis et al., 2020). Trained immunity is regarded as immunological memory in the innate immune system and likely mediated by epigenetic modifications (Netea et al., 2016). If non-specific trained innate immunity protects against SARS-CoV-2, this would provide an intriguing alternate to antigen-specific vaccination.

CONCLUSIONS

A vaccine will likely be required to end the global SARS-CoV-2 pandemic (Lurie et al., 2020). A successful vaccine will need to be safe, effective, durable, and deployable to large populations. There are currently at least 40 SARS-CoV-2 vaccine clinical trials ongoing. Most vaccine strategies aim to generate S-specific neutralizing antibodies, and preclinical studies have shown substantial protection against SARS-CoV-2 challenge in NHPs. Over the next several months, additional preclinical efficacy studies as well as studies on enhanced respiratory disease will likely be reported, and multiple large-scale clinical efficacy trials will be conducted.

ACKNOWLEDGMENTS

We acknowledge support from the Ragon Institute of MGH, MIT, and Harvard; Mark and Lisa Schwartz Foundation; Massachusetts Consortium on Pathogen

Readiness (MassCPR); Bill & Melinda Gates Foundation (INV-006131); Janssen Vaccines & Prevention BV; and the National Institutes of Health (OD024917, AI129797, AI124377, AI128751, and AI126603).

DECLARATION OF INTERESTS

Correspondence and requests for materials should be addressed to D.H.B. (dbarouch@bidmc.harvard.edu). D.H.B. is a co-inventor on provisional vaccine patents (62/969,008; 62/994,630) that have been licensed.

REFERENCES

- Akerström, S., Tan, Y.J., and Mirazimi, A. (2006). Amino acids 15–28 in the ectodomain of SARS coronavirus 3a protein induces neutralizing antibodies. *FEBS Lett.* *580*, 3799–3803.
- Altmann, D.M., and Boyton, R.J. (2020). SARS-CoV-2 T cell immunity: Specificity, function, durability, and role in protection. *Sci. Immunol.* *5*, eabd6160.
- Arons, M.M., Hatfield, K.M., Reddy, S.C., Kimball, A., James, A., Jacobs, J.R., Taylor, J., Spicer, K., Bardossy, A.C., Oakley, L.P., et al.; Public Health–Seattle and King County and CDC COVID-19 Investigation Team (2020). Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *N. Engl. J. Med.* *382*, 2081–2090.
- Barrow, G.I., Higgins, P.G., al-Nakib, W., Smith, A.P., Wenham, R.B.M., and Tyrrell, D.A.J. (1990). The effect of intranasal nedocromil sodium on viral upper respiratory tract infections in human volunteers. *Clin. Exp. Allergy* *20*, 45–51.
- Baum, A., Fulton, B.O., Wloga, E., Copin, R., Pascal, K.E., Russo, V., Giordano, S., Lanza, K., Negron, N., Ni, M., et al. (2020). Antibody cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual antibodies. *Science*, eabd0831.
- Bradburne, A.F., Bynoe, M.L., and Tyrrell, D.A. (1967). Effects of a “new” human respiratory virus in volunteers. *BMJ* *3*, 767–769.
- Callow, K.A. (1985). Effect of specific humoral immunity and some non-specific factors on resistance of volunteers to respiratory coronavirus infection. *J. Hyg. (Lond.)* *95*, 173–189.
- Cameron, M.J., Ran, L., Xu, L., Danesh, A., Bermejo-Martin, J.F., Cameron, C.M., Muller, M.P., Gold, W.L., Richardson, S.E., Poutanen, S.M., et al.; Canadian SARS Research Network (2007). Interferon-mediated immunopathological events are associated with atypical innate and adaptive immune responses in patients with severe acute respiratory syndrome. *J. Virol.* *81*, 8692–8706.
- Chan, J.F.-W., Yuan, S., Kok, K.-H., To, K.K.-W., Chu, H., Yang, J., Xing, F., Liu, J., Yip, C.C.-Y., Poon, R.W.-S., et al. (2020). A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet* *395*, 514–523.
- Chandrashekar, A., Liu, J., Martinot, A.J., McMahan, K., Mercado, N.B., Peter, L., Tostanoski, L.H., Yu, J., Maliga, Z., Nekorchuk, M., et al. (2020). SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science*, eabc4776.
- Channappanavar, R., and Perlman, S. (2017). Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin. Immunopathol.* *39*, 529–539.
- Channappanavar, R., Fehr, A.R., Zheng, J., Wohlford-Lenane, C., Abrahante, J.E., Mack, M., Sompallae, R., McCray, P.B., Jr., Meyerholz, D.K., and Perlman, S. (2019). IFN-I response timing relative to virus replication determines MERS coronavirus infection outcomes. *J. Clin. Invest.* *129*, 3625–3639.
- Chen, Z., Zhang, L., Qin, C., Ba, L., Yi, C.E., Zhang, F., Wei, Q., He, T., Yu, W., Yu, J., et al. (2005). Recombinant modified vaccinia virus Ankara expressing the spike glycoprotein of severe acute respiratory syndrome coronavirus induces protective neutralizing antibodies primarily targeting the receptor binding region. *J. Virol.* *79*, 2678–2688.
- Cheng, Y., Wong, R., Soo, Y.O., Wong, W.S., Lee, C.K., Ng, M.H., Chan, P., Wong, K.C., Leung, C.B., and Cheng, G. (2005). Use of convalescent plasma therapy in SARS patients in Hong Kong. *European journal of clinical microbiology & infectious diseases. European Society of Clinical Microbiology* *24*, 44–46.
- Corbett, K.S., Flynn, B., Foulds, K.E., Francica, J.R., Boyoglu-Barnum, S., Werner, A.P., Flach, B., O’Connell, S., Bock, K.W., Minai, M., et al. (2020). Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in Nonhuman Primates. *N. Engl. J. Med.*
- Curtis, N., Sparrow, A., Ghebreyesus, T.A., and Netea, M.G. (2020). Considering BCG vaccination to reduce the impact of COVID-19. *Lancet* *395*, 1545–1546.
- Deming, D., Sheahan, T., Heise, M., Yount, B., Davis, N., Sims, A., Suthar, M., Harkema, J., Whitmore, A., Pickles, R., et al. (2006). Vaccine efficacy in senescent mice challenged with recombinant SARS-CoV bearing epidemic and zoonotic spike variants. *PLoS Med.* *3*, e525.
- Du, L., He, Y., Jiang, S., and Zheng, B.-J. (2008a). Development of subunit vaccines against severe acute respiratory syndrome. *Drugs Today (Barc)* *44*, 63–73.
- Du, L., Zhao, G., Lin, Y., Sui, H., Chan, C., Ma, S., He, Y., Jiang, S., Wu, C., Yuen, K.-Y., et al. (2008b). Intranasal vaccination of recombinant adeno-associated virus encoding receptor-binding domain of severe acute respiratory syndrome coronavirus (SARS-CoV) spike protein induces strong mucosal immune responses and provides long-term protection against SARS-CoV infection. *J. Immunol.* *180*, 948–956.
- Enjuanes, L., Dediego, M.L., Alvarez, E., Deming, D., Sheahan, T., and Baric, R. (2008). Vaccines to prevent severe acute respiratory syndrome coronavirus-induced disease. *Virus Res.* *133*, 45–62.
- Folegatti, P.M., Ewer, K.J., Aley, P.K., Angus, B., Becker, S., Belli-Rammerstorfer, S., Bellamy, D., Bibi, S., Bittaye, M., Clutterbuck, E.A., et al.; Oxford COVID Vaccine Trial Group (2020). Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial. *Lancet*, S0140-6736(20)31604-4.
- Frieman, M.B., Chen, J., Morrison, T.E., Whitmore, A., Funkhouser, W., Ward, J.M., Lamirande, E.W., Roberts, A., Heise, M., Subbarao, K., and Baric, R.S. (2010). SARS-CoV pathogenesis is regulated by a STAT1 dependent but a type I, II and III interferon receptor independent mechanism. *PLoS Pathog.* *6*, e1000849.
- Gao, Q., Bao, L., Mao, H., Wang, L., Xu, K., Yang, M., Li, Y., Zhu, L., Wang, N., Lv, Z., et al. (2020). Development of an inactivated vaccine candidate for SARS-CoV-2. *Science* *369*, 77–81.
- Grifoni, A., Weiskopf, D., Ramirez, S.I., Mateus, J., Dan, J.M., Moderbacher, C.R., Rawlings, S.A., Sutherland, A., Premkumar, L., Jadi, R.S., et al. (2020). Targets of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and Unexposed Individuals. *Cell* *181*, 1489–1501.e15, e1415.
- Hansen, J., Baum, A., Pascal, K.E., Russo, V., Giordano, S., Wloga, E., Fulton, B.O., Yan, Y., Koon, K., Patel, K., et al. (2020). Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science*, eabd0827.
- He, Y., Zhu, Q., Liu, S., Zhou, Y., Yang, B., Li, J., and Jiang, S. (2005). Identification of a critical neutralization determinant of severe acute respiratory syndrome (SARS)-associated coronavirus: importance for designing SARS vaccines. *Virology* *334*, 74–82.
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T., Erichsen, S., Schiergens, T.S., Herrler, G., Wu, N.-H., Nitsche, A., et al. (2020). SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* *181*, 271–280.e8.
- Huang, X., Lu, B., Yu, W., Fang, Q., Liu, L., Zhuang, K., Shen, T., Wang, H., Tian, P., Zhang, L., and Chen, Z. (2009). A novel replication-competent vaccinia vector MVTT is superior to MVA for inducing high levels of neutralizing antibody via mucosal vaccination. *PLoS ONE* *4*, e4180.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., et al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* *395*, 497–506.
- Jackson, L.A., Anderson, E.J., Roupheal, N.G., Roberts, P.C., Makhene, M., Coler, R.N., McCullough, M.P., Chappell, J.D., Denison, M.R., Stevens, L.J., et al.; mRNA-1273 Study Group (2020). An mRNA Vaccine against SARS-CoV-2 - Preliminary Report. *N. Engl. J. Med.*

Jaume, M., Yip, M.S., Cheung, C.Y., Leung, H.L., Li, P.H., Kien, F., Dutry, I., Callendret, B., Escriou, N., Altmeyer, R., et al. (2011). Anti-severe acute respiratory syndrome coronavirus spike antibodies trigger infection of human immune cells via a pH- and cysteine protease-independent FcγR pathway. *J. Virol.* **85**, 10582–10597.

Ju, B., Zhang, Q., Ge, J., Wang, R., Sun, J., Ge, X., Yu, J., Shan, S., Zhou, B., Song, S., et al. (2020). Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**, 115–119.

Kim, E.S., Choe, P.G., Park, W.B., Oh, H.S., Kim, E.J., Nam, E.Y., Na, S.H., Kim, M., Song, K.H., Bang, J.H., et al. (2016). Clinical Progression and Cytokine Profiles of Middle East Respiratory Syndrome Coronavirus Infection. *J. Korean Med. Sci.* **31**, 1717–1725.

Kirchdoerfer, R.N., Wang, N., Pallesen, J., Wrapp, D., Turner, H.L., Cottrell, C.A., Corbett, K.S., Graham, B.S., McLellan, J.S., and Ward, A.B. (2018). Stabilized coronavirus spikes are resistant to conformational changes induced by receptor recognition or proteolysis. *Sci. Rep.* **8**, 15701.

Li, Q., Guan, X., Wu, P., Wang, X., Zhou, L., Tong, Y., Ren, R., Leung, K.S.M., Lau, E.H.Y., Wong, J.Y., et al. (2020). Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N. Engl. J. Med.* **382**, 1199–1207.

Liu, S.-J., Leng, C.-H., Lien, S.P., Chi, H.-Y., Huang, C.-Y., Lin, C.-L., Lian, W.-C., Chen, C.-J., Hsieh, S.-L., and Chong, P. (2006). Immunological characterizations of the nucleocapsid protein based SARS vaccine candidates. *Vaccine* **24**, 3100–3108.

Luo, M., Tao, P., Li, J., Zhou, S., Guo, D., and Pan, Z. (2008). Immunization with plasmid DNA encoding influenza A virus nucleoprotein fused to a tissue plasminogen activator signal sequence elicits strong immune responses and protection against H5N1 challenge in mice. *J. Virol. Methods* **154**, 121–127.

Lurie, N., Saville, M., Hatchett, R., and Halton, J. (2020). Developing Covid-19 Vaccines at Pandemic Speed. *N. Engl. J. Med.* **382**, 1969–1973.

Mathew, D., Giles, J.R., Baxter, A.E., Oldridge, D.A., Greenplate, A.R., Wu, J.E., Alanio, C., Kuri-Cervantes, L., Pampena, M.B., D'Andrea, K., et al.; UPenn COVID Processing Unit (2020). Deep immune profiling of COVID-19 patients reveals distinct immunotypes with therapeutic implications. *Science*, eabc8511.

McMichael, T.M., Currie, D.W., Clark, S., Pogosjans, S., Kay, M., Schwartz, N.G., Lewis, J., Baer, A., Kawakami, V., Lukoff, M.D., et al.; Public Health–Seattle and King County, EvergreenHealth, and CDC COVID-19 Investigation Team (2020). Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N. Engl. J. Med.* **382**, 2005–2011.

Mercado, N.B., Zahn, R., Wegmann, F., Loos, C., Chandrashekar, A., Yu, J., Liu, J., Peter, L., McMahan, K., Tostanoski, L.H., et al. (2020). Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus macaques. *Nature*.

Mulligan, M.J., Lyke, K.E., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S.P., Neuzil, K., Raabe, V., Bailey, R., Swanson, K.A., et al. (2020). Phase 1/2 Study to Describe the Safety and Immunogenicity of a COVID-19 RNA Vaccine Candidate (BNT162b1) in Adults 18 to 55 Years of Age: Interim Report. *medRxiv*. <https://doi.org/10.1101/2020.06.30.20142570>.

Munster, V.J., Feldmann, F., Williamson, B.N., van Doremalen, N., Pérez-Pérez, L., Schulz, J., Meade-White, K., Okumura, A., Callison, J., Brumbaugh, B., et al. (2020). Respiratory disease and virus shedding in rhesus macaques inoculated with SARS-CoV-2. *bioRxiv*, 2020.03.21.001628.

Muth, D., Corman, V.M., Meyer, B., Assiri, A., Al-Masri, M., Farah, M., Steinhagen, K., Lattwein, E., Al-Tawfiq, J.A., Albarrak, A., et al. (2015). Infectious Middle East Respiratory Syndrome Coronavirus Excretion and Serotype Variability Based on Live Virus Isolates from Patients in Saudi Arabia. *J. Clin. Microbiol.* **53**, 2951–2955.

Netea, M.G., Joosten, L.A.B., Latz, E., Mills, K.H.G., Natoli, G., Stunnenberg, H.G., O'Neill, L.A.J., and Xavier, R.J. (2016). Trained immunity: A program of innate immune memory in health and disease. *Science* **352**, aaf1098.

Ni, L., Ye, F., Cheng, M.L., Feng, Y., Deng, Y.Q., Zhao, H., Wei, P., Ge, J., Gou, M., Li, X., et al. (2020). Detection of SARS-CoV-2-Specific Humoral and Cellular Immunity in COVID-19 Convalescent Individuals. *Immunity* **52**, 971–977.e3, e973.

Pallesen, J., Wang, N., Corbett, K.S., Wrapp, D., Kirchdoerfer, R.N., Turner, H.L., Cottrell, C.A., Becker, M.M., Wang, L., Shi, W., et al. (2017).

Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike antigen. *Proc. Natl. Acad. Sci. USA* **114**, E7348–E7357.

Park, A., and Iwasaki, A. (2020). Type I and Type III Interferons - Induction, Signaling, Evasion, and Application to Combat COVID-19. *Cell Host Microbe* **27**, 870–878.

Pei, H., Liu, J., Cheng, Y., Sun, C., Wang, C., Lu, Y., Ding, J., Zhou, J., and Xiang, H. (2005). Expression of SARS-coronavirus nucleocapsid protein in *Escherichia coli* and *Lactococcus lactis* for serodiagnosis and mucosal vaccination. *Appl. Microbiol. Biotechnol.* **68**, 220–227.

Quinlan, B.D., Mou, H., Zhang, L., Guo, Y., He, W., Ojha, A., Parcells, M.S., Luo, G., Li, W., Zhong, G., et al. (2020). The SARS-CoV-2 receptor-binding domain elicits a potent neutralizing response without antibody-dependent enhancement. *bioRxiv*. <https://doi.org/10.1101/2020.04.10.036418>.

Ravichandran, S., Coyle, E.M., Klenow, L., Tang, J., Grubbs, G., Liu, S., Wang, T., Golding, H., and Khurana, S. (2020). Antibody signature induced by SARS-CoV-2 spike protein immunogens in rabbits. *Sci. Transl. Med.* **12**, eabc3539.

Roberts, A., Thomas, W.D., Guarnier, J., Lamirande, E.W., Babcock, G.J., Greenough, T.C., Vogel, L., Hayes, N., Sullivan, J.L., Zaki, S., et al. (2006). Therapy with a severe acute respiratory syndrome-associated coronavirus-neutralizing human monoclonal antibody reduces disease severity and viral burden in golden Syrian hamsters. *J. Infect. Dis.* **193**, 685–692.

Rockx, B., Kuiken, T., Herfst, S., Bestebroer, T., Lamers, M.M., Oude Munnink, B.B., de Meulder, D., van Amerongen, G., van den Brand, J., Okba, N.M.A., et al. (2020). Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model. *Science* **368**, 1012–1015.

Rogers, T.F., Zhao, F., Huang, D., Beutler, N., Burns, A., He, W.T., Limbo, O., Smith, C., Song, G., Woehl, J., et al. (2020). Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model. *Science*, eabc7520.

Roper, R.L., and Rehm, K.E. (2009). SARS vaccines: where are we? *Expert Rev. Vaccines* **8**, 887–898.

Sallard, E., Lescure, F.-X., Yazdanpanah, Y., Mentre, F., and Peiffer-Smadja, N. (2020). Type I interferons as a potential treatment against COVID-19. *Antiviral Res.* **178**, 104791.

Sekine, T., Perez-Potti, A., Rivera-Ballesteros, O., Strålin, K., Gorin, J.-B., Olsson, A., Llewellyn-Lacey, S., Kamal, H., Bogdanovic, G., Muschiol, S., et al. (2020). Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *bioRxiv*. <https://doi.org/10.1101/2020.06.29.174888>.

Shi, R., Shan, C., Duan, X., Chen, Z., Liu, P., Song, J., Song, T., Bi, X., Han, C., Wu, L., et al. (2020). A human neutralizing antibody targets the receptor-binding site of SARS-CoV-2. *Nature* **584**, 120–124.

Song, Z., Xu, Y., Bao, L., Zhang, L., Yu, P., Qu, Y., Zhu, H., Zhao, W., Han, Y., and Qin, C. (2019). From SARS to MERS, thrusting coronaviruses into the spotlight. *Viruses* **11**, 59.

Srinivasan, S., Cui, H., Gao, Z., Liu, M., Lu, S., Mkandawire, W., Narykov, O., Sun, M., and Korkin, D. (2020). Structural Genomics of SARS-CoV-2 Indicates Evolutionary Conserved Functional Regions of Viral Proteins. *Viruses* **12**, 360.

Stanifer, M.L., Kee, C., Cortese, M., Zumarán, C.M., Triana, S., Mukenhirn, M., Kraeusslich, H.-G., Alexandrov, T., Bartenschlager, R., and Boulant, S. (2020). Critical Role of Type III Interferon in Controlling SARS-CoV-2 Infection in Human Intestinal Epithelial Cells. *Cell Rep.* **32**, 107863.

Subbarao, K., McAuliffe, J., Vogel, L., Fahle, G., Fischer, S., Tatti, K., Packard, M., Shieh, W.J., Zaki, S., and Murphy, B. (2004). Prior infection and passive transfer of neutralizing antibody prevent replication of severe acute respiratory syndrome coronavirus in the respiratory tract of mice. *J. Virol.* **78**, 3572–3577.

Sui, J., Li, W., Roberts, A., Matthews, L.J., Murakami, A., Vogel, L., Wong, S.K., Subbarao, K., Farzan, M., and Marasco, W.A. (2005). Evaluation of human monoclonal antibody 80R for immunoprophylaxis of severe acute respiratory syndrome by an animal study, epitope mapping, and analysis of spike variants. *J. Virol.* **79**, 5900–5906.

Thieme, C.J., Anft, M., Paniskaki, K., Blazquez-Navarro, A., Doevelaar, A., Seibert, F.S., Hoelzer, B., Konik, M.J., Brenner, T., Tempfer, C., et al. (2020). The SARS-CoV-2 T-cell immunity is directed against the spike, membrane, and nucleocapsid protein and associated with COVID 19 severity. *medRxiv*. <https://doi.org/10.1101/2020.05.13.20100636>.

- van Doremalen, N., Lambe, T., Spencer, A., Belij-Rammerstorfer, S., Purushotham, J.N., Port, J.R., Avanzato, V.A., Bushmaker, T., Flaxman, A., Ulaszewska, M., et al. (2020). ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia in rhesus macaques. *Nature*.
- Vennema, H., de Groot, R.J., Harbour, D.A., Dalderup, M., Gruffydd-Jones, T., Horzinek, M.C., and Spaan, W.J. (1990). Early death after feline infectious peritonitis virus challenge due to recombinant vaccinia virus immunization. *J. Virol.* *64*, 1407–1409.
- Wallace, A., West, K., Rothman, A.L., Ennis, F.A., Lu, S., and Wang, S. (2013). Post-translational intracellular trafficking determines the type of immune response elicited by DNA vaccines expressing Gag antigen of Human Immunodeficiency Virus Type 1 (HIV-1). *Hum. Vaccin. Immunother.* *9*, 2095–2102.
- Walls, A.C., Xiong, X., Park, Y.-J., Tortorici, M.A., Snijder, J., Quispe, J., Cameroni, E., Gopal, R., Dai, M., Lanzavecchia, A., et al. (2019). Unexpected Receptor Functional Mimicry Elucidates Activation of Coronavirus Fusion. *Cell* *176*, 1026–1039.e15.
- Walls, A.C., Park, Y.-J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. (2020). Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell* *181*, 281–292.e6.
- Wan, Y., Shang, J., Sun, S., Tai, W., Chen, J., Geng, Q., He, L., Chen, Y., Wu, J., Shi, Z., et al. (2020). Molecular Mechanism for Antibody-Dependent Enhancement of Coronavirus Entry. *J. Virol.* *94*, 94.
- Wang, J., Wen, J., Li, J., Yin, J., Zhu, Q., Wang, H., Yang, Y., Qin, E., You, B., Li, W., et al. (2003). Assessment of immunoreactive synthetic peptides from the structural proteins of severe acute respiratory syndrome coronavirus. *Clin. Chem.* *49*, 1989–1996.
- Wang, S.F., Tseng, S.P., Yen, C.H., Yang, J.Y., Tsao, C.H., Shen, C.W., Chen, K.H., Liu, F.T., Liu, W.T., Chen, Y.M., and Huang, J.C. (2014). Antibody-dependent SARS coronavirus infection is mediated by antibodies against spike proteins. *Biochem. Biophys. Res. Commun.* *451*, 208–214.
- Wang, H., Zhang, Y., Huang, B., Deng, W., Quan, Y., Wang, W., Xu, W., Zhao, Y., Li, N., Zhang, J., et al. (2020). Development of an Inactivated Vaccine Candidate, BBIBP-CoV, with Potent Protection against SARS-CoV-2. *Cell*.
- Wölfel, R., Corman, V.M., Guggemos, W., Seilmaier, M., Zange, S., Müller, M.A., Niemeyer, D., Jones, T.C., Vollmar, P., Rothe, C., et al. (2020). Virological assessment of hospitalized patients with COVID-2019. *Nature* *581*, 465–469.
- Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.-L., Abiona, O., Graham, B.S., and McLellan, J.S. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* *367*, 1260–1263.
- Wu, F., Zhao, S., Yu, B., Chen, Y.-M., Wang, W., Song, Z.-G., Hu, Y., Tao, Z.-W., Tian, J.-H., Pei, Y.-Y., et al. (2020). A new coronavirus associated with human respiratory disease in China. *Nature* *579*, 265–269.
- Xiao, F., Tang, M., Zheng, X., Liu, Y., Li, X., and Shan, H. (2020). Evidence for Gastrointestinal Infection of SARS-CoV-2. *Gastroenterology* *158*, 1831–1833.e3.
- Yang, Z.Y., Kong, W.P., Huang, Y., Roberts, A., Murphy, B.R., Subbarao, K., and Nabel, G.J. (2004). A DNA vaccine induces SARS coronavirus neutralization and protective immunity in mice. *Nature* *428*, 561–564.
- Yang, X., Yu, Y., Xu, J., Shu, H., Xia, J., Liu, H., Wu, Y., Zhang, L., Yu, Z., Fang, M., et al. (2020). Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir. Med.* *8*, 475–481.
- Yasui, F., Kai, C., Kitabatake, M., Inoue, S., Yoneda, M., Yokochi, S., Kase, R., Sekiguchi, S., Morita, K., Hishima, T., et al. (2008). Prior immunization with severe acute respiratory syndrome (SARS)-associated coronavirus (SARS-CoV) nucleocapsid protein causes severe pneumonia in mice infected with SARS-CoV. *J. Immunol.* *181*, 6337–6348.
- Yu, J., Tostanoski, L.H., Peter, L., Mercado, N.B., McMahan, K., Mahrokhian, S.H., Nkolola, J.P., Liu, J., Li, Z., Chandrashekar, A., et al. (2020). DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science*, eabc6284.
- Yuan, A., Yang, H., Qi, H., Cui, J., Hua, W., Li, C., Pang, Z., Zheng, W., and Cui, G. (2015). IL-9 antibody injection suppresses the inflammation in colitis mice. *Biochem. Biophys. Res. Commun.* *468*, 921–926.
- Zhao, J., Zhao, J., and Perlman, S. (2010). T cell responses are required for protection from clinical disease and for virus clearance in severe acute respiratory syndrome coronavirus-infected mice. *J. Virol.* *84*, 9318–9325.
- Zhao, J., Zhao, J., Mangalam, A.K., Channappanavar, R., Fett, C., Meyerholz, D.K., Agnihothram, S., Baric, R.S., David, C.S., and Perlman, S. (2016). Airway Memory CD4(+) T Cells Mediate Protective Immunity against Emerging Respiratory Coronaviruses. *Immunity* *44*, 1379–1391.
- Zhu, F.-C., Guan, X.-H., Li, Y.-H., Huang, J.-Y., Jiang, T., Hou, L.-H., Li, J.-X., Yang, B.-F., Wang, L., Wang, W.-J., et al. (2020a). Immunogenicity and safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet*, S0140-6736(20)31605-6.
- Zhu, F.-C., Li, Y.-H., Guan, X.-H., Hou, L.-H., Wang, W.-J., Li, J.-X., Wu, S.-P., Wang, B.-S., Wang, Z., Wang, L., et al. (2020b). Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial. *Lancet* *395*, 1845–1854.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., et al.; China Novel Coronavirus Investigating and Research Team (2020c). A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J. Med.* *382*, 727–733.
- Zohar, T., and Alter, G. (2020). Dissecting antibody-mediated protection against SARS-CoV-2. *Nat. Rev. Immunol.* *20*, 392–394.
- Zost, S.J., Gilchuk, P., Case, J.B., Binshtein, E., Chen, R.E., Nkolola, J.P., Schäfer, A., Reidy, J.X., Trivette, A., Nargi, R.S., et al. (2020). Potently neutralizing and protective human antibodies against SARS-CoV-2. *Nature*.