



Ex-vivo Sensitivity of *Plasmodium falciparum* to Common Anti-malarial Drugs: The Case of Kéniéroba, a Malaria Endemic Village in Mali

Karim Traoré¹ · Seidina A. S. Diakité¹ · Sekou Bah² · Drissa S. Konaté¹ · Djeneba Dabita¹ · Ibrahim Sanogo¹ · Modibo Sangaré¹ · Souleymane Dama¹ · Bourama Keita¹ · Mory Doumbouya¹ · Merepen A. Guindo¹ · Seydou Doumbia¹ · Mahamadou Diakité¹

Published online: 18 June 2020
© The Author(s) 2020

Abstract

Background In 2006, the National Malaria Control Program in Mali recommended artemisinin-based combination therapy as the first-line treatment for uncomplicated malaria. Since the introduction of artemisinin-based combination therapy, few reports are available on the level of resistance of *Plasmodium falciparum* to the most common anti-malarial drugs in Mali.

Methods From 2016 to 2017, we assessed the ex-vivo drug sensitivity of *P. falciparum* isolates in Kéniéroba, a village located in a rural area of southern Mali. We collected *P. falciparum* isolates from malaria-infected children living in Kéniéroba. The isolates were tested for ex-vivo sensitivity to commonly used anti-malarial drugs, namely chloroquine, quinine, amodiaquine, mefloquine, lumefantrine, dihydroartemisinin, and piperazine. We used the 50% inhibitory concentration determination method, which is based on the incorporation of SYBR[®] Green into the parasite's genetic material.

Results *Plasmodium falciparum* isolates were found to have a reduced ex-vivo sensitivity to quinine (25.7%), chloroquine (12.2%), amodiaquine (2.7%), and mefloquine (1.3%). In contrast, the isolates were 100% sensitive to lumefantrine, dihydroartemisinin, and piperazine. A statistically significant correlation was found between 50% inhibitory concentration values of quinine and amodiaquine ($r=0.80$; $p<0.0001$).

Conclusions *Plasmodium falciparum* isolates were highly sensitive to dihydroartemisinin, lumefantrine, and piperazine and less sensitive to amodiaquine ($n=2$), mefloquine ($n=1$), and quinine ($n=19$). Therefore, our data support the previously reported increasing trend in chloroquine sensitivity in Mali.

1 Background

The resistance of *Plasmodium falciparum* to common anti-malarial drugs represents a serious hurdle for malaria control in endemic countries [1]. *Plasmodium falciparum* has developed resistance to low-cost and well-tolerated anti-malarial drugs such as chloroquine (CQ), amodiaquine (AQ), anti-folates, and mefloquine (MQ) [2–5]. This has led the World Health Organization and National Malaria Control Program (NMCP) to recommend artemisinin-based combination

therapy (ACT) for malaria treatment [6]. There are two main reasons for using ACT in the clinical care of malaria. First, ACT relies on the high efficacy of artemisinin, which is a drug known to suppress disease progression in *P. falciparum*-infected individuals at an earlier stage of infection. Second, combining artemisinin with other anti-malarial drugs will likely delay the emergence and spread of parasites that are resistant to artemisinin. This is an important consideration, given that *P. falciparum* isolates that are resistant to artemisinin-derivative drugs have been detected in Southeast Asia [7, 8]. Specifically, mutations in the *kelch* (*K13*) gene have been associated with resistance to artemisinin in vitro in Asia [9–11]. To our knowledge, such resistance to artemisinin has not been reported in Sub-Saharan Africa yet, and ACT is still very effective in that part of the continent [9, 12]. However, Sub-Saharan Africa remains under threat because of the widespread use of ACT and the increasing intercontinental human migrations. In addition, the circulation of sub-standard or counterfeit drugs combined with

✉ Karim Traoré
ktraore@icermali.org

¹ Malaria Research and Training Center, Mali International Center for Excellence in Research (Mali-ICER), University of Sciences, Techniques and Technologies of Bamako (USTTB), BP 1805, Bamako, Mali

² Hospital of Point-G/University of Sciences, Techniques and Technologies of Bamako (USTTB), Bamako, Mali

poor adherence of patients to antimalarial treatment may contribute to a rapid selection of resistant malaria parasites in Sub-Saharan Africa.

The Malian NMCP adopted ACT as the first-line treatment for uncomplicated malaria in 2006. Notably, artemether-lumefantrine (LUM) and AQ-artesunate are used for uncomplicated malaria treatment in Mali. In addition, three drugs namely, artesunate, artemether, and quinine (QN) were designated to be used in the clinical management of severe malaria via the intravenous route. It is therefore necessary to perform regular monitoring of antimalarial drug resistance in this setting. The ex-vivo drug susceptibility testing is one of the most efficient indirect approaches to assess the efficacy of anti-malarial drugs. Few data are available on in-vitro and ex-vivo efficacy of artemisinin alone or in association with other ACT drugs on isolates circulating in Mali. In 2017, Dama et al. reported a reduced ex-vivo susceptibility of *P. falciparum* after oral artemether-LUM treatment in Mali. This study did not include key anti-malarial drugs such as CQ, QN, AQ, MQ, and piperazine (PPQ). Therefore, additional studies are needed for a comprehensive analysis of drug resistance during malaria treatment to help national and international policy makers in providing evidence-based guidance about clinical management. Here, we report ex-vivo sensitivity of *P. falciparum* isolates to seven (7) common antimalarial drugs in Kénieroba, in southern Mali.

2 Materials and Methods

2.1 Collection of *Plasmodium falciparum* Isolates

From June 2016 to October 2017, *P. falciparum* isolates were collected from patients with malaria aged between 6 months and 18 years in Kénieroba, a village located in the Sudano-Guinean area of Mali (12° 6' 50" N and 8° 19' 58" W). Study participants were enrolled at the health center of Kénieroba. The inclusion criterion was a *P. falciparum* mono-infection with a parasitemia between 1000 and 100,000 parasites per microliter. We collected 2–3 mL of venous blood samples on citric acid and a dextrose collection tube (ACD tubes) from each patient before treatment initiation, according to the NMCP's recommendations.

2.2 Preparation of Anti-malarial Drugs

Plasmodium falciparum isolates were tested using seven common anti-malarial drugs: CQ, QN, AQ, MQ, LUM, dihydroartemisinin (DHA), and PPQ. All drugs were obtained from Sigma. A stock solution of CQ diphosphate, QN, MQ, and DHA were prepared in 70% ethanol. Amodiaquine and LUM were initially dissolved in methanol

while PPQ was dissolved in lactic acid 0.5% first and then in dimethyl sulfoxide. Two-fold serial dilutions were prepared using sterile distilled water and distributed in duplicate into 96-well flat-bottom plates. Final concentration ranges of 2.44–2500 nM for CQ, 4.88–5000 nM for QN, 1.22–1250 nM for MQ, 1.22–1250 nM for AQ, 0.34–350 nM for LUM, 0.10–100 nM for DHA, and 0.98–1000 nM for PPQ. Fifty microliters of each diluted anti-malarial drug were added in a 96-well plate in duplicates. The 96-well plate was first dried up in the ambient air, then maintained at 4 °C, and finally prepared for the culture within 2 weeks. The suitability of the prepared 96-well plate for in-vitro testing was continuously monitored using reference strain 3D7 [13, 14] during the test.

2.3 Culture of the *P. falciparum* Isolates

Briefly, 2–3 mL of whole blood was obtained by venipuncture from each patient with *P. falciparum* mono-infection and transported to the laboratory in Bamako, which is located within 2 h of our study site in Kenieroba. The plasma was removed after centrifugation of the whole blood. The cell pellet was washed three times with incomplete RPMI 1640 medium (Gibco™; Invitrogen Corporation, Carlsbad, California, USA) buffered with 25 mM of HEPES (5.95 g/L) followed by a centrifugation at 2000 rpm for 5 min. Parasites were tested directly without culture adaptation. The suspension of parasites was distributed in 96-well plates pre-loaded with antimalarial drugs (as described above). Culture plates were incubated at 37 °C and 5% CO₂ for 72 h. After this incubation (which corresponds to the schizont stage), a blood smear was prepared to confirm healthy growth of controls (drug-free parasites). Samples were then stored at – 20 °C overnight.

2.4 Malaria SYBR® Green I-Based Fluorescence Assay

The Malaria SYBR® Green I-Based Fluorescence Assay was performed as described in the modified version of the Smilkstein and colleagues paper [15] by Johnson et al. [16]. The plate was thawed for 1–2 h at room temperature to lyse the cells. Then, 100 µL of 0.002% SYBR® Green lysis buffer (10 mL of lysis buffer plus 2 µL of SYBR® Green) was added in each well. The plate was then covered with aluminum foil and kept under agitation using a plate shaker at room temperature for 30 min. The parasite growth was determined by measuring fluorescence of the SYBR® Green incorporated into nucleic acids of the parasites. The plate was read using a fluorometer plate reader using a 485-nm excitation filter and a 538-nm emission filter. The IC₅₀, defined as a drug concentration at which the SYBR® Green

signal was 50% of that measured in drug-free control wells, was calculated by using the In-Vitro Analysis and Reporting Tool (IVART) software.

The IC₅₀ threshold value measuring the parasite's sensitivity to a given drug was defined as reported in previous studies. These values were set at 100 nM for the CQ [17, 18], 30 nM for MQ [17–19], 80 nM for AQ [18, 20], 150 nM for LUM [18, 19, 21], 800 nM for QN [18–20], 12 nM for DHA [18], and 135 for PPQ [22, 23]. The SYBR[®] Green I-based method for determination of drug sensitivity concentrations provides results that are similar to other traditional methods, suggesting that this method can be routinely used to conduct surveillance for drug resistance with fresh or cultured parasites [24]. For example, Smilkstein and co-workers did not find a difference between the IC₅₀ of standard anti-malarial drugs determined by the mean of a radio-isotopic approach from those generated using the SYBR[®] Green method [25].

2.5 Statistical Analysis

We analyzed results from samples that had good ex-vivo growth, free of contamination during culture, and with a good fit on the log dose–response curve. The concentration of a given anti-malarial drug that can inhibit the growth of 50% (IC₅₀) of the parasites in culture was estimated from a dose–response curve by non-linear regression analysis using an online program previously described elsewhere [26, 27]. The program generated IC₅₀ estimates with associated 95% confidence intervals (CIs). Estimated values with insufficient precision based on the CI were discarded. Geometric mean IC₅₀ was calculated for each drug. Cross-susceptibility was analyzed using the Pearson correlation (*r*) generated by the software GraphPad Prism version 8. A two-sided *p* value ≤ 0.05 was set as the significance threshold. The strength of correlation (with a significant linear relationship) was classified as moderate, if “0.3 < *r* < 0.7” or strong, if “*r* ≥ 0.7”.

3 Results

A total of 100 parasite isolates were prepared for analysis. The IC₅₀ was successfully determined for 74 (74%) isolates for CQ, 74 (74%) for QN, 73 (73%) for AQ, 75 (75%) for MQ, 73 (73%) for LUM, 69 (69%) for DHA, and 73 (73%) for PPQ (Table 1). The geometric mean of the IC₅₀ was 46.07 nM with a 95% CI range between 4.0 and 724.3 for CQ. Similarly, it was 264.97 nM (39.90–1567.38) for QN, 25.5 nM (5.73–88) for AQ, 13.13 nM (1.82–36.16) for MQ, 10.14 nM (0.82–50.82) for LUM, 0.87 nM (0.34–3.03) for DHA, and 15.86 nM (3.72–35.72) for PPQ (Table 1). The proportion of resistant *P. falciparum*

Table 1 Geometric mean values of 50% inhibitory concentration (IC₅₀) of different anti-malarial drugs

Molecules	Number of experiments	IC ₅₀ geometric mean (95% CI)
CQ	74	46.07 (4.90–724.3)
QN	74	264.97 (39.90–1567.38)
AQ	73	25.5 (5.73–88)
MQ	75	13.13 (1.82–36.16)
LUM	73	10.14 (0.82–50.82)
DHA	69	0.87 (0.34–3.03)
PPQ	73	15.86 (3.72–35.72)

AQ amodiaquine, CI confidence interval, CQ chloroquine, DHA dihydroartemisinin, LUM lumefantrine, MQ mefloquine, PPQ piperazine, QN quinine

Table 2 Ex-vivo response of isolates of *Plasmodium falciparum* from Kenieroba to antimalarial drugs

Drugs	Cut-off values (nM)	Resistant isolates		Sensitive isolates	
		<i>N</i>	%	<i>N</i>	%
CQ	100	9	12.2	65	87.8
AQ	80	2	2.8	71	97.2
MQ	30	1	1.3	74	98.7
QN	800	19	25.7	55	74.3
LUM	150	0	0	73	100
PPQ	135	0	0	73	100
DHA	12	0	0	69	100

AQ amodiaquine, CQ chloroquine, DHA dihydroartemisinin, LUM lumefantrine, MQ mefloquine, PPQ piperazine, QN quinine

isolates, as defined in the methods section, were 12.2% (9/74) for CQ, 25.7% (19/74) for QN, 2.7% (2/73) for AQ, and 1.3% (1/75) for MQ. No resistant *P. falciparum* isolates to LUM, DHA, and PPQ were found in this study (Table 2).

A strong positive correlation was observed between the IC₅₀ values for QN and AQ (*r* = 0.80; *r*² = 0.64, *p* < 0.0001), suggesting an ex-vivo cross-resistance among these drugs. Moderate positive correlations were found between the IC₅₀ values of CQ vs AQ (*r* = 0.57, *p* < 0.0001), CQ vs QN (*r* = 0.49, *p* < 0.0001), DHA vs MFQ (*r* = 0.35, *p* = 0.002), LUM vs MQ (*r* = 0.5, *p* < 0.0001), DHA vs PPQ (*r* = 0.43, *p* < 0.0001), and MQ vs PPQ (*r* = 0.45, *p* < 0.0001). A weak positive correlation was observed between CQ vs MQ (*r* = 0.29; *r*² = 0.08, *p* = 0.008) and DHA vs LUM (*r* = 0.30, *p* = 0.012) (Table 3). In contrast, negative correlations were observed between IC₅₀ values of DHA vs QN (*r* = − 0.35, *p* = 0.002), LUM vs QN (*r* = − 0.36,

Table 3 Correlation between anti-malarial drug 50% maximal inhibitory concentration values

Drug pairing	Number of experiments	Correlation coefficient (<i>r</i>)	<i>p</i> value
AQ vs CQ	74	0.517	< 0.001
AQ vs DHA	68	- 0.28	0.01
AQ vs LUM	73	- 0.45	< 0.001
AQ vs MQ	74	- 0.04	0.72
AQ vs PPQ	74	0.07	0.55
AQ vs QN	73	0.80	< 0.001
CQ vs DHA	70	0.096	0.41
CQ vs LUM	73	0.036	0.76
CQ vs MQ	76	0.29	0.008
CQ vs PPQ	75	0.21	0.06
CQ vs QN	73	0.49	< 0.001
DHA vs LUM	67	0.30	0.01
DHA vs MQ	70	0.35	0.002
DHA vs PPQ	68	0.43	< 0.0001
DHA vs QN	67	- 0.35	0.002
LUM vs MQ	74	0.5	< 0.001
LUM vs PPQ	73	0.11	0.34
LUM vs QN	71	- 0.36	0.002
MQ vs PPQ	75	0.45	< 0.001
MQ vs QN	73	- 0.02	0.87
PPQ vs QN	72	- 0.10	0.37

Significant correlation values are highlighted in bold

AQ amodiaquine, CQ chloroquine, DHA dihydroartemisinin, LUM lumefantrine, MQ mefloquine, PPQ piperazine, QN quinine

$p = 0.002$), DHA vs AQ ($r = - 0.28$, $p = 0.01$), and AQ vs LUM ($r = - 0.45$, $p < 0.0001$).

4 Discussion

Ex-vivo assessment of the susceptibility of malaria parasites to antimalarial drugs remains an important component in the surveillance of the efficacy of anti-malarial drugs. As this method is largely independent on clinical factors, it provides information that complements clinical assessment of drug efficacy. The SYBR[®] Green method for assessing the outcome of the ex-vivo drug sensitivity test was revalidated and used to assess the responses of *P. falciparum* clinical isolates to seven anti-malarial drugs in Mali. To the best of our knowledge, this is the first study reporting the use of SYBR[®] Green method in Mali. We confirmed that this method is easy to use, reliable, and cheaper, as evidenced by the quick and successful evaluation of ex-vivo sensitivity *P. falciparum* isolates from Kéniéroba to a comprehensive panel of anti-malarial drugs, namely CQ, QN, AQ, MQ, LUM, DHA, and PPQ.

The prevalence of *P. falciparum* parasites with reduced susceptibility to QN was 25.7% in Kéniéroba. This prevalence was higher than those reported in Ghana, in 2011 (19.4% [6/31]) [28] and Senegal in 2017 (9.7% [3/31]) [29]. Globally, similar sensitivity of *P. falciparum* isolates to QN were found (mean IC₅₀ = 264.97 nM; 95% CI 39.90–1567.38) as in a study conducted in Cote d'Ivoire in 2008 (mean IC₅₀ = 272.12 nM) by Touré et al. [30]. In contrast, a higher IC₅₀ mean value for QN (355.37 nM) was reported by Kwansa-Bentum et al. in 2011 in Ghana [28]. To treat complicated malaria, QN by the intravenous route still remains the third-line treatment in Mali. Oral QN is also a recommended drug to treat uncomplicated malaria during the first trimester of pregnancy in Mali. As a reduced susceptibility in vitro is an early warning of impending resistance, a very close follow-up of the resistance to QN is necessary in Mali.

In the present study, 12.2% of the *P. falciparum* isolates exhibited a reduced susceptibility to CQ (IC₅₀ ≥ 100 nM). In 2004–6, a higher proportion of CQ-resistant *P. falciparum* isolates (60–69%) was found in Mali [31]; however, we do not know if current circulating *P. falciparum* isolates are now sensitive to CQ because of the reduced selective pressure on that drug after the introduction of ACT in Sub-Saharan Africa. Reports from Malawi and Kenya suggest that likelihood [32–34]. This trend was also found in studies conducted by French scientists, using samples from patients with malaria from Senegal, Mali, Ivory Coast, and Cameroon [35, 36]. The data indicate that it may be possible to re-introduce CQ as the first-line treatment of malaria in Mali because CQ is easily manufactured locally at a very low cost. Hence, we recommend routine monitoring of CQ resistance despite the widespread use of ACT.

Only 2.7% of the parasite isolates exhibited a reduced ex-vivo susceptibility to AQ (IC₅₀ ≥ 80 nM). This was interesting because AQ is one of the most commonly associated molecules with artemisinin in the ACT used in Mali. Higher prevalence of AQ-resistant parasites have been reported in 2017 in Senegal (28.1%) [29] and in India (8%) [37]. A strict adherence to the NMCP's malaria treatment guideline in conjunction with the World Health Organization guidelines [38] may considerably reduce the spread of the AQ resistance in Africa.

In our study, *P. falciparum* isolates displayed a very high sensitivity to MQ, LUM, DHA, and PPQ similar to previous findings in Mali [39] and in Senegal [40]. To determine the most suitable antimalarial therapeutic combinations and to prevent the emergence of resistant parasites in the study site, correlations between IC₅₀ values of the *P. falciparum* isolate were assessed. A positive correlation was found between the IC₅₀ values for: CQ vs AQ, CQ vs QN, and CQ vs MQ. The variation in the ex-vivo response to AQ (32%), QN (24%), and MQ (8%) could be explained by the variation in

response to CQ. However, the values of correlation coefficient remain low with no possible threat of a cross-resistance between CQ and AQ, CQ and QN, or CQ and MQ.

The artemisinin derivatives are recommended to be associated with LUM, AQ, PPQ, and MQ. In this study, we demonstrated a positive correlation between DHA vs LUM, DHA vs PPQ, and DHA vs MQ. Twelve percent, 9%, and 18% of the variation in the ex-vivo response to MQ, LUM, and PPQ could be explained by variations in response to DHA. The values of the coefficient of determination between DHA vs MQ, DHA vs LUM, and DHA vs PPQ were too low to explain the cross-resistance. A cross-resistance between LUM and artemisinin derivatives has been observed previously [41]. There was a significantly positive correlation between the ex-vivo activity of MQ and PPQ ($r=0.45$; $r^2=0.20$, $p<0.0001$). A significant association between MQ and PPQ was already estimated with parasite isolates from Africa ($r=0.45$) [42]. The correlation between MQ and PPQ might be partly explained by their similar mode of action. However, the values of the coefficient of determination remain low to fear cross-resistance between MQ and PPQ. A strong significant positive correlation was observed between QN and AQ ($r=0.80$; $r^2=0.64$, $p<0.0001$). Sixty-four percent (64%) of the variation in the ex-vivo response to QN could be explained by variations in response to AQ. The high value of the correlation coefficient raises a concern about a potential threat of cross-resistance between QN and AQ. From these data, we suggest that common mechanisms of action of these two drugs may explain the positive correlation between them, suggesting that parasites that have reduced susceptibility to QN would also be less susceptible to AQ.

Cross-resistances between CQ-AQ, CQ-QN, and QN-AQ have already been reported in previous studies [43–45]. Cross-resistance is thought to occur when a drug confers resistance to other drugs that have a similar mode of action or belong to the same chemical group. Cross-resistance affects essentially drugs belonging to the same chemical family or drugs with similar modes of action or modes of transportation [46–49]. Theoretically, parasites are more prone to be sensitive to a molecule whose IC_{50} is inversely proportional to that of a molecule the parasites display resistance. Data from our study revealed light to moderate, but significant negative correlations between sensitivities of *P. falciparum* isolates to AQ and DHA ($r=-0.28$, $p=0.01$), DHA and QN ($r=-0.35$, $p=0.01$), and AQ and LUM ($r=-0.45$, $p<0.001$). The negative correlation observed between DHA and AQ implies that the use of artemisinin in combination with AQ is very relevant in Mali. Our findings are reassuring considering the emergence of resistance to artemisinin derivatives [50–53].

5 Conclusions

Plasmodium falciparum isolates were highly sensitive to DHA, LUM, and PPQ. In addition, few isolates showed a reduced sensitivity to AQ and MQ (only one isolate). A much lower sensitivity to QN was also detected. Our data confirm an increasing trend in the sensitivity of the *P. falciparum* field isolate to CQ in Mali. This study provides valuable data that may considerably contribute to a better clinical management of patients with malaria and guide the decision-making process for an effective national strategy for malaria treatment.

Acknowledgements We thank the parents, guardians, and children who participated in this study, and the technical, clinical, and nursing staff for assistance. We are grateful to many colleagues at the Malaria Research and Training Center for providing critical reviews of the manuscript that have helped improve the content.

Author Contributions KT, SASD, and MD provided the study design. KT, SASD, IS, DSK, BK, and MD provided the sample and collected the data. KT and SASD performed the data analysis. KT wrote the manuscript. KT, SASD, DD, SD, MS, SB, MAG, SD, and MD edited and reviewed the manuscript.

Data Availability All data generated or analyzed during this study are included in the published article.

Compliance with Ethical Standards

Funding This study is supported by the University of Sciences, Techniques, and Technologies of Bamako (USTTB).

Conflict of interest Karim Traoré, Seidina A.S. Diakité, Sekou Bah, Drissa S. Konaté, Djeneba Dabita, Ibrahim Sanogo, Modibo Sangaré, Souleymane Dama, Bourama Keita, Mory Doumbouya, Merepen A. Guindo, Seydou Doumbia, and Mahamadou Diakité have no conflicts of interest that are directly relevant to the content of this article.

Ethics approval The study was approved by the Ethics Committee of the Faculty of Medicine and Pharmacy of the University of Sciences, Techniques, and Technologies of Bamako (USTTB), Mali.

Consent to participate All study participants signed a written consent or assent (for children) form to participate in this study.

Consent for publication All authors read and approved the final manuscript.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission

directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

References

- World Health Organization. World malaria report 2015. Geneva: World Health Organization; 2015.
- Packard RM. The origins of antimalarial-drug resistance. *N Engl J Med*. 2014;371(5):397–9.
- Wongsrichanalai C, et al. Epidemiology of drug-resistant malaria. *Lancet Infect Dis*. 2002;2(4):209–18.
- Spencer HC. Drug-resistant malaria: changing patterns mean difficult decisions. *Trans R Soc Trop Med Hyg*. 1985;79(6):748–58.
- Wernsdorfer WH, Payne D. The dynamics of drug resistance in *Plasmodium falciparum*. *Pharmacol Ther*. 1991;50(1):95–121.
- World Health Organization. Guidelines for the treatment of malaria. Geneva: World Health Organization; 2011.
- Phyo AP, et al. Emergence of artemisinin-resistant malaria on the western border of Thailand: a longitudinal study. *Lancet*. 2012;379(9830):1960–6.
- Amaratunga C, et al. Artemisinin-resistant *Plasmodium falciparum* in Pursat province, western Cambodia: a parasite clearance rate study. *Lancet Infect Dis*. 2012;12(11):851–8.
- Ariey F, et al. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature*. 2014;505(7481):50–5.
- Tacoli C, et al. Artemisinin resistance-associated K13 polymorphisms of *Plasmodium falciparum* in Southern Rwanda, 2010–2015. *Am J Trop Med Hyg*. 2016;95(5):1090–3.
- Zhang J, et al. In vitro susceptibility of *Plasmodium falciparum* isolates from the China–Myanmar border area to artemisinins and correlation with K13 mutations. *Int J Parasitol Drugs Drug Resist*. 2019;10:20–7.
- OMS. Rapport mondial sur l'efficacité des médicaments antipaludiques et la pharmacorésistance: 2000–2010. Genève, S.O.m.d.l.s., rapport mondial sur l'efficacité des médicaments antipaludiques et la pharmacorésistance: 2000–2010. Genève: Organisation mondiale de la santé; 2011.
- Walliker D, et al. Genetic analysis of the human malaria parasite *Plasmodium falciparum*. *Science*. 1987;236(4809):661–6.
- Wellems TE, et al. Chloroquine resistance not linked to *mdr*-like genes in a *Plasmodium falciparum* cross. *Nature*. 1990;345(6272):253–5.
- Smilkstein M, et al. Simple and inexpensive fluorescence-based technique for high-throughput antimalarial drug screening. *Antimicrob Agents Chemother*. 2004;48(5):1803–6.
- Johnson JD, et al. Assessment and continued validation of the malaria SYBR Green I-based fluorescence assay for use in malaria drug screening. *Antimicrob Agents Chemother*. 2007;51(6):1926–33.
- Brasseur P, et al. Emergence of mefloquine-resistant malaria in Africa without drug pressure. *Lancet*. 1990;336(8706):59.
- Pradines B, et al. Reduced in vitro susceptibility to artemisinin derivatives associated with multi-resistance in a traveller returning from South-East Asia. *Malar J*. 2011;10:268.
- Reynes JM, et al. In vitro responses of *Plasmodium falciparum* isolates to five antimalarial drugs in French Guiana during 1994 and 1995. *Mem Inst Oswaldo Cruz*. 1997;92(2):251–2.
- Dedet JP, et al. In vitro activity of various antimalarials (chloroquine, amodiaquine, quinine and mefloquine) against 32 isolates of *Plasmodium falciparum* in French Guiana. *Bull Soc Pathol Exot Filiales*. 1988;81(1):88–93.
- Munghthin M, et al. Association between the *pfmdr1* gene and in vitro artemether and lumefantrine sensitivity in Thai isolates of *Plasmodium falciparum*. *Am J Trop Med Hyg*. 2010;83(5):1005–9.
- Fall B, et al. Ex vivo susceptibility of *Plasmodium falciparum* isolates from Dakar, Senegal, to seven standard anti-malarial drugs. *Malar J*. 2011;10:310.
- Pascual A, et al. Multinormal in vitro distribution of *Plasmodium falciparum* susceptibility to piperazine and pyronaridine. *Malar J*. 2015;14:49.
- Bacon DJ, et al. Comparison of a SYBR Green I-based assay with a histidine-rich protein II enzyme-linked immunosorbent assay for in vitro antimalarial drug efficacy testing and application to clinical isolates. *Antimicrob Agents Chemother*. 2007;51(4):1172–8.
- Afari EA, et al. In vitro responses of *P. falciparum* parasites to chloroquine, amodiaquine and quinine in two ecological zones in Ghana. *Cent Afr J Med*. 1993;39(7):136–40.
- Le Nagard H, et al. Online analysis of in vitro resistance to anti-malarial drugs through nonlinear regression. *Comput Methods Progr Biomed*. 2011;104(1):10–8.
- Kaddouri H, et al. Assessment of the drug susceptibility of *Plasmodium falciparum* clinical isolates from Africa by using a *Plasmodium* lactate dehydrogenase immunodetection assay and an inhibitory maximum effect model for precise measurement of the 50-percent inhibitory concentration. *Antimicrob Agents Chemother*. 2006;50(10):3343–9.
- Kwansa-Bentum B, et al. *Plasmodium falciparum* isolates from southern Ghana exhibit polymorphisms in the SERCA-type PfATPase6 though sensitive to artesunate in vitro. *Malar J*. 2011;10:187.
- Diawara S, et al. Confirmation of *Plasmodium falciparum* in vitro resistance to monodesethylamodiaquine and chloroquine in Dakar, Senegal, in 2015. *Malar J*. 2017;16(1):118.
- Touré AO, et al. In vitro susceptibility of *P. falciparum* isolates from Abidjan (Cote d'Ivoire) to quinine, artesunate and chloroquine. *Sante*. 2008;18(1):43–7.
- Kaddouri H, et al. Baseline in vitro efficacy of ACT component drugs on *Plasmodium falciparum* clinical isolates from Mali. *Int J Parasitol*. 2008;38(7):791–8.
- Kublin JG, et al. Reemergence of chloroquine-sensitive *Plasmodium falciparum* malaria after cessation of chloroquine use in Malawi. *J Infect Dis*. 2003;187(12):1870–5.
- Laufer MK, et al. Return of chloroquine antimalarial efficacy in Malawi. *N Engl J Med*. 2006;355(19):1959–66.
- Mwai L, et al. Chloroquine resistance before and after its withdrawal in Kenya. *Malar J*. 2009;8:106.
- Gharbi M, et al. Longitudinal study assessing the return of chloroquine susceptibility of *Plasmodium falciparum* in isolates from travellers returning from West and Central Africa, 2000–2011. *Malar J*. 2013;12:35.
- Lu F, et al. Return of chloroquine sensitivity to Africa? Surveillance of African *Plasmodium falciparum* chloroquine resistance through malaria imported to China. *Parasit Vectors*. 2017;10(1):355.
- Sharma S, et al. Correlation of in vitro sensitivity of chloroquine and other antimalarials with the partner drug resistance to *Plasmodium falciparum* malaria in selected sites of India. *Indian J Med Microbiol*. 2017;35(4):485–90.
- World Health Organization. WHO briefing on malaria treatment guidelines and artemisinin monotherapies. Geneva: World Health Organization; 2006.
- Dama S, et al. Reduced ex vivo susceptibility of *Plasmodium falciparum* after oral artemether-lumefantrine treatment in Mali. *Malar J*. 2017;16(1):59.
- Fall B, et al. *Plasmodium falciparum* susceptibility to anti-malarial drugs in Dakar, Senegal, in 2010: an ex vivo and drug resistance molecular markers study. *Malar J*. 2013;12:107.

41. Pradines B, et al. In vitro activities of benflumetol against 158 Senegalese isolates of *Plasmodium falciparum* in comparison with those of standard antimalarial drugs. *Antimicrob Agents Chemother.* 1999;43(2):418–20.
42. Robert MG, et al. Absence of a high level of duplication of the plasmepsin II gene in Africa. *Antimicrob Agents Chemother.* 2018;62(11):e00374–e418.
43. Ringwald P, et al. In vitro culture and drug sensitivity assay of *Plasmodium falciparum* with nonserum substitute and acute-phase sera. *J Clin Microbiol.* 1999;37(3):700–5.
44. Pradines B, et al. In-vitro activity of pyronaridine and amodiaquine against African isolates (Senegal) of *Plasmodium falciparum* in comparison with standard antimalarial agents. *J Antimicrob Chemother.* 1998;42(3):333–9.
45. Ouedraogo JB, et al. In vitro sensitivity of *Plasmodium falciparum* to halofantrine compared with chloroquine, quinine and mefloquine in the region of Bobo-Dioulasso, Burkina Faso (West Africa). *Trop Med Int Health.* 1998;3(5):381–4.
46. Ringwald P, Bickii J, Basco LK. In vitro activity of dihydroartemisinin against clinical isolates of *Plasmodium falciparum* in Yaounde, Cameroon. *Am J Trop Med Hyg.* 1999;61(2):187–92.
47. East African Network for Monitoring Antimalarial Treatment (EANMAT). Monitoring antimalarial drug resistance within National Malaria Control Programmes: the EANMAT experience. *Trop Med Int Health.* 2001;6(11):891–8.
48. Randrianarivelojosia M, et al. In-vitro sensitivity of *Plasmodium falciparum* to chloroquine, halofantrine, mefloquine and quinine in Madagascar. *East Afr Med J.* 2002;79(5):237–41.
49. Yavo W, et al. In vitro susceptibility of *Plasmodium falciparum* to monodesethylamodiaquine, quinine, mefloquine and halofantrine in Abidjan (Cote d'Ivoire). *Afr Health Sci.* 2010;10(2):111–6.
50. Na-Bangchang K, et al. Declining in efficacy of a three-day combination regimen of mefloquine-artesunate in a multi-drug resistance area along the Thai-Myanmar border. *Malar J.* 2010;9:273.
51. Woitsch B, et al. Susceptibility to chloroquine, mefloquine and artemisinin of *Plasmodium vivax* in northwestern Thailand. *Wien Klin Wochenschr.* 2007;119(19–20 Suppl. 3):76–82.
52. Noedl H, et al. Artemisinin resistance in Cambodia: a clinical trial designed to address an emerging problem in Southeast Asia. *Clin Infect Dis.* 2010;51(11):e82–e8989.
53. Spring MD, et al. Dihydroartemisinin-piperaquine failure associated with a triple mutant including kelch13 C580Y in Cambodia: an observational cohort study. *Lancet Infect Dis.* 2015;15(6):683–91.