

Original Article

Effect of prostaglandins E₂ and F_{2α} on granulosa cell apoptosis in goat ovarian follicles

Sharma, A. K.¹ and Sharma, R. K.^{2*}

¹Ph.D. Student in Reproductive Physiology, Reproductive Physiology Laboratory, Department of Zoology, Kurukshetra University, Kurukshetra-136119, India; ²Reproductive Physiology Laboratory, Department of Zoology, Kurukshetra University, Kurukshetra-136119, India

*Correspondence: R. K. Sharma, Reproductive Physiology Laboratory, Department of Zoology, Kurukshetra University, Kurukshetra-136119, India. E-mail: rkshukz@gmail.com

(Received 14 Oct 2019; revised version 10 Dec 2019; accepted 14 Jan 2020)

Abstract

Background: The goat is a multi-purpose animal producing meat, milk, hide, fiber and manure, and hence plays a significant role in providing supplementary income and livelihood to the huge number of asset-poor ranchers and landless workers of rural India. **Aims:** The present study is aimed to investigate the *in vitro* effects of prostaglandins E₂ (PGE₂) and F_{2α} (PGF_{2α}) on ovarian granulosa cells of a goat. **Methods:** The healthy, slightly atretic and atretic antral follicles of goats with diameter ranging from 3-8 mm were cultured with PGE₂ and PGF_{2α} (1.0 µg/ml) along with control for 24 h. Histomorphological analysis was done in order to study the prostaglandins induced changes in granulosa cells of all the three categories of follicles. The acridine orange (AO) and methylene blue (MB) staining were used to study the apoptotic index in all the three categories of follicles and the collected data were analyzed by ANOVA with Duncan post Hoc test. **Results:** Prostaglandins E₂ revealed positive while PGF_{2α} showed a negative effect on the rescue of apoptosis in granulosa cells. The PGE₂ treated follicles revealed a reduction in the attributes of apoptosis in granulosa cells while PGF_{2α} showed an increased in the apoptotic characteristics. **Conclusion:** The present study depicted that granulosa cell viability *in vitro* is dependent on the continuous supply of survival and growth factors and prostaglandin of E series synthesis is a crucial step in the suppression of granulosa cell apoptosis while that of F series induces apoptosis.

Key words: Apoptosis, Granulosa cells, Prostaglandins, Ovaries

Introduction

Follicular atresia in the ovary is a degenerative process that involves apoptosis of granulosa cells and is regulated by gonadotropins, steroids, cytokines and growth factors (Manchanda *et al.*, 2001). Most of the follicles (<5 mm in diameter) undergo atresia and regression during the follicular development and the follicles that attain >5 mm in diameter are destined to ovulate in goats and sheep. The regulation of follicular atresia by gonadotropins is well understood but little is known about its control by non-gonadotropin signals (Manchanda *et al.*, 2001).

Prostaglandins are cyclic carboxylic acids, derived from arachidonic acid or closely related fatty acids which have potent and diverse physiological actions. The role of prostaglandins in the governance of follicular function of the ovary was first speculated based on sustaining that prohibition of prostaglandin synthesis by aspirin and indomethacin results in blocking of ovulation in rats (Armstrong and Grinwich, 1972; Orcyzk and Behrman, 1972). Similar findings were further observed in other species, including rabbits (Grinwich *et al.*, 1972; O'Grady *et al.*, 1972), mice (Lau *et al.*, 1974), goldfish (Stacey and Pandey, 1975), rhesus and marmoset monkeys (Wallach *et al.*, 1975; Maia *et al.*, 1978), and pigs (Ainsworth *et al.*, 1979). Another clue for the role of

prostaglandins at the follicular level was supported by the findings that intra-follicular levels of prostaglandins of both the series i.e. E and F, elevated considerably in these species shortly before ovulation (Armstrong *et al.*, 1974; Yang *et al.*, 1974; Ainsworth *et al.*, 1975; Bauminger and Lindner, 1975; Tsang *et al.*, 1979). Triebwasser *et al.* (1983) already demonstrated that prostaglandins E₂ (PGE₂) can mimic the action of follicle stimulating hormone (FSH) on granulosa cells at the early stages of development. Prostaglandins E₂ can trigger progesterone production in rat granulosa cells (Hiller *et al.*, 1978). Thus prostaglandins are effective lipid compounds bearing distinct hormone-like effects, and are synthesized in many tissues and play various female reproductive functions such as steroidogenesis, tissue remodeling, neovascularization of the luteinizing follicle and modulation of the oocyte and ovulation. They are essential for follicle rupture and oocyte release in most of the mammals including bovine, ovine, and equine, etc. The reproductive physiology of goat concerning prostaglandins is less understood compared to cattle and pigs. Hence, it is inescapable to investigate practically the effects of prostaglandins on apoptosis in granulosa cells of goat ovarian tissue. Moreover, the intra-ovarian role of PGs in apoptosis has not been addressed so far in small ruminants.

Materials and Methods

Collection and culture of ovarian follicles *in vitro*

No animals were personally handled or sacrificed for our research purpose, only the reproductive tissues (ovaries only) of Jamnapari breed of goat (*Capra hircus*) were collected from government-approved abattoirs of Chandigarh (30.7333° N, 76.7794° E) in normal saline at 4°C.

Then the follicles were manually separated with fine forceps. The healthy antral follicles, slightly atretic follicles and atretic antral follicles having a diameter ranging from 3-8 mm were selected on a morphometric basis including color, the turbidity of follicular fluid and vascularity (Sharma and Bhardwaj, 2009). The follicles were then cultured with prostaglandins PGE₂ product code number 14010 and Prostaglandins F_{2α} (PGF_{2α}) product code number 16010 (purchased from Cayman Chemical Co. 1180E, Ellsworth Road, Ann Arbor, Michigan 48108, USA) with the concentration of 1.0 µg/ml in DMEM (Dulbecco's modified eagle medium), supplemented with antibiotics (HiMedia) 200-unit having concentration of penicillin 100IU/MI and streptomycin 100 IU/MI in CO₂ incubator (5% CO₂, 95% humidity, 38°C) for 24 h as per experimental layout (Fig. 1).




Fig. 1: Experimental layout of the study

Histomorphological analysis

Histomorphological analysis was done according to the method of Pearse (1968). The cultured follicles after prostaglandin treatment were fixed in Bouin's fixative for 24 h. After fixation washing of follicles under tap water was done for 2 h and then dehydration through a graded series of ethanol (30-100%) followed by embedding in paraffin wax at 60°C. Similar blocks were prepared, then trimmed and sectioned serially at 5 µm thickness. After this, the trimmed sections were stretched on albumin coated slides and air-dried for staining. Dewaxing of the sections was done by xylene followed by hydration and dehydration using a series of alcohol (30-100%) for staining with hematoxylin and eosin (H&E) to study the histomorphological characters of

apoptosis in granulosa cells of ovarian follicles.

Preparation of granulosa cell suspension

The treated follicles were aspirated with the help of a 20-gauge needle on a 2 ml syringe containing phosphate buffer saline at pH 7.4. The cumulus-oocyte complexes were removed with the help of micropipettes using stereo-microscopes. The granulosa cell aggregates from the aspirates were allowed to centrifuge at 2000 rpm for 5 min. Then the supernatant was removed and the pellet formed was resuspended in phosphate buffer saline (PBS). This step was repeated thrice to remove the debris.

Apoptotic assay

Apoptotic analysis was done according to the method of Broaddus *et al.* (1996) and Dave *et al.* (2001). Granulosa cells apoptosis was evaluated by acridine orange (AO) staining and methylene blue (MB) staining, analyzed under the fluorescent and light microscope (Olympus, Japan). The cell suspension prepared from treated follicles was mixed with an equal quantity of AO and MB working solution which was prepared by dissolving 1 µL each of AO and MB in 1 ml PBS. Normal cells were identified by green fluorescence and unstained whereas apoptotic cells showed red fluorescence and blue stain. Quantification of healthy and apoptotic granulosa cells was done to measure percentage apoptosis.

Statistical analysis was done by applying ANOVA with Duncan Post Hoc test. Significance is implied at $P < 0.05$.

Results

Histomorphological analysis

The healthy antral follicles of control group cultured for 24 h were composed of granulosa cells of normal contour having clear morphology. The cytoplasm was finely granulated and lightly stained; the nucleus was centrally located and spherical in outline. The cells reacted positively with eosin stain and are spherical (Fig. 2A). Slightly atretic antral follicles of the same group revealed the apoptotic granulosa cells lodged in between healthy cells. These apoptotic cells were characterized by their small size, pyknotic nucleus, dark staining and marginated chromatin (Fig. 2B). In atretic antral follicles of the same group cultured for the same duration, the apoptotic granulosa cells were more in number with densely stained chromatin material, fragmented nuclei, and hyalinized cytoplasm with varying degree of vacuolization and membrane blebbing was observed (Fig. 2C). In most of the degenerating cells, the cytoplasmic contents were condensed and clumped. Heterochromatization and chromatolysis of granulosa cells were observed. The cell membrane was loose, wrinkled and folded (Fig. 2C).

Prostaglandins E₂ (1.0 µg/ml) treated healthy and slightly atretic antral follicles cultured for 24 h revealed chiefly healthy granulosa cells with intact cell

membranes and maintained cellular integrity. The appearance of pyknotic nuclei, degenerative granulosa cells were minimum in number (Figs. 2D and E). Prostaglandins E₂ treated atretic antral follicles cultured for 24 h demonstrated marked features of decreasing the attributes of apoptosis where most of the granulosa cells remained healthy with maintained cellular integrity. The appearance of pyknotic nuclei, vacuoles and condensed chromatin and degenerating granulosa cells were minimum (Fig. 2F). Prostaglandins E₂ helped in the suppression of apoptosis by decreasing the incidence of the occurrence of apoptotic characters within granulosa cells. While the PGF_{2α} (1.0 μg/ml) treated healthy and

slightly atretic follicles cultured for 24 h reported an increasing incidence of characteristics of apoptosis within granulosa cells (Figs. 2G-I). Whereas the number of degenerative granulosa cells was more when atretic antral follicles were treated with PGF_{2α} with the same dose for the same duration (Figs. 2G-I).

Fluorescence assay for detection and quantification of apoptosis

Moreover, the quantification of apoptosis revealed that PGE₂ decreased the frequency of apoptosis in granulosa cells (Table 1). Scrutiny of apoptosis using fluorescent dye i.e. AO is based upon the differential

Table 1: Apoptotic index in various experimental groups using AO and MB dye

Stains	Treatments	Healthy (%)	Slightly atretic (%)	Atretic (%)
AO	Control (n=10)	13.49 ± 1.89 ^b	18.79 ± 1.47 ^b	49.40 ± 3.87 ^b
	PGE ₁ (n=10)	3.81 ± 2.34 ^c	8.29 ± 3.24 ^c	19.69 ± 3.32 ^c
	PGF _{2α} (n=10)	25.17 ± 2.91 ^a	52.19 ± 2.27 ^a	89.21 ± 3.60 ^a
MB	Control (n=10)	18.53 ± 3.25 ^b	25.08 ± 1.85 ^b	62.63 ± 2.64 ^b
	PGE ₁ (n=10)	5.48 ± 1.70 ^c	16.56 ± 1.86 ^b	24.22 ± 2.37 ^c
	PGF _{2α} (n=10)	31.50 ± 2.03 ^a	67.59 ± 4.14 ^a	92.69 ± 0.94 ^a

Values are expressed as mean±SEM. ANOVA with Duncan Post Hoc test was used for analysis; different letters indicate significant differences among means of treatment at each duration (P<0.05) after Duncan's test. AO: Acridine orange, and MB: Methylene blue




Fig. 2: Histological appearance of granulosa cells in ovarian antral follicles showing morphological characteristics of apoptosis after PGE₂ treatment at 1.0 μg/ml (D, E, F) and PGF_{2α} treatment at 1.0 μg/ml (G, H, I) in comparison with control post haematoxylin and eosin staining (A, B, C) after 24 h at ×1000. The photograph reveals decreased and increased incidence of apoptotic characteristics. * Represents normal healthy cells with intact membrane, arrow head: Darkly stained pyknotic nuclei, m: Marginated chromatin, f: Fragmented chromatin, and v: Vacuolated cell




Fig. 3: Fluorescent photographs of granulosa cells of ovarian antral follicles after PGE₂ treatment at 1.0 µg/ml (D, E, F) and PGF_{2α} treatment at 1.0 µg/ml (G, H, I) in comparison with control (A, B, C) stained with fluorescent dye AO depicting apoptotic granulosa cells with bright red fluorescence and normal live cells with green fluorescence after 24 h culture at ×400




Fig. 4: Microphotographs showing apoptosis in granulosa cells of ovarian antral follicles using MB staining after PGE₂ treatment at 1.0 µg/ml (D, E, F) and PGF_{2α} treatment at 1.0 µg/ml (G, H, I) in comparison with control (A, B, C) depicting apoptotic granulosa cells with blue color and normal live cells with no color after 24 h culture at ×400

uptake of dye by live and apoptotic cells. Healthy cells with intact cell membranes and integrity appear green in color because of AO, whereas the apoptotic cells appear red due to the intercalation of AO dye. Prostaglandins E₂ supplementation decreased the percentage of granulosa cells showing red fluorescence in healthy, slightly and atretic antral follicles. Whereas PGF_{2α} increased the percentage of granulosa cells showing red color in healthy, slightly atretic and atretic antral follicles (Figs. 3A-I).

Methylene blue staining

Furthermore, the MB staining of cells clearly showed that PGE₂ has suppressed apoptosis in granulosa cells (Table 1). Analysis of granulosa cell apoptosis is based upon the differential staining of granulosa cells; the normal healthy cells appeared transparent whereas apoptotic cells appeared blue as dead cells took up the stain. Prostaglandins E₂ treatment decreased the percentage of granulosa cells showing blue color whereas, in PGF_{2α} treatment, most of the granulosa cells appeared blue (Figs. 4A-I).

Discussion

Although PG synthesis in granulosa cells during ovarian follicular development is regulated by a complex of interactions of different factors, the physiological role of these eicosanoids in the control of ovarian cells function by these factors is unknown. The present results illustrate that notable change in the responsiveness of granulosa cells when cultured with PGs for 24 h. The results of the present study revealed that PGE₂ suppressed the apoptosis in granulosa cells as already reported by Manchanda *et al.* (2001), who studied the role of prostaglandins in the suppression of apoptosis in hen granulosa cells and demonstrated that a prostaglandin synthesis is a necessary event in the suppression of granulosa cell apoptosis. Histomorphological analysis of PGE₂ and PGF_{2α} supplemented granulosa cells revealed that PGE₂ reduced the frequency of apoptotic attributes within the granulosa cells of atretic antral follicles while PGF_{2α} upregulated the process of apoptosis. The PGF_{2α} activates the apoptotic signaling cascades have already been documented in corpus luteum (Yadav *et al.*, 2005). Prostaglandins E₂ suppressed the rate of apoptosis in healthy, slightly atretic and atretic follicles. Though the mechanism by which PGE₂ mitigates the apoptotic action is not elucidated, some hypothetical roles of prostaglandin in ovarian follicular regulation have already been suggested by Armstrong (1981).

Growing theca cells release PGE₂, which stimulates the production of cyclic adenosine monophosphate (cAMP) in theca and granulosa cells. The cAMP subsequently initiates the growth of follicles by exerting gonadotropin like action on granulosa cells, which do not yet possess receptors for pituitary gonadotropins. Prostaglandins F_{2α}, of follicular origin, may initiate or participate in the processes of luteolysis and atresia,

through interactions of specific receptors (Armstrong, 1981).

Our findings are consistent with the study of Li *et al.* (1995), who documented PGs as central elements in cell signaling for mitogenesis induced by growth factors, and in oncogenic transformation. An increased PG production is associated with many tumor cells and is believed to control tumor growth and metastasis. Further, he demonstrated, compared to PGs of the F series, PGE₁ and PGE₂ are more effective in persuading transforming growth factor- α (TGF- α) induced DNA synthesis in hen granulosa cells. Prostaglandins of I series were also shown to increase in the cAMP production which was the main indicator of cell division in cultured granulosa cells *in vitro* but as compared to E series PGs of I series are lower stimulator for cAMP production (Goff *et al.*, 1978). The goat granulosa cells are possibly regulated by a similar mechanism for multiplication or regression.

Steroidogenesis in granulosa cells requires the continuous supply of stimulating factors, and PGE₂ is one of them; granulosa cells become more sensitive to PGE₂ in the presence of FSH (Goff *et al.*, 1983). Bowolaksono *et al.* (2008) who observed the production of PGs and their receptors in bovine corpus luteum cells also suggested that PGs have anti-apoptotic roles in bovine luteal steroidogenic cells.

The use of AO differential staining undoubtedly interpreted the apoptosis and DNA damage in granulosa cells. Acridine orange is fluorescent compound which intercalates between the adjacent base pairs of DNA and RNA and enhances its fluorescent intensity (Nafisi *et al.*, 2007). Acridine orange is permeable to cell membrane stain apoptotic cells as red and non-apoptotic cells as green. The AO differential staining techniques aptly assisted in the analysis of PGs suppression or induction of cascades of apoptosis within granulosa cells. In our study, PGE₂ is found to be capable of suppressing apoptosis, which is revealed through fluorescent assay. In PGE₂ administered group, green fluorescence suggestive of live cells is higher as compared to orange-reddish fluorescence (an indicator of apoptosis). These findings endorse the role of PG in follicular growth and development as already documented in rats (Goff and Armstrong, 1983).

In summary, the present study shows that prostaglandin of E series synthesis is a crucial step in the suppression of granulosa cell apoptosis while that of F series has an apoptotic role. The availability of such factors or the capacity of granulosa cells to reciprocate them may be essential determinants of the fate of goat ovarian follicles.

Acknowledgements

A. K. Sharma acknowledges the financial assistance provided in the form of JRF by UGC, New Delhi. The authors are also thankful to the Department of Zoology, Kurukshetra University, Kurukshetra for providing Laboratory facilities. The authors acknowledge the SAP-UGC for financial support.

Conflict of interest

There is no conflict of interest between the authors.

References

- Ainsworth, L; Baker, RD and Armstrong, DT** (1975). Pre-ovulatory changes in follicular fluid prostaglandin F levels in swine. *Prostaglandins*. 9: 915-925.
- Ainsworth, L; Tsang, BK; Downey, BR; Baker, RD; Marcus, GJ and Armstrong, DT** (1979). Effects of indomethacin on ovulation and luteal function in gilts. *Bio. Reprod.*, 21: 401-411.
- Armstrong, DT** (1981). Prostaglandins and follicular functions. *Reproduction*. 62: 283-291.
- Armstrong, DT and Grinwich, DL** (1972). Blockade of spontaneous and LH-induced ovulation in rats by indomethacin, an inhibitor of prostaglandin biosynthesis. *Prostaglandins*. 1: 21-28.
- Armstrong, DT; Moon, YS and Zamecnik, J** (1974). Evidence for a role of ovarian prostaglandins in ovulation. In: Mougald, NR (Ed.), *Gonadotropins and gonadal function*. (1st Edn.), Elsevier Inc., PP: 343-356.
- Bauminger, S and Lindner, HR** (1975). Perioovulatory changes in ovarian prostaglandin formation and their hormonal control in the rat. *Prostaglandins*. 9: 737-751.
- Bowolaksono, A; Nishimura, R; Hojo, T; Sakumoto, R; Acosta, TJ and Okuda, K** (2008). Anti-apoptotic roles of prostaglandin E2 and F2alpha in bovine luteal steroidogenic cells. *Bio. Reprod.*, 79: 310-317.
- Broadus, VC; Yang, L; Scavo, LM; Ernst, JD and Boylan, AM** (1996). Asbestos induces apoptosis of human and rabbit pleural mesothelial cells via reactive oxygen species. *J. Clin. Investig.*, 98: 2050-2059.
- Dave, U; Shousha, S and Westaby, D** (2001). Methylene blue staining: is it really useful in Barrett's esophagus? *Gastro. Endosco.*, 53: 333-335.
- Goff, AK and Armstrong, DT** (1983). Changes in responsiveness of rat granulosa cells to prostaglandin E2 and follicle-stimulating hormone during culture. *Can. J. Physio. Pharmacol.*, 61: 608-613.
- Goff, AK; Zamecnik, J; Ali, M and Armstrong, DT** (1978). Prostaglandin I2 stimulation of granulosa cell cyclic AMP production. *Prostaglandins*. 15: 875-879.
- Grinwich, DL; Kennedy, TG and Armstrong, DT** (1972). Dissociation of ovulatory and steroidogenic actions of luteinizing hormone in rabbits with indomethacin, an inhibitor of prostaglandin biosynthesis. *Prostaglandins*. 1: 89-96.
- Hillier, SG; Zeleznik, AJ and Ross, GT** (1978). Independence of steroidogenic capacity and luteinizing hormone receptor induction in developing granulosa cells. *Endocrinology*. 102: 937-946.
- Lau, IF; Saksena, SK and Chang, MC** (1974). Prostaglandins F and ovulation in mice. *Reproduction*. 40: 467-469.
- Li, J and Tsang, BK** (1995). Prostaglandins mediate the stimulation of deoxyribonucleic acid synthesis by transforming growth factor α in hen granulosa cells during ovarian follicular development. *Bio. Reprod.*, 52: 1050-1058.
- Maia, JrH; Barbosa, I and Coutinho, EM** (1978). Inhibition of ovulation in marmoset monkeys by indomethacin. *Fert. Ster.*, 29: 565-570.
- Manchanda, R; Kim, JM and Tsang, BK** (2001). Role of prostaglandins in the suppression of apoptosis in hen granulosa cells by transforming growth factor alpha. *Reproduction*. 122: 91-101.
- Nafisi, S; Saboury, AA; Keramat, N; Neault, JF and Tajmir-Riahi, HA** (2007). Stability and structural features of DNA intercalation with ethidium bromide, acridine orange and methylene blue. *J. Mol. Struct.*, 827: 35-43.
- O'Grady, JP; Caldwell, BV; Auletta, FJ and Speroff, L** (1972). The effects of an inhibitor of prostaglandin synthesis (indomethacin) on ovulation, pregnancy, and pseudopregnancy in the rabbit. *Prostaglandins*. 1: 97-106.
- Orczyk, GP and Behrman, HR** (1972). Ovulation blockade by aspirin or indomethacin-*in vivo* evidence for a role of prostaglandin in gonadotrophin secretion. *Prostaglandins*. 1: 3-20.
- Pearse, AGE** (1968). *Histochemistry, theoretical and applied: theoretical and applied*. 3rd Edn., Boston, USA.
- Sharma, RK and Bhardwaj, JK** (2009). Ultrastructural characterization of apoptotic granulosa cells in caprine ovary. *J. Micro.*, 236: 236-242.
- Stacey, NE and Pandey, S** (1975). Effects of indomethacin and prostaglandins on ovulation of goldfish. *Prostaglandins*. 9: 597-607.
- Triebwasser, WF; Clark, MR; LeMaire, WJ and Marsh, JM** (1978). Localization and *in vitro* synthesis of prostaglandins in components of rabbit preovulatory Graafian follicles. *Prostaglandins*. 16: 621-632.
- Tsang, BK; Ainsworth, L; Downey, BR and Armstrong, DT** (1979). Pre-ovulatory changes in cyclic AMP and prostaglandin concentrations in follicular fluid of gilts. *Prostaglandins*. 17: 141-148.
- Wallach, EE; de la Cruz, A; Hunt, J; Wright, KH and Stevens, VC** (1975). The effect of indomethacin on HMG-HCG induced ovulation in the rhesus monkey. *Prostaglandins*. 9: 645-658.
- Yadav, VK; Lakshmi, G and Medhamurthy, R** (2005). Prostaglandin F2 α -mediated activation of apoptotic signaling cascades in the corpus luteum during apoptosis involvement of caspase-activated DNase. *J. Bio. Chem.*, 280: 10357-10367.
- Yang, NS; Marsh, JM and LeMaire, WJ** (1974). Post ovulatory changes in the concentration of prostaglandins in rabbit Graafian follicles. *Prostaglandins*. 6: 37-44.