

Clinical trial to test the efficacy of melatonin in COVID-19

Dario Acuña-Castroviejo^{1,2}  | Germaine Escames^{1,2}  | Juan C. Figueira³ |
Pedro de la Oliva^{4,5} | Alberto M. Borobia^{5,6} | Carlos Acuña-Fernández⁷

¹Faculty of Medicine, Department of Physiology, Biomedical Research Center, Health Sciences Technology Park, University of Granada, Granada, Spain

²Ibs.Granada, San Cecilio University Hospital, Granada, Spain

³Intensive Care Medicine Department, La Paz University Hospital, Madrid, Spain

⁴Pediatric Intensive Care Department, La Paz University Hospital, Madrid, Spain

⁵School of Medicine, Universidad Autónoma de Madrid, Madrid, Spain

⁶Clinical Pharmacology Department, Clinical Trial Unit, La Paz University Hospital – IdiPAZ, Madrid, Spain

⁷Anaesthesiology and Reanimation Unit, University Hospital of Canary Islands, Santa Cruz de Tenerife, Spain

Correspondence

Dario Acuña-Castroviejo, Centro de Investigación Biomédica, Parque Tecnológico de Ciencias de la Salud, Avda. del Conocimiento s/n, 18016 Granada, Spain.

Email: dacuna@ugr.es

Funding information

Clinical trial is financed by Pharmamel, a start-up of the University of Granada founded by DA-C and GE.

Abstract

The pharmacological properties of melatonin are well known. However, there is noticeable the lack of clinical trials that confirm the efficacy, security, absence of side effects in the short and long term, and the effective doses of melatonin. This point is especially important in diseases with high morbidity and mortality including COVID-19. There is not treatment for COVID-19, and several anti-inflammatory and antiviral molecules are being tested, and different vaccines are in preparation. Although the SARS-CoV-2 pandemic is apparently improving, it is expected new re-surges next fall. Thus, looking for an effective treatment of COVID-19 is mandatory. Melatonin has significant anti-inflammatory, antioxidant, and mitochondrial protective effects, and its efficacy has been demonstrated in multiple experimental models of disease and in a clinical trial in sepsis. Because COVID-19 courses with a severe septic response, multiple reviews proposing melatonin as a treatment for COVID-19 have been published. Nevertheless, there is a lack of experimental and clinical data on the use of melatonin on SARS-CoV-2 infection. Accordingly, we designed a clinical trial with an injectable formulation of melatonin for intravenous perfusion in ICU patients suffering from COVID-19 that has been just approved by the Spanish Agency of Medicines and Medical Devices (AEMPS). The trial will allow by the first time understand the doses and efficacy of melatonin against COVID-19.

KEYWORDS

clinical trial, COVID-19, intravenous melatonin, SARS-CoV-2, sepsis

Whereas COVID-19 expands around the world, several candidate molecules including different antiviral drugs are being testing or they are under clinical trials. Also, different types of vaccines are being under research or testing in animals and humans. No definitive treatment, however, exists to date. Due to the rapid expanding SARS-CoV-2 infection, the need for effective drug treatment of COVID-19 is an increasing urgent matter.

COVID-19 is the disease caused by SARS-CoV-2 virus, an RNA β -coronavirus that, as other respiratory

coronavirus, infects the respiratory track via respiratory droplets, although fecal and/or oral transmission paths are not excluded. To date, there are 15 787 461 cases, 639 012 deaths, and 9 622 749 recovered, although these data increase daily. The rapid transmission of SARS-CoV-2 may depend of several factors, including the incubation period variability (from 3 to 14 days until 27 days) before the presence of symptoms, and an elevated percent of infected subjects without symptoms, who are potential transmitters of

the virus.¹ Main symptoms include fever, cough, difficulty breathing, chills, muscle pain, headache, and loss of taste or smell, although 80% of the cases are mild.² Infected males are twice as females, and patients with diabetes, chronic pulmonary disease, cardiovascular disease, hypertension, or cancer, and age population are more prone to suffer from coronavirus infection.³

The pathophysiology of COVID-19 involves the binding of the coronavirus to the angiotensin-converting enzyme 2 (ACE2), which is expressed in airway epithelial cells including alveolar ones⁴ although other tissues including intestinal epithelial cells also express ACE2 and they can be an alternative entrance of the virus to the host. Two subunits of the S spike, S1 and S2, of the virus are involved in this reaction; S1 attach to ACE2, and S2 mediates the fusion of virus with cell membrane. After that, the viral RNA is released into the cytosol leading to the translation of two proteins forming a replication/transcription complex that synthesized new RNA encoding additional proteins. Once new genomic RNA is formed, it assembles with nucleocapsid proteins and glycoproteins yielding new viral particles.⁵

ACE2 can cleave angiotensin I (AI) to produce Ang 1-9 and angiotensin II (AII) to produce Ang 1-7. Ang 1-7 antagonizes the action of AII through Mas receptor providing a protective role including anti-atrophy, anti-fibrosis, anti-inflammation, antioxidant, and vasodilatation effects, whereas Ang 1-9 has counter-regulatory effects.⁶ When SARS-CoV-2 binds to ACE2, an additional angiotensin-converting enzyme (ACE) forms AII from AI, which binds to AT1R, an AII type 1 receptor, releasing pro-atrophy, pro-fibrosis, pro-inflammation, pro-oxidant, and vasoconstriction events.^{4,6} Thus, ACE2 does not only allow the entry of SARS-CoV-2 and blunts the protection of Ang 1-7 but triggers the opposite effects activating the AT1R. Together, these responses break down the lung cells and yield a local pro-inflammatory innate immune response with macrophages and monocytes recruitment and release of pro-inflammatory cytokines and adaptive immune response through B and T lymphocytes, which can block the local infection.

If these immune responses are not long enough to blunt the infection, infiltration of inflammatory cells causes release of proteases and reactive oxygen species (ROS), damaging the alveoli and reducing the gas exchange and blood PO₂, favoring additional pulmonary infections.⁷ Additionally, a systemic response triggers a cytokine storm with high levels of IL-1 β , IL-6, IL-7, IL-10, IP-10, GCSF, MIP1 α , and TNF α , requiring the patients to be controlled in the ICU. This exaggerated immune response released by SARS-CoV-2 involves not only the activation of the NF- κ B pathway, responsibly for the production of the precursors of some of these cytokines including pro-IL-1 β , but also the activation of the NLRP3 inflammasome that in turn matures these cytokines through a caspase-1 activation.^{7,8}

The excessive inflammatory response during COVID-19 is the major cause of death in these patients, leading to the use of a series of immunomodulatory agents into clinical trials but also in an off-label manner. However, from a therapeutic point of view, we must choose drugs that not only blunts the NF- κ B/NLRP3 inflammasome connection during infection, but also that combat the oxidative stress response causing pulmonary and systemic damage.

In this context, we want to call the attention to melatonin. Recently, a series of reviews have been published suggesting the beneficial effects of melatonin on COVID-19, and proposing different doses of melatonin to be used. There are not, however, clinical data, and very low experimental data that demonstrated the efficacy of melatonin in viral infections, and none in SARS-CoV-2 infection, that support these proposed doses. Nevertheless, let me suggest here why melatonin may be of interest in the treatment of COVID-19 and what we need to do to support it.

It is known that melatonin has significant antioxidant and anti-inflammatory properties, and these may be of application against the free radical's generation and inflammation in COVID-19. In 1999, we showed for the first time a dose-dependent effect of melatonin to counteracted the shock septic and multi-organ dysfunction syndrome (MODS) in a model of sepsis induced by LPS administration to rats.⁹ The anti-septic effect of melatonin was more pronounced in aged rats, which showed major inflammatory response.¹⁰ In both cases, melatonin, at doses of 60 mg/kg bw. restored the normal metabolic status and the function of kidney and liver in septic rats with not side effects. We also showed for the first time that the mitochondria are the main intracellular targets of melatonin, which is able to maintaining their homeostasis in conditions of strong oxidative stress even in sepsis.^{11,12} Melatonin also restored the normal homeostasis in every organ and tissue of mice in the model of sepsis induced by CLP and significantly increased their survival,¹³ supporting the utility of melatonin in the treatment of septic shock.¹⁴ We also identified the action's mechanisms and therapeutic targets of melatonin in sepsis.¹⁵ Melatonin induces ROR α -dependent *bmal1* expression and, in turn, BMAL1-dependent NAMPT production, promoting the synthesis of NAD + that is used by SIRT-1 to deacetylate p65, displacing it from DNA and blocking its transcriptional activity, reducing the inflammatory response. In parallel, melatonin counteracts the mitochondrial oxidative damage as a consequence of the inflammatory reaction during sepsis; it closes the MPT and prevents the release of ROS and mtDNA to the cytosol, stopping the activation of the NLRP3 inflammasome. The subsequent lack of activation of the caspase-1 impedes the maturation of pro-cytokines including pro-IL-1 β , blocking the positive feedback effect of IL-1 β on its membrane receptor and further NF- κ B activation.¹⁵ Melatonin becomes in the unique anti-inflammatory molecule able to block the two main pathways of the

innate immunity, NF- κ B and NLRP3 inflammasome that, with its ability to recover the mitochondrial homeostasis, explains its outstanding anti-septic properties.^{11,12} The doses of melatonin used in these experiments in septic mice ranged from 90 to 120 mg/kg bw., which are equivalent to that of 60 mg/kg bw used in septic rats according to dose equivalent calculation.¹⁶ Interesting, the relationship between melatonin, clock genes, and inflammatory response was recently analyzed in septic patients, showing a relationship between the degree of the inflammatory and oxidative response, the expression of *bmall*, and the levels of melatonin.¹⁷ Because septic patients exhibit a process of chronodisruption, clock genes represent another target for melatonin in this disease. Lastly, melatonin recovers lungs from oxidative damage induced by age, protecting against acute and chronic pathologies such as emphysema and pneumonia, respectively, effects that may be applicable to reduce the local oxidative and inflammatory damage in the lungs of COVID-19 patients.¹⁸

The trinity of COVID-19, immunity, inflammation, and intervention, was defined.⁷ From this point of view, and from the data here presented, it is wise for us to ask an intervention with melatonin administration in COVID-19 patients to assess its efficacy and therapeutic doses. In a recent clinical trial (EudraCT: 2008-006782-83), we showed that the intravenous administration of 60 mg/d of a patented formulation of melatonin improved septic patients, reduced their mortality to zero, and their hospital stay by 40% (manuscript in preparation). The similar inflammatory response in sepsis and COVID-19, the higher incidence of the latter in aged subjects, and the similar mortality in both groups, 20%-25% in ICU patients, together with the demonstrated efficacy of melatonin against inflammation, led us to propose a clinical trial in ICU patients with SARS-CoV-2 infection. Melatonin or placebo (the same excipients than in the melatonin solution but without melatonin) will be intravenously administered every 6 hours starting at the moment at which the patient is included in the trial, at doses that are calculated accordingly.¹⁶ The trial (EudraCT: 2020-001808-42) has been just approved by the Spanish Agency of Medicines and Medical Devices (AEMPS), allowing us to identify the doses of melatonin that can be effective in this disease. The study is a phase II, single-center, double-blind, randomized placebo-controlled trial, with the objective to address the efficacy and safety of intravenous melatonin administration in ICU patients suffering from COVID-19. Besides written informed consent from patients or family members, inclusion criteria include aged above 18 years; confirmed disease with positive PCR; acute hypoxemic respiratory failure related to COVID-19; and ICU stay less than 7 days before to randomization with Murray score at the time of randomization greater or equal at ICU admission. Exclusion criteria include participation in other COVID-19 trial; liver enzymes five times higher the normal range; glomerular filtration rate less than 30 mL/

min/1.73 m²; pregnancy; autoimmune diseases; and terminal medical illness.

CONFLICT OF INTEREST

DA-C and GE are co-inventors of the patents "Durable preparation of an injectable of melatonin exhibiting long-term stability" (PCT/ES2015/070236), and "Injectable composition of melatonin for the treatment of viral diseases" (PCT/ES2020/070234).

AUTHOR CONTRIBUTION

DA-C and GE prepared the documentation of the clinical trial; JCF, PdIO, and AMB designed the protocol of the trial; DA-C, GE, and CA-F wrote the manuscript, and all authors revised its final form.

ORCID

Dario Acuña-Castroviejo  <https://orcid.org/0000-0002-9680-1560>

Germaine Escames  <https://orcid.org/0000-0003-1256-7656>

REFERENCES

1. <https://www.worldometers.info/coronavirus/>. Accessed July 24, 2020.
2. Symptoms of Novel Coronavirus (2019-nCoV) - United States Centers for Disease Control and Prevention (CDC).
3. Guan W-J, Ni Z-Y, Hu YU, et al. Clinical characteristics of coronavirus disease 2019 in China. *New Engl J Med*. 2020;382(18):1708-1720.
4. Zhang H, Penninger JM, Li Y, et al. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med*. 2020;64:586-590.
5. Walls AC, Park Y-J, Tortorici MA, et al. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell*. 2020;181:281-292.
6. Paz Ocaranza M, Riquelme JA, García L, et al. Counter-regulatory renin-angiotensin system in cardiovascular disease. *Nat Rev Cardiol*. 2020;17:116-129.
7. Tay MZ, Poh CM, Rénia L, et al. The trinity of COVID-19: immunity, inflammation and intervention. *Nat Rev Immunol*. 2020;20(6):363-374. [1038/s41577-020-0311-8](https://doi.org/10.1038/s41577-020-0311-8)
8. Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol*. 2020;20(6):355-362.
9. Crespo E, Macías M, Pozo D, et al. Melatonin inhibits expression of the inducible NO synthase II in liver and lung and prevents endotoxemia in lipopolysaccharide-induced multiple organ dysfunction syndrome in rats. *FASEB J*. 1999;13:1537-1546.
10. Escames G, López LC, Ortiz F, Ros E, Acuña-Castroviejo D. Age-dependent lipopolysaccharide-induced iNOS expression and multi-organ failure in rats: effects of melatonin treatment. *Expt Gerontol*. 2006;41:1165-1173.
11. Martin M, Macías M, Escames G, León J, Acuña-Castroviejo D. Melatonin but not vitamins C and E maintains glutathione homeostasis in t-butyl hydroperoxide-induced mitochondrial oxidative stress. *FASEB J*. 2000;14:1677-1679.

12. Escames G, León J, Macías M, Khaldy H, Acuña-Castroviejo D. Melatonin counteracts lipopolysaccharide-induced expression and activity of mitochondrial nitric oxide synthase in rats. *FASEB J*. 2003;17:932-934.
13. Escames G, López LC, Ortiz F, et al. Attenuation of cardiac mitochondrial dysfunction by melatonin in septic mice. *FEBS J*. 2007;274:2137-2147.
14. Escames G, Acuña-Castroviejo D, López LC, et al. Pharmacological utility of melatonin in the treatment of septic shock. *J Pharm Pharmacol*. 2006;58:1153-1165.
15. Garcia JA, Volt H, Venegas C, et al. Disruption of the NF-kappaB/NLRP3 connection by melatonin requires retinoid-related orphan receptor-alpha and blocks the septic response in mice. *FASEB J*. 2015;29:3863-3875.
16. Nair AB, Jacob S. A simple practice guide for dose conversion between animals and humans. *J Basic Clin Pharma*. 2016;7:27-31.
17. Acuna-Fernandez C, Marin JS, Diaz-Casado ME, et al. Daily changes in the expression of clock genes in sepsis and their relation with sepsis outcome and urinary excretion of 6-sulfatoxymelatonin. *Shock*. 2020;53:550-559.
18. Acuña-Castroviejo D, Carretero M, Doerrier C, et al. Melatonin protects lung mitochondria from aging. *Age (Dordr)*. 2012;34:681-692.

How to cite this article: Acuña-Castroviejo D, Escames G, Figueira JC, de la Oliva P, Borobia AM, Acuña-Fernández C. Clinical trial to test the efficacy of melatonin in COVID-19. *J Pineal Res*. 2020;69:e12683. <https://doi.org/10.1111/jpi.12683>