

## RESEARCH ARTICLE

# The role of DBP gene polymorphisms in the prevalence of new coronavirus disease 2019 infection and mortality rate

Lutfiye Karcioğlu Batur PhD  | Nezh Hekim PhD 

Department of Molecular Biology and Genetics, Faculty of Engineering and Natural Sciences, Biruni University, Istanbul, Turkey

**Correspondence**

Lutfiye Karcioğlu Batur, PhD, Department of Molecular Biology and Genetics, Faculty of Engineering and Natural Sciences, Biruni University, 10. Yıl Street Protokol Way No: 45, Istanbul 34010, Turkey.  
Email: lbatur@biruni.edu.tr

**Abstract**

Since December 2019, coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2, has given rise to emerging respiratory infections with pandemic diffusion. The vitamin D binding protein (DBP) with emphasis on its regulation of total and free vitamin D metabolite levels participate in various clinical conditions. The main goal of this study was to evaluate if there was any association between the DBP gene polymorphism at rs7041 and rs4588 loci and the prevalence of COVID-19 and its mortality rates caused among populations of 10 countries including Turkey. Positive significant correlations were found between the prevalence (per million) and mortality rates (per million), and GT genotype ( $P < .05$ ) while there was a negative significant correlation between prevalence (per million) and mortality rates (per million), and TT genotype at rs7041 locus among all populations ( $P < .05$ ). However, no significant correlation was found at rs4588 locus. GT genotype was found to confer this susceptibility to the populations of Germany, Mexico, Italy, Czech, and Turkey. The variations in the prevalence of COVID-19 and its mortality rates among countries may be explained by Vitamin D metabolism differed by the DBP polymorphisms of rs7041 and rs4588.

**KEYWORDS**

coronavirus disease 2019, polymorphism, rs4588, rs7041, vitamin D binding protein

## 1 | INTRODUCTION

Since December 2019, coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has given rise to emerging respiratory infections with a pandemics diffusion.<sup>1</sup> By 2 July 2020, the global number of confirmed cases of COVID-19 reached 10 534 765 with a mortality of 512 881.<sup>2</sup> In Turkey, 201 098 cases and 5150 deaths have been confirmed until 2nd July 2020.<sup>2</sup> A relationship was recently found between vitamin D levels and the number COVID-19 cases and the mortality rates caused by the coronavirus infection.<sup>3</sup>

The vitamin D binding protein (DBP) with emphasis on its regulation of total and free vitamin D metabolite levels participate in various clinical conditions. Nearly all DBP is produced in the liver, where its regulation is influenced by estrogen, glucocorticoids, and inflammatory cytokines but not by vitamin D itself. DBP is the most polymorphic protein known, and different DBP alleles can have

substantial impact on its biologic functions.<sup>4</sup> The two most common alleles—Gc1s (rs7041 locus) and Gc2 (rs4588 locus)—differ in their affinity with the vitamin D metabolites and have been variably associated with several clinical conditions.<sup>4</sup> Among these conditions, G allele at the rs7041 locus was found to be related with increased susceptibility to hepatitis C viral infection.<sup>5</sup> Moreover, the individuals having an AA genotype within rs4588 locus of the Gc2 polymorphic region showed a greater increase at 25(OH)D levels following vitamin D supplementation than those having the GG genotype.<sup>6</sup> A single-nucleotide polymorphism at rs4588 has been associated with susceptibility to the metabolic syndrome.<sup>7</sup> Therefore, we hypothesized that DBP polymorphisms may play a significant role in COVID-19.

The main goal of this study was to evaluate if there is any association between the DBP gene polymorphism at rs7041 and rs4588 loci and the prevalence of COVID-19 and its mortality rates caused among populations of ten countries including Turkey.

## 2 | MATERIALS AND METHODS

To test this hypothesis and to limit confounding bias (latitude, etc), we focused on the countries whose DBP polymorphisms at rs7041 and rs4588 loci were defined and the allele frequencies reported in five cohort and two systematic review and meta-analysis studies.<sup>8-15</sup> We searched the literature for DBP gene polymorphism in each country. We recorded the total number of cases of COVID-19 and per million population in each of the countries to find the prevalence, and the mortality rates caused by the coronavirus infection recorded at 2nd July 2020 (Table 1) according to World Health Organization Coronavirus disease (COVID-19) Situation Report—164.<sup>2</sup>

The allele frequencies of DBP polymorphisms at rs7041 and rs4588 loci of Turkish population were retrieved from a previous thesis study (unpublished results).<sup>16</sup> The ethical approval for analyzing the blood samples to examine the polymorphisms was obtained from Biruni University Non-Interventional Research Ethics Committee (Approval No: 2017/10-1). The written consent forms were obtained from all subjects who were informed about the study.

Blood samples were collected from 51 healthy Turkish individuals who applied for check-up, met the study criteria, and agreed to participate in the study. The selection criteria were not using any vitamin D supplements for the last 2 years, not having any health problem that would affect the vitamin D concentrations, and not being a black person. Total 25(OH)D concentrations were measured by a chemiluminescence microparticle immunoassay method (CMIA), using Architect 25-OH Vitamin D kit (5P02; Abbott Diagnosis) and i1000SR analyzer (Abbott Laboratories).

DNA isolation from the whole blood was performed by a quick-DNATM miniprep plus (Zymo Research) DNA isolation kit. Isolated DNAs were stored at  $-200^{\circ}\text{C}$  for further analysis. DNA quality and

concentration measurements were performed in 2  $\mu\text{L}$  DNA with the NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific). Using isolated genomic DNA, genotyping for the most common single nucleotide polymorphisms (SNPs) of DBP, rs4588, and rs7041, was performed using TaqMan probes for real-time PCR. The SNP assay coded by C\_8278879\_10 (Applied Biosystems TaqMan SNP Genotyping Assays Thermo Fisher Scientific) was used for genotyping rs4588, and SNP assay coded by C\_3133594\_30 (Thermo Fisher Scientific) was used for genotyping rs7041. The base sequences were CTTGTTAACCAGCTT TGCCAGTTC\*[G/T]TGGGTGTGGCATCAGGCAATTTTC and CTTTGC CAGTTCCTGGGTGTGGC\*[A/C]TCAGGCAATTTTGCTTTTAGTCGT, respectively. Real-time temperature cycle reaction conditions were adjusted according to the protocol of manufacturer and the literature.<sup>17</sup>

All data were analyzed by SPSS (statistical package for social sciences) for Windows 22 program. In the analysis of the data, first, the assumptions that must be met were tested to decide which tests (parametric/nonparametric tests) to apply. Shapiro-Wilk test, kurtosis and skewness values that are other assumptions of normal distribution, and histogram graph was used to decide the normality of the distribution. Considering the insufficient number of data in each group, it was decided that the data did not exhibit normal distribution. The relationship between independent variables was examined with the Spearman correlation coefficient ( $\rho$ ). In the interpretation of whether the obtained values are significant or not, 0.05 significance level was used as a criterion.

## 3 | RESULTS

The mean age of 51 healthy individuals from the Turkish population was  $39.39 \pm 12.30$ . Among them, 49.1% of individuals were male

**TABLE 1** Population diversities of rs7041 and rs4588 polymorphisms, prevalence of COVID-19, and mortality rates per country recorded at 2nd July 2020

Country	rs7041			rs4588			Prevalence <sup>a</sup>		Mortality <sup>a</sup>		Reference
	GG	GT	TT	AA	AC	CC	Total	Per million	Total	Per million	
China	7.30	42.3	50.4	8.8	44.9	46.3	81263	59.24	4648	3.23	Zhou et al <sup>8</sup>
Japan	8.80	31.9	59.3	2.7	43.8	53.6	18874	149.23	975	7.71	Khanna et al <sup>9</sup>
Nigeria	0.70	15.6	83.7	0.0	8.20	91.8	26484	128.48	603	2.93	Khanna et al <sup>9</sup>
Kenya	0.90	12.7	86.4	0.9	11.0	88.1	6673	124.1	149	2.77	Jones et al <sup>10</sup>
Mexico	24.1	50.0	25.9	1.8	46.4	51.8	226089	1753.54	27769	215.38	Rivera-Paredes et al <sup>11</sup>
Italy	25.5	56.9	17.6	5.9	39.2	54.9	240760	3982.01	34788	575.37	Jones et al <sup>10</sup>
Turkey	31.0	49.0	20.0	2.0	53.0	45.0	201098	2384	5150	61.06	Present data
Finland	63.6	32.5	3.9	10.8	48.7	40.5	7236	1305.81	328	59.19	Enlund-Cerullo et al <sup>13</sup>
Germany	31.9	48.2	19.9	8.7	42.0	49.3	194725	2323.56	8985	107.21	Terock et al <sup>15</sup>
Czechia	38.8	49.4	11.9	6.2	42.5	51.2	12046	1124.69	349	32.58	Pleva et al <sup>14</sup>

Abbreviations: COVID-19, coronavirus disease 2019; WHO, World Health Organization.

<sup>a</sup>Recorded on 2nd July 2020 from WHO Coronavirus disease (COVID-19) Situation Report—164.

( $n = 25$ ) and 50.9% were female ( $n = 26$ ). The mean concentration of total 25(OH)D was  $20.75 \pm 14.2$  ng/mL.

Population diversities of *rs7041* polymorphisms showed that the populations of China, Japan, Nigeria, and Kenya mostly have TT genotype, while the populations of Germany, Mexico, Italy, Czech, and Turkey mostly have GT genotype (Table 1). Population diversities of *rs4588* polymorphisms revealed that the populations of all countries except Finland and Turkey mostly have CC genotype, while Finn and Turkish populations have AC genotype at *rs4588* locus. The prevalence of COVID-19 and mortality rates per country recorded at 2nd July 2020 showed that Germany, Mexico, Italy, and Turkey had the highest number of COVID-19 cases and mortality rates per million of the populations of countries involved in the study (Table 1).

Correlation between *rs7041* and *rs4588* polymorphisms and prevalence of COVID-19 and mortality rates per country demonstrated that there were positive significant correlations between the prevalence (per million) and mortality rates (per million), and GT genotype ( $P < .05$ ), while there was a negative significant correlation between prevalence (per million) and mortality rates (per million), and TT genotype at *rs7041* locus among all populations ( $P < .05$ ). However, no significant correlation was found between the prevalence (per million) and mortality rates (per million), and the polymorphism at *rs4588* locus (Table 2).

## 4 | DISCUSSION

In the present study, TT genotype was found to confer COVID-19 susceptibility to the populations of China, Japan, Nigeria, and Kenya. GT genotype was found to confer this susceptibility to the populations of Germany, Mexico, Italy, Czech, and Turkey. The variations in the prevalence of COVID-19 and its mortality rates among countries may be explained by vitamin D metabolism differed by the DBP polymorphisms of *rs7041* and *rs4588*.

DBP is the most polymorphic protein known, which regulates the total and circulating free vitamin D metabolite levels in various clinical conditions. DBP alleles differ in their affinity with the vitamin D metabolites and can have substantial impact on various clinical conditions.<sup>4</sup> Polymorphisms in the *DBP* gene have been reported to be associated with vitamin D deficiency in different populations.<sup>12,18,19</sup> The general prevalence of vitamin D deficiency varies in the world, ranging from 7% to 77%.<sup>20</sup> Vitamin D deficiency is more common in the subtropical (including China) and mid-latitude (including Italy, Japan, Turkey) countries than the tropical (Mexico, Nigeria, and Kenya) and high-latitude countries.<sup>21</sup> In a study conducted in 1161 healthy subjects from the Turkish population, Ozturk et al<sup>22</sup> found the overall mean serum 25(OH)D level as  $16.61 \pm 6.90$  ng/mL and reported a high prevalence of vitamin D insufficiency or deficiency in all age groups. They also showed that the vitamin D deficiency is very common (75.54% with 25(OH)D  $< 20$  ng/mL) among adult Turkish population.<sup>22</sup> Altogether, these studies implicate that the vitamin D levels are mainly regulated by the genetic background of both healthy population and patients. In the present study, the genetic variations in the *DBP* gene, specifically SNP in *rs7041*

**TABLE 2** Correlation between *rs7041* and *rs4588* polymorphisms and prevalence of COVID-19 and mortality rates per country

Spearman's rho	GG	GT	TT	AA	AC	CC
Prevalence per million						
<i>r</i>	.61	.73	-.62	.12	.27	-.25
<i>P</i>	.06	.02	.04	.75	.45	.49
Mortality per million						
<i>r</i>	.60	.87	-.66	.27	.36	-.31
<i>P</i>	.07	.01	.04	.45	.31	.38

Note: Bold values indicate  $P < .05$ .

Abbreviation: COVID-19, coronavirus disease 2019.

$P < 0.05$  for bold values

locus, were found to be correlated with the prevalence of COVID-19 and mortality rates among countries.

Previous studies have identified a potential common relationship between the mean vitamin D levels in various European countries with COVID-19 cases per million population and its mortality.<sup>3,23</sup> Ilie et al<sup>3</sup> reported negative correlations between mean levels of vitamin D in European countries including Turkey and the number of COVID-19 cases per million population.<sup>3</sup> They suggest that Spain, Italy, and Switzerland are the most vulnerable group of the population in relation to COVID-19 since the vitamin D levels are severely low in the aging population, especially in these countries. They explained the crude association by the role of vitamin D in the prevention of COVID-19 infection or more probably by a potential protection of vitamin D from the more negative consequences of the infection.<sup>3</sup>

Vitamin D plays a major role in regulating the immune system, including immune responses to viral infection.<sup>24</sup> Interventional and observational epidemiological studies provide evidence that vitamin D deficiency may confer an increased risk of influenza and respiratory tract infection.<sup>25</sup> Cell culture experiments support the thesis that vitamin D has direct antiviral effects particularly against enveloped viruses. Though vitamin D's antiviral mechanism has not been fully established, it may be linked to vitamin D's ability to upregulate the antimicrobial peptides LL-37 and human beta-defensin 2.<sup>26</sup> Regarding the genetic susceptibility to a viral infection in vitamin D deficiency, we also observed significant correlations between *rs7041* polymorphism and prevalence of COVID-19 and mortality rates per country. However, no significant correlation was found between the prevalence (per million) and mortality rates (per million) at *rs4588* locus.

The pathology of COVID-19 involves a complex interaction between the SARS-CoV-2 and the body immune system. Calcitriol (1,25-dihydroxyvitamin D3) exerts pronounced impacts on ACE2/Ang(1-7)/MasR axis with enhanced expression of ACE2.<sup>27</sup> ACE2 is the host cell receptor responsible for mediating infection by SARS-CoV-2. ACE2 polymorphisms were recently described in human populations.<sup>28</sup> Another common polymorphism was found in *DBP* gene, which is a highly polymorphic gene. Allelic variants of the *DBP* gene have been studied extensively for their association with vitamin D deficiency<sup>29,30</sup> and viral infections.<sup>5</sup> Two of these variants corresponding to different allelic arrangements of *rs7041* and *rs4588* were reported to have a different

affinity to bind to vitamin-D3, and hence affect its serum concentration.<sup>31</sup> Different DBP isoforms influence the serum concentration/bioavailability of vitamin-D3.<sup>9</sup> From this perspective these isoforms might be correlated with an increased risk of viral infection in populations, as reported in the present study. If a necessary concentration of bioavailable vitamin D is not reached in certain genotypes, the immune system may modulate the body reaction to an infection in a severe way.

Low levels of total vitamin D, which are more common in black than in white, are associated with negative health outcomes in epidemiologic studies.<sup>32-34</sup> Powe et al.<sup>34</sup> reported lower mean levels of both total vitamin D and DBP in blacks than in whites due to the genetic polymorphisms.<sup>34</sup> Among homozygous participants, blacks and whites had similar levels of bioavailable vitamin D. Therefore, racial differences in the prevalence of common genetic polymorphisms provide a likely explanation for altered Vitamin D metabolism. In the present study, the genetic polymorphism, especially in *rs7041* locus of the DBP gene is probably associated with the increased risk of COVID-19 infection and its mortality among populations with the white race.

Vitamin D deficiency is known to impair the ability of macrophages to mature, to produce macrophage-specific surface antigens, to produce the lysosomal enzyme acid phosphatase, and to secrete H<sub>2</sub>O<sub>2</sub>, a function integral to their antimicrobial function.<sup>35</sup> Vitamin D has been also reported to modulate macrophages' response, preventing them from releasing too many inflammatory cytokines and chemokines,<sup>36</sup> which are frequently observed in COVID-19 cases.<sup>37</sup> Therefore, the correlation of the variations in DBP polymorphisms and the prevalence of COVID-19 with its mortality rate may depend on the modulatory effect of bioavailable Vitamin D levels of individuals, which is determined by the genetic background. However, the prevalence of SARS-COV-2 infection differs from the severity of COVID-19, by association with many factors such as public awareness, behaviors, and antiviral policy of each country except the host genetic factors. On the contrary, the severity of the disease induced by viral infection might be associated with the genetic host factors. More detailed and large sampled studies about the genetic variations in infected patients with different degrees of severity are needed to explain the underlying mechanism of Vitamin D metabolism in COVID-19 patients.

## CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

## DATA AVAILABILITY STATEMENT

The data related to COVID-19 that support the findings of this study are available in Coronavirus disease (COVID-19) Situation Report—164 of World Health Organization, at [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200702-covid-19-sitrep-164.pdf?sfvrsn=ac074f58\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200702-covid-19-sitrep-164.pdf?sfvrsn=ac074f58_2).

The data of 51 healthy Turkish individuals that support the findings of this study are available from the corresponding author upon reasonable request.

## ORCID

Lutfiye Karcioğlu Batur  <http://orcid.org/0000-0002-4803-9137>

Nezih Hekim  <https://orcid.org/0000-0003-2613-8676>

## REFERENCES

- Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708-1720.
- World Health Organization. Coronavirus disease (COVID-19) Situation Report—164. Retrieved from [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200702-covid-19-sitrep-164.pdf?sfvrsn=ac074f58\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200702-covid-19-sitrep-164.pdf?sfvrsn=ac074f58_2)
- Ilie PC, Stefanescu S, Smith L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res*. 2020;32(7):1195-1198.
- Bikle DD, Schwartz J. Vitamin D binding protein, total and free vitamin D levels in different physiological and pathophysiological conditions. *Front Endocrinol (Lausanne)*. 2019;10:317. <https://doi.org/10.3389/fendo.2019.00317>
- Xie CN, Yue M, Huang P, et al. Vitamin D binding protein polymorphisms influence susceptibility to hepatitis C virus infection in a high-risk Chinese population. *Gene*. 2018;679:405-411. <https://doi.org/10.1016/j.gene.2018.09.021>
- Mehramiz M, Khayyatzadeh SS, Esmaily H, et al. Associations of vitamin D binding protein variants with the vitamin D-induced increase in serum 25-hydroxyvitamin D. *Clin Nutr ESPEN*. 2019;29:59-64. <https://doi.org/10.1016/j.clnesp.2018.12.005>
- Karuwanarint P, Phonrat B, Tungtrongchitr A, et al. Vitamin D-binding protein and its polymorphisms as a predictor for metabolic syndrome. *Biomark Med*. 2018;12:465-473. <https://doi.org/10.2217/bmm-2018-0029>
- Zhou L, Zhang X, Chen X, et al. GC Glu416Asp and Thr420Lys polymorphisms contribute to gastrointestinal cancer susceptibility in a Chinese population. *Int J Clin Exp Med*. 2012;5(1):72-79.
- Khanna R, Nandy D, Senapati S. Systematic review and meta-analysis to establish the association of common genetic variations in vitamin D binding protein with chronic obstructive pulmonary disease. *Front Genet*. 2019;10:413. <https://doi.org/10.3389/fgene.2019.00413>
- Jones P, Lucock M, Chaplin G, et al. Distribution of variants in multiple vitamin D-related loci (DHCR7/NADSYN1, GC, CYP2R1, CYP11A1, CYP24A1, VDR, RXR $\alpha$  and RXR $\gamma$ ) vary between European, East-Asian and Sub-Saharan African-ancestry populations. *Genes Nutr*. 2020; 15(1):5. <https://doi.org/10.1186/s12263-020-00663-3>
- Rivera-Paredes B, Hidalgo-Bravo A, de la Cruz-Montoya A, et al. Association between vitamin D deficiency and common variants of Vitamin D binding protein gene among Mexican Mestizo and indigenous postmenopausal women. *J Endocrinol Invest*. 2020;43:935-946.
- Takiar R, Lutsey PL, Zhao D, et al. The associations of 25-hydroxyvitamin D levels, vitamin D binding protein gene polymorphisms, and race with risk of incident fracture-related hospitalization: twenty-year follow-up in a bi-ethnic cohort (the ARIC Study). *Bone*. 2015;78:94-101.
- Enlund-Cerullo M, Koljonen L, Holmlund-Suila E, et al. Genetic variation of the vitamin D binding protein affects vitamin D status and response to supplementation in infants. *J Clin Endocrinol Metab*. 2019; 104(11):5483-5498. <https://doi.org/10.1210/jc.2019-00630>
- Pleva L, Kovarova P, Faldynova L, et al. The rs1803274 polymorphism of the BCHE gene is associated with an increased risk of coronary instant stenosis. *BMC Cardiovasc Disord*. 2015;15:135. <https://doi.org/10.1186/s12872-015-0128-8>
- Terock J, Hannemann A, Van der Auwera S, et al. Posttraumatic stress disorder is associated with reduced vitamin D levels and functional polymorphisms of the vitamin D binding-protein in a population-based sample. *Prog Neuropsychopharmacol Biol Psychiatry*. 2020;96:109760.
- Karcioğlu Batur L. The Relationship Between the Genotype of Vitamin D Binding Protein and Plasma 25 OH D Concentration in Turkish

- Population. [PhD Thesis]. Istanbul, Turkey: Istanbul University-Cerrahpasa Institute of Graduate Education, Department of Medical Biology; 2019.
17. Peršić V, Rajević D, Markova-Car E, et al. Vitamin D-binding protein (rs4588) T/T genotype is associated with antero-septal myocardial infarction in coronary artery disease patients. *Ann Transl Med.* 2019; 7:374.
  18. Khan AH, Jafri L, Siddiqui A, Naureen G, Morris H, Moatter T. Polymorphisms in the GC gene for vitamin D binding protein and their association with vitamin D and bone mass in young adults. *J Coll Physicians Surg Pak.* 2019;29(8):715-719. <https://doi.org/10.29271/jcsp.2019.08.715>
  19. Shao B, Jiang S, Muyiduli X, et al. Vitamin D pathway gene polymorphisms influenced vitamin D level among pregnant women. *Clin Nutr.* 2018;37:2230-2237. <https://doi.org/10.1016/j.clnu.2017.10.024>
  20. Arabi A, El Rassi R, El-Hajj Fuleihan G. Hypovitaminosis D in developing countries-prevalence, risk factors and outcomes. *Nat Rev Endocrinol.* 2010;6:550-561.
  21. Kara M, Ekiz T, Ricci V, Kara Ö, Chang KV, Özçakar L. "Scientific Strabismus" or two related pandemics: coronavirus disease and vitamin D deficiency. *Br J Nutr.* 2020;12:1-6. <https://doi.org/10.1017/S0007114520001749>
  22. Ozturk ZA, Gol M, Turkbeyler IH. Prevalence of vitamin D deficiency in otherwise healthy individuals between the ages of 18 and 90 years in southeast Turkey. *Wien Klin Wochenschr.* 2017;129:854-855. <https://doi.org/10.1007/s00508-017-1241-8>
  23. Boucher BJ. Adjustments in analyses of vitamin D status, allowing for vitamin D determinants, for Covid-19 risks. *Diabetes Metab Res Rev.* 2020:e3375-e3375. <https://doi.org/10.1002/dmrr.3375>. [published online ahead of print, 2020 Jun 26].
  24. Beard JA, Bearden A, Striker R. Vitamin D and the anti-viral state. *J Clin Virol.* 2011;50(3):194-200. <https://doi.org/10.1016/j.jcv.2010.12.006>
  25. Miller J, Gallo RL. Vitamin D and innate immunity. *Dermatol Ther.* 2010;23:13-22.
  26. Leikina E, Delanoë-Ayari H, Melikov K, et al. Carbohydrate-binding molecules inhibit viral fusion and entry by crosslinking membrane glycoproteins. *Nat Immunol.* 2005;6(10):995-1001.
  27. Cui C, Xu P, Li G, et al. Vitamin D receptor activation regulates microglia polarization and oxidative stress in spontaneously hypertensive rats and angiotensin II-exposed microglial cells: role of renin-angiotensin system. *Redox Biol.* 2019;26:101295.
  28. Devaux CA, Rolain JM, Raoult D. ACE2 receptor polymorphism: Susceptibility to SARS-CoV-2, hypertension, multi-organ failure, and COVID-19 disease outcome. *J Microbiol Immunol Infect.* 2020; 53(3):425-435.
  29. Chishimba L, Thickett DR, Stockley RA, Wood AM. The vitamin D axis in the lung: a key role for vitamin D-binding protein. *Thorax.* 2010;65:456-462. <https://doi.org/10.1136/thx.2009.128793>
  30. Wood AM, Bassford C, Webster D, et al. Vitamin D-binding protein contributes to COPD by activation of alveolar macrophages. *Thorax.* 2011;66:205-210. <https://doi.org/10.1136/thx.2010.140921>
  31. Janssens W, Bouillon R, Claes B, et al. Vitamin D deficiency is highly prevalent in COPD and correlates with variants in the vitamin D-binding gene. *Thorax.* 2010;65:215-220. <https://doi.org/10.1136/thx.2009.120659>
  32. Holick MF. Vitamin D deficiency. *N Engl J Med.* 2007;357:266-281.
  33. Adams JS, Hewison M. Update in vitamin D. *J Clin Endocrinol Metab.* 2010;95:471-478.
  34. Powe CE, Evans MK, Wenger J, et al. Vitamin D-binding protein and vitamin D status of black Americans and white Americans. *N Engl J Med.* 2013;369(21):1991-2000. <https://doi.org/10.1056/NEJMoa1306357>
  35. Abu-Amer Y, Bar-Shavit Z. Impaired bone marrow-derived macrophage differentiation in vitamin D deficiency. *Cell Immunol.* 1993;151:356-368.
  36. Helming L. 1 $\alpha$ ,25-Dihydroxyvitamin D3 is a potent suppressor of interferon gamma-mediated macrophage activation. *Blood.* 2005;106:4351-4358.
  37. Nikolich-Zugich J, Knox KS, Rios CT, Natt B, Bhattacharya D, Fain MJ. SARS-CoV-2 and COVID-19 in older adults: what we may expect regarding pathogenesis, immune responses, and outcomes. *GeroScience.* 2020;42(2):505-514.

**How to cite this article:** Karcioğlu Batur L, Hekim N. The role of DBP gene polymorphisms in the prevalence of new coronavirus disease 2019 infection and mortality rate. *J Med Virol.* 2021;93:1409–1413. <https://doi.org/10.1002/jmv.26409>