

LETTER TO THE EDITOR

COVID-19 and AIDS: Outcomes from the coexistence of two global pandemics and the importance of chronic antiretroviral therapy

To the Editor,

The outbreak of coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome 2 (SARS-CoV-2), has created a worldwide pandemic, raising multitude of challenges and uncertainties regarding disease management specific to immunocompromised patient populations.¹ Despite over ~16 million confirmed cases of COVID-19 worldwide as of July 25, 2020, there is a dearth of reports describing the co-infection of human immunodeficiency virus (HIV) and SARS-CoV-2.² There are various unknowns regarding the simultaneous impact from these two diseases, despite the fact that the majority of patients with acquired immunodeficiency syndrome (AIDS) are immunocompromised and have comorbidities, which can predispose them to severe clinical symptoms, and poorer prognosis, from COVID-19.^{3,4} A large majority of reports indicate that HIV patients with normal CD4 T-cell counts and suppressed viral loads and receiving regular, combined antiretroviral therapy (cART) do not present with a severe clinical course of COVID-19 and may not be at an increased risk of developing SARS-CoV-2 infection.⁵⁻⁹ These favorable indications for HIV patients may be the byproduct of a potential protective factor conferred by antiretroviral therapy, which has also been used successfully in previous coronavirus epidemics.^{10,11}

SARS-CoV-2 and HIV carry numerous similarities as they are both RNA viruses capable of undergoing mutations and recombination under selection pressure within the host.¹² During the previous SARS-CoV epidemic, the HIV-1 protease inhibitor nelfinavir was found to strongly inhibit the cytopathic effect induced by SARS-CoV infection.¹⁰ Lopinavir was also found to show efficacy both in vivo and in vitro, reducing viral titers and shortening disease progression in case of MERS-CoV-infected animals.¹¹ HIV-1 protease inhibitors including lopinavir/ritonavir and darunavir have been found as efficacious treatment options in patients coinfecting with COVID-19 and HIV by reducing SARS-CoV-2 viral loads and accelerating recovery.¹³⁻¹⁵ However, recently, remdesivir, a broad-spectrum antiviral drug, has also shown promising results in suppressing SARS-CoV-2 replication in a double-blind, randomized, placebo-controlled trial, where adults hospitalized with COVID-19 that received remdesivir showed reduced respiratory tract involvement.¹⁶ Similarly, tenofovir has also been found to be highly efficacious against SARS-CoV-2 through potently inhibiting SARS-CoV-2 RNA-dependent RNA polymerase.¹⁷ However, it is important to recognize

that clinical evidence regarding the role of antiretroviral therapy, specifically, lopinavir/ritonavir is still emerging. The current COVID-19 treatment guidelines issued by the NIH recommend against the use of lopinavir/ritonavir protease inhibitors (AIII) for the treatment of COVID-19, and is restricted to clinical trials.^{18,19}

Various large clinical case series have corroborated the growing evidence that patients with HIV/SARS-CoV-2 co-infection do not have excess morbidity and mortality when compared to non-HIV COVID-19 patients. In a study of 47 coinfecting patients reported by Gervasoni et al,⁹ HIV patients that were hospitalized with SARS-CoV-2 infection generally had favorable outcomes and did not experience severe symptoms requiring intensive care unit admission or mechanical ventilation. Another study by Blanco et al¹³ reporting 5 patients coinfecting with HIV/SARS-CoV-2 in Spain, showed that patients who were previously receiving ART before admission for COVID-19 experienced less severe symptoms, which did not require intensive care unit admission or mechanical ventilation. Härter et al⁵ reported similar outcomes in a study of 33 HIV patients infected with SARS-CoV-2 who were all on antiretroviral therapy at the time of COVID-19 diagnosis. This study also did not support excess mortality and morbidity among HIV patients with COVID-19 when patients were virally suppressed on antiretroviral therapy. Finally, Shalev et al also reported that HIV patients with COVID-19 shared similar clinical characteristics and outcomes to other hospitalized cohorts in their report of 31 coinfecting patients.⁶ However, contradictory results have also emerged. Vizcarra et al¹⁶ reported 51 HIV-infected individuals diagnosed with COVID-19 of which six were critically ill and two died. Notably, previous administration of antiretroviral therapy, CD4 T-cell counts, CD4/CD8 ratio, and pre-existing comorbidities were not significantly different in recovered patients vs still-admitted individuals in this study.²⁰ The authors in this study concluded that HIV-infected individuals should not be considered protected from SARS-CoV-2 or to have lower risk of severe disease. Another study by Suwanwongse et al²¹ reported that 7/9 HIV/SARS-CoV-2 coinfecting patients from New York City died from COVID-19-related respiratory failure, despite low HIV viral loads and previous ART regimen. The largest study by Del Amo et al²² reported 236 HIV/COVID-19 patients from a cohort of 77 590 HIV-positive patients on cART. Out of the 236 patients, 15 were admitted to the ICU, 20 died, and 151 were hospitalized with mild to moderate symptoms. The study notes that HIV-positive patients who received tenofovir/emtricitabine (TDF/FTC) as part of cART regimen

were found to have a lower risk for COVID-19 and related hospitalization than those receiving other therapies. This finding conforms with various other molecular docking and case series studies, which demonstrate more favorable outcomes for HIV/COVID-19 patients on tenofovir-based regimens.^{11,15} However, this study does not include non-AIDS comorbidities, such as CD4 + T-cell counts, HIV viral RNA loads, and other key confounders, which may influence the results. Further, this study did not report whether or not the patients receiving TDF/FTC had fewer comorbidities than those receiving other forms of ART.

Given the limited available evidence and mixed results from various studies as noted above, it is difficult to draw effective conclusions regarding the outcomes for HIV patients when infected with SARS-CoV-2 at this time and therefore more data/studies are needed to make meaningful conclusions. It is also important to recognize that these studies contain limited sample sizes, geographical limitations, and inconsistencies in information, which creates a challenge in stratifying the data to understand the impact of antiretroviral therapy in addition to taking into account individual patient factors and comorbid conditions. Thus, to further elucidate the role of cART on SARS-CoV-2, future large cohort studies are warranted with detailed meta-analysis. Nevertheless, these early studies place a spotlight on the unanswered role of antiretroviral therapy in pre-exposure prophylaxis for these patients in preventing incidence of severe clinical symptoms in these patients. Based on the trend of favorable prognosis as seen in the available reports, and the lack of large cohorts of HIV/SARS-CoV-2 infections, it is not unreasonable to believe that HIV patients who have no other comorbidities could be protected from severe COVID-19 clinical symptoms by following a strict chronic antiretroviral regimen.

Further, confounding factors may be present in the form of host factors (immune activation and viral genotypes), which can significantly alter the clinical outcomes in coinfecting patients. These factors are often unreported in the majority of studies describing HIV/SARS-CoV-2 co-infection but may still be significant influencers of clinical outcomes. Next, infection with certain HIV subtypes may lead to rapid or slow disease progression, host genetics such as human leukocyte antigen (HLA) can also influence differential clinical outcomes. In regard to immune activation, ART intervention may not completely stop the immune-activation process, and the type of ART may differentially affect immune activation. Thus, depending on the immune activation status of HIV patients coinfecting with SARS-CoV-2, a variety of clinical outcomes can be expected, which could partially explain the discrepancies of clinical outcomes among patients with HIV/SARS-CoV-2 co-infections in all previously reported studies. Therefore, further large systematic studies are warranted to further determine the contribution of these specific genetic/host factors that influence the clinical outcomes.


Another possible factor to consider as an explanation for the low incidence of severe outcomes is the impact of social and psychological factors.²³ Are HIV patients following stricter social distancing measures and self-quarantining due to a greater perceived risk of contracting severe SARS-CoV-2 infection? It is also important to consider the

backgrounds and demographics of the population represented by HIV patients, which includes drug users, poor patients living in crowded urban slums, migrant workers, sex workers, and prisoners, who may live in environments in which it may be difficult to practice successful social distancing and contact isolation. The challenges of universal HIV testing and now with COVID-19 testing also remains unsolved, which can impact the reported incidence and prevalence.

Although more studies are needed, it is undeniable that the overwhelming majority of available evidence regarding HIV/COVID-19 coinfecting patients overwhelmingly demonstrates the positive effects of chronic antiretroviral therapy. Individuals who are unaware of their HIV diagnosis and do not have access to antiretroviral therapy may be at heightened risk of experiencing increased morbidity and mortality if infected with SARS-CoV-2. Given the benefits of chronic antiretroviral therapy in reported patients thus far, it becomes increasingly important to sustain the HIV care continuum. Healthcare centers and HIV clinics should work to continue regular follow-up visits for patients and ensure that antiretroviral therapy is adequately rationed for HIV patients and distributed in a timely manner to protect this vulnerable population.

FUNDING INFORMATION

Research reported in this publication was partially supported by National Institutes of Health R01 AI129745 and Frances E. Lageschulte and Evelyn B. Weese New Frontiers in Medical Research Fund to SNB and R01DA042524, R01DA050169, P51OD011133 to M.M. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Raj H. Patel¹ 

Arpan Acharya²

Mahesh Mohan³

Siddappa N. Byrareddy^{2,4,5} 

¹Edward Via College of Osteopathic Medicine, Monroe, Louisiana

²Department of Pharmacology and Experimental Neuroscience, University of Nebraska Medical Centre, Omaha, Nebraska

³Texas Biomedical Research Institute, Southwest National Primate Research Center, San Antonio, Texas

⁴Department of Pathology and Microbiology, University of Nebraska Medical Center, Omaha, Nebraska

⁵Department of Biochemistry and Molecular Biology, University of Nebraska Medical Centre, Omaha, Nebraska

Correspondence

Siddappa N. Byrareddy, Department of Pharmacology and Experimental Neuroscience, University of Nebraska Medical Centre, Omaha, NE 68198-5880.

Email: sid.byrareddy@unmc.edu

ORCID

Raj H. Patel  <http://orcid.org/0000-0003-1896-1961>

Siddappa N. Byrareddy  <https://orcid.org/0000-0002-6889-4640>

REFERENCES

1. Harapan H, Itoh N, Yufika A, et al. Coronavirus disease 2019 (COVID-19): a literature review. *J Infect Public Health*. 2020;13(5):667-673. <https://doi.org/10.1016/j.jiph.2020.03.019>
2. Lesko CR, Bengtson AM. HIV and SARS-CoV-2: intersecting epidemics with many unknowns. *Am J Epidemiol*. 1975;229:1726. <https://doi.org/10.1093/aje/kwaa158>
3. Dauby N. Potential impact of COVID-19 in people living with HIV: experience from previous 21st century coronaviruses epidemics. *AIDS*. 2020;34(8):1255-1256.
4. Siordia JA Jr. Epidemiology and clinical features of COVID-19: a review of current literature. *J Clin Virol*. 2020;127:104357.
5. Härter G, Spinner CD, Roeder J, et al. COVID-19 in people living with human immunodeficiency virus: a case series of 33 patients [published online ahead of print May 11, 2020]. *Infection*. 2020.
6. Shalev N, Scherer M, LaSota ED, et al. Clinical characteristics and outcomes in people living with HIV hospitalized for COVID-19 [published online ahead of print May 30, 2020]. *Clin Infect Dis*. <https://doi.org/10.1093/cid/ciaa635>
7. Benkovic S, Kim M, Sin E. 4 Cases: HIV and SARS-CoV-2 Co-infection in patients from Long Island, New York [published online ahead of print May 19, 2020]. *J Med Virol*. <https://doi.org/10.1002/jmv.26029>
8. Wu Q, Chen T, Zhang H. Recovery from COVID-19 in two patients with coexisted HIV infection [published online ahead of print May 13, 2020]. *J Med Virol*. <https://doi.org/10.1002/jmv.26006>
9. Gervasoni C, Meraviglia P, Riva A, et al. Clinical features and outcomes of HIV patients with coronavirus disease 2019 [published online ahead of print May 14, 2020]. *Clin Infect Dis*. <https://doi.org/10.1093/cid/ciaa579>
10. YamamotoNorio, Yang Rongge, Yoshinaka Yoshiyuki, Amari Shinji, Nakano Tatsuya, Cinatl Jindrich, Rabenau Holger, Doerr Hans Wilhelm, Hunsmann Gerhard, Otaka Akira, Tamamura Hirokazu, Fujii Nobutaka, Yamamoto Naoki. HIV protease inhibitor nelfinavir inhibits replication of SARS-associated coronavirus. *Biochemical and Biophysical Research Communications*. 2004;318(3):719-725. <https://doi.org/10.1016/j.bbrc.2004.04.083>
11. Chan JF, Yao Y, Yeung ML, et al. Treatment with lopinavir/ritonavir or interferon-beta1b improves outcome of MERS-CoV infection in a nonhuman primate model of common marmoset. *J Infect Dis*. 2015; 212(12):1904-1913.
12. De Groot AS. How the SARS vaccine effort can learn from HIV-speeding towards the future, learning from the past. *Vaccine*. 2003; 21(27-30):4095-4104.
13. Blanco JL, Ambrosioni J, Garcia F, et al. COVID-19 in patients with HIV: clinical case series. *Lancet HIV*. 2020;7(5):e314-e316.
14. Chen J, Cheng X, Wang R, Zeng X. Computed tomography imaging of an HIV-infected patient with coronavirus disease 2019 (COVID-19). *J Med Virol*. 2020. <https://doi.org/10.1002/jmv.25879>
15. Cao B, Wang Y, Wen D, et al. A trial of lopinavir-ritonavir in adults hospitalized with severe COVID-19. *N Engl J Med*. 2020;382(19): 1787-1799.
16. Grein J, Ohmagari N, Shin D, et al. Compassionate use of remdesivir for patients with severe COVID-19. *N Engl J Med*. 2020;382(24): 2327-2336. <https://doi.org/10.1056/nejmoa2007016>
17. Elfiky AA. Ribavirin, remdesivir, sofosbuvir, galidesivir, and tenofovir against SARS-CoV-2 RNA dependent RNA polymerase (RdRp): a molecular docking study. *Life Sci*. 2020;253:117592.
18. COVID-19 Treatment Guidelines Panel. Coronavirus disease 2019 (COVID-19) treatment Guidelines. National Institutes of Health. <https://www.covid19treatmentguidelines.nih.gov/antiviral-therapy/lopinavir-ritonavir-and-other-hiv-protease-inhibitors/>. Accessed July 24, 2020.
19. Marimuthu J, Kumar BS, Aravind Gandhi P. HIV and SARS CoV-2 co-infection: a retrospective, record based, case series from South India [published online ahead of print Jul 7, 2020]. *J Med Virol*. <https://doi.org/10.1002/jmv.26271>
20. Vizcarra P, Pérez-Eliás MJ, Quereda C, et al. Description of COVID-19 in HIV-infected individuals: a single-centre, prospective cohort. *Lancet HIV*. 2020;S2352-3018(20):30164-30168.
21. Suwanwongse K, Shabarek N. Clinical features and outcome of HIV/SARS-CoV-2 coinfecting patients in The Bronx, New York city [published online ahead of print May 28, 2020]. *J Med Virol*. <https://doi.org/10.1002/jmv.26077>
22. Del Amo J, Polo R, Moreno S, et al. Incidence and severity of COVID-19 in HIV-positive persons receiving antiretroviral therapy: a cohort study [published online ahead of print Jun 26, 2020]. *Ann Intern Med*. <https://doi.org/10.7326/M20-3689>
23. Singu S, Acharya A, Challagundla K, Byrareddy SN. Impact of social determinants of health on the emerging COVID-19 pandemic in the United States. *Front Public Health*. 2020;8:406.