



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Management and Outcomes of Patients With STEMI During the COVID-19 Pandemic in China



Dingcheng Xiang, MD,<sup>a,\*</sup> Xin Xiang, PhD,<sup>b,\*</sup> Wei Zhang, PhD,<sup>c</sup> Shaodong Yi, MD,<sup>a</sup> Jinxia Zhang, MD,<sup>a</sup> Xiaolong Gu, MD,<sup>a</sup> Yawei Xu, MD,<sup>d</sup> Kai Huang, MD,<sup>e</sup> Xi Su, MD,<sup>f</sup> Bo Yu, MD,<sup>g</sup> Yan Wang, MD,<sup>h</sup> Weiyi Fang, MD,<sup>i</sup> Yong Huo, MD,<sup>j</sup> Junbo Ge, MD<sup>k</sup>

## ABSTRACT

**BACKGROUND** ST-segment elevation myocardial infarction (STEMI) is a fatal cardiovascular emergency requiring rapid reperfusion treatment. During the coronavirus disease-2019 (COVID-19) pandemic, medical professionals need to strike a balance between providing timely treatment for STEMI patients and implementing infection control procedures to prevent nosocomial spread of COVID-19 among health care workers and other vulnerable cardiovascular patients.

**OBJECTIVES** This study evaluates the impact of the COVID-19 outbreak and China Chest Pain Center's modified STEMI protocol on the treatment and prognosis of STEMI patients in China.

**METHODS** Based on the data of 28,189 STEMI patients admitted to 1,372 Chest Pain Centers in China between December 27, 2019 and February 20, 2020, the study analyzed how the COVID-19 outbreak and China Chest Pain Center's modified STEMI protocol influenced the number of admitted STEMI cases, reperfusion strategy, key treatment time points, and in-hospital mortality and heart failure for STEMI patients.

**RESULTS** The COVID-19 outbreak reduced the number of STEMI cases reported to China Chest Pain Centers. Consistent with China Chest Pain Center's modified STEMI protocol, the percentage of patients undergoing primary percutaneous coronary intervention declined while the percentage of patients undergoing thrombolysis increased. With an average delay of approximately 20 min for reperfusion therapy, the rate of in-hospital mortality and in-hospital heart failure increased during the outbreak, but the rate of in-hospital hemorrhage remained stable.

**CONCLUSIONS** There were reductions in STEMI patients' access to care, delays in treatment timelines, changes in reperfusion strategies, and an increase of in-hospital mortality and heart failure during the COVID-19 pandemic in China. (J Am Coll Cardiol 2020;76:1318-24) © 2020 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Listen to this manuscript's audio summary by Editor-in-Chief Dr. Valentin Fuster on JACC.org.

From the <sup>a</sup>Department of Cardiology, General Hospital of Southern Theatre Command of PLA, Guangzhou, China; <sup>b</sup>Graduate School of Arts and Sciences, Harvard University, Cambridge, Massachusetts; <sup>c</sup>Department of Biostatistics, School of Public Health, Fudan University, Shanghai, China; <sup>d</sup>Department of Cardiology, 10th Hospital of Tongji University, Shanghai, China; <sup>e</sup>Department of Cardiology, Union Hospital of Huazhong University of Science and Technology, Wuhan, China; <sup>f</sup>Department of Cardiology, Wuhan Asia Heart Hospital, Wuhan, China; <sup>g</sup>Department of Cardiology, Key Laboratories of Education Ministry for Myocardial Ischemia Mechanism and Treatment, 2nd Affiliated Hospital of Harbin Medical University, Harbin, China; <sup>h</sup>Department of Cardiology, Xiamen Heart Center, Xiamen, China; <sup>i</sup>Department of Cardiology, Shanghai Dongfang Hospital, Shanghai, China; <sup>j</sup>Department of Cardiology, Peking University First Hospital, Beijing, China; and the <sup>k</sup>Department of Cardiology, Zhongshan Hospital, Fudan University, Shanghai, China. \*Drs. Dingcheng Xiang and Xin Xiang contributed equally to this work. This study is supported by Ministry of Science and Technology of the People's Republic of China (2016YFC1301201). The authors have reported that they have no relationships relevant to the contents of this paper to disclose.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the JACC [author instructions page](#).

Manuscript received May 3, 2020; revised manuscript received June 17, 2020, accepted June 18, 2020.

**S**T-segment elevation myocardial infarction (STEMI), usually resulting from thrombotic occlusion of a coronary artery, is a fatal cardiovascular emergency requiring early diagnosis and rapid reperfusion therapy. The Chest Pain Center (CPC) was developed to accelerate this process (1,2). The growing coronavirus disease-2019 (COVID-19) pandemic caused by severe acute respiratory syndrome-coronavirus-2 poses a severe challenge to the care of STEMI patients. On the one hand, performing primary percutaneous coronary intervention (PCI)—the preferred reperfusion strategy recommended by most STEMI guidelines (3)—for a patient with unconfirmed COVID-19 status is a high-risk procedure that may expose health care workers and other hospitalized cardiovascular patients, who are particularly vulnerable if infected by severe acute respiratory syndrome-coronavirus-2 (4-7). On the other hand, the screening and infectious control procedures required to reduce the nosocomial spread of COVID-19 may substantially delay primary PCI and negatively impact patient prognosis (8,9). The scarcity of personal protective equipment and rapid testing for COVID-19 further exacerbates the problem, as most medical facilities could not afford to engage full personal protective equipment for all STEMI patients with confirmed COVID-19 status.

As a result, medical professionals across the globe have been debating about the extent to which to initiate COVID-19 screening protocols for STEMI patients and to adjust treatment procedures to prevent nosocomial infection. Organizations or experts in the United States, Italy, Australia, and New Zealand have mostly recommended continuing with existing primary PCI protocols for STEMI patients except for confirmed COVID-19 patients and persons under investigation or cases in which primary PCI could not be performed within required time frames (10-13). In contrast, experts in China, Iran, Palestine, and Jordan recommended prioritizing thrombolytic treatment for most patients with unconfirmed COVID-19 status (14-16).

However, there is no empirical study evaluating the impact of the COVID-19 pandemic and these different protocols on the prognosis of STEMI patients by the time of the writing of this paper, apart from 2 observational studies reporting 38% to 40% reduction in cardiac catheterization laboratory STEMI activities in Spain and United States and 1 small sample study reporting significant primary PCI delays in Hong Kong (8,17,18).

We draw on data from the China Chest Pain Center (CCPC) Database to evaluate the impact of the COVID-19 pandemic on the care of STEMI patients

and provide much needed empirical evidence for this challenge. Like those in other countries, CCPC's regular STEMI protocol recommends primary PCI as the standard of care for STEMI patients (1). Anticipating infectious control measures needed during the COVID-19 outbreak, the CCPC Executive Committee (CCPCEC) modified its STEMI protocol on January 23, 2020. This modified protocol recommended thrombolysis as the preferred reperfusion strategy for patients with unconfirmed COVID-19 status in areas affected by the outbreak to ensure timely reperfusion and prevent nosocomial infection (15). In this paper, we use data from the CCPC Database to evaluate how the COVID-19 pandemic and the related changes in STEMI protocols have affected access to cardiac care, reperfusion strategies, reperfusion efficiency, and prognosis of STEMI patients.

SEE PAGE 1325

## METHODS

**DATA SOURCE AND SAMPLE.** All data included in this study come from the CCPC Database, which was established in 2015. All Chinese hospitals with CPCs are required to register in this database and report the basic information, diagnosis and treatment procedures, and timelines of all patients presenting with acute chest pain. The CCPCEC uses the data in this database to accredit CPCs in China and monitor their operation.

The CPC of the General Hospital of the Southern Theatre Command created the first modified STEMI protocol—the “Treatment Protocol for Novel Coronavirus Pneumonia Patients with STEMI”—on January 23, 2020, the same day that the Chinese government announced the decision to lock down the epicenter Wuhan. CCPCEC immediately distributed and recommended this protocol to its network of 4,420 (accredited and unaccredited) CPCs across the country. On February 10, a refined version of this protocol was formally published as the “Chinese Expert Consensus on the Operation of Chest Pain Centers during the COVID-19 Pandemic” (15), jointly signed by the Chest Pain Committee of the Chinese College of Physicians, Chinese College of Cardiovascular Physicians, CCPCEC, and the Headquarters of the CCPC. The CCPC hosted 2 online training sessions on this protocol on February 7 and February 10, 2020, reaching a total of 24,705 participants.

In this study, we treated January 24, 2020, as the start time of the COVID-19 outbreak in China as well

## ABBREVIATIONS AND ACRONYMS

- CCPC** = China Chest Pain Center
- CCPCEC** = China Chest Pain Center Executive Committee
- CI** = confidence interval
- COVID-19** = coronavirus disease-2019
- CPC** = Chest Pain Center
- FMC-to-N** = time from first medical contact to needle (the beginning of thrombolysis)
- FMC-to-W** = time from first medical contact to wire crossing
- IQR** = interquartile range
- OR** = odds ratio
- PCI** = percutaneous coronary intervention
- STEMI** = ST-segment elevation myocardial infarction
- S-to-FMC** = time from symptom onset to first medical contact

**TABLE 1 Sample Baseline Characteristics**

	Hubei Sample			Non-Hubei Sample		
	Pre-Outbreak	Outbreak	p Value	Pre-Outbreak	Outbreak	p Value
Total STEMI cases reported	626	236	—	15,729	11,598	—
Included cases*	564	220	0.740	14,070	10,296	0.838
S-to-FMC ≤12	425 (75.35)	158 (71.82)		11,151 (79.25)	8,126 (78.92)	
Male	427 (75.7)	170 (77.3)	0.644	10,592 (75.3)	7,816 (76.0)	0.245
Age, yrs	62.86 ± 12.33	61.59 ± 13.10	0.203	62.91 ± 12.95	62.37 ± 12.81	0.001
Pattern of patient arrival			0.352			0.005
Walk-in	283 (50.2)	126 (57.3)		7,740 (55.0)	5,562 (54.0)	
EMS	51 (9.0)	16 (7.3)		1,498 (10.7)	1,150 (11.2)	
Transfer	208 (36.9)	71 (32.3)		4,536 (32.2)	3,420 (33.2)	
In-hospital onset	22 (3.9)	7 (3.2)		296 (2.1)	163 (1.6)	
S-to-FMC, h†	2.57 (1.12-7.67)	3.22 (1.34-7.86)	0.114	2.33 (1.05-6.16)	2.53 (1.15-6.38)	<0.001
FMC-to-N, min†	33.0 (20.0-52.0)	45.0 (27.0-83.0)	0.098	35.0 (24.0-62.0)	38.0 (26.0-70.0)	<0.001
FMC-to-W, min†	101.5 (74.0-187.0)	108.5 (75.5-204.5)	0.245	93.0 (69.0-154.0)	99.0 (73.0-159.0)	<0.001
Effective reperfusion	58 (68.2)	38 (67.9)	0.962	3,070 (79.4)	2,165 (78.7)	0.496
Timely reperfusion	214 (59.44)	66 (51.56)	0.121	6,261 (65.13)	4,224 (60.10)	<0.001
In-hospital mortality	26 (4.6)	16 (7.3)	0.137	566 (4.0)	480 (4.7)	0.015
In-hospital heart failure	78 (14.2)	40 (18.4)	0.139	1,794 (13.2)	1,404 (14.0)	0.081
Hemorrhage	4 (0.9)	4 (2.3)	0.176	108 (1.1)	68 (0.9)	0.320

Values are n, n (%), mean ± SD, or median (interquartile range). \*Cases are included if they were admitted during the sampling period, had complete key time point information, and had complete outcome information. †The z estimates are from the Mann-Whitney U test

EMS = emergency medical services; FMC-to-N = time from first medical contact to needle (the beginning of thrombolysis); FMC-to-W = time from first medical contact to wire crossing; S-to-FMC = time from symptom onset to first medical contact; STEMI = ST-segment elevation myocardial infarction.

as the implementation of the revised protocol and analyzed changes in the treatment and prognosis of STEMI patients before and after this point. We included in our analysis all STEMI cases admitted to the 1,372 accredited CPCs from the 4 weeks before and the 4 weeks after January 24, 2020 (December 27, 2019 to February 20, 2020). We excluded cases when the records for key time points were incomplete and when transfer to other medical facilities prevented the collection of outcome data. Our research protocol was approved by the Medical Ethics Committee of the General Hospital of Southern Theatre Command of PLA.

**KEY VARIABLES.** The key independent variables are the COVID-19 outbreak period (whether the patient was admitted between January 24, 2020 and February 20, 2020), Hubei (whether the patient was admitted to a CPC in Hubei province, the epicenter of the COVID-19 outbreak in China that accounted for 80.78% of the country's confirmed cases as of April 26, 2020), and the interaction of these 2 variables. We analyzed the impact of these independent variables on 4 groups of key outcome variables: 1) indicators of STEMI patients' access to care: the weekly total number of admitted STEMI cases at each accredited CPC and the time from symptom onset to first medical contact (S-to-FMC); 2) reperfusion strategy: primary PCI, thrombolysis, or conservative therapy; 3)

indicators of STEMI reperfusion efficiency: time from first medical contact to wire crossing (FMC-to-W) for patients undergoing primary PCI, time from first medical contact to needle (the beginning of thrombolysis) (FMC-to-N) for patients undergoing thrombolytic therapy, effective reperfusion (successful thrombolysis or Thrombolysis In Myocardial Infarction flow grade 3), and timely reperfusion (FMC-to-N ≤30 min or FMC-to-W ≤120 min); and 4) indicators of patient prognosis: in-hospital mortality, in-hospital heart failure, and in-hospital hemorrhage. We included patient age and sex as control variables for all models of individual-level variables.

**STATISTICAL PROCEDURES.** Continuous variables are summarized as mean ± SD for symmetric distributions and median (interquartile range [IQR]) for skewed distributions. Discrete variables are presented as frequency (percentage). We conducted Student's *t*-tests (for continuous variables) and chi-square tests (for discrete variables) to compare the characteristics of the outbreak sample and those of the pre-outbreak sample for Hubei and for other regions, respectively. We then estimated multivariate linear regression models for continuous outcome variables and logistic regression models for dichotomous outcome variables, except for using a mixed-effects model for weekly STEMI cases to account for dependence among the weekly numbers reported by

**TABLE 2 Results of Regression Models**

Model	Outcome Variable	n	Outbreak	Outbreak × Hubei
1	Hospitalized STEMI cases (per CPC per week)	7,145	-0.55 (-0.71 to -0.40)*	-0.50 (-1.37 to 0.37)
2	S-to-FMC, h	23,133	0.12 (-0.10 to 0.33)	0.88 (-0.48 to 2.24)
3	Primary PCI	21,205	0.76 (0.71 to 0.81)*	0.58 (0.38 to 0.89)†
4	Thrombolysis	21,205	1.67 (1.50 to 1.84)*	4.78 (2.45 to 9.34)*
5	Timely reperfusion	17,107	0.81 (0.76 to 0.86)*	0.90 (0.60 to 1.36)
6	Effective reperfusion	16,759	0.96 (0.85 to 1.08)	1.03 (0.49 to 2.13)
7	Total ischemic time, h	16,525	0.09 (-0.12 to 0.29)	-0.06 (-1.45 to 1.32)
8	FMC-to-W, min	15,280	4.44 (0.43 to 8.44)†	16.20 (-11.76 to 44.17)
9	FMC-to-N, min	2,602	4.54 (0.72 to 8.36)†	17.56 (-6.01 to 41.12)
10	In-hospital mortality	25,118	1.21 (1.07 to 1.37)‡	1.42 (0.73 to 2.76)
11	In-hospital heart failure	24,342	1.10 (1.02 to 1.18)†	1.30 (0.85 to 2.00)
12	In-hospital hemorrhage	18,031	0.87 (0.64 to 1.19)	2.93 (0.70 to 12.27)

Values are n or odds ratio/coefficient (95% confidence interval). Each row in the table represents a regression model. Models 2-12 included age and sex as control variables. \*p < 0.001. †p < 0.05. ‡p < 0.01.  
 CPC = Chest Pain Center; PCI = percutaneous coronary intervention; other abbreviations as in Table 1.

the same CPC. We also produced descriptive plots based on weekly means of outcome variables to provide an intuitive understanding of trends in the raw data. All statistical analysis was conducted using SAS software version 9 (SAS Institute, Cary, North Carolina).

**RESULTS**

A total of 28,189 STEMI cases were reported by 1,372 accredited CPCs during the 8-week sampling period. Among these cases, 25,150 (89.22%) met the inclusion criteria of having complete information regarding key time points and patient outcome. The final COVID-19 sample included slightly more male patients and slightly younger patients (Table 1). There were also statistically significant differences in key time points, proportion of effective reperfusion and timely reperfusion, and patient prognosis, which will be analyzed in the following 4 sections.

**COVID-19 AND STEMI PATIENTS' ACCESS TO CARE.**


**Hospitalized STEMI cases.** There was an approximately 26% drop in the weekly total number of hospitalized STEMI cases during the COVID-19 outbreak nationwide, and an approximately 62% drop in Hubei (Table 1). Regression analysis (Table 2) confirmed the negative and significant effect of the COVID-19 outbreak on the number of cases reported by each CCPC each week (coefficient: -0.55; 95% confidence interval [CI]: -0.71 to -0.40; p < 0.001), but the interaction term was not statistically significant (coefficient: -0.50; 95% CI: -1.37 to 0.37; p = 0.25). Our model estimated that each CCPC admitted 2.47 patients (95% CI: 1.78 to 3.16 patients) (Hubei) and 3.24 patients (95% CI: 3.12 to 3.36

patients) (non-Hubei) each week during the COVID-19 outbreak period, in contrast to 3.53 patients (95% CI: 3.02 to 4.04 patients) (Hubei) and 3.79 patients (95% CI: 3.69 to 3.89 patients) (non-Hubei) per center per week before the outbreak (Central Illustration).

**Time from symptom onset to first medical contact.** During the COVID-19 outbreak, the percentage of reported STEMI cases with FMC within 12 h of symptom onset dropped slightly in Hubei (Table 1), but this trend was not statistically significant in our regression analysis (Table 2). Our model estimated that mean S-to-FMC during the COVID-19 outbreak period was 6.75 h (IQR: 5.61 to 7.89 h) (Hubei) and 5.48 h (IQR: 5.30 to 5.66 h) (non-Hubei), in contrast to 5.66 h (IQR: 4.99 to 6.32 h) (Hubei) and 5.32 h (IQR: 5.16 to 5.48 h) (non-Hubei) before the outbreak.

**COVID-19 AND CHANGES IN REPERFUSION STRATEGIES AND EFFICIENCY FOR STEMI PATIENTS. Reperfusion strategies.**

The percentage of STEMI patients undergoing primary PCI dropped by one-half, whereas the percentage of thrombolysis increased sharply in Hubei during the COVID-19 outbreak (Central Illustration). There were similar but more moderate trends in the non-Hubei sample. Regression analysis (Table 2) confirmed these trends. The COVID-19 outbreak significantly reduced the probability of primary PCI (odds ratio [OR]: 0.76; 95% CI: 0.71 to 0.81; p < 0.001), with a much larger effect in Hubei province (interaction term OR: 0.57; 95% CI: 0.37 to 0.88; p < 0.001). In contrast, the COVID-19 outbreak had a substantial positive effect on the probability of thrombolysis (OR: 1.66; 95% CI: 1.50 to 1.84; p < 0.001), particularly in Hubei (interaction term OR: 4.78; 95% CI: 2.45 to 9.33; p < 0.001). These



results are consistent with the modified CCPC STEMI protocol that recommended prioritizing thrombolysis. **Timely reperfusion.** The proportion of STEMI patients receiving timely reperfusion therapy dropped

from 59.44% to 51.56% in Hubei during the COVID-19 outbreak and from 65.13% to 60.10% in other provinces. Regression analysis confirmed the negative impact of being admitted during the COVID-19

outbreak period (OR: 0.80; 95% CI: 0.76 to 0.86;  $p < 0.001$ ), but the interaction term between the COVID-19 outbreak period and Hubei province was not statistically significant (Table 2).

Further analysis of FMC-to-W and FMC-to-N revealed that the COVID-19 outbreak delayed primary PCI for 20.82 min in Hubei and 4.43 min in other provinces; the delay for thrombolytic treatment was 22.60 min in Hubei and 4.49 min in other provinces (Table 2). Overall, the impact of the COVID-19 outbreak on total ischemic time was not statistically significant in Hubei or in other provinces.

**Effective reperfusion.** Being admitted during the COVID-19 impact period had no significant effect on the probability of receiving effective reperfusion (defined as successful thrombolysis or reaching TIMI flow grade 3 after primary PCI), in Hubei or in other provinces (Table 2).

**IN-HOSPITAL MORTALITY AND HEART FAILURE RATES.** The rates of in-hospital mortality and in-hospital heart failure increased from 4.6% to 7.3% and from 14.2% to 18.4% in Hubei during the outbreak period (Table 1), while the increase for other provinces was much smaller. Regression analysis confirmed that being admitted during the COVID-19 outbreak period was associated with higher likelihood of in-hospital mortality (OR: 1.21; 95% CI: 1.07 to 1.37;  $p = 0.003$ ) and in-hospital heart failure (OR: 1.10; 95% CI: 1.02 to 1.18;  $p = 0.020$ ), but the interaction term between the COVID-19 outbreak period and Hubei province was not statistically significant (Table 2). The COVID-19 outbreak did not have an impact on the rate of in-hospital hemorrhage, in Hubei or in other parts of the country (Table 2).

## DISCUSSION

The decline in hospitalized STEMI cases and the slight-to-moderate increases in S-to-FMC, FMC-to-N and FMC-to-W during the COVID-19 outbreak suggest that the outbreak reduced STEMI patients' probability of accessing care and receiving timely reperfusion. These changes were more pronounced in Hubei province, the epicenter of the outbreak in China. Consistent with CCPC's modified STEMI protocol, there was an increase in the proportion of STEMI cases undergoing thrombolysis and a decrease in that of primary PCI during the outbreak, with no significant change in the proportion of STEMI cases receiving effective reperfusion therapy. The in-hospital mortality and in-hospital heart failure rate of STEMI patients increased during the COVID-19

outbreak in China, while the rate of in-hospital hemorrhage remained stable in Hubei as well as in other parts of the country.

Our findings provide much needed empirical evidence for health care professionals searching for a balance between optimizing timely treatment for STEMI patients and protecting health care workers and vulnerable cardiovascular patients from the risk of nosocomial COVID-19 infection. Despite the inevitable delays in treatment timeline due to mandatory infection control procedures and changes in reperfusion strategies during the outbreak, the proportion of patients receiving effective reperfusion remained stable.

**STUDY LIMITATIONS.** Our study has 3 important limitations. First and foremost, because there was little time lag between the start of China's response to the COVID-19 outbreak and the release of CCPC's modified STEMI protocol, this observational study cannot fully dissociate the effects of the CCPC's modified protocol from the other aspects of the COVID-19 outbreak. In other words, we cannot conclude the extent to which the switch to prioritizing thrombolysis for patients with unclear COVID status contributed to the deterioration of STEMI prognosis or attenuated the negative impact of other factors during the outbreak. Because most unobserved factors—such as mild patients delaying or avoiding hospital visits during the outbreak—tend to worsen the prognosis of hospitalized patients entering our sample, the results that we currently observe is likely an overestimation of the overall negative impact of the COVID-19 outbreak.

The second limitation is that we were unable to assess the extent to which CCPC's modified protocol protected health care workers and other vulnerable cardiovascular patients from COVID-19 infection due to the lack of complete data on nosocomial infection. Nonetheless, the information that we do have suggest that the protocol was likely effective in this regard. As of April 27, 2020, the CPCs in our network had reported only 2 cases of nosocomial infection related to procedures of primary PCI or thrombolysis during this pandemic, both occurring in catheterization labs during primary PCI in Wuhan, Hubei, before we released the modified protocol on January 23, 2020.

Last, but not least, the analysis presented in this paper did not account for the influence of the spring festival holiday, a major 2-week festival that coincided with the first 2 weeks of the COVID-19 outbreak period. The spring festival significantly influences



patient behavior, public transportation, and hospital operation. Because this study did not include a spring festival period control group, it cannot dissociate the impacts of the spring festival on STEMI treatment and prognosis from those of the COVID-19 outbreak in China.

## CONCLUSIONS

The COVID-19 outbreak led to a substantial drop in the number of admitted STEMI cases as well as delays in patients' access to care. Consistent with CCPC's recommendations, the proportion of patients undergoing primary PCI decreased and that of thrombolysis increased, but the proportion of patients receiving effective reperfusion therapy remained stable. The outbreak was associated with moderately increased in-hospital mortality and in-hospital heart failure, while the probability of in-hospital hemorrhage did not change.

**ACKNOWLEDGMENTS** The authors thank the staff members of all the CPCs in China for their hard work in caring for STEMI patients, and the staff of CCPC Headquarters for their full support in promoting the modified the STEMI protocol and in collecting data for this analysis. The authors are particularly grateful

for Wei Sun for providing invaluable advice for the statistical analysis process.

**ADDRESS FOR CORRESPONDENCE:** Dr. Dingcheng Xiang, Department of Cardiology, General Hospital of Southern Theatre Command of PLA, Liuhua Road 111, Guangzhou 510010, China. E-mail: [dcxiang@foxmail.com](mailto:dcxiang@foxmail.com). OR Dr. Yong Huo, Department of Cardiology, Peking University First Hospital, 8# St. Xishiku, Beijing, China. E-mail: [drhuoyong@163.com](mailto:drhuoyong@163.com).

## PERSPECTIVES

**COMPETENCY IN MEDICAL KNOWLEDGE:** The COVID-19 outbreak negatively affected STEMI patient's access to care, treatment timeline, and prognosis.

**TRANSLATIONAL OUTLOOK:** Future studies should compare the treatment and prognosis of STEMI patients in countries implementing different pandemic STEMI protocols to further evaluate the effects of different strategies.

## REFERENCES

- Zhang Y, Yu B, Han Y, et al. Protocol of the China ST-segment elevation myocardial infarction (STEMI) Care Project (CSCAP): a 10-year project to improve quality of care by building up a regional STEMI care network. *BMJ Open* 2019;9:e026362.
- Xiang D, Yi S. Chest pain centers in China: Current status and prospects. *Cardiol Plus* 2017;2:18-21.
- Ibanez B, James S, Agewall S, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation. *Eur Heart J* 2018;39:119-77.
- Rodríguez-Leor O, Cid-Alvarez B. STEMI care during COVID-19: losing sight of the forest for the trees. *J Am Coll Cardiol Case Rep* 2020 Apr 23 [E-pub ahead of print].
- Driggin E, Madhavan MV, Bikdeli B, et al. Cardiovascular considerations for patients, health care workers, and health systems during the coronavirus disease 2019 (COVID-19) pandemic. *J Am Coll Cardiol* 2020;75:2352-71.
- Zheng Y-Y, Ma Y-T, Zhang J-Y, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol* 2020;17:259-60.
- Ardati AK, Mena Lora AJ. Be prepared. *Circ Cardiovasc Qual Outcomes* 2020;13:e006661.
- Tam C-CF, Cheung K-S, Lam S, et al. Impact of coronavirus disease 2019 (COVID-19) outbreak on ST-segment-elevation myocardial infarction care in Hong Kong, China. *Circ Cardiovasc Qual Outcomes* 2020;13:e006631.
- Daniels MJ, Cohen MG, Bavry AA, Kumbhani DJ. Reperfusion of STEMI in the COVID-19 era - business as usual? *Circulation* 2020;141:1948-50.
- Szerlip M, Anwaruddin S, Aronow HD, et al. Considerations for cardiac catheterization laboratory procedures during the COVID-19 pandemic. *Catheter Cardiovasc Interv* 2020 Mar 25 [E-pub ahead of print].
- Zaman S, Maclsaac AI, Jennings GL, et al. Cardiovascular disease and COVID-19: Australian and New Zealand consensus statement. *Med J Aust* 2020 Jul 31 [E-pub ahead of print].
- Uccio FSD, Valente S, Colivicchi F, et al. ANMCO position paper: the network organization for the management of patients with acute coronary syndrome during the COVID-19 pandemic. *G Ital Cardiol* 2020;21:332-5.
- Mahmud E, Dauerman HL, Welt FG, et al. Management of acute myocardial infarction during the COVID-19 pandemic: a position statement from the Society for Cardiovascular Angiography and Interventions (SCAI), the American College of Cardiology (ACC), and the American College of Emergency Physicians (ACEP). *J Am Coll Cardiol* 2020;76:1375-84.
- Daralammouri Y, Azamta M, Hamayel H, et al. Recommendations for safe and effective practice of interventional cardiology during COVID-19 pandemic: expert opinion from Jordan and Palestine. *Palest Med Pharm J* 2020;5:65-73.
- Xiang D, Huo Y, Ge J. Expert consensus on operating procedures at chest pain centers in China during the coronavirus infectious disease-19 epidemic. *Cardiol Plus* 2020;5:21-32.
- Sadeghipour P, Talasaz AH, Estami V, et al. Management of ST-segment-elevation myocardial infarction during the coronavirus disease 2019 (COVID-19) outbreak: Iranian "247" National Committee's position paper on primary percutaneous coronary intervention. *Catheter Cardiovasc Interv* 2020 Apr 22 [E-pub ahead of print].
- Garcia S, Albaghdadi MS, Meraj PM, et al. Reduction in ST-segment elevation cardiac catheterization laboratory activations in the United States during COVID-19 pandemic. *J Am Coll Cardiol* 2020;75:2871-2.
- Rodríguez-Leor O, Cid-Alvarez B, Ojeda S, et al. The impact of the COVID-19 pandemic on healthcare activities in interventional cardiology units in Spain. *REC Interv Cardiol* 2020;2:82-9.

**KEY WORDS** chest pain center, COVID-19, primary percutaneous coronary intervention, ST-segment elevation myocardial infarction, thrombolysis