



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Editorial

Functional Limitations Post-COVID-19: A Comprehensive Assessment Strategy



Limitaciones funcionales post COVID-19: Una estrategia de evaluación integral

In late 2019, the new SARS-CoV-2 coronavirus began to spread rapidly, causing respiratory diseases of different severity.¹ Approximately 80% of cases are asymptomatic or mild, and about 20% cause severe respiratory compromises that need hospital management, eventually requiring invasive mechanical ventilation and life support,² which can lead to permanent sequelae and even death.¹

Muscle weakness is one of the most frequent problems in the Intensive Care Unit (ICU) and is often generated because of critical conditions that pose a vital risk. Intensive care unit-acquired weakness (ICUAW) is one factor related to muscle weakness and its prevalence, in people who survive a critical illness, is close to 40%.³ Critical illness survivors experience marked disability and deficiencies in physical and cognitive function that may even persist for years after their initial ICU stay.⁴ Post-ICU-acquired disability is associated with rising health costs and the use of more medical care, reduced health-related quality of life, and prolonged unemployment.⁵

Surviving older adults have limited activities of daily life (ADLs), reaching more significant limitations in basic ADLs such as walking, bathing, or dressing.⁶ Disability in ADLs is associated with increased risk of institutionalization, more significant health care expenditures, more hospitalizations, and higher mortality rates.⁴ Furthermore, post-critical illness patients present cognitive impairments and mental problems such as anxiety or depression that make returning to normal life more difficult.⁷

Recent SARS-CoV-2 experience has revealed the need for a multidisciplinary rehabilitation approach, especially for those patients with advanced age, obesity, comorbidities, and organ failure.⁸ Although the evaluation of patients with post-COVID-19 sequelae must include an exhaustive and complete medical evaluation, we consider that a specific assessment focused on functional rehabilitation must contemplate at least these five dimensions: (1) ADLs; (2) cognitive function; (3) physical function; (4) respiratory function; and (5) quality of life.

The dimension of ADLs and physical capacity must be assessed due to the consequences of extended bed rest and/or mechanical ventilation and, in some cases, the use of sedatives. The importance of ADLs lies in the fact that they allow patients to be easily stratified by level of dependency and/or activity limitations and to guide rehabilitation strategies by groups with similar needs.⁴ The ADLs and physical capacity evaluation are also a widely used

recommendation for assessing the health status of people in the ageing process (those most severely affected by COVID-19). There are tools to assess ADLs, such as the Barthel index or the Functional Independence Measure (FIM),^{9,10} as well as tests to assess physical fitness such as the six-minute walk test (6MWT) or the sit-to-stand test (STST) which have already been used in patients with COVID-19.^{10,11}

The cognitive dimension must be evaluated by the consequences due to the use of medicaments and/or presence of delirium. Hypoglycemia, hyperglycemia, delirium, and in-hospital acute stress symptoms have been identified as prognostic factors of persistent cognitive impairment after critical illness.⁷ Additionally, there is strong evidence that patients with delirium in the ICU are at a higher risk of long-term outcomes of cognitive dysfunction.³ The Mini-Mental State Examination (MMSE) and the Montreal Cognitive Assessment (MoCA) are some tools commonly used that also has the advantage of being able to be used through telemedicine platforms.¹²

The respiratory dimension must be evaluated due to the damage provoked by mechanical ventilation and the pathophysiological response to the virus. People who had received intensive care and had abnormal pulmonary function tests showed a restrictive pattern of pulmonary impairments at six months.⁴ On the other hand, there is a risk of ventilator-induced lung injury and subsequent lung fibrosis in persons that had received mechanical ventilation.⁴ The spirometry and the diffusion capacity of the lungs for carbon monoxide (DL_{CO}) determination have been used to assess the respiratory function in patients with COVID-19. Particularly, DL_{CO} has been shown to be diminished in about half of the patients.¹³

The quality of life dimension is necessary to assess because represent the perceptions, comfort, and feelings experienced during and after COVID-19 by the individual. We must not forget that one of the most crucial purposes of rehabilitation is to improve the quality of life by maintaining, improving, and reacquiring activities of daily living.⁷ There are some quality of life questionnaires, such as the Short Form-36 (SF-36) or the European Quality of Life-5 Dimensions (E5-QD) that have been used in this type of patient.^{10,14}

Another critical point in the assessment of post-COVID-19 patients is to consider the muscle function (peripheral/respiratory) that involves at least three dimensions. The peripheral muscles are crucial for carrying out the ADLs, for the physical capacity, and

the respiratory muscles are important for the respiratory pump function. Their evaluation has been included in rehabilitation programmes in the current context.^{9,10}

Additionally, other factors would be to influence the functional rehabilitation of post-COVID-19 patients and may affect performance during therapeutic interventions; for example, pain, sleep quality, nutritional status, mood, and necessity of returning to work, among other factors. These must be assessed according to the needs of each person.

The proposed model contemplates four different levels of severity: minimal, mild, moderate and severe functional limitations, which allow for adjusting the intervention according to the functional limitations present in each one. It is essential to classify the people into these groups because not all people will have the same impairments, mainly because the limitations to perform physical, cognitive, or functional tasks depend not only on the magnitude of the post-COVID 19 damage but also on the previous state of health, age, and comorbidities. We believe that in this way, the objectives can be better customized to the needs of each person.

Although there are other compromised systems, we think that these dimensions are the minimum to evaluate in a rehabilitation programme. However, if the patient requires it, other important aspects that influence rehabilitation must be evaluated, such as nutritional support, psychosocial support, swallowing, or referral to other medical specialities.

Finally, we call on the professionals involved in rehabilitation to evaluate at least these five dimensions so that we can carry out interventions adjusted to the reality of each patient and thus improve their quality of life.

Funding

Rodrigo Torres-Castro is funded by a grant from the National Agency for Research and Development (ANID) / Scholarship Program / DOCTORADO BECAS CHILE / 2018 – 72190117.

Conflict of interests

The authors declare no conflicts of interests.

References

1. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708–20. <http://dx.doi.org/10.1056/NEJMoa2002032>.
2. World Health Organization. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). 2020. Available from: [https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-.-\(covid-19\)](https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-.-(covid-19)) [cited 19.05.20].

3. Fan E, Cheek F, Chlan L, Gosselink R, Hart Nicholas, Herridge MS, et al. An official American Thoracic Society Clinical Practice guideline: the diagnosis of intensive care unit-acquired weakness in adults. *Am J Respir Crit Care Med*. 2014;190:1437–46. <http://dx.doi.org/10.1164/rccm.201411-2011ST>.
4. Ohtake PJ, Lee AC, Scott JC, Hinman RS, Ali NA, Hinkson CR, et al. Physical impairments associated with post-intensive care syndrome: systematic review based on the World Health Organization's International Classification of Functioning, Disability and Health Framework. *Phys Ther*. 2018;98:631–45. <http://dx.doi.org/10.1093/ptj/pzy059>.
5. Hermans G, Van den Berghe G. Clinical review: intensive care unit acquired weakness. *Crit Care*. 2015;19:274. <http://dx.doi.org/10.1186/s13054-015-0993-7>.
6. Iwashyna TJ, Ely EW, Smith DM, Langa KM. Long-term cognitive impairment and functional disability among survivors of severe sepsis. *JAMA*. 2010;304:1787–94. <http://dx.doi.org/10.1001/jama.2010.1553>.
7. Inoue S, Hatakeyama J, Kondo Y, Hifumi T, Sakuramoto H, Kawasaki T, et al. Post-intensive care syndrome: its pathophysiology, prevention, and future directions. *Acute Med Surg*. 2019;6:233–46. <http://dx.doi.org/10.1002/ams2.415>.
8. Brugliera L, Spina A, Castellazzi P, Cimino P, Tettamanti A, Houdayer E, et al. Rehabilitation of COVID-19 patients. *J Rehabil Med*. 2020;52:jrm00046. <http://dx.doi.org/10.2340/16501977-2678>.
9. Pancera S, Galeri S, Porta R, Pietta I, Bianchi LNC, Carrozza MC, et al. Feasibility and efficacy of the pulmonary rehabilitation program in a rehabilitation center: case report of a young patient developing severe COVID-19 acute respiratory distress syndrome. *J Cardiopulm Rehabil Prev*. 2020;40:205–8. <http://dx.doi.org/10.1097/HCR.0000000000000529>.
10. Liu K, Zhang W, Yang Y, Zhang J, Li Y, Chen Y. Respiratory rehabilitation in elderly patients with COVID-19: a randomized controlled study. *Complement Ther Clin Pract*. 2020;39:101166. <http://dx.doi.org/10.1016/j.ctcp.2020.101166>.
11. Simonelli C, Paneroni M, Fokom AG, Saleri M, Speltoni I, Favero I, et al. How the COVID-19 infection tsunami revolutionized the work of respiratory physiotherapists: an experience from Northern Italy. *Monaldi Arch Chest Dis*. 2020;90(2.). <http://dx.doi.org/10.4081/monaldi.2020.1085>.
12. Hantke NC, Gould C. Examining older adult cognitive status in the time of COVID-19. *J Am Geriatr Soc*. 2020;68:1387–9. <http://dx.doi.org/10.1111/jgs.16514>.
13. Frija-Masson J, Debray MP, Gilbert M, Lescure FX, Travert F, Borie R, et al. Functional characteristics of patients with SARS-CoV-2 pneumonia at 30 days post infection. *Eur Respir J*. 2020;2020:2001754. <http://dx.doi.org/10.1183/13993003.01754->.
14. Ping W, Zheng J, Niu X, Guo C, Zhang J, Yang H, et al. Evaluation of health-related quality of life using EQ-5D in China during the COVID-19 pandemic. *PLOS ONE*. 2020;15:e0234850. <http://dx.doi.org/10.1371/journal.pone.0234850>.

Rodrigo Torres-Castro^{a,*}, Lilian Solis-Navarro^a,
Mercè Sitjà-Rabert^b, Jordi Vilaró^b

^a Department of Physical Therapy, Faculty of Medicine, University of Chile, Santiago, Chile

^b Blanquerna School of Health Sciences, Global Research on Wellbeing (GRoW), Universitat Ramon Llull, Barcelona, Spain

* Corresponding author.

E-mail address: klgorodrigotorres@gmail.com (R. Torres-Castro).