


# Extracellular Vesicle-Related Thrombosis in Viral Infection

This article was published in the following Dove Press journal:  
*International Journal of General Medicine*

Shosaku Nomura<sup>1</sup>  
Takehito Taniura<sup>2</sup>  
Tomoki Ito <sup>1</sup>

<sup>1</sup>First Department of Internal Medicine, Kansai Medical University, Hirakata, Japan; <sup>2</sup>Division of Internal Medicine, Daiwa Hospital, Suita, Japan

**Abstract:** Although the outcomes of viral infectious diseases are remarkably varied, most infections cause acute diseases after a short period. Novel coronavirus disease 2019, which recently spread worldwide, is no exception. Extracellular vesicles (EVs) are small circulating membrane-enclosed entities shed from the cell surface in response to cell activation or apoptosis. EVs transport various kinds of bioactive molecules between cells, including functional RNAs, such as viral RNAs and proteins. Therefore, when EVs are at high levels, changes in cell activation, inflammation, angioplasty and transportation suggest that EVs are associated with various diseases. Clinical research on EVs includes studies on the coagulatory system. In particular, abnormal enhancement of the coagulatory system through EVs can cause thrombosis. In this review, we address the functions of EVs, thrombosis, and their involvement in viral infection.

**Keywords:** viral infection, thrombosis, extracellular vesicle, exosome, microvesicle

## Introduction

Most viruses are small sized (typically 0.02 to 0.3  $\mu\text{m}$ ), unlike large and megaviruses whose maximum lengths can reach 1  $\mu\text{m}$  in length at the maximum.<sup>1–3</sup> The reproduction of a virus depends on bacterial, plant, and animal cells including those in humans.<sup>4</sup> Viruses are classified according to genomic properties and structures as well as their reproduction method, and not according to the disease that each virus causes.<sup>5</sup> Replication in DNA viruses typically occurs in the nucleus of the host cell, whereas RNA virus replication typically occurs in the cytoplasm.<sup>5,6</sup> There are exceptions, however. For example, H1N1 (an RNA virus) cannot reproduce in the host's cytoplasm, whereas vaccinia (a DNA virus) does not need the nucleus to reproduce. After reproduction of a complete virus particle, the host cell typically perishes and the virus is released to infect other host cells.<sup>4–6</sup> Although the outcomes of viral infectious diseases are remarkably varied, most infections cause acute disease after a short period.<sup>7</sup> Novel coronavirus disease 2019 (COVID-19), which recently spread worldwide, is no exception.<sup>8,9</sup>

Extracellular vesicles (EVs) are small circulating membrane-enclosed entities shed from cell surface in response to cell activation or apoptosis.<sup>10</sup> Although detailed understanding of EVs is lacking, information about EVs has been accumulating.<sup>11–13</sup> EVs measure 0.01–4  $\mu\text{m}$  and are generated by various processes.<sup>11,12</sup> EVs transport various types of molecules between cells, such as viral RNAs and proteins.<sup>14–16</sup> Therefore, when EVs are at high levels, cell activation, inflammation, angioplasty and states involving transportation occur, indicating the association of EVs with

Correspondence: Shosaku Nomura  
First Department of Internal Medicine,  
Kansai Medical University, 2-3-1  
Shinmachi, Hirakata, Osaka 573-1010,  
Japan  
Tel + 81 72 804 2754  
Fax + 81 72 804 2041  
Email shosaku-n@mbp.ocn.ne.jp

various diseases.<sup>11,14,17–19</sup> Clinical research of EVs includes studies on the coagulatory system. In particular, abnormal enhancement of the coagulatory system by EVs can cause thrombosis and disseminated intravascular coagulation (DIC). Such an abnormality after a viral infectious disease can become a significant clinical problem. In this review, we address the functions of EVs, thrombosis, and their involvement in viral infection.

## Classification of EVs

EVs are classified into three groups by size, namely exosomes, microvesicles (MVs), and apoptotic bodies (ABs) (Table 1).<sup>11</sup> Exosomes are EVs with diameters of 30–200 nm, and these membrane-bound vesicles can be precipitated by ultracentrifugation at 100,000×g.<sup>11,12,15,20</sup> Most exosomes carry specific proteins reflecting the characteristics of the origin cell.<sup>21–23</sup> They form through multi-vesicular bodies (MVBs) that fuse with the cell membrane for release.<sup>15,24,25</sup> Exosomes are released from a cell via a mechanism of endosomal complexes required for transport (Figure 1).<sup>26</sup>

MVs are EVs of 10–1000 nm in size.<sup>11</sup> MVs are also called microparticles (MPs).<sup>10,11,27</sup> The differences between MVs and exosomes other than their size are the processes of formation and secretion.<sup>15,28</sup> MVs are generated by the surface of the cell breaking off, which is controlled by cell activation.<sup>29</sup> The plasma membrane includes several kinds of phospholipids.<sup>11,30</sup> The internal leaflet contains aminophosphatides (eg, phosphatidylserine:PS) for a negative charge.<sup>30</sup> MV biogenesis, which occurs via blebbing, is a fragmentation phenomenon whereby nascent MVs are released into the extracellular space via pinching off from the plasma membrane.<sup>31,32</sup> MVs contain distinct protein and lipid components from the plasma membrane.<sup>32</sup> During cell activation, the charge of the leaflet changes the structure of the normal lipid

layer, and PS is exposed to MV (Figure 1).<sup>33</sup> This leads to procoagulant activity in the MVs.<sup>34</sup>

EVs generated by apoptosis are called apoptotic bodies (ABs).<sup>11,35,36</sup> During its final phase of apoptotic death, the cell divides into several Ab,<sup>32</sup> the size of which are 1000–3000 nm.<sup>37,38</sup> Similar to MVs, ABs are formed by PS moving to the cell surface.<sup>33–35</sup> ABs may contain a wide variety of cellular components, such as micronuclei, chromatin remnants, cytosol portions, degraded proteins, DNA fragments, and even intact organelles.<sup>32</sup> The main difference between ABs and MVs is the existence of materials derived from the nucleus such as histones and DNA fragments (Figure 1).<sup>39,40</sup>

## Function of EVs

EVs can carry activated coagulation factors, by expressing phosphatides on their surface.<sup>11</sup> Therefore, it is thought that the existence of EVs is related to several diseases, indicating a coagulatory promotion tendency.<sup>11,27</sup> Because EVs promote intravascular coagulation to support thrombin generation, they may be linked with coagulation abnormalities.<sup>27</sup> The procoagulant activity observed on platelet-derived EV surfaces is 50 to 100-fold higher than that observed on activated platelets.<sup>41</sup> This suggests that the coagulatory promotion by EVs is an important defense mechanism for bleeding risk.<sup>11</sup> EVs also carry tissue factor (TF) that is important to activate the coagulation system.<sup>42</sup>


The main characteristics of atherosclerosis are the adhesion of monocytes to the endothelium and movement of monocytes into the subendothelium.<sup>27</sup> When a monocyte is activated by platelet-derived EVs and adheres to the endothelium, several inflammatory cytokines are generated.<sup>43</sup> Furthermore, stimulation or activation of the endothelium by EVs increases the expression of adhesion molecules on the endothelial surface.<sup>43,44</sup> Atherosclerotic lesions can develop and progress in severity via the apoptosis of endothelial cells,

**Table 1** Population and Characteristics of EVs

	Exosome	MV (MP)	AB
Size	30–200 nm	100–1000 nm	1000–3000 nm
Shape	Homogeneous	Variable	Variable
Origine	MBV fusion with the plasma membrane	Budding from the plasma membrane	Budding from the plasma membrane
Markers	Tetraspanins (CD9, CD63, CD81) Alix, TSG101, HSP70	Annexin V, Integrin, Selectin, CD40 ligand, metalloproteinase	Annexin V, DNA fragment Caspase 3, Histones
Isolation	Ultracentrifugation (100,000g)	Ultracentrifugation (10,000–100,000g)	Ultracentrifugation (6000–10,000g)
P-Act	Weak	Powerful	Powerful

**Notes:** Data from Nomura.<sup>11</sup> All vesicles preparations are heterogeneous with different protocols allowing the enrichment of one type over another.

**Abbreviations:** MV, microvesicle; MP, microparticle; AB, apoptotic body; MBV, multivesicular body; TSG101, tumor susceptibility gene 101; HSP70, heat shock protein 70; P-Act, procoagulant activity.



**Figure 1** Convergence of EV and virus biogenesis. Original cell owns the endocytic and secretory pathways. Viruses share effectors of EV production for their assembly and release. Exosomes produced in the MVB and shed MV (MP) budding of the plasma membrane. Apoptotic cell finally releases shedding AB.

**Abbreviations:** MV, microvesicle; MP, microparticle; AB, apoptotic body; N, nucleus; MBV, multivesicular body; ER, endoplasmic reticulum.

a process induced by a substantial number of EV-dependent coagulatory factors.<sup>11,27,45</sup>

EV appear to constitute a new system of cell-cell communication.<sup>16,46,47</sup> EVs have various important functions such as coagulatory promotion, immunosuppression, and angioplasty.<sup>48,49</sup> EVs may be the most suitable mechanism through which cells communicate with others.<sup>15,48</sup> For example, EVs produced by one kind of cell stimulate another specific cell.<sup>50,51</sup> EVs carry tetraspanin protein and may employ a mechanism that can return in a specific organization.<sup>52,53</sup> Another mechanism involves fusion with the cell membrane, which results in the transfer of mRNA, micro(mi)RNA, proteins, and signaling molecules by EVs.<sup>54,55</sup> The existence of miRNAs has been found in EVs released with biological fluid of patients with various viral infectious diseases.<sup>13,56–58</sup> EVs might play a crucial role in dissemination of pathogens as well as host-derived molecules during infection.<sup>58</sup> Therefore, EVs may be strongly involved in progression of the post-viral condition and the origin of complications.<sup>59</sup>


## Viral Infection and Thrombosis

Ebola, H1N1 influenza, cytomegalovirus, chickenpox – herpes zoster, hepatitis C virus, human immunodeficiency

virus (HIV), coxsackie virus B3, herpes simplex virus-1, dengue, and Junin virus are accompanied by thrombotic complications and bleeding.<sup>60–71</sup> There are at least two major factors underlying the onset of thrombosis that are associated with these viral infections (Figure 2).<sup>72,73</sup> One factor influencing the blood vessel system is the viruses themselves, because they can influence monocytes, neutrophils and the blood vessel endothelium and also induce the expression of TF.<sup>74</sup> Some viruses can also directly influence platelets and enhance PS expression during their infections.<sup>75–77</sup> Plasma from influenza patients may also contain MVs with TF activity.<sup>78</sup> This leads to strong thrombosis onset, by enhancing the activity of the exogenous coagulatory system. The second mechanism influences the immune system. The viral infection influences the immune system, and inflammation with the cell which immunity decreased spreads, and an imbalance between coagulation and anti-coagulation occurs.<sup>79</sup> This mechanism is related to interactions between the virus and Toll-like receptors (TLRs).<sup>70,79</sup>

## Ebola Virus (EBOV)

EBOV is a negative-sense RNA virus that causes a severe disease characterized by high fever, diarrhea, and unexpected



**Figure 2** EV mediated thrombotic event pathway.

**Abbreviations:** EV, extracellular vesicle; MV, microvesicle; MP, microparticle; EC, endothelial cell; PAI-1, plasminogen activator inhibitor-1; VWF, von Willebrand factor.

onset of vomiting.<sup>80</sup> Additionally, this virus causes a serious procoagulatory abnormality with liver damage.<sup>80</sup> EBOV infection induces TF expression in infected cells and serum D-dimer rises, finally causing DIC.<sup>81</sup> EBOV disrupts the functions of dendritic cells coordinating a T cell increase.<sup>82,83</sup> Therefore, it is thought that EBOV infectious disease develops a thrombotic tendency by failure of the immune system and enhancement of coagulation.<sup>70,84</sup>

## Influenza A Virus (IAV)

Influenza A virus (IAV) is a negative-sense RNA virus that commonly causes death.<sup>85</sup> Cardiovascular system disorders, such as acute myocardial infarction, deep vein thrombosis, and pulmonary embolism, are observed.<sup>86–90</sup> Disorder of the blood vessel wall barrier by IAV may contribute to the development of pulmonary damage in patients with influenza.<sup>91,92</sup> H5N1-infected chickens show microthrombosis and thrombocytopenia.<sup>93</sup> IAV infectious disease causes a high level of plasma von Willebrand factor (vWF) and a decrease of a disintegrin-like and metalloproteinase with thrombospondin type 1 motifs 13 (ADAMTS13), which may result in clot-related microangiopathy.<sup>94,95</sup> Therefore, IAV infectious disease is thought to cause immunity and aggravation of the wall system, which results in inflammatory induction and coagulatory enhancement.

## Human Immunodeficiency Virus (HIV)

Human immunodeficiency virus (HIV) is a positive-sense single-stranded enveloped RNA virus of the Retroviridae family.<sup>96</sup> One cause of death in HIV-infected patients is cardiovascular disease (CVD), and functional disorder in the blood

walls caused by HIV reproduction is a considerable determinant of it.<sup>96–98</sup> Additionally, the immune reaction and inflammation caused by HIV infection may be cardiovascular risks.<sup>99,100</sup> The death rate is strongly related to interleukin (IL)-6, high sensitivity C-reactive protein (hsCRP), and D-dimer. In particular, IL-6 and hsCRP are related to the development of acquired immunodeficiency syndrome.<sup>101,102</sup> Funderburg et al<sup>103,104</sup> reported that TF expression on the surface of monocytes increases in HIV patients. Additionally, Harley et al<sup>105</sup> reported that T-cell kinetics and activity of the thrombin-PAR1 signaling axis are increased by proinflammatory cytokines during HIV infection and contribute to adaptive immunoreactions. Furthermore, young patients with HIV infection have a high level of vWF and a low level of ADAMTS13, which are related to stroke.<sup>106</sup> Therefore, HIV infection has a thrombotic risk through various mechanisms.<sup>107,108</sup>

## Hepatitis C Virus (HCV)

Hepatitis C virus (HCV) is a positive-sense, single-stranded RNA virus of the Flaviviridae family. HCV infection carry the risk of thrombosis, which increases with expression of TF, fibrinolysis interference, and increased platelet aggregation and activation.<sup>109</sup> HCV viral RNA activates TLR-3 in endothelial cells (ECs), leading to inflammation and expression of tumor necrosis factor (TNF)- $\alpha$ .<sup>110</sup> Damaged ECs can affect immune cells through CXCL12 chemokine expression.<sup>111</sup> Through these mechanisms, increased TNF- $\alpha$  induces expression of TF and downregulation of thrombomodulin (TM).<sup>112</sup> Hodowanec et al<sup>113</sup> confirmed that elevation of TF expression and enhancement of coagulatory activity increase in some chronic HCV patients. Similarly, during chronic HCV infectious disease, wall functional disorder caused by plasma vWF levels increases.<sup>114</sup>

## Coxsackievirus (CoxV)

Coxsackievirus (CoxV) is negative-sense RNA virus of the Picornaviridae family.<sup>70</sup> It is accompanied by a considerable risk of thromboembolism caused by myocarditis.<sup>115</sup> Additionally, myocarditis coagulopathy and hepatic necrosis occur, and ventricular clot formation increases significantly because of platelet activation.<sup>116</sup>

## Herpes Simplex Virus (HSV)

Herpes simplex virus (HSV) is double-stranded, linear DNA genome virus of the herpesvirus family.<sup>70</sup> Infection of blood vessel ECs by HSV increases TF activity and reduces TM

expression.<sup>69,117</sup> Sutherland et al<sup>118</sup> reported that HSV type 1 (HSV-1) is a cofactor for PAR-1 that induces TF and glycoprotein C, causing thrombin production. HSV-1 and HSV-2 move FX to the cell surface via internal mechanism phosphatides.<sup>117–119</sup> It is thought that these viruses drive thrombin generation on the cell surface via this mechanism.<sup>117</sup>

## Epstein-Barr Virus (EBV)

Epstein-Barr virus (EBV) belongs to the herpesvirus family and has double-stranded DNA. Its name was changed to human herpesvirus type 4 (HHV-4), but the former name is still widely used.<sup>70</sup> EBV triggers autoantibody-producing autoimmunity in response to various autoimmune diseases.<sup>120</sup> Patients with EBV experience portal vein thrombosis caused by hypercoagulable syndrome.<sup>121</sup>

## Cytomegalovirus (CMV)

Cytomegalovirus (CMV) has double-stranded DNA with the generic name of herpesvirus characterized by forming a characteristic inclusion body of the observable “eyes of the owl” state in the nucleus of the host cell under an optical microscope. CMV transforms monocytes and induces production of inflammatory cytokines. Therefore, the incidence of thrombosis in patients with acute CMV infection is common.<sup>121,122</sup>

## Corona Virus (CoV)

Corona virus (CoV) causes disease in mammals and birds, and has a single strand plus chain RNA genome.<sup>123</sup> In 2003, a CoV epidemic caused by severe acute respiratory syndrome CoV (SARS-CoV-1) emerged in China.<sup>124</sup> SARS-CoV-1 was associated with severe thrombotic complications.<sup>124</sup> Several SARS-CoV-1 cases exhibited pulmonary embolism and deep vein thrombosis.<sup>125</sup> The characteristics of SARS-CoV-1 are a prolonged prothrombin time, prolonged activated partial thromboplastin time, elevated D-dimer, and worsening thrombocytopenia.<sup>126</sup> These findings are consistent with DIC. Furthermore, interestingly, thrombopoietin levels increase in SARS-CoV-1 patients at the convalescent phase compared with normal controls.<sup>127</sup> These findings have also been reported in patients with septic DIC.<sup>128</sup>

Another CoV infection was reported in 2012.<sup>129</sup> This CoV was responsible for “Middle East respiratory syndrome” (MERS-CoV).<sup>129</sup> MERS-CoV is similar to SARS-CoV-1 and associated with thrombotic complications. Specifically, DIC is one of the major complications reported in fatal MERS-CoV cases.<sup>130</sup>

In 2019, another CoV caused a global pandemic. CoV disease 2019 (COVID-19) is a novel CoV strain disease.<sup>8,9</sup> COVID-19 patients suffer from severe respiratory or systemic manifestations. Therefore, COVID-19 is also called SARS-CoV-2.<sup>131</sup> Thrombotic complications also emerge in patients with COVID-19.<sup>132–135</sup> COVID-19 patients have elevated D-dimer levels, prolonged prothrombin times, and thrombocytopenia similar to DIC.<sup>132,136–138</sup> Both elevated D-dimer levels and thrombocytopenia can be explained by excessive activation of platelets and the coagulation cascade. In contrast, phagocytosis or direct viral targeting by the immune system can also cause these abnormal clinical findings. Thus, viral infections elicit systemic inflammatory immune responses resulting in imbalanced coagulation.<sup>70</sup>

## EVs and Coagulatory Abnormalities During Viral Infection

EVs from a virally infected cell include not only the virus, but also information on the host.<sup>139</sup> Therefore, EVs are involved in interactions between a virus and various cells of the host. As we have mentioned, the function and role of EVs are varied.<sup>15,27,46,47</sup> Most importantly, EVs that participate in post-viral coagulatory abnormalities may be MVs (or MPs), because it appears that most characteristic functions of MVs promote coagulation depending on TF (Figure 2). In addition, MVs can substantially increase procoagulant activity on platelet surfaces, thereby contributing to coagulant abnormality during viral infection.<sup>11,41</sup> In contrast, exosomes contain many functional features, one of which concerns the immune system (Figure 2).<sup>56,139–142</sup> For example, lymphatic exosomes promote dendritic cell migration along guidance cues, thereby regulating the immune system.<sup>142</sup> Therefore, it is possible that exosomes do not influence the coagulatory system.

EVs are involved in some aspects of viral infectious disease.<sup>143</sup> After infection, for example, EBOV packages the protein moiety characteristic of this virus in EVs. Apoptosis of immune cells in the host is guided by inflammatory cytokines.<sup>144</sup> This drives failure of the homeostasis mechanism in blood vessels, leading to coagulatory abnormalities such as vein clots and DIC. Airway epithelial cells release EVs that neutralize human influenza virus.<sup>145</sup> This mechanism of EVs might play an important role in defense against respiratory viruses.<sup>146</sup> EVs after HIV infection have a significant influence on the immune system.<sup>147–149</sup> CMV-related EVs are useful to reinforce infectivity of CMV.<sup>143,150</sup> Human CMV infection is controlled by T cell-mediated immunity and CMV infects ECs.<sup>151,152</sup> One of the functions

of EC-derived EVs after CMV infection might be contributing to innate surveillance.<sup>151</sup>

One role played by post-viral EVs exosomes is their participation in maintaining the pathological state and promoting the spread of viral infection. The main roles of MVs and MPs appear to be in coagulation abnormality.<sup>59,153–161</sup> Regarding COVID-19, reports involving EVs are still rare.<sup>162–165</sup> However, a similar role of exosomes and MVs is assumed in COVID-19 as other virus infectious diseases.<sup>166–170</sup> In particular, concerning thrombosis, there is no doubt that COVID-19 causes more symptoms in comparison with the past virus infectious disease.<sup>171</sup> Accumulation of important reports of EVs in conjunction with COVID-19 is expected in the future.

## Conclusions

We have addressed the functions of EVs in thrombosis, and their involvement in viral infections. Clinical research on EVs should include studies on the coagulatory system. In particular, abnormal enhancement of the coagulatory system by EVs can cause thrombosis and DIC (Figure 2). There are reports about EVs in some virus infectious diseases. Apoptosis of immune cells in the host is guided by inflammatory cytokines. As a role of post-viral EVs, exosomes participate in pathological maintenance and infection spreading. The main roles of MVs appear to be in coagulation abnormalities. Although many viral infections involve EVs, the role of EVs in COVID-19 is unclear. Nevertheless, thrombosis is a major problem that affects the prognosis of COVID-19 patients. Accumulation of important reports of EVs in conjunction with COVID-19 is expected in the future.

## Acknowledgments

This review was supported in part by grants (15K08657 and 19K07948 to S.N.) from the Ministry of Education, Culture, Science and Technology of Japan. We thank Edanz Group for editing a draft of this manuscript.

## Disclosure

The authors declare no conflicts of interest regarding the publication of this study.

## References

- La Scola B, Audic S, Robert C, et al. A giant virus in amoebae. *Science*. 2003;299(5615):2033. doi:10.1126/science.1081867
- Iyer LM, Balaji S, Koonin EV, Aravind L. Evolutionary genomics of nucleocapsid cytoplasmic large DNA viruses. *Virus Res*. 2006;117(1):156–184.
- Koonin EV, Yutin N. Evolution of the large nucleocytoplasmic DNA viruses of eukaryotes and convergent origins of viral gigantism. *Adv Virus Res*. 2019;103:167–202.
- Ali H, Ali AA, Atta MS, Cepica A. Common, emerging, vector-borne and infrequent abortogenic virus infections of cattle. *Transbound Emerg Dis*. 2012;59(1):11–25.
- Poirier EZ, Vignuzzi M. Virus population dynamics during infection. *Curr Opin Virol*. 2017;23:82–87.
- Tan B, Gao SJ. RNA epitranscriptomics: regulation of infection of RNA and DNA viruses by N<sup>6</sup>-methyladenosine (m<sup>6</sup>A). *Rev Med Virol*. 2018;28(4):e1983.
- Mayo MA, Pringle CR. Virus taxonomy – 1997. *J Gen Virol*. 1998;79(Pt 4):649–657.
- Chen Y, Liu Q, Guo D. Emerging coronaviruses: genome structure, replication, and pathogenesis. *J Med Virol*. 2020;92(4):418–423.
- Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med*. 2020;382(13):1199–1207.
- Nomura S, Ozaki Y, Ikeda Y. Function and role of microparticles in various clinical settings. *Thromb Res*. 2008;123(1):8–23. doi:10.1016/j.thromres.2008.06.006
- Nomura S. Extracellular vesicles and blood diseases. *Int J Hematol*. 2017;105(4):392–405. doi:10.1007/s12185-017-2180-x
- Raposo G, Stoovogel W. Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol*. 2013;200(4):373–383. doi:10.1083/jcb.201211138
- Ciardello C, Cavallini L, Spinelli C, et al. Focus on extracellular vesicles: new frontiers of cell-to-cell communication in cancer. *Int J Mol Sci*. 2016;17(2):175–191. doi:10.3390/ijms17020175
- Valadi H, Ekström K, Bossios A, et al. Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat Cell Biol*. 2007;9(6):654–659. doi:10.1038/ncb1596
- They C, Ostrowski M, Segura E. Membrane vesicles as conveyors of immune responses. *Nat Rev Immunol*. 2009;9(8):581–593.
- Hoehn EN, Cremer T, Gallo RC, Margolis LB. Extracellular vesicles and viruses: are they close relatives? *Proc Natl Acad Sci U S A*. 2016;113(33):9155–9161.
- Huang ME, Leonard JN. A platform for actively loading cargo RNA to elucidate limiting steps in EV-mediated delivery. *J Extracell Vesicles*. 2016;5:31027.
- Iraci N, Leonardi T, Gessler F, Vega B, Pluchino S. Focus on extracellular vesicles: physiological role and signalling properties of extracellular membrane vesicles. *Int J Mol Sci*. 2016;17(2):171.
- Alberro A, Sáenz-Cuesta M, Muñoz-Culla M, et al. Inflammation and frailty status do not result in an increased extracellular vesicle concentration in circulation. *Int J Mol Sci*. 2016;17(7):1168.
- Conde-Vancells J, Rodríguez-Suárez E, Embade N, et al. Characterization and comprehensive proteome profiling of exosomes secreted by hepatocytes. *J Proteome Res*. 2008;7(12):5157–5166.
- Mathivanan S, Ji H, Simpson RJ. Exosomes: extracellular organelles important in intercellular communication. *J Proteomics*. 2010;73(10):1907–1920.
- Alvarez-Erviti L, Seow Y, Yin H, Betts C, Likhani S, Wood MJ. Delivery of siRNA to the mouse brain by systemic injection of targeted exosomes. *Nat Biotechnol*. 2011;29(4):341–345.
- Baietti MF, Zhang Z, Mortier E, et al. Syndecan-synternin-ALIX regulates the biogenesis of exosomes. *Nat Cell Biol*. 2012;14(7):677–685.
- György B, Szabó TG, Pásztói M, et al. Membrane vesicles, current state-of-the-art: emerging role of extracellular vesicles. *Cell Mol Life Sci*. 2011;68(16):2667–2688.

25. Muhsin-Sharafaldine MR, Saunderson SC, Dunn AC, Faed JM, McLellan AD. Procoagulant and immunogenic properties of melanoma exosomes, microvesicles and apoptotic vesicles. *Oncotarget*. 2016;7(35):56279–56294.
26. Colombo M, Moita C, van Niel G, et al. Analysis of ESCRT functions in exosome biogenesis, composition and secretion highlights the heterogeneity of extracellular vesicles. *J Cell Sci*. 2013;126(Pt 24):5553–5565.
27. Nomura S. Microparticle and atherothrombotic diseases. *J Atheroscler Thromb*. 2016;23(1):1–9. doi:10.5551/jat.32326
28. Jayachandran M, Miller VM, Heit JA, Owen WG. Methodology for isolation, identification and characterization of microvesicles in peripheral blood. *J Immunol Methods*. 2012;375(1–2):207–214.
29. Sabatier F, Camoin-Jau L, Anfosso F, Sampol J, Dignat-George F. Circulating endothelial cells, microparticles and progenitors: key players towards the definition of vascular competence. *J Cell Mol Med*. 2009;13(3):454–471.
30. Angelillo-Scherrer A. Leukocyte-derived microparticles in vascular homeostasis. *Circ Res*. 2012;110(2):356–369.
31. Tricarico C, Clancy J, D'Souza-Schorey C. Biology and biogenesis of shed microvesicles. *Small GTPases*. 2017;8(4):220–232.
32. Battistelli M, Falcieri E. Apoptotic bodies: particular extracellular vesicles involved in intercellular communication. *Biology (Basel)*. 2020;9(1):21.
33. Leventis PA, Grinstein S. The distribution and function of phosphatidylserine in cellular membranes. *Annu Rev Biophys*. 2010;39:407–427.
34. Kaplan ZS, Jackson SP. The role of platelets in atherothrombosis. *Hematology Am Soc Hematol Educ Program*. 2011;2011:51–61.
35. Akers JC, Gonda D, Kim R, Carter BS, Chen CC. Biogenesis of extracellular vesicles (EV): exosomes, microvesicles, retrovirus-like vesicles, and apoptotic bodies. *J Neurooncol*. 2013;113(1):1–11.
36. Black LV, Saunderson SC, Coutinho FP, et al. The CD169 sialoadhesin molecule mediates cytotoxic T-cell responses to tumour apoptotic vesicles. *Immunol Cell Biol*. 2015;94(5):430–438.
37. Thery C, Boussac M, Véron P, et al. Proteomic analysis of dendritic cell-derived exosomes: a secreted subcellular compartment distinct from apoptotic vesicles. *J Immunol*. 2001;166(12):7309–7318.
38. Bilyy RO, Shkandina T, Tomin A, et al. Macrophages discriminate glycosylation patterns of apoptotic cell-derived microparticles. *J Biol Chem*. 2012;287(1):496–503.
39. Hristov M, Erl W, Linder S, Weber PC. Apoptotic bodies from endothelial cells enhance the number and initiate the differentiation of human endothelial progenitor cells in vitro. *Blood*. 2004;104(9):2761–2766.
40. Turiak L, Misjak P, Szabo TG, et al. Proteomic characterization of thymocyte-derived microvesicles and apoptotic bodies in BALB/C mice. *J Proteome*. 2011;74(10):2025–2033.
41. Sinauridze EI, Kireev DA, Popenko NY, et al. Platelet microparticle membranes have 50- to 100-fold higher specific procoagulant activity than activated platelets. *Thromb Haemost*. 2007;97(3):425–434.
42. Wolberg AS, Monroe DM, Roberts HR, Hoffman MR. Tissue factor de-encryption: ionophore treatment induces changes in tissue factor activity by phosphatidylserine-dependent and -independent mechanisms. *Blood Coagul Fibrinolysis*. 1999;10(4):201–210.
43. Nomura S, Tandon NN, Nakamura T, Cone J, Kambayashi J. High-shear-stress-induced activation of platelets and microparticles enhances expression of cell adhesion molecules in THP-1 and endothelial cells. *Atherosclerosis*. 2001;158(2):277–287.
44. Barry OP, Praticò D, Savani RC, FitzGerald GA. Modulation of monocyte-endothelial cell interactions by platelet microparticles. *J Clin Invest*. 1998;102(1):136–144.
45. Mallat Z, Hugel B, Ohan J, Lesèche G, Freyssinet JM, Tedgui A. Shed membrane microparticles with procoagulant potential in human atherosclerotic plaques: a role for apoptosis in plaque thrombogenicity. *Circulation*. 1999;99(3):348–353.
46. Colombo M, Raposo G, Théry C. Biogenesis, secretion, and intercellular interactions of exosomes and other extracellular vesicles. *Annu Rev Cell Dev Biol*. 2014;30:255–289.
47. Yáñez-Mó M, Siljander PR, Andreu Z, et al. Biological properties of extracellular vesicles and their physiological functions. *J Extracell Vesicles*. 2015;4:27066.
48. Al-Nedawi K, Meehan B, Micallef J, et al. Intercellular transfer of the oncogenic receptor EGFRvIII by microvesicles derived from tumour cells. *Nat Cell Biol*. 2008;10(5):619–624.
49. Hood JL, San RS, Wickline SA. Exosomes released by melanoma cells prepare sentinel lymph nodes for tumor metastasis. *Cancer Res*. 2011;71(11):3792–3801.
50. Tomasoni S, Longaretti L, Rota C, et al. Transfer of growth factor receptor mRNA via exosomes unravels the regenerative effect of mesenchymal stem cells. *Stem Cells Dev*. 2013;22(5):772–780.
51. Borges FT, Reis LA, Schor N. Extracellular vesicles: structure, function, and potential clinical uses in renal diseases. *Braz J Med Biol Res*. 2013;46(10):824–830.
52. Chen TS, Lai RC, Lee MM, Choo AB, Lee CN, Lim SK. Mesenchymal stem cell secretes microparticles enriched in pre-microRNAs. *Nucleic Acids Res*. 2010;38(1):215–224.
53. Mathivanan S, Simpson RJ. ExoCarta: a compendium of exosomal proteins and RNA. *Proteomics*. 2009;9(21):4997–5000.
54. Hemler ME. Tetraspanin proteins mediate cellular penetration, invasion, and fusion events and define a novel type of membrane microdomain. *Annu Rev Cell Dev Biol*. 2003;19:397–422.
55. Tian T, Wang Y, Wang H, Zhu Z, Xiao Z. Visualizing of the cellular uptake and intracellular trafficking of exosomes by live-cell microscopy. *J Cell Biochem*. 2010;111(2):488–496.
56. Janas T, Janas MM, Sapon K, Janas T. Mechanisms of RNA loading into exosomes. *FEBS Lett*. 2015;589(13):1391–1398.
57. Kosaka N, Yoshioka Y, Hagiwara K, Tominaga N, Katsuda T, Ochiya T. Trash or treasure: extracellular microRNAs and cell-to-cell communication. *Front Genet*. 2013;4:173.
58. Schorey JS, Cheng Y, Singh PP, Smith VL. Exosomes and other extracellular vesicles in host-pathogen interactions. *EMBO Rep*. 2015;16(1):24–43.
59. Alenquer M, Amorim MJ. Exosome biogenesis, regulation, and function in viral infection. *Viruses*. 2015;7(9):5066–5083.
60. Miller HC, Stephan M. Hemorrhagic varicella: a case report and review of the complications of varicella in children. *Am J Emerg Med*. 1993;11(6):633–638.
61. Uthman IW, Gharavi AE. Viral infections and antiphospholipid antibodies. *Semin Arthritis Rheum*. 2002;31(4):256–263.
62. Geisbert TW, Jahrling PB. Exotic emerging viral diseases: progress and challenges. *Nat Med*. 2004;10(12 Suppl):S110–S121.
63. Squizzato A, Gerdes VE, Buller HR. Effects of human cytomegalovirus infection on the coagulation system. *Thromb Haemost*. 2005;93(3):403–410.
64. Wang CC, Chang CT, Lin CL, Lin IC, Kao CH. Hepatitis C virus infection associated with an increased risk of deep vein thrombosis: a population-based cohort study. *Medicine (Baltimore)*. 2015;94(38):e1585.
65. Bunce PE, High SM, Nadjafi M, Stanley K, Liles WC, Christian MD. Pandemic H1N1 influenza infection and vascular thrombosis. *Clin Infect Dis*. 2011;52(2):e14–e17.
66. Kiser KL, Badowski ME. Risk factors for venous thromboembolism in patients with human immunodeficiency virus infection. *Pharmacotherapy*. 2010;30(12):1292–1302.
67. Matta F, Yaekoub AY, Stein D. Human immunodeficiency virus infection and risk of venous thromboembolism. *Am J Med Sci*. 2008;336(5):402–406.

68. Kishimoto C, Ochiai H, Sasayama S. Intracardiac thrombus in murine Coxsackievirus B3 myocarditis. *Heart Vessels*. 1992;7(2):76–81.
69. Friedman MG, Phillip M, Dagan R. Virus-specific IgA in serum, saliva, and tears of children with measles. *Clin Exp Immunol*. 1989;75(1):58–63.
70. Hidaka Y, Sakai Y, Toh Y, Mori R. Glycoprotein C of herpes simplex virus type 1 is essential for the virus to evade antibody-independent complement-mediated virus inactivation and lysis of virus-infected cells. *J Gen Virol*. 1991;72(Pt 4):915–921.
71. Key NS, Bach RR, Vercellotti GM, Moldow CF. Herpes simplex virus type I does not require productive infection to induce tissue factor in human umbilical vein endothelial cells. *Lab Invest*. 1993;68(6):645–651.
72. Subeamaniam S, Scharrer I. Procoagulant activity during viral infections. *Front Biosci*. 2018;23:1060–1081.
73. Yu Y, Shen Y, Li J, et al. Viral infection related venous thromboembolism: potential mechanism and therapeutic targets. *Ann Palliat Med*. 2020;9(3):1257–1263.
74. van Gorp EC, Suharti C, Ten Cate H, et al. Review: infectious diseases and coagulation disorders. *J Infect Dis*. 1999;180(1):176–186.
75. Koupenova M, Vitseva O, Mackay CR, et al. Platelet-TLR7 mediates host survival and platelet count during viral infection in the absence of platelet-dependent thrombosis. *Blood*. 2014;124(5):791–802.
76. Campbell RA, Schwertz H, Hottz ED, et al. Human megakaryocytes possess intrinsic antiviral immunity through regulated induction of IFITM3. *Blood*. 2019;133(19):2013–2026.
77. Koupenova M, Corkrey HA, Vitseva O, et al. The role of platelet in mediating a response to human influenza infection. *Nat Commun*. 2019;10(1):1780.
78. Rondina M, Tatsumi K, Bastarache JA, Mackman N. Microvesicle tissue factor activity and interleukin-8 levels are associated with mortality in patients with influenza A/H1N1 infection. *Crit Care Med*. 2016;44(7):e574–e578.
79. van der Poll T, Levi M. Crosstalk between inflammation and coagulation: the lessons of sepsis. *Curr Vasc Pharmacol*. 2012;10(5):632–638.
80. Leroy EM, Baize S, Volchkov VE, et al. Human asymptomatic Ebola infection and strong inflammatory response. *Lancet*. 2000;355(9222):2210–2215.
81. Mohamadzadeh M, Chen L, Schmaljohn AL. How Ebola and Marburg viruses battle the immune system. *Nat Rev Immunol*. 2007;7(7):556–567.
82. Wong G, Kobinger GP, Qiu X. Characterization of host immune responses in Ebola virus infections. *Expert Rev Clin Immunol*. 2014;10(6):781–790.
83. Mahanty S, Bray M. Pathogenesis of filoviral haemorrhagic fevers. *Lancet Infect Dis*. 2004;4(8):487–498.
84. Antoniak S. The coagulation system in host defense. *Res Pract Thromb Haemost*. 2018;2(3):549–557.
85. Short KR, Veldhuis Kroeze EJB, Reperant LA, Richard M, Kuiken T. Influenza virus and endothelial cells: a species specific relationship. *Front Microbiol*. 2014;5:653.
86. Barnes M, Heywood AE, Mahimbo A, Rahman B, Newall AT, Macintyre CR. Acute myocardial infarction and influenza: a meta-analysis of case-control studies. *Heart*. 2015;101(21):1738–1747.
87. Corrales-Medina VF, Madjid M, Musher DM. Role of acute infection in triggering acute coronary syndromes. *Lancet Infect Dis*. 2010;10(2):83–92.
88. Ludwig A, Lucero-Obusan C, Schirmer P, Winston C, Holodniy M. Acute cardiac injury events  $\leq 30$  days after laboratory-confirmed influenza virus infection among U.S. veterans, 2010–2012. *BMC Cardiovasc Disord*. 2015;15:109.
89. Marsden PA. Inflammation and coagulation in the cardiovascular system: the contribution of influenza. *Circ Res*. 2006;99(11):1152–1153.
90. Rothberg MB, Haessler SD, Brown RB. Complications of viral influenza. *Am J Med*. 2008;121(4):258–264.
91. Armstrong SM, Darwish I, Lee WL. Endothelial activation and dysfunction in the pathogenesis of influenza A virus infection. *Virulence*. 2013;4(6):537–542.
92. Feng Y, Hu L, Lu S, et al. Molecular pathology analyses of two fatal human infections of avian influenza A(H7N9) virus. *J Clin Pathol*. 2015;68(1):57–63.
93. Muramoto Y, Ozaki H, Takada A, et al. Highly pathogenic H5N1 influenza virus causes coagulopathy in chickens. *Microbiol Immunol*. 2006;50(1):73–81.
94. Akiyama R, Komori I, Hiramoto R, Isonishi A, Matsumoto M, Fujimura Y. H1N1 influenza (Swine flu)-associated thrombotic microangiopathy with a markedly high plasma ratio of von Willebrand factor to ADAMTS13. *Intern Med*. 2011;50(6):643–647.
95. Tsujii N, Nogami K, Yoshizawa H, et al. Influenza-associated thrombotic microangiopathy with unbalanced von Willebrand factor and a disintegrin and metalloproteinase with a thrombospondin type 1 motif, member 13 levels in a heterozygous protein S-deficient boy. *Pediatr Int*. 2016;58(9):926–929.
96. Blum A, Hadas V, Burke M, Yust I, Kessler A. Viral load of the human immunodeficiency virus could be an independent risk factor for endothelial dysfunction. *Clin Cardiol*. 2005;28(3):149–153.
97. Baker JV, Duprez D. Biomarkers and HIV-associated cardiovascular disease. *Curr Opin HIV AIDS*. 2010;5(6):511–516.
98. Palella FJ Jr., Phair JP. Cardiovascular disease in HIV infection. *Curr Opin HIV AIDS*. 2011;6(4):266–271.
99. Shen YM, Frenkel EP. Thrombosis and a hypercoagulable state in HIV-infected patients. *Clin Appl Thromb Hemost*. 2004;10(3):277–280.
100. O'Brien MP, Zafar MU, Rodriguez JC, et al. Targeting thrombogenicity and inflammation in chronic HIV infection. *Sci Adv*. 2019;5(6):eaav5463. doi:10.1126/sciadv.aav5463
101. Kuller LH, Tracy R, Bellosso W, et al. Inflammatory and coagulation biomarkers and mortality in patients with HIV infection. *PLoS Med*. 2008;5(10):e203. doi:10.1371/journal.pmed.0050203
102. Rodger AJ, Fox Z, Lundgren JD, et al. Activation and coagulation biomarkers are independent predictors of the development of opportunistic disease in patients with HIV infection. *J Infect Dis*. 2009;200(6):973–983. doi:10.1086/605447
103. Funderburg NT, Mayne E, Sieg SF, et al. Increased tissue factor expression on circulating monocytes in chronic HIV infection: relationship to *in vivo* coagulation and immune activation. *Blood*. 2010;115(2):161–167. doi:10.1182/blood-2009-03-210179
104. Funderburg NT, Zidar DA, Shive C, et al. Shared monocyte subset phenotypes in HIV-1 infection and in uninfected subjects with acute coronary syndrome. *Blood*. 2012;120(23):4599–4608. doi:10.1182/blood-2012-05-433946
105. Hurley A, Smith M, Karpova T, et al. Enhanced effector function of CD8(+) T cells from healthy controls and HIV-infected patients occurs through thrombin activation of protease-activated receptor 1. *J Infect Dis*. 2013;207(4):638–650. doi:10.1093/infdis/jis730
106. Allie S, Stanley A, Bryer A, Meiring M, Combrinck MI. High levels of von Willebrand factor and low levels of its cleaving protease, ADAMTS13, are associated with stroke in young HIV-infected patients. *Int J Stroke*. 2015;10(8):1294–1296. doi:10.1111/ijss.12550
107. Ellwanger JH, Veit TD, Chies JAB. Exosomes in HIV infection: a reviewer and critical look. *Infect Genet Evol*. 2017;53:146–154. doi:10.1016/j.meegid.2017.05.021



108. Welch JL, Stapleton JT, Okeoma CM. Vehicles of intercellular communication: exosomes and HIV-1. *J Gen Virol.* 2019;100(3):350–366. doi:10.1099/jgv.0.001193
109. Gonzalez-Reimers E, Quintero-Platt G, Martin-Gonzalez C, Perez-Hernandez O, Romero-Acevedo L, Santolaria-Fernandez F. Thrombin activation and liver inflammation in advanced hepatitis C virus infection. *World J Gastroenterol.* 2016;22(18):4427–4437. doi:10.3748/wjg.v22.i18.4427
110. Pircher J, Czermak T, Merkle M, et al. Hepatitis C virus induced endothelial inflammatory response depends on the functional expression of TNFalpha receptor subtype 2. *PLoS One.* 2014;9(11):e113351. doi:10.1371/journal.pone.0113351
111. Wald O, Pappo O, Safadi R, et al. Involvement of the CXCL12/CXCR4 pathway in the advanced liver disease that is associated with hepatitis C virus or hepatitis B virus. *Eur J Immunol.* 2004;34(4):1164–1174. doi:10.1002/eji.200324441
112. Falasca K, Mancino P, Ucciferri C, et al. Inflammatory cytokines and S-100b protein in patients with hepatitis C infection and cryoglobulinemias. *Clin Invest Med.* 2007;30(5):E167–E176. doi:10.25011/cim.v30i5.2892
113. Hodowanec AC, Lee RD, Brady KE, et al. A matched cross-sectional study of the association between circulating tissue factor activity, immune activation and advanced liver fibrosis in hepatitis C infection. *BMC Infect Dis.* 2015;15:190. doi:10.1186/s12879-015-0920-1
114. Barone M, Viggiani MT, Amoruso A, et al. Endothelial dysfunction correlates with liver fibrosis in chronic HCV infection. *Gastroenterol Res Pract.* 2015;2015:682174. doi:10.1155/2015/682174
115. Lee CJ, Huang YC, Yang S, et al. Clinical features of coxsackievirus A4, B3 and B4 infections in children. *PLoS One.* 2014;9(2):e87391. doi:10.1371/journal.pone.0087391
116. Antoniak S, Mackman N. Coagulation, protease-activated receptors, and viral myocarditis. *J Cardiovasc Transl Res.* 2014;7(2):203–211. doi:10.1007/s12265-013-9515-7
117. Sutherland MR, Raynor CM, Leenknecht H, Wright JF, Prydzial EL. Coagulation initiated on herpesviruses. *Proc Natl Acad Sci U S A.* 1997;94(25):13510–13514. doi:10.1073/pnas.94.25.13510
118. Sutherland MR, Friedman HM, Prydzial EL. Thrombin enhances herpes simplex virus infection of cells involving protease-activated receptor 1. *J Thromb Haemost.* 2007;5(5):1055–1061. doi:10.1111/j.1538-7836.2007.02441.x
119. Sutherland MR, Ruf W, Prydzial EL. Tissue factor and glycoprotein C on herpes simplex virus type 1 are protease-activated receptor 2 cofactors that enhance infection. *Blood.* 2012;119(15):3638–3645. doi:10.1182/blood-2011-08-376814
120. Toussiot E, Roudier J. Epstein-Barr virus in autoimmune diseases. *Best Pract Res Clin Rheumatol.* 2008;22(5):883–896. doi:10.1016/j.berh.2008.09.007
121. Hyakutake M, Steinberg E, Disla E, Heller M. Concomitant infection with Epstein-Barr virus and cytomegalovirus infection leading to portal vein thrombosis. *J Emerg Med.* 2019;57(2):e49–e51. doi:10.1016/j.jemermed.2019.02.010
122. Justo D, Finn T, Atzmony L, Guy N, Steinvil A. Thrombosis associated with acute cytomegalovirus infection: a meta-analysis. *Eur J Intern Med.* 2011;22(2):195–199. doi:10.1016/j.ejim.2010.11.006
123. Simmonds P, Aiewsakun P. Virus classification – where do you draw the line? *Arch Virol.* 2018;163(8):2037–2046. doi:10.1007/s00705-018-3938-z
124. Wong RSM, Wu A, To KF, et al. Haematological manifestations in patients with severe acute respiratory syndrome: retrospective analysis. *BMJ.* 2003;326(7403):1358–1362. doi:10.1136/bmj.326.7403.1358
125. Chong PY, Chui P, Ling AE, et al. Analysis of deaths during the severe acute respiratory syndrome (SARS) epidemic in Singapore: challenges in determining a SARS diagnosis. *Arch Pathol Lab Med.* 2004;128(2):195–204. doi:10.1043/1543-2165-(2004)128<195:AODDTS>2.0.CO;2
126. Ng KHL, Wu AKL, Cheng VCC, et al. Pulmonary artery thrombosis in a patient with severe acute respiratory syndrome. *Postgrad Med J.* 2005;81(956):e3. doi:10.1136/pgmj.2004.030049
127. Yang M, Ng MHL, Li CK, et al. Thrombopoietin levels increased in patients with severe acute respiratory syndrome. *Thromb Res.* 2008;122(4):473–477. doi:10.1016/j.thromres.2007.12.021
128. Shimizu M, Konishi A, Nomura S. Examination of biomarker expressions in sepsis-related DIC patients. *Int J Gen Med.* 2018;11:353–361. doi:10.2147/IJGM.S173684
129. Assiri A, Al-Tawfiq JA, Al-Rabeeh AA, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis.* 2013;13(9):752–761. doi:10.1016/S1473-3099(13)70204-4
130. Singh SK. Middle east respiratory syndrome virus pathogenesis. *Semin Respir Crit Care Med.* 2016;37(4):572–577. doi:10.1055/s-0036-1584796
131. Han H, Yang L, Liu R, et al. Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med.* 2020.
132. Giannis D, Ziogas IA, Gianni P. Coagulation disorders in coronavirus infected patients: COVID-19, SARS-Cov-1, MERS-CoV and lessons from the past. *J Clin Virol.* 2020;127:104362. doi:10.1016/j.jcv.2020.104362
133. Lodigiani C, Iapichino G, Carenzo L, et al. Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res.* 2020;191:9–14. doi:10.1016/j.thromres.2020.04.024
134. Lacour T, Semaan C, Genet T, Ivanov F. Insights for increased risk of failed fibrinolytic therapy and stent thrombosis associated with COVID-19 in ST-segment elevation myocardial infarction patients. *Catheter Cardiovasc Interv.* 2020.
135. Bikkdeli B, Madhavan MV, Jimenez D, et al. COVID-19 and thrombotic or thromboembolic disease; implications for prevention, antithrombotic therapy, and follow-up. *J Am Coll Cardiol.* 2020;75:2950–2973.
136. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost.* 2020;18(4):844–847. doi:10.1111/jth.14768
137. Griffin DO, Jensen A, Khan M, et al. Pulmonary embolism and increased levels of d-dimer in patients with coronavirus disease. *Emerg Infect Dis.* 2020;26(8).
138. Terpos E, Ntanasis-Stathopoulos I, Elalamy I, et al. Hematological findings and complications of COVID-19. *Am J Hematol.* 2020.
139. Schwab A, Meyering SS, Lepence B, et al. Extracellular vesicles from infected cells: potential for direct pathogenesis. *Front Microbiol.* 2015;6:1132. doi:10.3389/fmicb.2015.01132
140. Rodrigues M, Fan J, Lyon C, Wan M, Hu Y. Role of extracellular vesicles in viral and bacterial infections: pathogenesis, diagnosis, and therapeutics. *Theranostics.* 2018;8(10):2709–2721. doi:10.7150/thno.20576
141. Zhang W, Jiang X, Bao J, Wang Y, Liu H, Tang L. Exosomes in pathogen infections: a bridge to deliver molecules and link functions. *Front Immunol.* 2018;9:90. doi:10.3389/fimmu.2018.00090
142. Brown M, Johnson LA, Leone DA, et al. Lymphatic exosomes promote dendritic cell migration along guidance cues. *J Cell Biol.* 2018;217(6):2205–2221. doi:10.1083/jcb.201612051

143. Anderson MR, Kashanchi F, Jacobson S. Exosomes in viral disease. *Neurotherapeutics*. 2016;13(3):535–546. doi:10.1007/s13311-016-0450-6
144. Pleet ML, DeMarino C, Stonier SW, et al. Extracellular vesicles and Ebola virus: a new mechanism of immune evasion. *Viruses*. 2019;11(5):410. doi:10.3390/v11050410
145. Kesimer M, Scull M, Brighton B, et al. Characterization of exosome-like vesicles released from human tracheobronchial ciliated epithelium: a possible role in innate defense. *FASEB J*. 2009;23(6):1858–1868. doi:10.1096/fj.08-119131
146. Suptawiwat O, Ruangrung K, Boonarkart C, et al. Microparticle and anti-influenza activity in human respiratory secretion. *PLoS One*. 2017;12(8):e0183717. doi:10.1371/journal.pone.0183717
147. Konadu KA, Huang MB, Roth W, et al. Isolation of exosomes from the plasma of HIV-1 positive individuals. *J Vis Exp*. 2016;107:e53495.
148. Chettimada S, Lorenz DR, Misra V, et al. Exosome markers associated with immune activation and oxidative stress in HIV patients on antiretroviral therapy. *Sci Rep*. 2018;8(1):7227. doi:10.1038/s41598-018-25515-4
149. Pérez PS, Romaniuk MA, Duette GA, et al. Extracellular vesicles and chronic inflammation during HIV infection. *J Extracell Vesicles*. 2019;8(1):1687275. doi:10.1080/20013078.2019.1687275
150. Plazolles N, Humbert JM, Vachot L, Verrier B, Hocke C, Halary F. Pivotal advance: the promotion of soluble DC-SIGN release by inflammatory signals and its enhancement of cytomegalovirus-mediated Cis- infection of myeloid dendritic cells. *J Leukoc Biol*. 2011;89(3):329–342. doi:10.1189/jlb.0710386
151. Walker JD, Maier CL, Pober JS. Cytomegalovirus-infected human endothelial cells can stimulate allogeneic CD4+ memory T cells by releasing antigenic exosomes. *J Immunol*. 2009;182(3):1548–1559. doi:10.4049/jimmunol.182.3.1548
152. Kuo HT, Ye X, Sampaio MS, Reddy P, Bunnapradist S. Cytomegalovirus serostatus pairing and deceased donor kidney transplant outcomes in adult recipients with antiviral prophylaxis. *Transplantation*. 2010;90(10):1091–1098.
153. Meckes DG, Raab-Traub N. Microvesicles and viral infection. *J Virol*. 2011;85(24):12844–12854.
154. Wurdinger T, Gatsen NN, Balaj L, Kaur B, Breakefield XO, Pegtel DM. Extracellular vesicles and their convergence with viral pathway. *Adv Virol*. 2012;2012:767694.
155. Delabranche X, Berger A, Boisramé-Helms J, Meziani F. Microparticles and infectious disease. *Med Mal Infect*. 2012;42(8):335–343.
156. Coakley C, Maizels RM, Buck AH. Exosome and other extracellular vesicles: the new communicators in parasite infections. *Trends Parasitol*. 2015;31(10):477–489.
157. Schorey JS, Harding CV. Extracellular vesicles and infectious diseases: new complexity to an old story. *J Clin Invest*. 2016;126(4):1181–1189.
158. Raab-Traub N, Dittmer DP. Viral effects on the content and function of extracellular vesicles. *Nat Rev Microbiol*. 2017;15(9):559–572.
159. Altan-Bonnet N, Perales C, Domingo E. Extracellular vesicles: vesicles of en bloc viral transmission. *Virus Res*. 2019;265:143–149.
160. Urbanelli L, Buratta S, Tancini B, et al. The role of extracellular vesicles in viral infection and transmission. *Vaccines (Basel)*. 2019;7(3):102.
161. Goncalves-Alves E, Saferding V, Schliehe C, et al. MicroRNA-155 controls T helper cell activation during viral infection. *Front Immunol*. 2019;10:1367.
162. AmraneDjedidi R, Rousseau A, Larsen AK, et al. Extracellular vesicles derived from pancreatic cancer cells BXP3 or breast cancer cells MCF7 induce a permanent procoagulant shift to endothelial cells. *Thromb Res*. 2020;187:170–179.
163. Deffune E, Prudenciatti A, Moroz A. Mesenchymal stem cell (MSc) secretome: a possible therapeutic strategy for intensive-care COVID-19 patients. *Med Hypotheses*. 2020;142:109769.
164. O'Driscoll L. Extracellular vesicles from mesenchymal stem cells as a COVID-19 treatment. *Drug Discov Today*. 2020.
165. Bari E, Ferrarotti I, Saracino L, Perteghella S, Torre ML, Corsico AG. Mesenchymal stromal cell secretome or severe COVID-19 infections: premises for the therapeutic use. *Cells*. 2020;9(4):924.
166. Kumar S, Zhi K, Mukherji A, Gerth K. Repurposing antiviral protease inhibitors using extracellular vesicles for potential therapy of COVID-19. *Viruses*. 2020;12(5):E486.
167. Schmedes CM, Grover SP, Hisada YM, et al. Increased circulating extracellular vesicle tissue factor activity during orthohanta-virus infection is associated with intravascular coagulation. *J Infect Dis*. 2019.
168. Felsenstein S, Herbert JA, McNamara PS, Hedrich CM. COVID-19: immunology and treatment options. *Clin Immunol*. 2020;108448.
169. Myer AD, Rishmawi AR, Kamucheka R, et al. Effect of blood flow on platelets, leukocytes, and extracellular vesicles in thrombosis of simulated neonatal extracorporeal circulation. *J Thromb Haemost*. 2020;18(2):399–410.
170. Saenz-Pipaon G, San Martin P, Planell N, et al. Functional and transcriptomic analysis of extracellular vesicles identifies calprotectin as a new prognostic marker in peripheral arterial disease (PAD). *J Extracell Vesicles*. 2020;9:1729646.
171. Zou B, She J, Wang Y, Ma X. Venous thrombosis and arteriosclerosis obliterans of lower extremities in a very severe patient with 2019 novel coronavirus disease: a case report. *J Thromb Thrombolysis*. 2020.

## International Journal of General Medicine

### Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies

across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>