



HHS Public Access

Author manuscript

Diabetologia. Author manuscript; available in PMC 2020 October 01.

Published in final edited form as:

Diabetologia. 2019 October ; 62(10): 1811–1822. doi:10.1007/s00125-019-4940-z.

Environmental neglect: endocrine disruptors as underappreciated but potentially modifiable diabetes risk factors

Robert M. Sargis^{1,2}, Rebecca A. Simmons^{3,4,5}

¹Division of Endocrinology, Diabetes, and Metabolism Department of Medicine, University of Illinois at Chicago, 835 S. Wolcott, Suite E625; M/C 640, Chicago, IL 60612, USA

²ChicAGO Center for Health and Environment (CACHET), University of Illinois at Chicago, Chicago, IL, USA

³Division of Neonatology, Children's Hospital of Philadelphia, Philadelphia, PA, USA

⁴Center for Research on Reproduction and Women's Health, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

⁵Center of Excellence in Environmental Toxicology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

Abstract

Type 2 diabetes prevalence is increasing dramatically across the globe, imposing a tremendous toll on individuals and healthcare systems. Reversing these trends requires comprehensive approaches to address both classical and emerging diabetes risk factors. Recently, environmental toxicants acting as endocrine-disrupting chemicals (EDCs) have emerged as novel metabolic disease risk factors. EDCs implicated in diabetes pathogenesis include various inorganic and organic molecules of both natural and synthetic origin, including arsenic, bisphenol A, phthalates, polychlorinated biphenyls and organochlorine pesticides. Indeed, evidence implicates EDC exposures across the lifespan in metabolic dysfunction; moreover, specific developmental windows exhibit enhanced sensitivity to EDC-induced metabolic disruption, with potential impacts across generations. Importantly, differential exposures to diabetogenic EDCs are likely to also contribute to racial/ethnic and economic disparities. Despite these emerging links, clinical practice guidelines fail to address this underappreciated diabetes risk factor. Comprehensive approaches to stem the tide of diabetes must include efforts to address its environmental drivers.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <http://www.springer.com/gb/open-access/authors-rights/aam-terms-v1>

Corresponding Author: Robert M. Sargis, Division of Endocrinology, Diabetes, and Metabolism Department of Medicine, University of Illinois at Chicago, 835 S. Wolcott, Suite E625; M/C 640, Chicago, IL 60612, USA rsargis@uic.edu.

Contribution statement

Both authors were responsible for drafting the article and revising it critically for important intellectual content. Both authors approved the version to be published.

Duality of interest

RMS declares he has received honoraria from CVS/Health. RAS has no duality of interest.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of a an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Keywords

Beta cell; Bisphenol A; Diabetes; Endocrine-disrupting chemical; Endocrine disruptor; Environmental justice; Glucose; Insulin; Life course development; Review

Introduction

The global burden of diabetes is projected to reach a staggering 629 million individuals by 2045 [1], and the WHO envisages metabolic disorders will be a major cause of death by 2030, placing substantial economic stresses on healthcare systems worldwide [2]. Indeed, 12% of global health expenditures are already estimated to be spent on diabetes [1] and, in the USA alone, annual diabetes-associated economic costs are calculated to be \$327 billion [3]. Most of the diabetes health burden arises from type 2 diabetes, which accounts for 90–95% of all diabetes cases. As both disease prevalence and treatment costs continue to rise dramatically [4], comprehensive approaches to reverse these trends are desperately needed. Essential to this effort are strategies that simultaneously address the myriad risk factors contributing to diabetes pathogenesis. While this certainly includes efforts to tackle the contributions of excess food intake and sedentary lifestyles, it is also essential to identify other modifiable risk factors that promote diabetes development.

One emerging but underappreciated realm of risk is our ubiquitous exposure to environmental chemicals through food and water, skin contact and the air we breathe. Indeed, exposures to endocrine-disrupting chemicals (EDCs) are associated with an increased risk of various endocrine-related disorders in both human and animal studies [5]. The Endocrine Society classifies exogenous chemicals as EDCs based upon their ability to disrupt any aspect of hormone secretion or action [6]. Over the last two decades, exposure to many EDCs from multiple sources has been associated with increased risk of diabetes, obesity and other metabolic disorders (Fig. 1). These metabolism-disrupting EDCs include a variety of inorganic and organic molecules of natural or synthetic origin, including many to which humans are frequently exposed, including arsenic, bisphenol A (BPA) and phthalates, as well as legacy compounds, which have profound chemical and biological stability and persist in the environment despite regulatory action to eliminate or curtail their use, such as polychlorinated biphenyls (PCBs) and organochlorine pesticides. With increasing evidence linking exposure to various EDCs with diabetes in human populations, as well as diabetes-associated pathophysiological defects in cell-based and animal studies [7–10] (Fig. 2), the time has come to incorporate environmental interventions into comprehensive strategies to reduce the individual and societal burden of this devastating disease.

Exposures during sensitive developmental windows

While EDC exposures occur across the lifespan, specific developmental windows are recognised as periods of unique vulnerability (Fig. 3). The period from conception to birth is a time of cellular replication and differentiation, functional organ system maturation and rapid growth. These processes are exquisitely sensitive to perturbations in the intrauterine metabolic milieu that exert long-lasting effects on offspring. Animal and epidemiology data demonstrate that several critical periods during development influence the later-life onset of

type 2 diabetes, including preconception and gestation, early infancy, the adiposity rebound period between 5–7 years of age, and puberty. Until recently, most studies examining developmental programming were focused on inadequate nutrition or uteroplacental insufficiency; however, it is now apparent that developmental exposure to EDCs may also increase the incidence of type 2 diabetes [7, 11].

Perhaps the prototypical EDC is BPA, a common contaminant used widely in polycarbonate and other plastics, epoxy resins lining cans and the thermal paper of cash register receipts. BPA binds both oestrogen receptor α (ER α) and β (ER β), although with less affinity than 17 β -oestradiol (E2). Humans are ubiquitously exposed to BPA, with low levels detectable in the blood and urine of virtually all individuals across multiple populations [12–14]. Importantly, human fetuses and children are likely to be at heightened risk of endocrine disruption from BPA, since BPA clearance enzymes are not fully functional at these life stages [15]. Consequently, blood and urine levels of BPA are higher in fetuses than in their pregnant mothers [13], and in infants and children compared with adults [12]. Importantly, while exact human exposures are difficult to estimate, animal studies demonstrate that levels below those considered safe for human health are sufficient to induce metabolic dysfunction [16].

Multiple animal studies have explored the impact of developmental BPA exposure on metabolic programming; however, controversy persists regarding the effects of maternal BPA exposure on offspring. Differences across studies, include species and strain, developmental stage during exposure, ages at metabolic interrogation, levels of exposure, housing, diet, control conditions and exposure routes have all contributed to a lack of consensus on the gestational effects of BPA. Particularly robust debate has focused upon the importance of the route of exposure (oral vs subcutaneous) as well as evidence of non-monotonic dose–response relationships demonstrating that BPA exposures at or below the current US Environmental Protection Agency’s reference dose often yield metabolic disruptions that are either not observed or different at higher doses [17–19]. Acknowledging these challenges, when experimental conditions are strictly maintained, reproducible and stable metabolic disruptions are consistently observed in the offspring of BPA-exposed dams [20].

Some of the first evidence that developmental EDC exposure alters metabolic health came from studies exposing pregnant rats to BPA. Exposure resulted in dose- and sex-specific increases in offspring body weights that persisted into adulthood [21]. Since then, multiple studies have examined links between perinatal EDC exposure and the later-life development of metabolic dysfunction, including obesity. Fewer studies have examined the effects of EDC on beta cell development and function. Importantly, all three oestrogen receptors (ER α , ER β and the G-protein coupled ER [GPER]) are present in rodent and human beta cells, where they play essential roles in islet survival and function [22]; thus, beta cells are primed for disruption by ER-active EDCs like BPA. Indeed, fetal offspring of C57BL/6 dams fed a BPA-containing diet from embryonic day 7.5 (E7.5) to E18.5 exhibited altered endocrine pancreas development, with disruptions in the number and relative cellular composition of islets [23]. Using similar exposures, male offspring of pregnant mice exposed subcutaneously to BPA exhibited enhanced insulin secretion at 17 weeks of age,

suggesting BPA-induced insulin resistance; moreover, by 28 weeks, the mice exhibited hyperglycaemia without alterations in glucose-stimulated insulin secretion (GSIS) [24]. Of note, follow-up studies using a similar exposure window and a higher BPA dose demonstrated that the lower BPA dose, but not the higher dose, decreased GSIS at postnatal day 30, despite increased beta cell mass, which was attributed to increased expression of pro-proliferative genes and decreased expression of genes regulating beta cell death [25]. Intriguingly, beta cell mass was actually decreased in BPA-treated animals by postnatal day 120. The discrepancies in the two studies are likely to be related to the different postnatal ages examined; however, it remains unclear why GSIS was impaired in young animals but was normal at later ages.

To better mimic exposures during human development we examined the impact of dietary exposure of mice to environmentally relevant BPA doses from 2 weeks prior to pregnancy to the end of lactation using a low phytoestrogen control diet and BPA-free cages; serum BPA levels approximated those in humans [17]. Male offspring of exposed dams, but not females, developed abnormal glucose tolerance associated with insulin secretory defects, including increased basal rates of insulin release at the higher BPA dose that was suggestive of insulin resistance. Interestingly, differences in insulin release using a glucose gradient were not observed between control males and those that received a higher dose of BPA, nor was KCl-induced insulin release in islets altered by the higher BPA dose, suggesting an intact insulin secretory apparatus. In contrast, male offspring exposed to the lower BPA dose exhibited similar basal rates of insulin release; however, maximal GSIS was impaired [17]. These data suggest that the metabolic consequences of developmental BPA exposure exhibit complex dose–response relationships.

An area of intense interest in the developmental programming of metabolism is the multigenerational heritability of metabolic traits after ancestral EDC exposure. To determine whether BPA-induced metabolic disruptions persisted into F2 offspring, BPA-exposed F1 females were mated to unexposed males [17]. Like F1 offspring, F2 males of dams exposed to a higher dose of BPA were glucose intolerant with preserved GSIS, while F2 males of dams exposed to a lower dose of BPA exhibited blunted GSIS. In addition, lower dose BPA exposure significantly reduced beta cell mass and increased beta cell death in F1 males, which persisted into the F2 generation, and transcriptomic analyses indicated significant dose-specific changes in genes regulating inflammation and mitochondrial function [18]. Collectively, these results demonstrate multigenerational dose- and sex-specific metabolic effects of developmental BPA exposure.

These data fit within a broadening context of multigenerational and transgenerational effects of multiple EDC classes on various endocrine and metabolic tissues [7, 26–29]. Recently, studies have demonstrated that changes in the expression and methylation of imprinted genes in the brain persist across three generations after BPA exposure in mice [30]. In examining the transgenerational (F3) effects of BPA on metabolic health, we found that F3 adult males exposed ancestrally to either the lower or higher BPA dose had increased body weight with preserved glucose tolerance [19]. F3 male offspring of dams exposed to the lower BPA dose had reduced beta cell mass and smaller islets that exhibited enhanced GSIS. There was no effect of BPA in F3 females. These studies show that maternal BPA exposure

resulted in fewer metabolic defects in F3 than F1 and F2 offspring, and these were sex- and dose-specific. Interestingly, while these data suggest a decay in EDC effects over generations, the relationships may be more complex for other exposures. For example, the fungicide and booster biocide tributyltin (TBT) has been shown to exert transgenerational effects on adipose tissue, with some phenotypes that amplify over successive generations [31].

Mechanisms of EDC action

There are a number of possible mechanisms by which EDCs can influence health outcomes. These include interference with the activity of nuclear receptors, noncanonical steroid hormone receptors and orphan receptors, as well as disruptions in the enzymatic pathways regulating steroid biosynthesis and/or metabolism. For example, BPA can bind multiple oestrogen receptor subtypes. Moreover, several studies indicate that BPA also promotes signalling via extranuclear signal transduction pathways, including extracellular signal-regulated kinase (ERK)/mitogen-activated protein kinase (MAPK) and Akt [32], and BPA also exerts oestradiol-like effects by binding to ER α and thereby activating extranuclear pathways [33]. Thus, it is increasingly clear that the diverse effects of BPA are likely to arise from interactions across multiple signal transduction pathways [34]. Adding complexity is the fact that epidemiological data have linked BPA to disruptions in androgen-dependent developmental outcomes [35], despite few mechanistic data demonstrating that BPA alters androgen receptor activity or interferes with androgen-dependent extranuclear receptors. Finally, there is some evidence that BPA may also disrupt glucocorticoid receptor signalling [36–38]. Because endocrine signalling adapts to tonic activation through feedback circuits and because metabolic phenotypes arise from the integrated crosstalk of multiple signalling cascades that act additively, antagonistically or synergistically, the diversity of BPA actions illustrates how this and other EDCs may exhibit non-monotonic dose–response relationships in which phenotypic changes at low doses are not necessarily observed with higher exposures.

While significant literature linking developmental EDC exposures to metabolic disorders focuses on oestrogen receptor actions, robust data also indicate that perinatal exposures to some EDCs alter the expression or activity of the nuclear receptor peroxisome proliferator-activated receptor γ (PPAR γ) and its heterodimeric partner retinoid X receptor (RXR), which play essential roles in adipocyte development and function. Interestingly, BPA has been shown to increase PPAR γ expression [39]. While no studies have shown that BPA directly binds to PPAR γ , brominated BPA analogues, also found in high quantities in the environment, bind to the receptor at low concentrations [40]. The prototypical obesogen TBT binds and activates both PPAR γ and RXR at very low concentrations, augmenting adipocyte differentiation, promoting adipocyte dysfunction and resulting in transgenerational disruptions in metabolic homeostasis [26, 31, 41–43].

Our recent studies suggest that mitochondrial dysfunction could be an early event in the BPA-induced impairment of beta cells [18, 19]. Low-dose exposure of pregnant dams impairs beta cell mitochondria function, blunts GSIS, and reduces beta cell mass in male offspring [18]. These effects persisted into the second, but not the third, generation [19]. In

another study, perinatal BPA exposure decreased activity of Complexes I and III of the electron transport chain in the livers of suckling rats, with later-life development of microvesicular liver steatosis and enhanced lipogenic gene expression [44]. Beyond BPA, other common EDCs have also been shown to disrupt mitochondrial function after developmental exposure, including arsenic [45]. Collectively, these data underscore the complexity of endocrine disruption with regard to metabolic outcomes. Unlike pharmaceutical agents that are designed to hit specific molecular targets, EDCs often interfere with multiple pathways, with specific effects likely dependent on both the dose and timing of exposure as well as the target and the background hormonal milieu. However, to date, while multiple studies have demonstrated that EDCs exhibit multiple modes of action, very few studies have definitively ascribed specific mechanisms to particular phenotypes. This greatly complicates our understanding of the molecular mechanisms of these agents and argues for comprehensive unbiased approaches for uncovering the network of metabolic toxicities induced by these agents.

Epigenetic mechanisms

With studies demonstrating transgenerational effects of EDCs on metabolic health, attention has turned to how EDCs directly influence the epigenome. Multiple studies report associations between developmental exposures to EDCs and epigenetic changes in key metabolic tissues, including beta cells, liver, adipose and muscle [7]. One of the earliest studies examining this phenomenon demonstrated that developmental BPA exposure shifts the coat colour of Agouti mice through stable epigenetic changes [46]. In general, imprinted genes are particularly vulnerable to de novo epigenetic modifications. Dams exposed to BPA from 2 weeks prior to mating to the end of gestation yielded offspring with loss of imprinting and biparental expression of *Igf2*, increased embryonic *Igf2* gene expression and enhanced methylation of the *Igf2* differentially methylated region 1 [17]. Extension of these studies demonstrated that *Igf2* differentially methylated region 1 (DMR1) hypermethylation persisted in islets of F1 and F2 offspring of BPA-exposed dams with an associated increased expression of *Igf2* [18]. While *Igf2* has multiple functions, it is a key regulator of beta cell development, and aberrant early-life *Igf2* imprinting could potentially disrupt beta cell development.

Epigenetic alterations have also been described for several other metabolically active EDCs [7]. Phthalates are a family of phthalic acid diesters that are commonly found in consumer products, resulting in ubiquitous exposure in the USA. Limited but growing evidence in human studies indicate that phthalate exposure is associated with DNA methylation changes of imprinted genes in cord blood and placenta [47]. Animal studies demonstrate associations between phthalates and global and site-specific methylation [48]; however, studies were performed in mixed cell populations, and changes in DNA methylation were usually modest.

In addition to BPA and phthalates, other EDCs linked to epigenetic modifications include the metabolism-disrupting chemicals arsenic [49] and TBT [26]. Importantly, epigenetic modifications are cell- and tissue-specific, and findings from one tissue or cell type may not serve as a proxy for another. Moreover, even single cell types are often heterogeneous, and this lack of uniformity is likely to reflect differences in cell state or cell-specific functions.

For example, not all beta cells function similarly in an islet, and this may be reflected in epigenetic differences. Demonstrating that epigenetic modifications induce changes in gene expression that, in turn, cause abnormal phenotypes is experimentally challenging; however, it is imperative to link exposure-associated changes to metabolic dysregulation.

Beyond offspring: impact of gestational exposures on mothers

While environmental exposures during gestation and lactation can clearly impact the long-term metabolic trajectory of offspring, the consequences of exposures during this window on mothers is less appreciated. Gestational diabetes (GDM) imposes unique pregnancy-associated risks, including gestational hypertension, preeclampsia, future diabetes and obstetric complications. Assessments of EDC effects on GDM in humans remain in their infancy; however, emerging data suggest potential associations. Commonly found in cosmetics, pharmaceuticals and food, parabens were associated with glucose levels during pregnancy in a cohort of women at high risk of GDM [50], while the widely used antibacterial triclosan was nearly significantly associated with GDM in another cohort [51]. In the Infant Development and Environment Study, monoethyl phthalate levels were associated with increased GDM risk [52]. Among the inorganic EDCs, arsenic has been associated with GDM in a host of studies. In the Maternal-Infant Research on Environmental Chemicals Study, the risk of GDM was nearly fourfold greater in the highest vs lowest tertile of dimethylarsinic acid levels [53]. Overall, arsenic has been associated with GDM in many [54–57] but not all studies [58]. In addition to disruptions during gestation, the dynamic changes that happen in the pancreatic islet under the influence of pregnancy hormones suggest that environmental insults that occur during this window may permanently reprogram beta cells and thereby alter their long-term function. Indeed, gestational BPA exposure was shown to not only induce metabolic dysfunction in adult offspring [59], but exposed dams also exhibited glucose intolerance later in life [60]. While more data with comprehensive exposure assessments throughout pregnancy are needed, these data highlight pregnancy as a unique window of susceptibility to EDC-induced metabolic dysfunction.

Exposures in later life and metabolic dysfunction

Robust evidence now links exposure to a variety of EDCs with diabetes risk (reviewed extensively in refs. [7–10]). The majority of cross-sectional and prospective epidemiological analyses have been conducted in adult populations, while most animal studies have employed exposure paradigms using adult mice. Across these studies, there is great diversity in the spectrum of environmental toxicants linked to diabetes, with multiple potential mechanisms of metabolic toxicity (Fig. 2). EDCs linked to diabetes and/or alterations in insulin–glucose dynamics in human studies include arsenic, BPA, dioxins, organochlorine pesticides, PCBs and phthalates, among others [61–72]. In animal studies, disruptions in glucose homeostasis have been linked to arsenic, atrazine, BPA, cadmium, dioxins, malathion, particulate matter air pollution, PCBs, phthalates, persistent organic pollutants, tolylfluanid, and tributyltin [73–93]. Importantly, the diversity of EDCs now linked with metabolic dysfunction, coupled with the frequency with which human populations are exposed, now raise important questions with regard to how these diabetogenic agents

interact with each other, as well as with traditional diabetes risk factors, to augment disease risk [94].

Indeed, diabetogenic EDC exposures do not exist within a vacuum; rather, exposures occur concurrently with other metabolic stressors. Recent animal models demonstrate that excess energy intake induced by provision of a high-fat or high-fat/high-sucrose diet augments the metabolic dysfunction induced by BPA [90, 95, 96], DDT [97], perfluoroalkyl substances [98], tolylfluanid [79] and phthalates [99]. Intriguingly, some studies suggest an EDC–diet potentiation that evolves over time as with dichlorodiphenyldichloroethylene (DDE) [100]. In contrast, other data suggest that changes in diet may precipitate adverse EDC effects; this has been suggested for PCBs, which may specifically impair glucose tolerance after diet-induced weight loss [91, 101]. Importantly, since many EDCs are lipophilic, high-fat diets are likely to augment exposure, whereas reductions in adipose mass with weight loss release EDCs into the circulation [102–105]. Whether this mobilisation from adipose tissue antagonises subsequent weight loss remains under debate.

Environmental diabetogens and health justice

It has long been recognised that diabetes disproportionately afflicts racial and ethnic minorities, as well as those with lower incomes. Despite this knowledge, these disparities have been amplified over time [106]. In addition to higher rates of disease, African-Americans and Latinx/Hispanics are more likely to suffer microvascular complications, including nephropathy [107], neuropathy/amputations [108] and retinopathy [109]. Furthermore, diabetes-related mortality is also higher in these communities [110]. While various social factors are posited to contribute to diabetes disparities, less appreciated is the additional contribution of environmental injustice to this heightened risk. In our recent analysis, five classes of EDCs associated with diabetic phenotypes in cell-based and animal studies (PCBs, organochlorine pesticides, chemical constituents of air pollution, BPA and phthalates) were shown to be linked with incident diabetes in prospective epidemiological studies [111]. Importantly, for each of these EDC classes, data indicated that racial and ethnic minorities, as well as those with lower incomes, are more highly exposed [111]. In addition, studies suggest that racial and ethnic minorities may be at heightened risk for EDC-induced metabolic dysfunction. In a meta-analysis, exposure to higher vs lower levels of both PCBs and organochlorine pesticides was associated with greater diabetes risk in non-white compared with white populations [65]. In a study of phthalates and metabolic control, phthalate-associated insulin resistance was observed in Mexican Americans and African-Americans but not in non-Hispanic whites [112]. In another study, phthalates were associated with worse glucose control and more severe insulin resistance in African-Americans and Mexican Americans compared with non-Hispanic whites [113]. A recent analysis examining the links between phthalates and GDM suggested potentially higher phthalate-associated risk among Asians [52]. Finally, in a study of overall mortality, the hazard ratio for death per unit increase in particulate matter air pollution was higher in Hispanics and much greater in blacks than in whites [114]. Whether the enhanced diabetes- and mortality-associated risks of environmental toxicants in communities with heightened diabetes risk arises from higher baseline exposures, gene–environment interactions or environment-amplified social risk factors remains to be clarified; however, regardless of the

origin of this enhanced risk, addressing environmental potentiators of diabetes is essential for reducing disease disparities and improving health equity.

From abdication to engagement

With mounting evidence that EDCs contribute to diabetes pathogenesis, it is incumbent upon the diabetes care community to address the environmental drivers of the disease; however, current clinical practice guidelines wholly neglect environmental toxicants as diabetes risk factors [115]. While data are required to definitively demonstrate that exposure reduction improves outcomes, emerging evidence provides a framework for action [116]. Importantly, paradigms for environmental risk reduction are available in other fields, such as chronic obstructive lung disease [117] and asthma [118], which acknowledge the health threat posed by air pollution, an exposure also linked to diabetes [119, 120]. In addition, leadership from the American College of Obstetricians and Gynecologists, International Federation of Gynecology and Obstetrics, and the Endocrine Society provide further guidance on areas for risk reduction [5, 121, 122]. The urgency for such action is underscored by data from the European Union estimating that five EDCs alone add €18–29 billion annually to obesity- and diabetes-associated healthcare costs [123]. Conversely, data from the Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study indicate that 25% reductions in only four target compounds (PCB 153, monoethylphthalate, DDE and perfluorononanoic acid) would reduce diabetes prevalence by 13% in Europe (over 150,000 cases), with associated cost savings of €4.51 billion annually [124]. Given the narrow focus of these analyses relative to the total burden of human chemical exposure, reducing contact with diabetogenic chemicals has potential to substantially lower the individual and societal toll of diabetes and its complications. Capitalising on this, however, requires both improved clinical guidance, as well as transformations in public policy to incorporate diabetes and its associated costs into the development of environmental policies [125].

Challenges and controversies

Despite consensus statements regarding EDC risks from the Organisation for Economic Co-operation and Development in the European Union and the Endocrine Society, some controversies persist. These include robust discussion regarding methodologies used in biomonitoring studies, including the validity of ELISA for measuring BPA concentrations in bodily tissues. Importantly, more recent studies have employed more sensitive and reliable techniques to measure very low levels of EDCs. It has also been suggested that low levels of unconjugated BPA in bodily tissues and fluids arise from contamination of collection materials and/or deconjugation of BPA metabolites during storage. To address this concern, multiple human and animal studies have detected BPA in a variety of biological specimens using alternative methods, and storage vials are now BPA-free. Furthermore, high-quality animal studies are now designed to limit background EDC exposure from diet, caging, bedding and water bottles. The route of exposure used in animal studies also remains hotly debated. While differences in exposure routes can affect circulating BPA levels, several studies have shown that there are fewer differences in route-dependent BPA metabolism and excretion than previously thought [126]. Collectively, these issues mandate rigorously designed epidemiological and animal studies. Importantly, recent clinical studies have

demonstrated that administering BPA to humans alters glucose–insulin homeostasis [127, 128]. While these data validate the robust cell-based, animal and epidemiological evidence linking BPA to metabolic disruptions, they also raise an important ethical question: Will such direct evidence of harm be required for each of the thousands of chemicals to which humans are exposed before action is taken to protect public health?

Conclusions

Burgeoning evidence now implicates exposure to a variety of environmental toxicants in the pathogenesis of diabetes. These exposures occur across the lifespan; however, certain developmental periods are uniquely sensitive windows during which metabolism can be permanently disrupted in both the exposed individual and subsequent generations. Importantly, several communities at heightened diabetes risk are also exposed to higher levels of chemicals linked to the disease. Despite this evidence, clinical practice remains blind to EDCs as diabetes drivers. While more work is required to address knowledge gaps regarding environmental exposures and diabetes risk, the weight of the evidence now mandates action to empower individuals and governments to address environmental contributors to diabetes risk.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding

Work in the authors' laboratories is supported by the National Institutes of Health (R01 ES028879, P30 ES027792 and R01 ES025159 supporting RMS; R01 ES023284, R01 ES028206 and P30 ES013508 supporting RAS).

Abbreviations

BPA	Bisphenol A
EDC	Endocrine-disrupting chemical
GDM	Gestational diabetes
GSIS	Glucose-stimulated insulin secretion
PCBs	Polychlorinated biphenyls
PPARγ	Peroxisome proliferator-activated receptor γ
RXR	Retinoid X receptor

References

- [1]. International Diabetes Federation (2017) IDF Diabetes Atlas, 8th Edition. Available from www.idf.org/e-library/epidemiology-research/diabetes-atlas/134-idf-diabetes-atlas-8th-edition.html. Accessed 15 Jan 2018

- [2]. World Health Organization (2016) Global report on diabetes. World Health Organization, Geneva
- [3]. American Diabetes Association (2018) Economic costs of diabetes in the U.S. in 2017. *Diabetes care* 41(5): 917–928. 10.2337/dci18-0007 [PubMed: 29567642]
- [4]. Squires E, Duber H, Campbell M, et al. (2018) Health care spending on diabetes in the U.S., 1996–2013. *Diabetes care* 41(7): 1423–1431. 10.2337/dc17-1376 [PubMed: 29748431]
- [5]. Gore AC, Chappell VA, Fenton SE, et al. (2015) EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals. *Endocrine reviews* 36(6): E1–E150. 10.1210/er.2015-1010 [PubMed: 26544531]
- [6]. Zoeller RT, Brown TR, Doan LL, et al. (2012) Endocrine-disrupting chemicals and public health protection: a statement of principles from The Endocrine Society. *Endocrinology* 153(9): 4097–4110. 10.1210/en.2012-1422 [PubMed: 22733974]
- [7]. Heindel JJ, Blumberg B, Cave M, et al. (2017) Metabolism disrupting chemicals and metabolic disorders. *Reprod Toxicol* 68: 3–33. 10.1016/j.reprotox.2016.10.001 [PubMed: 27760374]
- [8]. Mimoto MS, Nadal A, Sargis RM (2017) Polluted Pathways: mechanisms of metabolic disruption by endocrine disrupting chemicals. *Curr Environ Health Rep* 4(2): 208–222. 10.1007/s40572-017-0137-0 [PubMed: 28432637]
- [9]. Sargis RM (2014) The hijacking of cellular signaling and the diabetes epidemic: mechanisms of environmental disruption of insulin action and glucose homeostasis. *Diabetes & metabolism journal* 38: 1–12. 10.4093/dmj.2014.38.1.1 [PubMed: 24627822]
- [10]. Neel BA, Sargis RM (2011) The paradox of progress: environmental disruption of metabolism and the diabetes epidemic. *Diabetes* 60(7): 1838–1848. 60/7/1838 [pii] 10.2337/db11-0153 [PubMed: 21709279]
- [11]. Tuduri EM L; Dos Santos RS; Quesada I; Fuentes E; Alonso-Magdalena P (2019) Timing of exposure to bisphenol-A: implications for diabetes development. *Front Endocrinol (Lausanne)* 9: 648 10.3389/fendo.2018.00648
- [12]. Calafat AM, Ye X, Wong LY, Reidy JA, Needham LL (2008) Exposure of the U.S. population to bisphenol A and 4-tertiary-octylphenol: 2003–2004. *Environ Health Perspect* 116(1): 39–44. 10.1289/ehp.10753 [PubMed: 18197297]
- [13]. Ikezuki Y, Tsutsumi O, Takai Y, Kamei Y, Taketani Y (2002) Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum Reprod* 17(11): 2839–2841 [PubMed: 12407035]
- [14]. Schonfelder G, Wittfoht W, Hopp H, Talsness CE, Paul M, Chahoud I (2002) Parent bisphenol A accumulation in the human maternal-fetal-placental unit. *Environ Health Perspect* 110(11): A703–707. 10.1289/ehp.110-1241091 [PubMed: 12417499]
- [15]. Nahar MS, Liao C, Kannan K, Dolinoy DC (2013) Fetal liver bisphenol A concentrations and biotransformation gene expression reveal variable exposure and altered capacity for metabolism in humans. *J Biochem Mol Toxicol* 27(2): 116–123. 10.1002/jbt.21459 [PubMed: 23208979]
- [16]. Vom Saal FS, Nagel SC, Coe BL, Angle BM, Taylor JA (2012) The estrogenic endocrine disrupting chemical bisphenol A (BPA) and obesity. *Molecular and cellular endocrinology* 354(1–2): 74–84. 10.1016/j.mce.2012.01.001 [PubMed: 22249005]
- [17]. Susiarjo M, Xin F, Bansal A, et al. (2015) Bisphenol a exposure disrupts metabolic health across multiple generations in the mouse. *Endocrinology* 156(6): 2049–2058. 10.1210/en.2014-2027 [PubMed: 25807043]
- [18]. Bansal A, Rashid C, Xin F, et al. (2017) Sex- and dose-specific effects of maternal bisphenol a exposure on pancreatic islets of first- and second-generation adult mice offspring. *Environ Health Perspect* 125(9): 097022. 10.1289/EHP1674
- [19]. Bansal A, Li C, Xin F, et al. (2018) Transgenerational effects of maternal bisphenol: a exposure on offspring metabolic health. *Journal of developmental origins of health and disease*: 1–12. 10.1017/S2040174418000764
- [20]. Rubin BS CM; Soto AM (2019) The case for BPA as an obesogen: contributors to the controversy. *Front Endocrinol (Lausanne)* 10: 30 10.3389/fendo.2019.00030 [PubMed: 30787907]

- [21]. Rubin BS, Murray MK, Damassa DA, King JC, Soto AM (2001) Perinatal exposure to low doses of bisphenol A affects body weight, patterns of estrous cyclicity, and plasma LH levels. *Environ Health Perspect* 109(7): 675–680. 10.1289/ehp.01109675 [PubMed: 11485865]
- [22]. Tiano JP, Mauvais-Jarvis F (2012) Importance of oestrogen receptors to preserve functional beta-cell mass in diabetes. *Nat Rev Endocrinol* 8(6): 342–351. 10.1038/nrendo.2011.242 [PubMed: 22330739]
- [23]. Whitehead R, Guan H, Arany E, Cernea M, Yang K (2016) Prenatal exposure to bisphenol A alters mouse fetal pancreatic morphology and islet composition. *Horm Mol Biol Clin Investig* 25(3): 171–179. 10.1515/hmbci-2015-0052
- [24]. Garcia-Arevalo M, Alonso-Magdalena P, Rebelo Dos Santos J, Quesada I, Carneiro EM, Nadal A (2014) Exposure to bisphenol-A during pregnancy partially mimics the effects of a high-fat diet altering glucose homeostasis and gene expression in adult male mice. *PLoS one* 9(6): e100214. 10.1371/journal.pone.0100214
- [25]. Garcia-Arevalo M, Alonso-Magdalena P, Servitja JM, et al. (2016) Maternal exposure to bisphenol-A during pregnancy increases pancreatic beta-cell growth during early life in male mice offspring. *Endocrinology* 157(11): 4158–4171. 10.1210/en.2016-1390 [PubMed: 27623287]
- [26]. Chamorro-Garcia R, Diaz-Castillo C, Shoucri BM, et al. (2017) Ancestral perinatal obesogen exposure results in a transgenerational thrifty phenotype in mice. *Nat Commun* 8(1): 2012. 10.1038/s41467-017-01944-z [PubMed: 29222412]
- [27]. Veiga-Lopez A, Pu Y, Gingrich J, Padmanabhan V (2018) Obesogenic Endocrine disrupting chemicals: identifying knowledge gaps. *Trends Endocrinol Metab* 29(9): 607–625. 10.1016/j.tem.2018.06.003 [PubMed: 30017741]
- [28]. Mennigen JA, Thompson LM, Bell M, Tellez Santos M, Gore AC (2018) Transgenerational effects of polychlorinated biphenyls: 1. Development and physiology across 3 generations of rats. *Environ Health* 17(1): 18. 10.1186/s12940-018-0362-5 [PubMed: 29458364]
- [29]. Skinner MK (2016) Endocrine disruptors in 2015: Epigenetic transgenerational inheritance. *Nat Rev Endocrinol* 12(2): 68–70. 10.1038/nrendo.2015.206 [PubMed: 26585656]
- [30]. Drobna Z, Henriksen AD, Wolstenholme JT, et al. (2018) Transgenerational Effects of Bisphenol A on Gene Expression and DNA Methylation of Imprinted Genes in Brain. *Endocrinology* 159(1): 132–144. 10.1210/en.2017-00730 [PubMed: 29165653]
- [31]. Chamorro-Garcia R, Sahu M, Abbey RJ, Laude J, Pham N, Blumberg B (2013) Transgenerational inheritance of increased fat depot size, stem cell reprogramming, and hepatic steatosis elicited by prenatal exposure to the obesogen tributyltin in mice. *Environ Health Perspect* 121(3): 359–366. 10.1289/ehp.1205701 [PubMed: 23322813]
- [32]. Ptak A, Gregoraszczyk EL (2012) Bisphenol A induces leptin receptor expression, creating more binding sites for leptin, and activates the JAK/Stat, MAPK/ERK and PI3K/Akt signalling pathways in human ovarian cancer cell. *Toxicology letters* 210(3): 332–337. 10.1016/j.toxlet.2012.02.003 [PubMed: 22343039]
- [33]. Xu X, Lu Y, Zhang G, et al. (2014) Bisphenol A promotes dendritic morphogenesis of hippocampal neurons through estrogen receptor-mediated ERK1/2 signal pathway. *Chemosphere* 96: 129–137. 10.1016/j.chemosphere.2013.09.063 [PubMed: 24231043]
- [34]. Acconcia F, Pallottini V, Marino M (2015) Molecular mechanisms of action of BPA. *Dose Response* 13(4): 1559325815610582. 10.1177/1559325815610582
- [35]. Fernandez MF, Arrebola JP, Jimenez-Diaz I, et al. (2016) Bisphenol A and other phenols in human placenta from children with cryptorchidism or hypospadias. *Reprod Toxicol* 59: 89–95. 10.1016/j.reprotox.2015.11.002 [PubMed: 26602963]
- [36]. Prasanth GK, Divya LM, Sadasivan C (2010) Bisphenol-A can bind to human glucocorticoid receptor as an agonist: an in silico study. *Journal of applied toxicology : JAT* 30(8): 769–774. 10.1002/jat.1570 [PubMed: 20669259]
- [37]. Sargis RM, Johnson DN, Choudhury RA, Brady MJ (2010) Environmental endocrine disruptors promote adipogenesis in the 3T3-L1 cell line through glucocorticoid receptor activation. *Obesity* 18(7): 1283–1288. 10.1038/oby.2009.419 [PubMed: 19927138]
- [38]. Wang J, Sun B, Hou M, Pan X, Li X (2013) The environmental obesogen bisphenol A promotes adipogenesis by increasing the amount of 11 β -hydroxysteroid dehydrogenase type 1 in the

- adipose tissue of children. *International journal of obesity* 37(7): 999–1005. 10.1038/ijo.2012.173 [PubMed: 23090578]
- [39]. Somm E, Schwitzgebel VM, Toulotte A, et al. (2009) Perinatal exposure to bisphenol A alters early adipogenesis in the rat. *Environ Health Perspect* 117(10): 1549–1555. 10.1289/ehp.11342 [PubMed: 20019905]
- [40]. Riu A, Grimaldi M, le Maire A, et al. (2011) Peroxisome proliferator-activated receptor gamma is a target for halogenated analogs of bisphenol A. *Environ Health Perspect* 119(9): 1227–1232. 10.1289/ehp.1003328 [PubMed: 21561829]
- [41]. Grun F, Watanabe H, Zamanian Z, et al. (2006) Endocrine-disrupting organotin compounds are potent inducers of adipogenesis in vertebrates. *Molecular endocrinology* 20(9): 2141–2155. me.2005–0367 [pii] 10.1210/me.2005-0367 [PubMed: 16613991]
- [42]. Shoucri BM, Hung VT, Chamorro-Garcia R, Shioda T, Blumberg B (2018) Retinoid X receptor activation during adipogenesis of female mesenchymal stem cells programs a dysfunctional adipocyte. *Endocrinology* 159(8): 2863–2883. 10.1210/en.2018-00056 [PubMed: 29860300]
- [43]. Regnier SM, El-Hashani E, Kamau W, Zhang X, Massad NL, Sargis RM (2015) Tributyltin differentially promotes development of a phenotypically distinct adipocyte. *Obesity* 23(9): 1864–1871. 10.1002/oby.21174 [PubMed: 26243053]
- [44]. Jiang Y, Xia W, Zhu Y, et al. (2014) Mitochondrial dysfunction in early life resulted from perinatal bisphenol A exposure contributes to hepatic steatosis in rat offspring. *Toxicology letters* 228(2): 85–92. 10.1016/j.toxlet.2014.04.013 [PubMed: 24768697]
- [45]. Chandravanshi LP, Gupta R, Shukla RK (2018) Developmental neurotoxicity of arsenic: involvement of oxidative stress and mitochondrial functions. *Biol Trace Elem Res* 186(1): 185–198. 10.1007/s12011-018-1286-1 [PubMed: 29502250]
- [46]. Dolinoy DC, Huang D, Jirtle RL (2007) Maternal nutrient supplementation counteracts bisphenol A-induced DNA hypomethylation in early development. *Proceedings of the National Academy of Sciences of the United States of America* 104(32): 13056–13061. 10.1073/pnas.0703739104
- [47]. Tindula G, Murphy SK, Grenier C, et al. (2018) DNA methylation of imprinted genes in Mexican-American newborn children with prenatal phthalate exposure. *Epigenomics* 10(7): 1011–1026. 10.2217/epi-2017-0178 [PubMed: 29957030]
- [48]. Pogribny IP, Tryndyak VP, Boureiko A, et al. (2008) Mechanisms of peroxisome proliferator-induced DNA hypomethylation in rat liver. *Mutat Res* 644(1–2): 17–23. 10.1016/j.mrfmmm.2008.06.009 [PubMed: 18639561]
- [49]. Meakin CJ, Martin EM, Szilagy J, Nylander-French LA, Fry RC (2019) Inorganic arsenic as an endocrine disruptor: modulation of the glucocorticoid receptor pathway in placental cells via CpG methylation. *Chem Res Toxicol*. 10.1021/acs.chemrestox.8b00352
- [50]. Bellavia A, Chiu YH, Brown FM, et al. (2019) Urinary concentrations of parabens mixture and pregnancy glucose levels among women from a fertility clinic. *Environ Res* 168: 389–396. 10.1016/j.envres.2018.10.009 [PubMed: 30384233]
- [51]. Ouyang F, Tang N, Zhang HJ, et al. (2018) Maternal urinary triclosan level, gestational diabetes mellitus and birth weight in Chinese women. *Sci Total Environ* 626: 451–457. 10.1016/j.scitotenv.2018.01.102 [PubMed: 29353787]
- [52]. Shaffer RM, Ferguson KK, Sheppard L, et al. (2019) Maternal urinary phthalate metabolites in relation to gestational diabetes and glucose intolerance during pregnancy. *Environment international* 123: 588–596. 10.1016/j.envint.2018.12.021 [PubMed: 30622083]
- [53]. Ashley-Martin J, Dodds L, Arbuckle TE, et al. (2018) Association between maternal urinary speciated arsenic concentrations and gestational diabetes in a cohort of Canadian women. *Environment international* 121(Pt 1): 714–720. 10.1016/j.envint.2018.10.008 [PubMed: 30321846]
- [54]. Xia X, Liang C, Sheng J, et al. (2018) Association between serum arsenic levels and gestational diabetes mellitus: A population-based birth cohort study. *Environ Pollut* 235: 850–856. 10.1016/j.envpol.2018.01.016 [PubMed: 29348076]
- [55]. Marie C, Leger S, Guttman A, et al. (2018) Exposure to arsenic in tap water and gestational diabetes: A French semi-ecological study. *Environ Res* 161: 248–255. 10.1016/j.envres.2017.11.016 [PubMed: 29169099]

- [56]. Farzan SF, Gossai A, Chen Y, Chasan-Taber L, Baker E, Karagas M (2016) Maternal arsenic exposure and gestational diabetes and glucose intolerance in the New Hampshire birth cohort study. *Environ Health* 15(1): 106 10.1186/s12940-016-0194-0 [PubMed: 27825389]
- [57]. Shapiro GD, Dodds L, Arbuckle TE, et al. (2015) Exposure to phthalates, bisphenol A and metals in pregnancy and the association with impaired glucose tolerance and gestational diabetes mellitus: The MIREC study. *Environment international* 83: 63–71. 10.1016/j.envint.2015.05.016 [PubMed: 26101084]
- [58]. Munoz MP, Valdes M, Munoz-Quezada MT, et al. (2018) Urinary inorganic arsenic concentration and gestational diabetes mellitus in pregnant women from Arica, Chile. *Int J Environ Res Public Health* 15(7). 10.3390/ijerph15071418
- [59]. Alonso-Magdalena P, Vieira E, Soriano S, et al. (2010) Bisphenol A exposure during pregnancy disrupts glucose homeostasis in mothers and adult male offspring. *Environ Health Perspect* 118(9): 1243–1250. 10.1289/ehp.1001993 [PubMed: 20488778]
- [60]. Alonso-Magdalena P, Garcia-Arevalo M, Quesada I, Nadal A (2015) Bisphenol-A treatment during pregnancy in mice: a new window of susceptibility for the development of diabetes in mothers later in life. *Endocrinology* 156(5): 1659–1670. 10.1210/en.2014-1952 [PubMed: 25830705]
- [61]. Maull EA, Ahsan H, Edwards J, et al. (2012) Evaluation of the association between arsenic and diabetes: a National Toxicology Program workshop review. *Environ Health Perspect* 120(12): 1658–1670. 10.1289/ehp.1104579 [PubMed: 22889723]
- [62]. Sung TC, Huang JW, Guo HR (2015) Association between arsenic exposure and diabetes: a meta-analysis. *Biomed Res Int* 2015: 368087. 10.1155/2015/368087
- [63]. Wang W, Xie Z, Lin Y, Zhang D (2014) Association of inorganic arsenic exposure with type 2 diabetes mellitus: a meta-analysis. *Journal of epidemiology and community health* 68(2): 176–184. 10.1136/jech-2013-203114 [PubMed: 24133074]
- [64]. Ranciere F, Lyons JG, Loh VH, et al. (2015) Bisphenol A and the risk of cardiometabolic disorders: a systematic review with meta-analysis of the epidemiological evidence. *Environ Health* 14: 46 10.1186/s12940-015-0036-5 [PubMed: 26026606]
- [65]. Song Y, Chou EL, Baecker A, et al. (2016) Endocrine-disrupting chemicals, risk of type 2 diabetes, and diabetes-related metabolic traits: A systematic review and meta-analysis. *J Diabetes* 8(4): 516–532 [PubMed: 26119400]
- [66]. Jensen TK, Timmermann AG, Rossing LI, et al. (2014) Polychlorinated biphenyl exposure and glucose metabolism in 9-year-old Danish children. *The Journal of clinical endocrinology and metabolism* 99(12): E2643–E2651. 10.1210/jc.2014-1683
- [67]. Suarez-Lopez JR, Lee DH, Porta M, Steffes MW, Jacobs DR Jr. (2015) Persistent organic pollutants in young adults and changes in glucose related metabolism over a 23-year follow-up. *Environ Res* 137: 485–494. 10.1016/j.envres.2014.11.001 [PubMed: 25706918]
- [68]. Tang M, Chen K, Yang F, Liu W (2014) Exposure to organochlorine pollutants and type 2 diabetes: a systematic review and meta-analysis. *PLoS one* 9(10): e85556. 10.1371/journal.pone.0085556
- [69]. Taylor KW, Novak RF, Anderson HA, et al. (2013) Evaluation of the association between persistent organic pollutants (POPs) and diabetes in epidemiological studies: a national toxicology program workshop review. *Environ Health Perspect* 121(7): 774–783. 10.1289/ehp.1205502 [PubMed: 23651634]
- [70]. Shoshtari-Yeganeh B, Zarean M, Mansourian M, et al. (2019) Systematic review and meta-analysis on the association between phthalates exposure and insulin resistance. *Environ Sci Pollut Res Int*. 10.1007/s11356-019-04373-1
- [71]. Sun Q, Cornelis MC, Townsend MK, et al. (2014) Association of urinary concentrations of bisphenol A and phthalate metabolites with risk of type 2 diabetes: a prospective investigation in the Nurses' Health Study (NHS) and NHSII cohorts. *Environ Health Perspect* 122(6): 616–623. 10.1289/ehp.1307201 [PubMed: 24633239]
- [72]. Lind PM, Zethelius B, Lind L (2012) Circulating levels of phthalate metabolites are associated with prevalent diabetes in the elderly. *Diabetes care* 35(7): 1519–1524. 10.2337/dc11-2396 [PubMed: 22498808]

- [73]. Rajesh P, Sathish S, Srinivasan C, Selvaraj J, Balasubramanian K (2013) Phthalate is associated with insulin resistance in adipose tissue of male rat: role of antioxidant vitamins. *J Cell Biochem* 114(3): 558–569. 10.1002/jcb.24399 [PubMed: 22991202]
- [74]. Indumathi D, Jayashree S, Selvaraj J, et al. (2013) Effect of bisphenol-A on insulin signal transduction and glucose oxidation in skeletal muscle of adult male albino rat. *Hum Exp Toxicol* 32(9): 960–971. 10.1177/0960327112470273 [PubMed: 23424204]
- [75]. Jayashree S, Indumathi D, Akilavalli N, Sathish S, Selvaraj J, Balasubramanian K (2013) Effect of bisphenol-A on insulin signal transduction and glucose oxidation in liver of adult male albino rat. *Environ Toxicol Pharmacol* 35(2): 300–310. 10.1016/j.etap.2012.12.016 [PubMed: 23376180]
- [76]. Han JC, Park SY, Hah BG, et al. (2003) Cadmium induces impaired glucose tolerance in rat by down-regulating GLUT4 expression in adipocytes. *Arch Biochem Biophys* 413(2): 213–220. [PubMed: 12729619]
- [77]. Liu PC, Matsumura F (1995) Differential effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the “adipose-type” and “brain-type” glucose transporters in mice. *Molecular pharmacology* 47(1): 65–73 [PubMed: 7530807]
- [78]. Enan E, Matsumura F (1994) 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-induced changes in glucose transporting activity in guinea pigs, mice, and rats in vivo and in vitro. *J Biochem Toxicol* 9(2): 97–106 [PubMed: 8071950]
- [79]. Regnier SM, Kirkley AG, Ruiz D, et al. (2018) Diet-dependence of metabolic perturbations mediated by the endocrine disruptor tolylfluanid. *Endocr Connect* 7(1): 159–168. 10.1530/EC-17-0320 [PubMed: 29187361]
- [80]. Regnier SM, Kirkley AG, Ye H, et al. (2015) Dietary exposure to the endocrine disruptor tolylfluanid promotes global metabolic dysfunction in male mice. *Endocrinology* 156(3): 896–910. 10.1210/en.2014-1668 [PubMed: 25535829]
- [81]. Paul DS, Walton FS, Saunders RJ, Styblo M (2011) Characterization of the impaired glucose homeostasis produced in C57BL/6 mice by chronic exposure to arsenic and high-fat diet. *Environ Health Perspect* 119(8): 1104–1109. 10.1289/ehp.1003324 [PubMed: 21592922]
- [82]. Kirkley AG, Carmean CM, Ruiz D, et al. (2018) Arsenic exposure induces glucose intolerance and alters global energy metabolism. *Am J Physiol Regul Integr Comp Physiol* 314(2): R294–R303. 10.1152/ajpregu.00522.2016 [PubMed: 29118024]
- [83]. Hill DS, Wlodarczyk BJ, Mitchell LE, Finnell RH (2009) Arsenate-induced maternal glucose intolerance and neural tube defects in a mouse model. *Toxicol Appl Pharmacol* 239(1): 29–36. 10.1016/j.taap.2009.05.009 [PubMed: 19446573]
- [84]. Sun Q, Yue P, Deiluiis JA, et al. (2009) Ambient air pollution exaggerates adipose inflammation and insulin resistance in a mouse model of diet-induced obesity. *Circulation* 119(4): 538–546. 10.1161/CIRCULATIONAHA.108.799015 [PubMed: 19153269]
- [85]. Panahi P, Vosough-Ghanbari S, Pournourmohammadi S, et al. (2006) Stimulatory effects of malathion on the key enzymes activities of insulin secretion in langerhans islets, glutamate dehydrogenase and glucokinase. *Toxicol Mech Methods* 16(4): 161–167. 10.1080/15376520500191623 [PubMed: 20021041]
- [86]. Batista TM, Alonso-Magdalena P, Vieira E, et al. (2012) Short-term treatment with bisphenol-a leads to metabolic abnormalities in adult male mice. *PloS one* 7(3): e33814. 10.1371/journal.pone.0033814
- [87]. Zuo Z, Chen S, Wu T, et al. (2011) Tributyltin causes obesity and hepatic steatosis in male mice. *Environ Toxicol* 26(1): 79–85. 10.1002/tox.20531 [PubMed: 19760618]
- [88]. Huang CF, Yang CY, Chan DC, et al. (2015) Arsenic Exposure and glucose intolerance/insulin resistance in estrogen-deficient female mice. *Environ Health Perspect* 123(11): 1138–1144. 10.1289/ehp.1408663 [PubMed: 25859628]
- [89]. Lim S, Ahn SY, Song IC, et al. (2009) Chronic exposure to the herbicide, atrazine, causes mitochondrial dysfunction and insulin resistance. *PloS one* 4(4): e5186. 10.1371/journal.pone.0005186

- [90]. Moon MK, Jeong IK, Jung Oh T, et al. (2015) Long-term oral exposure to bisphenol A induces glucose intolerance and insulin resistance. *The Journal of endocrinology* 226(1): 35–42. 10.1530/JOE-14-0714 [PubMed: 25972359]
- [91]. Baker NA, Karounos M, English V, et al. (2013) Coplanar polychlorinated biphenyls impair glucose homeostasis in lean C57BL/6 mice and mitigate beneficial effects of weight loss on glucose homeostasis in obese mice. *Environ Health Perspect* 121(1): 105–110. 10.1289/ehp.1205421 [PubMed: 23099484]
- [92]. Gray SL, Shaw AC, Gagne AX, Chan HM (2013) Chronic exposure to PCBs (Aroclor 1254) exacerbates obesity-induced insulin resistance and hyperinsulinemia in mice. *J Toxicol Environ Health A* 76(12): 701–715. 10.1080/15287394.2013.796503 [PubMed: 23980837]
- [93]. Huang CF, Yang CY, Tsai JR, Wu CT, Liu SH, Lan KC (2018) Low-dose tributyltin exposure induces an oxidative stress-triggered JNK-related pancreatic beta-cell apoptosis and a reversible hypoinsulinemic hyperglycemia in mice. *Sci Rep* 8(1): 5734 10.1038/s41598-018-24076-w [PubMed: 29636531]
- [94]. Sargis RM (2015) Metabolic disruption in context: clinical avenues for synergistic perturbations in energy homeostasis by endocrine disrupting chemicals. *Endocrine Disruptors* 3(1): e1080788–1080781-1080787. 10.1080/23273747.2015.1080788
- [95]. Wei J, Lin Y, Li Y, et al. (2011) Perinatal exposure to bisphenol A at reference dose predisposes offspring to metabolic syndrome in adult rats on a high-fat diet. *Endocrinology* 152(8): 3049–3061. 10.1210/en.2011-0045 [PubMed: 21586551]
- [96]. Ding S, Fan Y, Zhao N, et al. (2014) High-fat diet aggravates glucose homeostasis disorder caused by chronic exposure to bisphenol A. *The Journal of endocrinology* 221(1): 167–179. 10.1530/JOE-13-0386 [PubMed: 24501380]
- [97]. La Merrill M, Karey E, Moshier E, et al. (2014) Perinatal exposure of mice to the pesticide DDT impairs energy expenditure and metabolism in adult female offspring. *PloS one* 9(7): e103337. 10.1371/journal.pone.0103337
- [98]. Wan HT, Zhao YG, Leung PY, Wong CK (2014) Perinatal exposure to perfluorooctane sulfonate affects glucose metabolism in adult offspring. *PloS one* 9(1): e87137. 10.1371/journal.pone.0087137
- [99]. Strakovsky RS, Lezmi S, Shkoda I, Flaws JA, Helferich WG, Pan YX (2015) In utero growth restriction and catch-up adipogenesis after developmental di (2-ethylhexyl) phthalate exposure cause glucose intolerance in adult male rats following a high-fat dietary challenge. *J Nutr Biochem* 26(11): 1208–1220. 10.1016/j.jnutbio.2015.05.012 [PubMed: 26188368]
- [100]. Howell GE 3rd, Mulligan C, Meek E, Chambers JE (2015) Effect of chronic p,p'-dichlorodiphenyldichloroethylene (DDE) exposure on high fat diet-induced alterations in glucose and lipid metabolism in male C57BL/6H mice. *Toxicology* 328: 112–122. 10.1016/j.tox.2014.12.017 [PubMed: 25541407]
- [101]. Baker NA, Shoemaker R, English V, et al. (2015) Effects of adipocyte aryl hydrocarbon receptor deficiency on PCB-induced disruption of glucose homeostasis in lean and obese mice. *Environ Health Perspect* 123(10): 944–950. 10.1289/ehp.1408594 [PubMed: 25734695]
- [102]. Dirtu AC, Dirinck E, Malarvannan G, et al. (2013) Dynamics of organohalogenated contaminants in human serum from obese individuals during one year of weight loss treatment. *Environ Sci Technol* 47(21): 12441–12449. 10.1021/es400657t
- [103]. Dirinck E, Dirtu AC, Jorens PG, Malarvannan G, Covaci A, Van Gaal LF (2015) Pivotal role for the visceral fat compartment in the release of persistent organic pollutants during weight loss. *The Journal of clinical endocrinology and metabolism* 100(12): 4463–4471. 10.1210/jc.2015-2571 [PubMed: 26469381]
- [104]. Hue O, Marcotte J, Berrigan F, et al. (2006) Increased plasma levels of toxic pollutants accompanying weight loss induced by hypocaloric diet or by bariatric surgery. *Obes Surg* 16(9): 1145–1154. 10.1381/096089206778392356 [PubMed: 16989697]
- [105]. Charlier C, Desai C, Plomteux G (2002) Human exposure to endocrine disrupters: consequences of gastroplasty on plasma concentration of toxic pollutants. *Int J Obes Relat Metab Disord* 26(11): 1465–1468. 10.1038/sj.ijo.0802144 [PubMed: 12439648]

- [106]. Beckles GL, Chou CF; Centers for Disease Control and Prevention (CDC) (2013) Diabetes - United States, 2006 and 2010. *MMWR Surveill Summ* 62 (3): 99–104.
- [107]. Ferdinand KC, Nasser SA (2015) Racial/ethnic disparities in prevalence and care of patients with type 2 diabetes mellitus. *Curr Med Res Opin* 31(5): 913–923. 10.1185/03007995.2015.1029894 [PubMed: 25772230]
- [108]. Peek ME (2011) Gender differences in diabetes-related lower extremity amputations. *Clin Orthop Relat Res* 469(7): 1951–1955. 10.1007/s11999-010-1735-4 [PubMed: 21161738]
- [109]. Nsiah-Kumi P, Ortmeier SR, Brown AE (2009) Disparities in diabetic retinopathy screening and disease for racial and ethnic minority populations—a literature review. *J Natl Med Assoc* 101(5): 430–437 [PubMed: 19476196]
- [110]. Rosenstock S, Whitman S, West JF, Balkin M (2014) Racial disparities in diabetes mortality in the 50 most populous US cities. *J Urban Health* 91(5): 873–885. 10.1007/s11524-013-9861-4 [PubMed: 24532483]
- [111]. Ruiz D, Becerra M, Jagai JS, Ard K, Sargis RM (2018) Disparities in Environmental exposures to endocrine-disrupting chemicals and diabetes risk in vulnerable populations. *Diabetes care* 41(1): 193–205. 10.2337/dc16-2765 [PubMed: 29142003]
- [112]. Trasande L, Spanier AJ, Sathyanarayana S, Attina TM, Blustein J (2013) Urinary phthalates and increased insulin resistance in adolescents. *Pediatrics* 132(3): e646–655. 10.1542/peds.2012-4022 [PubMed: 23958772]
- [113]. Huang T, Saxena AR, Isganaitis E, James-Todd T (2014) Gender and racial/ethnic differences in the associations of urinary phthalate metabolites with markers of diabetes risk: National Health and Nutrition Examination Survey 2001–2008. *Environ Health* 13(1): 6 10.1186/1476-069X-13-6 [PubMed: 24499162]
- [114]. Di Q, Wang Y, Zanobetti A, et al. (2017) Air pollution and mortality in the medicare population. *N Engl J Med* 376(26): 2513–2522. 10.1056/NEJMoa1702747 [PubMed: 28657878]
- [115]. American Diabetes Association (2019) Standards of medical care in diabetes: 2019. *Diabetes care* 42(Suppl 1): S1–S186. 10.2337/dc19-S001 [PubMed: 30559224]
- [116]. Sargis RM, Heindel JJ, Padmanabhan V (2019) Interventions to address environmental metabolism-disrupting chemicals: changing the narrative to empower action to restore metabolic health. *Front Endocrinol (Lausanne)* 10: 33 10.3389/fendo.2019.00033 [PubMed: 30778334]
- [117]. Vogelmeier CF, Criner GJ, Martinez FJ, et al. (2017) Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Lung Disease 2017 Report. GOLD Executive Summary. *Am J Respir Crit Care Med* 195(5): 557–582. 10.1164/rccm.201701-0218PP [PubMed: 28128970]
- [118]. National Asthma Education Prevention Program (2007) Expert Panel Report 3 (EPR-3): Guidelines for the Diagnosis and Management of Asthma—Summary Report 2007. *J Allergy Clin Immunol* 120(5 Suppl): S94–138. 10.1016/j.jaci.2007.09.043 [PubMed: 17983880]
- [119]. Wang B, Xu D, Jing Z, Liu D, Yan S, Wang Y (2014) Effect of long-term exposure to air pollution on type 2 diabetes mellitus risk: a systemic review and meta-analysis of cohort studies. *Eur J Endocrinol* 171(5): R173–R182. 10.1530/EJE-14-0365 [PubMed: 25298376]
- [120]. Zhao Z, Lin F, Wang B, Cao Y, Hou X, Wang Y (2016) Residential proximity to major roadways and risk of type 2 diabetes mellitus: a meta-analysis. *Int J Environ Res Public Health* 14(1). 10.3390/ijerph14010003
- [121]. Di Renzo GC, Conry JA, Blake J, et al. (2015) International Federation of Gynecology and Obstetrics opinion on reproductive health impacts of exposure to toxic environmental chemicals. *Int J Gynaecol Obstet* 131(3): 219–225. 10.1016/j.ijgo.2015.09.002 [PubMed: 26433469]
- [122]. American College of Obstetricians and Gynecologists (2013) Exposure to toxic environmental agents. *Fertil Steril* 100(4): 931–934. 10.1016/j.fertnstert.2013.08.043 [PubMed: 24070500]
- [123]. Legler J, Fletcher T, Govarts E, et al. (2015) Obesity, diabetes, and associated costs of exposure to endocrine-disrupting chemicals in the European Union. *The Journal of clinical endocrinology and metabolism* 100(4): 1278–1288. 10.1210/jc.2014-4326 [PubMed: 25742518]
- [124]. Trasande L, Lampa E, Lind L, Lind PM (2017) Population attributable risks and costs of diabetogenic chemical exposures in the elderly. *Journal of epidemiology and community health* 71(2): 111–114. 10.1136/jech-2016-208006 [PubMed: 27789757]

- [125]. Shaikh S, Jagai JS, Ashley C, Zhou S, Sargis RM (2018) Underutilized and under threat: environmental policy as a tool to address diabetes risk. *Curr Diab Rep* 18(5): 25 10.1007/s11892-018-0993-5 [PubMed: 29582168]
- [126]. Vandenberg LN, Maffini MV, Sonnenschein C, Rubin BS, Soto AM (2009) Bisphenol-A and the great divide: a review of controversies in the field of endocrine disruption. *Endocrine reviews* 30(1): 75–95. 10.1210/er.2008-0021 [PubMed: 19074586]
- [127]. Stahlhut RW, Myers JP, Taylor JA, Nadal A, Dyer JA, Vom Saal FS (2018) Experimental BPA exposure and glucose-stimulated insulin response in adult men and women. *J Endocr Soc* 2(10): 1173–1187. 10.1210/js.2018-00151 [PubMed: 30302422]
- [128]. Hagobian TA, Bird A, Stanelle S, Williams D, Schaffner A, Phelan S (2019) Pilot study on the effect of orally administered bisphenol A on glucose and insulin response in nonobese adults. *J Endocr Soc* 3(3): 643–654. 10.1210/js.2018-00322 [PubMed: 30842988]
- [129]. Kahn SE, Hull RL, Utzschneider KM (2006) Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* 444(7121): 840–846. 10.1038/nature05482 [PubMed: 17167471]





Fig. 1. Sources of EDCs. A diverse array of chemicals from various sources have been linked to metabolic dysfunction in cell-based, animal and epidemiological studies. These metabolism-disrupting EDCs include both inorganic and organic compounds of both natural and synthetic origin. Humans are exposed through the use, production and environmental dissemination of these chemicals in food production, industrial activity, and home and personal care, as well as through medical care. This figure is available as part of a downloadable slideset

**Fig. 2.**

EDCs associated with type 2 diabetes pathogenesis and their proposed mechanisms of action. Type 2 diabetes arises from a combination of reduced insulin sensitivity (solid black arrow) and progressive beta cell failure (dashed black arrow). The blue circles, labelled a to e, represent individual states. Starting at state 'a', (in the middle of the normoglycaemia curve, green shading), if a person with these levels of insulin sensitivity and insulin secretion were to exercise, lose weight and get better sleep, they would slide down the curve to 'b'. Unfortunately, our society is instead less active, consuming excess food and sleeping less, driving individuals from state 'a' to state 'c'. This situation is, however, untenable in the long run. As one's beta cells begin to fail, they fall off the curve (dashed line) to 'd' (impaired glucose tolerance, yellow shading); this condition then further deteriorates to 'e' (type 2 diabetes, red). Several EDCs have been linked to altered insulin sensitivity (darker red text box), beta cell disruptions (darker blue text box), or both. Multiple mechanisms have been ascribed to EDC-induced beta cell dysfunction and altered functioning of insulin-responsive tissues (lighter blue and lighter red text boxes). Data supporting EDC-mediated diabetes pathogenesis are derived from: cell-, islet- and tissue-based studies; animal models; and epidemiological and clinical studies (indicated by the superscript numbers; purple text box). BPA, bisphenol A; DDE, dichlorodiphenyldichloroethylene; DDT, dichlorodiphenyltrichloroethane; GR, glucocorticoid receptor; OC, organochlorine; PBDEs, polybrominated diphenyl ethers; PCBs, polychlorinated biphenyls; PFASs, per- and polyfluoroalkyl substances; PCDD/Fs, polychlorinated dibenzo-*p*-dioxins and furans; PM, particulate matter; POPs, persistent organic pollutants; PPARs, peroxisome proliferator-

activated receptors; TCDD, 2,3,7,8-tetrachlorodibenzodioxin. Data compiled from [7–10].
Figure adapted by permission from Springer Nature [129], ©2006 Nature Publishing Group.
This figure is available as part of a downloadable slideset

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Fig. 3.

Influence of EDCs across the lifespan. EDCs and other environmental toxicants are an important component of the overall environmental milieu (blue) that interacts with genetic susceptibility to influence diabetes pathogenesis across the lifespan (green arrow). Importantly, exposures during sensitive windows can disrupt developmental programming and result in long-term metabolic dysfunction in both the exposed individual as well as future generations (black arrows). This figure is available as part of a downloadable slideset