



HHS Public Access

Author manuscript

J Biomol NMR. Author manuscript; available in PMC 2020 September 04.

Published in final edited form as:

J Biomol NMR. 2020 July ; 74(6-7): 355–363. doi:10.1007/s10858-020-00323-z.

Site-Specific Resolution of Anionic Residues in Proteins Using Solid-State NMR Spectroscopy

Jianping Li[†], Ampon Sae Her[†], Nathaniel J. Traaseth^{*}

Department of Chemistry, New York University, New York, NY 10003

Abstract

NMR spectroscopy is commonly used to infer site-specific acid dissociation constants (pK_a) since the chemical shift is sensitive to the protonation state. Methods that probe atoms nearest to the functional groups involved in acid/base chemistry are the most sensitive for determining the protonation state. In this work, we describe a magic-angle-spinning (MAS) solid-state NMR approach to measure chemical shifts on the side chain of the anionic residues aspartate and glutamate. This method involves a combination of double quantum spectroscopy in the indirect dimension and REDOR dephasing to provide a sensitive and resolved view of these amino acid residues that are commonly involved in enzyme catalysis and membrane protein transport. To demonstrate the applicability of the approach, we carried out measurements using a microcrystalline soluble protein (ubiquitin) and a membrane protein embedded in lipid bilayers (EmrE). Overall, the resolution available from the double quantum dimension and confidence in identification of aspartate and glutamate residues from the REDOR filter make this method the most convenient for characterizing protonation states and deriving pK_a values using MAS.

Keywords

magic-angle-spinning; solid-state NMR; acid dissociation constants; microcrystalline proteins; membrane proteins; EmrE

INTRODUCTION

Charged amino acids play key roles in biological functions, including protein folding and binding interactions^{1,2}. In addition, acid/base chemistry at charged residues are involved in enzyme catalysis such as protein degradation and proton transport across cellular membranes^{3–7}. Since the chemical shift is a sensitive reporter of the electrostatic environment, NMR spectroscopy has been a valuable experimental approach to probe site-specific protonation states and acid dissociation constants (pK_a)^{8–18}. The most effective NMR methods for inferring charged states rely on direct measurements of nuclei at the site of acid/base chemistry. For example, solution NMR methods have been developed for determining pK_a values for the anionic residues aspartate and glutamate by recording ¹³C chemical shifts of the side chain carboxyl carbon in the indirect dimension (i.e., HCABGCO

^{*} Author for correspondence: Nathaniel J. Traaseth, 100 Washington Square East, New York, NY 10003, Phone: (212) 992-9784, traaseth@nyu.edu.

[†] These authors contributed equally to this work.

pulse sequence¹⁹). This experiment was employed to determine asymmetric protonation sites at a conserved aspartate residue in the HIV protease dimer when bound to an asymmetric inhibitor¹⁹. Although this experiment has been used with success for small and medium sized soluble proteins, application of the method to macromolecular systems is limited due to the requirement of multiple magnetization transfers that reduce sensitivity. Furthermore, this solution NMR experiment is not applicable for solid-like samples such as microcrystalline proteins or membrane proteins in liposomes, which are ideally studied using solid-state NMR spectroscopy.

In this article, we present a magic-angle-spinning (MAS) solid-state NMR method to directly observe the carboxyl groups of aspartate and glutamate using double quantum spectroscopy²⁰. A new pulse sequence is presented by combining double quantum spectroscopy with REDOR dephasing to sensitively distinguish aspartate and glutamate side chains from those of asparagine and glutamine. As proof of concept, we applied the method to microcrystalline ubiquitin and a mutant of the multidrug efflux pump EmrE reconstituted into lipid bilayers. This method offers improved sensitivity and resolution relative to available methods for performing pH titration experiments for deriving pK_a values directly at the side chain carboxyl carbon.

MATERIALS AND METHODS

Preparation of Ubiquitin and Crystallization

Uniformly labeled $^{13}\text{C}/^{15}\text{N}$ ubiquitin protein was expressed and purified as previously published^{21–23}. In brief, ubiquitin was expressed in BL21(DE3) *E. coli* bacteria in the presence of uniformly labeled $^{13}\text{C}_6$ glucose and ^{15}N ammonium chloride in minimal media (M9). Purification involved resuspending bacterial cells in 25 mM Tris buffer, pH 7.5 and 0.04 mg/mL lysozyme. Following sonication for 30 mins, the lysate was centrifuged for 20 min at a speed of 48,000 x g at 4 °C. The supernatant was dialyzed against 50 mM sodium acetate buffer (pH 4.1) overnight at 4 °C and heated at 70 °C for 10 mins. Precipitated protein was removed by centrifugation for 40 min at a speed of 70,000 x g at 4 °C. The supernatant was concentrated and purified using size exclusion chromatography with a Superdex 75 column (GE Healthcare). For preparation of the solid-state NMR samples, 25 mg/ml ubiquitin (10 mg total) in 20 mM sodium citrate at pH 4.3 was crystallized by dropwise addition of 2-methyl-2,4-pentanediol (MPD) to a final concentration of 60%. The sample was incubated overnight at 4 °C. The microcrystalline samples were packed into a 3.2 mm MAS rotor with sample spacers to prevent dehydration.

Preparation of EmrE Samples

Protein expression and purification of the E14Q mutant of EmrE (EmrE^{E14Q}) has been described previously^{3,24–26}. In brief, uniformly labeled $^{13}\text{C}/^{15}\text{N}$ EmrE^{E14Q} protein was obtained by expression of BL21(DE3) *E. coli* bacteria in the presence of uniformly labeled $^{13}\text{C}_6$ glucose and ^{15}N ammonium chloride in minimal media (M9). EmrE^{E14Q} was expressed as a fusion construct with maltose binding protein and purified using amylose affinity chromatography and size exclusion chromatography in n-dodecyl- β -D-maltopyranoside (DDM, Anatrace). Purified EmrE^{E14Q} was reconstituted in 1,2-di-*O*-

tetradecyl-sn-glycero-3-phosphocholine (*O*-14:0-PC) (Avanti Polar Lipids) by removing DDM detergent using Bio-Beads SM-2 resin (Bio-Rad). Proteoliposomes were pelleted by ultra-centrifugation for 12 hours at 436,000 x g using a TLA-100 rotor (Beckman-Coulter) and packed into a 3.2 mm MAS rotor using sample spacers to prevent dehydration. Proteoliposomes were in 150 mM sodium phosphate and 20 mM sodium chloride and were buffer exchanged to give a range of pH values from 1.7 to 11.0.

Solid State NMR Spectroscopy

All NMR experiments were carried out using an Agilent DD2 NMR spectrometer operating at a ^1H frequency of 600 MHz (14.1 T) using a 3.2 mm triple resonance MAS probe manufactured by Black Fox, LLC. The sample temperature was set to 5 °C and -5 °C for ubiquitin and EmrE^{E14Q}, respectively. The MAS rate was 8333 ± 5 Hz. The ^1H - ^{13}C cross polarization time was set to 0.5 msec (ubiquitin) or 1.45 msec (EmrE^{E14Q}) using a tangent ramp²⁷ on ^1H with field strengths of $\omega/2\pi = \sim 66.7$ kHz (^1H , middle of pulse) and $\omega/2\pi = 50$ kHz (^{13}C). For REDOR dephasing²⁸, ^{15}N inversion was achieved by composite 90° - 180° - 90° pulses^{29,30,31} using a 5 μs 90° pulse. For double quantum experiments, an SPC-5 pulse train of 0.48 msec was utilized for each of the conversion and reconversion steps. During the SPC-5 element²⁰, continuous wave ^1H decoupling was applied at a strength of $\omega/2\pi = 100$ kHz. All other ^1H decoupling periods utilized SPINAL-64³² at $\omega/2\pi = 100$ kHz. The indirect dimension spectral width was 8.333 kHz for the double quantum indirect dimension and 100 kHz for the direct ^{13}C dimension. The ^{13}C offset was 99.81 ppm and the ^{15}N offset was 120.03 ppm. Note that the relevant aspartate and glutamate peaks in the double quantum dimension were aliased and subsequently corrected using the circular shift function in NMRPipe³³. Spectral referencing in the ^{13}C dimension was carried out by setting the CH_2 resonance of adamantane to 40.48 ppm³⁴.

RESULTS AND DISCUSSION

Statistical Distribution of Anionic Amino Acid Side Chains

Single quantum 2D spectroscopy such as $^{13}\text{C}/^{13}\text{C}$ correlations can be used to distinguish aspartate and glutamate side chain carboxyl groups in the $\beta 1$ immunoglobulin binding domain of protein G³⁵. We wondered whether an alternative approach to single quantum correlations, through the use of double quantum spectroscopy, would provide improved resolution to probe functionally important anionic residues. The use of double quantum spectroscopy typically involves evolution of double quantum coherences in indirect dimensions, which are correlated with single quantum coherences in the direct dimension (i.e., DQSQ experiment). Since the double quantum shifts appear at the additive chemical shift frequencies of the two nuclei in the double quantum coherence, we envisioned that the indirect dimension of the DQSQ experiment would nicely resolve side chain carboxyl groups of aspartate and glutamate residues from backbone carbonyl sites that often share overlapping ^{13}C resonances. To test this theory, we analyzed statistical distributions of chemical shifts^{36,37} and plotted these in the form of a DQSQ spectrum (Figure 1). This plot shows that the DQSQ experiment provides frequency isolation for the side chain amino acids of aspartate, glutamate, asparagine, and glutamine relative to the backbone carbonyl sites. Namely, the indirect double quantum dimension evolves under the additive chemical

shifts of $\delta_{\text{C}\beta} + \delta_{\text{C}\gamma}$ for aspartate and asparagine and $\delta_{\text{C}\gamma} + \delta_{\text{C}\delta}$ for glutamate and glutamine, which separates these resonances from the backbone sites corresponding to $\delta_{\text{C}\gamma} + \delta_{\text{C}\alpha}$ (Figure 1). More significantly, the evolution of the additive chemical shifts in the indirect dimension increases the effective resolution between protonated and deprotonated side chain chemical shifts for aspartate and glutamate. Specifically, the average chemical shift differences between charged and uncharged sites in the two dimensions of the DQSQ experiment are 6.2 and 3.2 ppm for aspartate and 7.6 and 4.1 ppm for glutamate. This can be compared to chemical shift differences of 3.0 and 3.2 ppm for aspartate and 3.5 and 4.1 ppm for glutamate in a single quantum based 2D experiment. From this analysis, DQSQ spectroscopy is anticipated to improve the spectral dispersion of aspartate and glutamate side chains to provide insight into protonation states of proteins during a pH titration to determine $\text{p}K_a$ values.

Combination of REDOR Filter with DQSQ Spectroscopy to Distinguish Aspartate and Glutamate from Asparagine and Glutamine

While the 2D DQSQ experiment is expected to provide enhanced resolution for observing side chain carboxyl sites between protonated and deprotonated states, the statistical distribution of chemical shifts in Figure 1 does not inherently distinguish aspartate and glutamate from asparagine and glutamine side chains for all protonation states. However, these residue types differ in bonding at the terminal carbon: $-\text{NH}_2$ in asparagine and glutamine or $-\text{OH}$ in aspartic acid and glutamic acid (Figure 1). Thus, the application of ^{15}N REDOR pulses can be used to distinguish asparagine and glutamine from aspartic acid and glutamic acid through a loss in signal intensity. A suitable pulse sequence that combines DQSQ and REDOR is shown in Figure 2. This sequence, referred to as DQSQ-REDOR, utilizes SPC-5²⁰ for double quantum conversion and reconversion back to the single quantum coherence for detection followed by REDOR dephasing²⁸ on the ^{15}N channel. The only requisite for the DQSQ-REDOR experiments is a uniformly $^{13}\text{C}/^{15}\text{N}$ labeled protein, which is the same sample as those used to obtain resonance assignments. Therefore, this experiment has the potential for enhancing resolution for aspartate and glutamate residues while selectively reducing overlapping resonances with the goal of studying biologically important mechanisms. Note that filtering approaches have been previously used to reduce spectral complexity both at backbone^{38,39} and side chain sites⁹. It should also be noted that the REDOR period was also placed before the DQSQ element; however, this produced poorer dephasing relative to DQSQ-REDOR.

Application of DQSQ-REDOR to Microcrystalline Ubiquitin

To test the effectiveness of DQSQ and DQSQ-REDOR to resolve side chain anionic residues, we applied these experiments to uniformly $^{13}\text{C}/^{15}\text{N}$ microcrystalline ubiquitin as a model protein where the primary sequence contained several asparagine, aspartate, glutamine, and glutamate residues (Figure 3A). The DQSQ spectrum of ubiquitin displayed several isolated peaks corresponding to these residue side chains that were predicted from the statistical distribution of chemical shifts (Figure 3B). Namely, residues Glu18, Asn25, Gln31, Asp32, Asp39, Gln41, Asp58, and Gln62 in ubiquitin were identified from previously reported assignments^{21,40}. Note that five other glutamate residues in the primary

sequence were not observed at 5 °C and with the employed DQ mixing time, which is consistent with a prior study⁴⁰ and likely stems from the disordered nature of these side chains. For the residues that were observed, the DQSQ spectrum showed dispersed side chains of asparagine, aspartate, glutamine, and glutamate and the glycine backbone from other backbone carbonyls within ubiquitin. Based on our experimental result and the statistical distribution of chemical shifts in Figure 1, we anticipate the DQSQ experiment will be a valuable method in mechanistic studies for probing side chains of essential asparagine, aspartate, glutamine, and glutamate residues.

Next, we explored whether the DQSQ-REDOR pulse sequence (Figure 2) would be a suitable technique to distinguish side chain carboxyls within aspartate and glutamate from side chain carbonyls within asparagine and glutamine residues. Since one-bond distances for $C\gamma-N^{\delta 2}$ (asparagine) and $C^{\delta}-N^{\epsilon 2}$ (glutamine) are 1.3 Å, a REDOR dephasing period of ~1.2 msec would lead to efficient dephasing for an ordered site. Using this dephasing time, we collected two DQSQ-REDOR datasets corresponding to experiments with and without ¹⁵N REDOR 180° pulses (Figure 3C, top and bottom). These spectra showed significant reduction of backbone signal intensity when REDOR pulses were applied (S) compared to when REDOR pulses were turned off (S_0). Quantification of these spectra showed that most asparagine and glutamine side chains displayed intensity ratios (S/S_0) less than ~0.25 (Figure 3D), which validated the method. However, the S/S_0 values were not effectively dephased for all non-carboxyl sites at a dephasing period of 1.2 msec (e.g., Gln62), which could stem from mobility of the side chain leading to a reduced effective dipolar coupling or imperfect REDOR dephasing. To identify mobile sites, we also employed longer dephasing time periods of 2.2, 3.8 and 4.6 msec. These data showed that progressively longer REDOR dephasing times reduced S/S_0 for Gln62 (Figure 3C, 3D), while maintaining S/S_0 values for aspartate and glutamate side chains greater than ~0.8. The observation of Gln62 as a mobile side chain is potentially required for its function in polyubiquitination⁴¹. We also observed that several side chain asparagine and glutamine residues (e.g., Asn25, Gln31, Gln41) maintained S/S_0 values less than ~0.25, which means these residues were efficiently dephased at the shortest and longest dephasing times. Thus, for asparagine and glutamine sites that are well ordered, the DQSQ-REDOR method can be used to distinguish anionic residues, which increases the signal to noise relative to experiments with longer dephasing times. Notably, a previous method that combined REDOR dephasing and DARR spectroscopy to observe side chain aspartate and glutamate residues required a dephasing period of 3.8 msec to completely dephase asparagine and glutamine side chains⁹. Thus, shorter REDOR dephasing times available with DQSQ-REDOR will achieve higher signal to noise ratios for confidently identifying aspartate and glutamate residues. However, we emphasize that highly mobile residues that are not detected in cross-polarization based techniques also cannot be studied using the DQSQ-REDOR method.

Lastly, we compared our DQSQ-REDOR method with the 2D ¹³C/¹³C DARR experiment⁴² for ubiquitin (Figure 4). The DARR spectrum was obtained by using a 20 msec mixing time while the DQSQ-REDOR spectrum was carried out with 1.2 msec of dephasing (same spectrum as in Figure 3C). As expected, the sensitivity of DARR was higher than that of the DQSQ-REDOR method since DARR does not employ double quantum or REDOR filters. The two resolvable signals in the DARR spectrum showed 1.4-fold and 2.9-fold better

sensitivity for Asp32 and Glu18, respectively, for the same total time to collect each dataset. However, as seen in Figure 4, the DARR spectrum was quite congested and most of the aspartate residues were not clearly resolvable. Therefore, while a DARR spectrum provided for better sensitivity, the major advantage of DQSQ-REDOR is the reduced spectral overlap that enabled identification of aspartate and glutamate residues in a straightforward manner.

Application of DQSQ Method to a Membrane Protein in Lipid Bilayers

pH titrations are commonly employed to derive pK_a values to discern roles of carboxyl residues in biological function. To demonstrate the value of DQSQ-REDOR for determining a pK_a value using MAS solid-state NMR, we applied these experiments to the E14Q mutant of the membrane protein EmrE (EmrE^{E14Q}) in lipid bilayers. This mutant was selected to serve as a simpler model system than wild-type EmrE due to the role of pH dependent changes in structure and dynamics stemming from acid/base chemistry at Glu14^{3,4,25}. Similar to the ubiquitin sample, EmrE^{E14Q} contains asparagine and glutamine (Asn2, Gln14, Gln81, Asn102) and aspartate and glutamate residues (Glu25 and Asp84) that serve to test the robustness of the method for distinguishing these residue types within a membrane protein (Figure 5A).

Uniformly ¹³C/¹⁵N EmrE^{E14Q} was reconstituted in ether-14:0-PC (*O*-14:0-PC) to ensure lipid stability during the pH titration^{24,43}. Proteoliposomes were initially prepared at a pH value of 5.0. DQSQ-REDOR experiments were collected using dephasing times of 1.2, 2.2 and 4.0 msec in the presence and absence of ¹⁵N REDOR pulses. At a dephasing time of 1.2 msec with REDOR pulses on, several peaks including the side chains of glutamine and the backbone of glycine residues were found to be significantly attenuated in intensity (Figure 5B). In fact, many of the asparagine and glutamine signals gave very low signal intensities relative to a reference DQSQ spectrum (Figure 5C). The relative intensities remained low in intensity with longer dephasing times of 2.2 and 4.0 msec, which is observed in the quantification of intensity ratios (S/S_0) (Figure 5D). To the contrary, S/S_0 values for Glu25 and Asp84 remained greater than ~0.75 at all mixing times (Figure 5D). These results indicate effective dephasing at a short time period of 1.2 msec and demonstrates the effectiveness of the method for distinguishing side chain carboxyl sites of aspartate and glutamate from side chain carbonyl sites of asparagine and glutamine. Lastly, note that detection of aspartate and glutamate signals using DQSQ-REDOR with a 1.2 msec dephasing time retained excellent sensitivity relative to DQSQ, as evidenced by the intensity ratios of ~0.8 for Glu25 and Asp84 when comparing signal intensities of DQSQ-REDOR and the DQSQ experiment (S/S_{DQSQ}) (Figure 5E).

We next performed a pH titration on EmrE^{E14Q} as proof of principle of the method to detect pH dependent chemical shift changes. For these measurements, we used DQSQ-REDOR at a dephasing time of 2.2 msec for each pH value within the range 1.7 to 11.0 (Figure 6). However, we note that it is possible to perform the titration experiments using only DQSQ once an aspartate or glutamate side chain of interest has been identified and assigned using DQSQ-REDOR and triple resonance experiments, respectively. The pH titration showed that Glu25 experienced chemical shift perturbations in the fast exchange regime such that the peak moved by ~4.5 ppm in the direct dimension and ~8.5 ppm in the indirect dimension.

No other peaks in the spectrum had similarly large chemical shift perturbations over this pH range. The observed pH dependent chemical shifts of Glu25 were in agreement with previously reported chemical shifts of model tripeptides containing a glutamate residue where the deprotonated chemical shifts are downfield ($C\delta = 183.8$ ppm; $C\gamma = 36.1$) relative to the protonated side chain ($C\delta = 179.7$ ppm; $C\gamma = 32.7$)³⁶. Using the chemical shift changes in both the single quantum and double quantum dimensions, we fitted an apparent pK_a of Glu25 to be 4.4 ± 0.3 using a modified Henderson-Hasselbalch equation³. Note that we did not observe changes in chemical shifts for Asp84 over the pH range of 3.0 to 11.0, implying this residue has a pK_a value below ~ 3.0 and would be deprotonated under all physiologically relevant pH conditions.

In summary, DQSQ offers excellent separation of the side chains of asparagine, aspartate, glutamine, and glutamate residues relative to backbone amide sites. Inclusion of a REDOR period into this sequence enabled aspartate and glutamate side chains to be easily distinguished from asparagine and glutamine side chain residues that often overlap within the DQSQ spectrum. The DQSQ-REDOR offers excellent sensitivity since it can be applied with a short REDOR dephasing time of ~ 1.2 msec that is sufficient for identifying anionic side chains through the comparison of signal intensities in the presence and absence of REDOR dephasing pulses. Thus, DQSQ-REDOR is anticipated to be a valuable method for characterizing catalytically important anionic residues with MAS solid-state NMR spectroscopy.

CONCLUSION

We proposed and validated a DQSQ-based pulse sequence for distinguishing asparagine, aspartate, glutamine, and glutamate side chains in proteins. Inclusion of a REDOR dephasing period into this sequence (i.e., DQSQ-REDOR) was effective at distinguishing aspartate and glutamate side chains from asparagine and glutamine. The DQSQ-REDOR method has an intrinsic resolution advantage due to the double quantum chemical shift indirect dimension that better resolves protonated side chains of aspartate and glutamate from their deprotonated states. Since ^{13}C chemical shifts of carboxyl sites are sensitive to the protonation state, this method can be employed in microcrystalline proteins and membrane proteins to infer charged states of anionic side chains as well as in pH titrations to quantify pK_a values.

ACKNOWLEDGEMENT

The NMR methodology was supported by NSF grant MCB 1902449 (to N.J.T.). Application to EmrE was funded by NIH grant R01 AI108889 (to N.J.T.). A.S.H. was supported from a Dean's Dissertation Fellowship from New York University.

REFERENCES:

1. Tollinger M, Crowhurst KA, Kay LE & Forman-Kay JD Site-specific contributions to the pH dependence of protein stability. *Proc Natl Acad Sci U S A* 100, 4545–50 (2003). [PubMed: 12671071]

2. Wells JA, Powers DB, Bott RR, Graycar TP & Estell DA Designing substrate specificity by protein engineering of electrostatic interactions. *Proc Natl Acad Sci U S A* 84, 1219–23 (1987). [PubMed: 3547407]
3. Gayen A, Leninger M & Traaseth NJ Protonation of a glutamate residue modulates the dynamics of the drug transporter EmrE. *Nat Chem Biol* 12, 141–5 (2016). [PubMed: 26751516]
4. Morrison EA, Robinson AE, Liu Y & Henzler-Wildman KA Asymmetric protonation of EmrE. *J Gen Physiol* 146, 445–61 (2015). [PubMed: 26573622]
5. Masureel M et al. Protonation drives the conformational switch in the multidrug transporter LmrP. *Nat Chem Biol* 10, 149–55 (2014). [PubMed: 24316739]
6. Fluman N, Ryan CM, Whitelegge JP & Bibi E Dissection of mechanistic principles of a secondary multidrug efflux protein. *Mol Cell* 47, 777–87 (2012). [PubMed: 22841484]
7. Fluman N & Bibi E Bacterial multidrug transport through the lens of the major facilitator superfamily. *Biochim Biophys Acta* 1794, 738–47 (2009). [PubMed: 19103310]
8. Jehle S et al. Spectral editing: selection of methyl groups in multidimensional solid-state magic-angle spinning NMR. *J Biomol NMR* 36, 169–77 (2006). [PubMed: 17031530]
9. Schmidt-Rohr K, Fritzsching KJ, Liao SY & Hong M Spectral editing of two-dimensional magic-angle-spinning solid-state NMR spectra for protein resonance assignment and structure determination. *J Biomol NMR* 54, 343–53 (2012). [PubMed: 23053913]
10. Reggie L, Lopez JJ, Collinson I, Glaubitz C & Lorch M Dynamic nuclear polarization-enhanced solid-state NMR of a ¹³C-labeled signal peptide bound to lipid-reconstituted Sec translocon. *J Am Chem Soc* 133, 19084–6 (2011). [PubMed: 22040139]
11. Nielsen JE Chapter 5 - Analyzing Protein NMR pH-Titration Curves in *Annual Reports in Computational Chemistry*, Vol. 4 (eds. Wheeler RA & Spellmeyer DC) 89–106 (Elsevier, 2008).
12. Oregioni A, Stieglitz B, Kelly G, Rittinger K & Frenkiel T Determination of the pKa of the N-terminal amino group of ubiquitin by NMR. *Sci Rep* 7, 43748 (2017). [PubMed: 28252051]
13. Pielak RM & Chou JJ Influenza M2 proton channels. *Biochimica et Biophysica Acta (BBA) - Biomembranes* 1808, 522–529 (2011). [PubMed: 20451491]
14. Hu J et al. Histidines, heart of the hydrogen ion channel from influenza A virus: toward an understanding of conductance and proton selectivity. *Proc Natl Acad Sci U S A* 103, 6865–70 (2006). [PubMed: 16632600]
15. Hu F, Schmidt-Rohr K & Hong M NMR detection of pH-dependent histidine-water proton exchange reveals the conduction mechanism of a transmembrane proton channel. *J Am Chem Soc* 134, 3703–13 (2012). [PubMed: 21974716]
16. Bartik K, Redfield C & Dobson CM Measurement of the individual pKa values of acidic residues of hen and turkey lysozymes by two-dimensional ¹H NMR. *Biophys J* 66, 1180–4 (1994). [PubMed: 8038389]
17. McIntosh LP et al. The pKa of the general acid/base carboxyl group of a glycosidase cycles during catalysis: a ¹³C-NMR study of bacillus circulans xylanase. *Biochemistry* 35, 9958–66 (1996). [PubMed: 8756457]
18. Oda Y et al. Individual ionization constants of all the carboxyl groups in ribonuclease HI from *Escherichia coli* determined by NMR. *Biochemistry* 33, 5275–84 (1994). [PubMed: 7909691]
19. Wang YX et al. Solution NMR evidence that the HIV-1 protease catalytic aspartyl groups have different ionization states in the complex formed with the asymmetric drug KNI-272. *Biochemistry* 35, 9945–50 (1996). [PubMed: 8756455]
20. Hohwy M, Rienstra CM, Jaroniec CP & Griffin RG Fivefold symmetric homonuclear dipolar recoupling in rotating solids: Application to double quantum spectroscopy. *The Journal of Chemical Physics* 110, 7983–7992 (1999).
21. Zech SG, Wand AJ & McDermott AE Protein structure determination by high-resolution solid-state NMR spectroscopy: application to microcrystalline ubiquitin. *J Am Chem Soc* 127, 8618–26 (2005). [PubMed: 15954766]
22. Banigan JR & Traaseth NJ Utilizing afterglow magnetization from cross-polarization magic-angle-spinning solid-state NMR spectroscopy to obtain simultaneous heteronuclear multidimensional spectra. *J Phys Chem B* 116, 7138–44 (2012). [PubMed: 22582831]

23. Lazar GA, Desjarlais JR & Handel TM De novo design of the hydrophobic core of ubiquitin. *Protein Sci* 6, 1167–78 (1997). [PubMed: 9194177]
24. Banigan JR, Leninger M, Her AS & Traaseth NJ Assessing Interactions Between a Polytopic Membrane Protein and Lipid Bilayers Using Differential Scanning Calorimetry and Solid-State NMR. *J Phys Chem B* 122, 2314–2322 (2018). [PubMed: 29457729]
25. Cho MK, Gayen A, Banigan JR, Leninger M & Traaseth NJ Intrinsic conformational plasticity of native EmrE provides a pathway for multidrug resistance. *J Am Chem Soc* 136, 8072–80 (2014). [PubMed: 24856154]
26. Gayen A, Banigan JR & Traaseth NJ Ligand-induced conformational changes of the multidrug resistance transporter EmrE probed by oriented solid-state NMR spectroscopy. *Angew Chem Int Ed Engl* 52, 10321–4 (2013). [PubMed: 23939862]
27. Baldus MG, D. G.; Hediger, S.; Meier, B. H. Efficient 15N- 13C Polarization Transfer by Adiabatic-Passage Hartmann-Hahn Cross Polarization. *Journal of Magnetic Resonance* 118, 140–144 (1996).
28. Gullion T & Schaefer J Rotational-echo double-resonance NMR. 1989. *J Magn Reson* 213, 413–7 (2011). [PubMed: 22152360]
29. Levitt MH Short perspective on “NMR population inversion using a composite pulse” by Levitt MH and Freeman R [J. Magn. Reson. 33 (1979) 473–476]. *J Magn Reson* 213, 274–5 (2011). [PubMed: 21903438]
30. Levitt MH & Freeman R NMR population inversion using a composite pulse. *Journal of Magnetic Resonance* (1969) 33, 473–476 (1979).
31. Sinha N, Schmidt-Rohr K & Hong M Compensation for pulse imperfections in rotational-echo double-resonance NMR by composite pulses and EXORCYCLE. *J Magn Reson* 168, 358–65 (2004). [PubMed: 15140448]
32. Fung BM, Khitrin AK & Ermolaev K An improved broadband decoupling sequence for liquid crystals and solids. *J Magn Reson* 142, 97–101 (2000). [PubMed: 10617439]
33. Delaglio F et al. NMRPipe: a multidimensional spectral processing system based on UNIX pipes. *J Biomol NMR* 6, 277–93 (1995). [PubMed: 8520220]
34. Morcombe CR & Zilm KW Chemical shift referencing in MAS solid state NMR. *J Magn Reson* 162, 479–86 (2003). [PubMed: 12810033]
35. Schmidt HL, Shah GJ, Sperling LJ & Rienstra CM NMR Determination of Protein pK(a) Values in the Solid State. *J Phys Chem Lett* 1, 1623–1628 (2010). [PubMed: 20563223]
36. Platzer G, Okon M & McIntosh LP pH-dependent random coil (1)H, (13)C, and (15)N chemical shifts of the ionizable amino acids: a guide for protein pK a measurements. *J Biomol NMR* 60, 109–29 (2014). [PubMed: 25239571]
37. Ulrich EL et al. BioMagResBank. *Nucleic Acids Res* 36, D402–8 (2008). [PubMed: 17984079]
38. Banigan JR, Gayen A & Traaseth NJ Combination of (1)(5)N reverse labeling and afterglow spectroscopy for assigning membrane protein spectra by magic-angle-spinning solid-state NMR: application to the multidrug resistance protein EmrE. *J Biomol NMR* 55, 391–9 (2013). [PubMed: 23539118]
39. Traaseth NJ & Veglia G Frequency-selective heteronuclear dephasing and selective carbonyl labeling to deconvolute crowded spectra of membrane proteins by magic angle spinning NMR. *J Magn Reson* 211, 18–24 (2011). [PubMed: 21482162]
40. Igumenova TI et al. Assignments of carbon NMR resonances for microcrystalline ubiquitin. *J Am Chem Soc* 126, 6720–7 (2004). [PubMed: 15161300]
41. Suryadinata R et al. Molecular and structural insight into lysine selection on substrate and ubiquitin lysine 48 by the ubiquitin-conjugating enzyme Cdc34. *Cell Cycle* 12, 1732–44 (2013). [PubMed: 23656784]
42. Takegoshi K, Nakamura S & Terao T 13C–1H dipolar-assisted rotational resonance in magic-angle spinning NMR. *Chemical Physics Letters* 344, 631–637 (2001).
43. Leninger M, Sae Her A & Traaseth NJ Inducing conformational preference of the membrane protein transporter EmrE through conservative mutations. *Elife* 8(2019).





Figure 1.

Projection of chemical shifts onto a DQSQ spectrum using the chemical shift statistics from the BMRB database³⁷ and Platzer *et al.*³⁶. Errors reflect the standard deviation from the chemical shift assignments of these residues. Chemical shifts of aspartate and glutamate side chains were taken from Platzer *et al.*³⁶ because both protonated and deprotonated chemical shifts were reported. The structures of the side chains of aspartate, asparagine, glutamate, and glutamine are color coded with the DQSQ plot.

**Figure 2.**

DQSQ-REDOR pulse sequence. All ^{15}N REDOR dephasing pulses were applied as composition pulses ($90^\circ-180^\circ-90^\circ$) with the indicated phases. A 4-step phase cycle was implemented as follows: $\phi_1 = [x, y, -x, -y]$, $\phi_2 = [y, -x, -y, x]$, $\phi_3 = [y, y, -y, -y]$, $\phi_4 = [x, x, -x, -x]$, and $\phi_{\text{receiver}} = [x, -y, -x, y]$. τ_r is the rotor period (120 μsec) and $t_{z\text{filter}}$ is the z-filter time (120 μsec).




Figure 3.

DQSQ and DQSQ-REDOR applied to microcrystalline ubiquitin. (A) Primary sequence of ubiquitin with aspartate, asparagine, glutamate, and glutamine are color coded with residue types as in Figure 1. (B) DQSQ and (C) DQSQ-REDOR with dephasing periods of 1.2 msec and 2.2 msec. Top spectra in panel C were collected with application of ^{15}N dephasing pulses, while the bottom spectra were collected with the same time period without ^{15}N REDOR pulses. All peak labels are color coded with residue types as in Figure 1. (D) Intensity ratios at different dephasing times of select side chains (Asn, Asp, Gln, Glu) and backbone peaks (Gly) in the presence of ^{15}N pulses (S) divided by those in the absence of ^{15}N pulses (S_0).




Figure 4. Comparison of (A) DQSQ-REDOR with 1.2 msec ^{15}N dephasing and (B) DARR $^{13}\text{C}/^{13}\text{C}$ correlation with 20 msec mixing time on a microcrystalline sample of uniformly ^{13}C , ^{15}N ubiquitin. Note that the total time to collect the spectra were the same.




Figure 5. (A) Primary sequence of EmrE^{E14Q} with aspartate, asparagine, glutamate, and glutamine are color coded with residue types as in Figure 1. (B) DQSQ-REDOR and (C) DQSQ spectra of uniformly labeled ¹³C/¹⁵N EmrE^{E14Q} in *O*-14:0-PC liposomes at a pH value of 5.0. The REDOR dephasing times in panel B are indicated on top of each spectrum. (D) Intensity ratios (S/S_0) of select side chains (Asn, Asp, Gln, Glu) and backbone peaks (Gly) in the presence of ¹⁵N pulses (S) divided by those in the absence of ¹⁵N pulses (S_0). (E) Intensity ratios (S/S_{DQSQ}) of select side chains (Asn, Asp, Gln, Glu) and backbone peaks (Gly) in the presence of ¹⁵N REDOR pulses (S) divided by those in the DQSQ spectrum (S_{DQSQ}).




Figure 6.

Spectra and subsequent pK_a fitting of pH-dependent chemical shift changes for Glu25 within EmrE^{E14Q}. Left: DQSQ spectra with 2.2 msec REDOR dephasing for Glu25 over the indicated pH values. Center and right: experimental data points of single quantum chemical shifts in the direct dimension (middle) and double quantum chemical shifts in the indirect dimension (right). The continuous line is the best global fit using a modified Henderson-Hasselbalch equation³ that yielded an apparent pK_a value of 4.4 ± 0.3 for Glu25.