



Cite this: *Med. Chem. Commun.*,  
2019, 10, 1958

## A focused structure–activity relationship study of psoralen-based immunoproteasome inhibitors†

Eva Shannon Schiffrer,<sup>a</sup> Izidor Sosič,<sup>id</sup><sup>a</sup> Andrej Šterman,<sup>id</sup><sup>a</sup> Janez Mravljak,<sup>id</sup><sup>a</sup>  
Irena Mlinarič Raščan,<sup>b</sup> Stanislav Gobec<sup>a</sup> and Martina Gobec<sup>id</sup><sup>\*b</sup>

The immunoproteasome is a multicatalytic protease that is predominantly expressed in cells of hematopoietic origin. Its elevated expression has been associated with autoimmune diseases, various types of cancer, and inflammatory diseases. The development of immunoproteasome-selective inhibitors with non-peptidic scaffolds remains a challenging task. Here, we describe a focused series of psoralen-based inhibitors of the  $\beta 5i$  subunit of the immunoproteasome with different substituents placed at position 4'. The most promising compound was further evaluated through changes at position 3 of the psoralen ring. Despite a small decrease in the inhibitory potency in comparison with the parent compound, we were able to improve the selectivity against other subunits of both the immunoproteasome and the constitutive proteasome. The most potent compounds discriminated between both proteasome types in cell lysates and also showed a decrease in cytokine secretion in peripheral blood mononuclear cells.

Received 17th July 2019,  
Accepted 8th September 2019

DOI: 10.1039/c9md00365g

rsc.li/medchemcomm

### Introduction

The heart of the ubiquitin–proteasome system, the 26S proteasome, is responsible for the regulation of various cell processes and the maintenance of protein homeostasis in cells. It is the most important non-lysosomal proteolytic complex degrading misfolded or damaged proteins.<sup>1</sup> The 26S proteasome consists of a 20S core and 19S regulatory sections, with the 20S core comprising three active subunits with distinct catalytic activities, *i.e.*  $\beta 1$  (caspase-like),  $\beta 2$  (trypsin-like), and  $\beta 5$  (chymotrypsin-like).<sup>2,3</sup> Two major types of 20S cores exist in vertebrates: the constitutive proteasome (cCP), expressed in all cell types, and the immunoproteasome (iCP), expressed mostly in hematopoietic cells. The latter can also be found in other cell types upon induction by the tumour necrosis factor- $\alpha$  and interferon- $\gamma$  during acute immune and inflammatory responses.<sup>4,5</sup> Of note, in the iCP, the catalytically active subunits  $\beta$  are replaced by their  $\beta 1i$ ,  $\beta 2i$ , and  $\beta 5i$  counterparts.

Extensive research in the proteasome field in the last two decades resulted in the development of non-selective proteasome inhibitors, such as bortezomib and carfilzomib, used in the treatment of multiple myeloma and mantle-cell


lymphoma.<sup>6</sup> However, the clinical use of such conventional, non-selective proteasome inhibitors is often limited due to the side effects caused by the non-selective inhibition of protein degradation.<sup>7</sup> While the cCP is expressed in all eukaryotic cells, the iCP expression is mostly upregulated during the course of disease processes,<sup>8–10</sup> leading to the assumption that its selective inhibition is expected to cause fewer side effects. The latest studies showed that co-inhibition of two subunits of the iCP is required to show beneficial effects in autoimmune diseases, *i.e.* either by combining  $\beta 1i$ - and  $\beta 5i$ -selective inhibitors in a synergistic approach<sup>11</sup> or with a single molecule that is able to simultaneously inhibit two catalytically active subunits of the iCP.<sup>12</sup> The excellent work by the groups of Overkleeft,<sup>13,14</sup> Groll,<sup>15</sup> Lin,<sup>16,17</sup> and Johnson<sup>12,18</sup> yielded selective inhibitors of all subunits of the iCP (Fig. 1) and these represent both very valuable molecular tools in the studies of iCP biology and lead compounds towards new drugs (Fig. 1). It is noteworthy that KZR-616, a  $\beta 1i$ - and  $\beta 5i$ -targeting compound, is currently being evaluated in clinical studies for the treatment of autoimmune and inflammatory diseases.<sup>12,19</sup>

From a structural perspective, the majority of these inhibitors of the iCP have a peptidic backbone, and are thus prone to poor metabolic stability and low bioavailability leading to lack of oral exposure.<sup>20,21</sup> With an attempt to circumvent these limitations, several lines of research by various groups are devoted to the development of non-peptidic inhibitors of the iCP (Fig. 2).<sup>22–26</sup> Despite this, the number of such inhibitors is scarce and, interestingly, only inhibitors of the  $\beta 5i$  subunit were discovered so far.

<sup>a</sup> Faculty of Pharmacy, Chair of Pharmaceutical Chemistry, University of Ljubljana, Aškerčeva 7, 1000 Ljubljana, Slovenia

<sup>b</sup> Faculty of Pharmacy, Chair of Clinical Biochemistry, University of Ljubljana, Aškerčeva 7, 1000 Ljubljana, Slovenia. E-mail: martina.gobec@ffa.uni-lj.si; Tel: +386 1 476 9636

† Electronic supplementary information (ESI) available: General chemistry methods, all biochemistry experimental data, spectroscopic analyses, and representative NMR, HPLC, and HRMS spectra. See DOI: 10.1039/c9md00365g




**Fig. 1** Examples of peptidic subunit-selective inhibitors of iCP; original labelling of compounds by the authors is used here. For more details, the corresponding references are given with each inhibitor.

Previously, we reported a series of psoralens as selective inhibitors of the  $\beta$ 5i subunit.<sup>22</sup> These compounds were discovered through a structure-based virtual screening of the ZINC drug-like compound library. A library of psoralen-based compounds was subsequently synthesised for the initial


structure–activity relationship (SAR) studies. The best non-covalent inhibitor was transformed into a covalent inhibitor through the addition of an electrophilic warhead on position 3 of the psoralen ring. The resulting compounds exhibited selectivity to  $\beta$ 5i compared to the  $\beta$ 2i and  $\beta$ 1i subunits, as well as compared to all cCP subunits. The most selective covalent inhibitor, a psoralen with an oxathiazolone as the electrophilic warhead (psoralen-based inhibitor, Fig. 2) and a phenyl group at position 4', exhibited more than 80-fold selectivity to  $\beta$ 5i compared to the other iCP and cCP subunits.<sup>22</sup> Of note, oxathiazolone as a covalent warhead had been used previously in inhibitors of *Mycobacterium tuberculosis* proteasome<sup>27</sup> and oxathiazolone-based compounds were also described as iCP-selective by the same group.<sup>25</sup>

Herein, we present a focused SAR study of psoralen-based  $\beta$ 5i inhibitors with oxathiazolone at position 3 (Fig. 3). A series of compounds with various substituents at position 4' of the psoralen ring is described and characterised to evaluate the effect of these modifications on the activity,



**Fig. 2** A representative set of non-peptidic selective inhibitors of the iCP; original labelling of compounds by the authors is used here for the bottom three compounds. For more details, the corresponding references are given with each inhibitor.

R = cycloalkyl, substituted phenyl, heteroaromate



**Fig. 3** Schematic representation of the work disclosed in this study. The numbering system for the psoralen ring is shown for clarity.

selectivity, and biochemical characteristics of psoralen-based compounds. The best performing 4'-substituted inhibitor was further modified at position 3 to evaluate the influence of different electrophilic groups at this position.

## Synthesis

The target oxathiazolone-based psoralens were synthesised using a straightforward synthetic route (Scheme 1). The 1-substituted-2-bromo-ethan-1-ones **2–7** were either commercially available or prepared from corresponding ethan-1-ones by bromination with copper(II) bromide (**2**, **3**, and **4**) or bromine (**5**, **6**, and **7**). These were then reacted with ethyl 7-hydroxy-4-methyl-2-oxo-2*H*-chromene-3-carboxylate (**1**), which was prepared under anhydrous conditions from 1-(2,4-dihydroxyphenyl)ethan-1-one and diethyl malonate using a microwave reactor, to yield alkylated compounds **8–17**.

To obtain the psoralen ring (compounds **18–27**), cyclisation under basic conditions was performed. The temperature needed to be maintained at 100 °C and the reaction times were carefully controlled as extensive heating resulted in the formation of undesired side products and low yields. Compounds **18–27** were then treated with Boc<sub>2</sub>O, pyridine and ammonium acetate to yield 4-methyl-3-carboxamide substituted psoralens **28–37**, which were finally reacted with chlorocarbonylsulfonyl chloride under carefully controlled anhydrous conditions. The formation of oxathiazolone-based compounds **38–47** was confirmed with


the disappearance of the amide signals in the <sup>1</sup>H NMR spectra and the observance of oxathiazolone carbonyl signals in the 170–176 ppm range in the <sup>13</sup>C NMR spectra.

The psoralens with the cyclohexyl moiety at position 4' with variation at position 3 were synthesised through a different, but nevertheless, straightforward synthetic route (Scheme 2). Compound **7** was reacted with ethyl 2-(7-hydroxy-4-methyl-2-oxo-2*H*-chromen-3-yl)acetate (**48**) or ethyl 3-(7-hydroxy-4-methyl-2-oxo-2*H*-chromen-3-yl)propanoate (**49**) as described previously,<sup>22</sup> to yield alkylated compounds **50** and **51**. To obtain the psoralen ring (compounds **52** and **53**), cyclisation was performed in the same manner as described in Scheme 1. Variation at position 3 was performed through an EDC-mediated coupling with *N*-hydroxysuccinimide, 2-(methylamino)acetonitrile, or 2-aminoacetonitrile.


## Biochemical evaluation

The synthesized compounds with various substituents at position 4' and oxathiazolone at position 3 (**38–47**) were first evaluated for their inhibition of β5i activity using succinyl-Leu-Leu-Val-Tyr-7-amino-4-methylcoumarin (Suc-LLVY-AMC) as a fluorogenic substrate (Table 1).

All compounds showed sub-micromolar inhibitory activity, with compounds **42**, **44**, and **47** showing the strongest β5i inhibition with IC<sub>50</sub> values of 141 nM, 174 nM, and 106 nM, respectively (Table 1). The position of the methoxy group on













**Scheme 1** Synthesis of psoralen-based compounds with an oxathiazolone warhead. Reagents and conditions: (a) diethyl malonate, KOtBu, molecular sieves (3 Å), MW (100 °C, 50 W, 15 min); (b) EtOAc, CuBr<sub>2</sub>, 85 °C, 24 h (for **2**, **3**, and **4**) or MeOH, Br<sub>2</sub>, 0 °C to rt, 48 h (for **5**, **6**, and **7**); (c) **1**, dioxane, K<sub>2</sub>CO<sub>3</sub>, KI, 100 °C, 24 h; (d) propan-2-ol, 1 M NaOH, 100 °C, 5–20 min; (e) dioxane, Boc<sub>2</sub>O, pyridine, ammonium acetate, rt, 24 h; (f) anhydrous THF, chlorocarbonylsulfonyl chloride, 80 °C, 1.5–2 h.



**Scheme 2** Synthesis of psoralens with the cyclohexyl moiety at position 4' with variation at position 3. Reagents and conditions: (a) 75% H<sub>2</sub>SO<sub>4</sub>, diethyl 2-acetylsuccinate (for **48**) or diethyl 2-acetylglutarate (for **49**), rt, 24 h; (b) MeOH, Br<sub>2</sub>, 0 °C to rt, 48 h; (c) **48** or **49**, dioxane, K<sub>2</sub>CO<sub>3</sub>, KI, 100 °C, 24 h; (d) propan-2-ol, 1 M NaOH, 100 °C, 5–20 min; (e) **52** or **53**, THF, *N*-hydroxysuccinimide, EDC, Et<sub>3</sub>N, 0 °C to rt, 24 h; (f) **52** or **53**, THF, 2-(methylamino)acetonitrile, EDC, Et<sub>3</sub>N, 0 °C to rt, 24 h; (g) **52** or **53**, THF, 2-aminoacetonitrile, EDC, Et<sub>3</sub>N, 0 °C to rt, 24 h.

**Table 1** Structures and *in vitro* inhibitory potencies of compounds 38–47 against the  $\beta 5i$  subunit of human iCP


Compound	R	IC <sub>50</sub> for $\beta 5i$ (nM)
38		227 ± 32
39		226 ± 21
40		351 ± 4
41		200 ± 13
42		141 ± 6
43		365 ± 5
44		174 ± 5
45		433 ± 13
46		264 ± 3
47		106 ± 4

the 4'-phenyl ring did not influence the activity as exemplified with compounds 38 and 39, which showed the same inhibition of  $\beta 5i$ . When methyl was placed at the *meta* position (compound 43) instead of the methoxy group (compound 39), a decrease in the inhibitory activity was observed (Table 1). Interestingly, the compound with bromine at the *para* position (40, IC<sub>50</sub> ( $\beta 5i$ ) = 351 nM) was less active than the corresponding *para* fluoro (41, IC<sub>50</sub> ( $\beta 5i$ ) = 200 nM) and *para* ethyl (44, IC<sub>50</sub> ( $\beta 5i$ ) = 174 nM) derivatives. In a series of cycloalkyl derivatives 45–47, a clear trend of improved  $\beta 5i$  inhibition with increasing size of substituents at position 4' of the psoralen ring was observed (Table 1). Additionally, the time-dependence of  $\beta 5i$  inhibition by compounds 42, 44, and 47 was determined by IC<sub>50</sub> shift assay (Table S3<sup>†</sup>); *i.e.* a clear increase in the IC<sub>50</sub> values was observed when the compounds were not pre-incubated with the enzyme prior to the addition of the substrate. Moreover, the irreversibility of oxathiazolones 42, 44, and 47 was determined by the rapid dilution assay (Table S4<sup>†</sup>). In Fig. 4, representative time course curves obtained in the rapid dilution assays for compound 47 and DPLG-3, one of the most selective and reversible  $\beta 5i$  inhibitors known,<sup>17</sup> are shown. We should state here that DPLG-3 is not commercially available, so we resynthesised it in our lab (see the ESI,<sup>†</sup> Scheme S1).

To assess the selectivity of these psoralen-based inhibitors 38–47, the compounds were evaluated against all remaining subunits of the iCP and the cCP (Table S1,<sup>†</sup> data presented as residual activities in the presence of 10  $\mu$ M of each compound). All the compounds showed preferential inhibition of  $\beta 5i$  over the  $\beta 2i$  and  $\beta 1i$  subunits of the human iCP, as well as over all three subunits of the human cCP. Due to poor solubility of the compounds in higher concentrations that did not allow precise IC<sub>50</sub> determinations for inhibition of other subunits, the degree of selectivity for compounds 38–47 can only be assumed; *e.g.* for compounds 42, 44, and 47 (which were the most potent  $\beta 5i$  inhibitors in the series), the selectivity can be estimated to be more than 100-fold. This represents an improvement in comparison with a previously described oxathiazolone-based psoralen (cpd. 42, Fig. 2, 80-fold selectivity, ref. 22), which was, however, more active with an IC<sub>50</sub> value for  $\beta 5i$  inhibition of 13 nM.

The most promising inhibitor, compound 47, was further modified at position 3 to evaluate the effect of different electrophilic groups at this position on the inhibitory properties of psoralens (compounds 54–59, Table 2). The nitrile-based compounds 56–59 showed no inhibition of  $\beta 5i$ , whereas only one succinimidyl-ester exhibited inhibitory properties, albeit to a moderate extent (55, IC<sub>50</sub> ( $\beta 5i$ ) = 1830 nM). These results show that oxathiazolone-based compounds are more preferable to achieve  $\beta 5i$  inhibition.

Next, we incubated compounds 44 and 47 with cell lysates of HeLa and THP-1 cells to estimate the capability of the compounds to inhibit the iCP in the presence of other cytosolic components. THP-1 cells were used as they are derived from acute monocytic leukemia and express high levels of iCP, whereas HeLa cells predominantly contain cCP.<sup>28</sup> Both compounds showed better inhibition of  $\beta 5i$  activity in THP-1 cell lysates than in HeLa cell lysates (Table S2<sup>†</sup>); although, similarly, as with previously described psoralens,<sup>22</sup> approximately 10-fold reductions in potency (44: IC<sub>50</sub> [THP-1 lysates] = 1.53  $\mu$ M, 47: IC<sub>50</sub> [THP-1 lysates] = 1.01  $\mu$ M) were observed when compared with those of *in vitro* assays with purified human iCP (Table 1). Unfortunately, we were not able to solve issues with poor cellular permeability



**Fig. 4** Recovery of iCP activity in the rapid dilution assay. To a 100-fold concentration of iCP used in the kinetic assay, a 10-fold IC<sub>50</sub> concentration of the selected inhibitor (either 45 nM DPLG-3 or 1060 nM compound 47) was added. The reaction mix was immediately (0 min), or after pre-incubation (30 min), diluted 100-fold with the substrate. For the reversible inhibitor DPLG-3 (A), almost a full recovery of iCP activity was observed, while in the case of compound 47 (B) only a 22% recovery was observed, indicating the irreversible mode of inhibition.







of psoralen-based compounds as worse inhibition of  $\beta 5$  activity was observed for both derivatives in intact cells (Table S2,† only RA values are given due to issues with solubility at higher concentrations of 44 and 47).

Despite poor permeability, we did test the ability of compounds 44 and 47 (both compounds at 10  $\mu\text{M}$  with 24-hour incubation) to inhibit cytokine secretion. Here, we used DPLG-3 as a positive control. In lipopolysaccharide (LPS) stimulated peripheral blood mononuclear cells, we determined that all compounds, to a small extent, affected the secretion of IL-6 and IL-10 (Fig. 5), while other assayed cytokines (IL-1 $\beta$ , IL-8, IL-12p70, and TNF $\alpha$ ) were not affected (Table S5†). The results obtained for DPLG-3 and compounds 44 and 47 are not only in line with permeability issues for compounds 44 and 47, but also corroborate previously determined data<sup>12</sup> that besides  $\beta 5$ i, another iCP subunit must be inhibited to achieve notable cellular effects in terms of decreasing the pro-inflammatory cytokine production.

## Conclusions

Discovery of non-peptidic and selective inhibitors of the iCP is an extremely challenging task, so such incremental steps, as shown in this study, are needed to optimize a given compound class. We showed that small changes in the structures can lead to further improvements in their selectivity for the inhibition of the  $\beta 5$ i subunit of iCP. In addition, we calculated parameters to evaluate pharmacokinetics and drug-likeness with the SwissADME Web tool<sup>29</sup> (ESI,† section 9); these indicate slightly better overall characteristics and predicted water solubility for structures with saturated rings attached to psoralen. However, problems with poor cellular permeability remain

**Table 2** Structures and *in vitro* inhibitory potencies of compounds 54–59 against the  $\beta 5$ i subunit of human iCP

Compound	R	RA (%) at 10 $\mu\text{M}$ or IC <sub>50</sub> for $\beta 5$ i (nM)
54		62 $\pm$ 3
55		IC <sub>50</sub> = 1830 $\pm$ 380
56		101 $\pm$ 7
57		111 $\pm$ 16
58		99 $\pm$ 9
59		102 $\pm$ 9

unsolved; therefore, more pronounced changes in the psoralen structure are needed in the future to tackle this problem. Of note, our results confirmed the recently disclosed data,<sup>12</sup> in which simultaneous inhibition of two iCP subunits is necessary to achieve a significant anti-inflammatory effect. Therefore, two separate issues, *i.e.* permeability and simultaneous inhibition of two iCP subunits, will represent our main focus during the design of next generation non-peptidic iCP inhibitors.

## Experimental

Synthetic procedures and analyses for a representative set of compounds are reported in the main text. General chemistry methods, all biochemistry experimental data, spectroscopic analyses, and representative NMR, HRMS, and HPLC spectra are available in the ESI.†


### Procedure for the synthesis of ethyl 7-hydroxy-4-methyl-2-oxo-2H-chromene-3-carboxylate 1

To a mixture of 1-(2,4-dihydroxyphenyl)ethan-1-one (10 mmol, 1 equiv.), diethyl malonate (2.3 equiv.) and KOtBu (0.22 equiv.), activated molecular sieves (pore diameter of 3 Å) were added. The mixture was reacted in a microwave reactor at 100 °C and 50 W for 15 min. Compound 1 was purified by column chromatography (*n*-hexane/ethyl acetate = 4:1, gradient to 1:1).

Off-white solid; yield 33%; <sup>1</sup>H NMR (400 MHz, acetone-*d*<sub>6</sub>)  $\delta$  1.34 (t, *J* = 7.1 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>), 2.43 (s, 3H, Ar-CH<sub>3</sub>), 4.35 (q, *J* = 7.1 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 6.76 (d, *J* = 2.4 Hz, 1H, Ar-H), 6.91 (dd, *J* = 8.8 Hz, 2.4 Hz, 1H, Ar-H), 7.73 (d, *J* = 8.8 Hz, 1H, Ar-H), 9.68 (br s, 1H, OH). HRMS (*m/z*) (ESI): calcd for C<sub>13</sub>H<sub>12</sub>O<sub>5</sub> [M + H]<sup>+</sup>: 249.0763, found: 249.0765.

### General procedure for the synthesis of 1-substituted-2-bromoethan-1-ones 2–4

The corresponding ethan-1-ones (starting amount ranging from 6.7 to 7.9 mmol, 1 equiv.) were dissolved in EtOAc (50



**Fig. 5** The effect of compounds 44, 47, and DPLG-3 on the secretion of cytokines. The results present values relative to LPS treated cells. The cells were pre-treated for 1 h with DPLG (50 nM), compound 44 (10  $\mu\text{M}$ ), or 47 (10  $\mu\text{M}$ ). Afterwards, LPS was added and the concentrations of cytokines were determined in the supernatants after additional 24 h of treatment. The results are expressed as means of duplicates  $\pm$  SEM of two independent experiments, \* is *p* < 0.05.

mL). Copper(II) bromide (1.7 equiv.) was added slowly and the reaction mixture was stirred at 85 °C for 24 h. The solvent was evaporated under reduced pressure and the solid residue was washed with EtOAc and dried under reduced pressure to obtain a crude product that was used in the next step without further purification.

**2-Bromo-1-(4-ethylphenyl)ethan-1-one 4.** Brown oil; yield 100% (crude, unpurified product); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.21 (t, *J* = 7.0 Hz, 3H, CH<sub>3</sub>), 2.70 (q, *J* = 7.0 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 4.91 (s, 2H, CH<sub>2</sub>), 7.40 (d, *J* = 7.7 Hz, 2H, Ar-H), 7.94 (d, *J* = 7.7 Hz, 2H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>10</sub>H<sub>11</sub>BrO [M + H]<sup>+</sup>: 227.0072, found: 227.0069.

#### General procedure for the synthesis of 1-substituted-2-bromoethan-1-ones 5–7

The corresponding ethan-1-ones (starting amount ranging from 2.4 to 10.2 mmol, 1 equiv.) were dissolved in MeOH (volume ranging from 2 to 8 mL). The solution was cooled to 0 °C and bromine (1 equiv.) was added dropwise. The mixture was stirred for 3 h at a temperature below 15 °C and then H<sub>2</sub>O (2 mL) was added. The mixture was stirred for additional 24–48 h at rt. H<sub>2</sub>O (25 mL) was subsequently added and the aqueous layer was extracted with EtOAc (3 × 25 mL). The combined organic phases were washed with an aqueous solution of Na<sub>2</sub>CO<sub>3</sub> (10%, 50 mL), dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure to obtain crude products that were used in the next step without further purification.

**2-Bromo-1-cyclohexylethan-1-one 7.** Orange oil; yield 90% (crude, unpurified product); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.15–1.48 (m, 5H, cyclohexyl-CH<sub>2</sub>), 1.62–1.70 (m, 1H, cyclohexyl-CH), 1.75–1.82 (m, 2H, cyclohexyl-CH<sub>2</sub>), 1.84–1.92 (m, 2H, cyclohexyl-CH<sub>2</sub>), 2.71 (*sym* m, 1H, cyclohexyl-CH), 3.97 (s, 2H, COCH<sub>2</sub>Br). HRMS (*m/z*) (ESI): calcd for C<sub>8</sub>H<sub>14</sub>OBr [M + H]<sup>+</sup>: 206.0922, found: 206.0918.

#### General procedure for the synthesis of compounds 8–17, 50 and 51

To a solution of 7-hydroxycoumarin 1, 48 or 49 (starting amount ranging from 0.93 to 2.0 mmol, 1 equiv.) in dioxane (50 mL), K<sub>2</sub>CO<sub>3</sub> (4 equiv.) and KI (0.1 equiv.) were added. After 5 min, the corresponding α-haloketone (1.5 equiv.) was added and the reaction mixture was stirred at 100 °C for 24 h. The solvent was then removed under reduced pressure, followed by the addition of H<sub>2</sub>O (100 mL) to the residue. The aqueous phase was extracted with EtOAc (3 × 100 mL), and the combined organic layers were washed with brine (200 mL), dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure. Compounds 8–17, 50 and 51 were subsequently purified by crystallization from EtOH.

**Ethyl 7-(2-(4-ethylphenyl)-2-oxoethoxy)-4-methyl-2-oxo-2H-chromene-3-carboxylate 14.** Yellow solid; yield 74%; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.22 (t, *J* = 7.5 Hz, 3H, Ar-CH<sub>2</sub>CH<sub>3</sub>), 1.30 (t, *J* = 7.1 Hz, 3H, COOCH<sub>2</sub>CH<sub>3</sub>), 2.44 (s, 3H, Ar-CH<sub>3</sub>), 2.72 (q, *J* = 7.5 Hz, 2H, Ar-CH<sub>2</sub>CH<sub>3</sub>), 4.33 (q, *J* = 7.1 Hz, 2H, COOCH<sub>2</sub>CH<sub>3</sub>), 5.77 (s, 2H, CH<sub>2</sub>), 7.09–7.15 (m, 2H, Ar-H),

7.44 (d, *J* = 8.0 Hz, 2H, Ar-H), 7.83 (d, *J* = 9.0 Hz, 1H, Ar-H), 7.97 (d, *J* = 8.0 Hz, 2H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>23</sub>H<sub>22</sub>O<sub>6</sub> [M + H]<sup>+</sup>: 395.1495, found: 395.1491.

**Ethyl 7-(2-(cyclohexyl-2-oxoethoxy)-4-methyl-2-oxo-2H-chromene-3-carboxylate 17.** Orange solid; yield 78%; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.15–1.34 (m, 8H, CH<sub>2</sub>CH<sub>3</sub> and cyclohexyl-CH<sub>2</sub>), 1.62–1.88 (m, 5H, cyclohexyl-CH<sub>2</sub>), 2.43 (s, 3H, Ar-CH<sub>3</sub>), 2.53–2.60 (m, 1H, cyclohexyl-CH), 4.33 (d, *J* = 7.1 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 5.16 (s, 2H, CH<sub>2</sub>), 6.96–7.03 (m, 2H, Ar-H), 7.81 (d, *J* = 8.8 Hz, 1H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>21</sub>H<sub>24</sub>O<sub>6</sub> [M + H]<sup>+</sup>: 373.1651, found: 373.1651.

Spectroscopic analyses of compounds 50 and 51 were previously reported.<sup>22</sup>

#### General procedure for the synthesis of 4-methyl-3-carboxyalkyl substituted psoralens 18–27, 52 and 53

To a suspension of the desired *O*-alkylated compound 8–17, 48 or 49 (starting amount ranging from 0.56 to 1.45 mmol, 1 equiv.) in propan-2-ol (volume ranging from 20–30 mL), an aqueous solution of NaOH (1 M, 10 equiv.) was added. The reaction mixture was stirred at 100 °C for 5–20 min. After the reaction was complete (monitored by TLC), propan-2-ol was evaporated under reduced pressure. The aqueous residue was acidified with 5 M HCl to pH 1, and the precipitate that formed was filtered off. In cases where there was no precipitate, the aqueous layer was extracted with EtOAc (2 × 50 mL), and the combined organic layers were washed with brine (100 mL), dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure to obtain a crude product that was used in the next step without further purification.

**3-(4-Ethylphenyl)-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-carboxylic acid 24.** Brown oil; yield 100% (crude, unpurified product); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.20 (t, *J* = 7.5 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>), 2.45 (s, 3H, Ar-CH<sub>3</sub>), 2.68 (q, *J* = 7.5 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 7.31 (s, 1H, Ar-H), 7.40–7.47 (m, 4H, Ar-H), 8.26 (s, 1H, Ar-H), 8.49 (s, 1H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>21</sub>H<sub>16</sub>O<sub>5</sub> [M + H]<sup>+</sup>: 349.1076, found: 349.1081.

**3-Cyclohexyl-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-carboxylic acid 27.** Orange solid; yield 100% (crude, unpurified product); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.26–1.31 (m, 4H, cyclohexyl-CH<sub>2</sub>), 1.42–1.49 (m, 4H, cyclohexyl-CH<sub>2</sub>), 2.05–2.07 (m, 2H, cyclohexyl-CH<sub>2</sub>), 2.43 (s, 3H, Ar-CH<sub>3</sub>), 2.82–2.87 (m, 1H, cyclohexyl-CH), 7.57 (s, 1H, Ar-H), 7.82 (d, *J* = 0.8 Hz, 1H, Ar-H), 7.93 (s, 1H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>19</sub>H<sub>18</sub>O<sub>5</sub> [M + H]<sup>+</sup>: 327.1232, found: 327.1235.

Spectroscopic analyses of compounds 52 and 53 were previously reported.<sup>22</sup>

#### General procedure for the synthesis of 4-methyl-3-carboxamide substituted psoralens 28–37

To a suspension of the desired 4-methyl-3-carboxyalkyl substituted psoralen 18–27 (starting amount ranging from 0.29 to 2.10 mmol, 1 equiv.) in dioxane (volume ranging from 4–20 mL), Boc<sub>2</sub>O (2 equiv.), pyridine (4 equiv.) and ammonium acetate (6 equiv.) were added. The reaction

mixture was stirred at rt for 24 h. The solvent was evaporated under reduced pressure and pyridine was removed by co-evaporation with toluene. CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added and the organic phase was washed with water (2 × 50 mL). The organic layer was washed with brine (100 mL), dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure. The compounds were purified by column chromatography (EtOAc/*n*-hexane = 2 : 1).

**3-(4-Ethylphenyl)-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-carboxamide 34.** Orange solid; yield 6%; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.32 (t, *J* = 7.6 Hz, 3H, CH<sub>2</sub>CH<sub>3</sub>), 2.75 (q, *J* = 7.6 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 2.90 (s, 3H, Ar-CH<sub>3</sub>), 7.36–7.41 (m, 2H, CONH<sub>2</sub>), 7.53 (s, 1H, Ar-H), 7.54–7.57 (m, 2H, Ar-H), 7.72–7.76 (m, 2H, Ar-H), 7.84 (s, 1H, Ar-H), 8.20 (s, 1H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>21</sub>H<sub>17</sub>NO<sub>4</sub> [M + H]<sup>+</sup>: 348.1236, found: 348.1241.

**3-Cyclohexyl-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-carboxamide 37.** White solid; yield 27%; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.42–1.52 (m, 4H, cyclohexyl-CH<sub>2</sub>), 1.73–1.83 (m, 4H, cyclohexyl-CH<sub>2</sub>), 2.05–2.09 (m, 2H, cyclohexyl-CH<sub>2</sub>), 2.54 (s, 3H, Ar-CH<sub>3</sub>), 2.83–2.91 (m, 1H, cyclohexyl-CH), 7.67–7.69 (m, 1H, CONH<sub>2</sub>), 7.71 (s, 1H, Ar-H), 7.87–7.89 (m, 2H, Ar-H and CONH<sub>2</sub>), 8.11 (s, 1H, Ar-H). HRMS (*m/z*) (ESI): calcd for C<sub>19</sub>H<sub>19</sub>NO<sub>4</sub> [M + H]<sup>+</sup>: 326.1392, found: 326.1395.

#### General procedure for the synthesis of oxathiazolone-based compounds 38–47

To a suspension of the desired 4-methyl-3-carboxamide substituted psoralen 28–37 (starting amount ranging from 0.09 to 0.47 mmol, 1 equiv.) in anhydrous THF (volume ranging from 3 to 5 mL), chlorocarbonylsulfonyl chloride (4 equiv.) was added under argon at rt. The resulting solution was stirred at 80 °C for 1.5–2 h. After the reaction was complete (monitored by TLC), the solvent was removed under reduced pressure. Compounds 38–47 were subsequently purified by crystallization from a mixture of EtOAc/*n*-hexane and column chromatography (EtOAc/*n*-hexane = 1 : 3).

**5-(3-(4-Ethylphenyl)-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-yl)-1,3,4-oxathiazol-2-one 44.** Yellow solid; yield 16%; mp 67.5–68.8 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.32 (t, *J* = 7.6 Hz, CH<sub>2</sub>CH<sub>3</sub>), 2.65 (s, 3H, Ar-CH<sub>3</sub>), 2.75 (q, *J* = 7.6 Hz, 2H, CH<sub>2</sub>CH<sub>3</sub>), 7.36–7.41 (m, 2H, Ar-H), 7.53–7.56 (m, 2H, Ar-H), 7.57 (s, 1H, Ar-H), 7.85 (s, 1H, Ar-H), 8.12 (s, 1H, Ar-H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 172.93, 158.17, 156.21, 153.57, 151.41, 149.65, 144.69, 143.35, 128.90 (2C), 127.75, 127.64 (2C), 125.16, 122.39, 117.64, 115.50, 112.71, 100.55, 28.72, 17.16, 15.60. HRMS (*m/z*) (ESI): calcd for C<sub>22</sub>H<sub>16</sub>NO<sub>5</sub>S [M + H]<sup>+</sup>: 406.07437, found: 406.07355. Purity by HPLC: 100%.

**5-(3-Cyclohexyl-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-yl)-1,3,4-oxathiazol-2-one 47.** Off-white solid; yield 45%; mp 124.8–125.8 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.28–1.39 (m, 1H, cyclohexyl-CH<sub>2</sub>), 1.47–1.55 (m, 4H, cyclohexyl-CH<sub>2</sub>), 1.83–1.94 (m, 3H, cyclohexyl-CH<sub>2</sub>), 2.13–2.15 (m, 2H, cyclohexyl-CH<sub>2</sub>), 2.70 (s, 3H, Ar-CH<sub>3</sub>), 2.76–2.86 (m, 1H, cyclohexyl-CH),

7.49 (s, 1H, Ar-H), 7.50 (d, *J* = 0.8 Hz, 1H, Ar-H), 7.91 (s, 1H, Ar-H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 173.00, 158.08, 156.29, 153.72, 151.23, 142.57, 126.54, 126.03, 116.97, 114.80, 112.38, 111.47, 100.31, 34.07, 33.09 (2C), 26.46 (2C), 26.18, 17.21. HRMS (*m/z*) (ESI): calcd for C<sub>20</sub>H<sub>17</sub>NO<sub>5</sub>S [M + H]<sup>+</sup>: 384.0906, found: 384.0899. Purity by HPLC: 96%.

#### General procedure for the synthesis of coumarin derivatives 48 and 49

To a solution of resorcinol (45.41 mmol, 1 equiv.) in H<sub>2</sub>SO<sub>4</sub> (75%, 50 mL), diethyl 2-acetylsuccinate or diethyl 2-acetylglutarate (1.1 equiv.) was added. The reaction mixture was stirred at room temperature for 24 h. After the reaction was complete, the solution was poured onto crushed ice. The solid that formed was filtered off, washed with water and dried. Spectroscopic analyses of compounds 48 and 49 were previously reported.<sup>22</sup>

#### General procedure for the synthesis of psoralens with the cyclohexyl moiety at position 4' with variation at position 3 (compounds 54–59)

The starting compound 52 or 53 (1 equiv.) was transferred to a dry flask. A suitable nucleophilic reagent (*N*-hydroxysuccinimide (1 equiv.), 2-(methylamino)acetonitrile (1.25 equiv.) or 2-aminoacetonitrile (2 equiv.)) was added. THF (5 mL) was added under an argon atmosphere at 0 °C. After stirring for 5 min at 0 °C, EDC (1.25 equiv.) and Et<sub>3</sub>N (2.5 equiv.) were added. The reaction mixture was stirred at room temperature for 24 h. The solvent was evaporated and HCl (0.01 M, 20 mL) was added. The reaction mixture was extracted with EtOAc (3 × 30 mL). The combined organic layers were dried with anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated under reduced pressure. The compounds were purified by column chromatography (EtOAc or EtOAc/*n*-hexane = 2 : 1).

**2,5-Dioxopyrrolidin-1-yl 3-(3-cyclohexyl-5-methyl-7-oxo-7H-furo[3,2-*g*]chromen-6-yl)propanoate 55.** White solid; yield 32%; mp 150.1–151.3 °C; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.25–1.30 (m, 1H, cyclohexyl-H), 1.36–1.50 (m, 4H, cyclohexyl-H), 1.70–1.85 (m, 3H, cyclohexyl-H), 2.02–2.09 (m, 2H, cyclohexyl-H), 2.58 (s, 3H, Ar-CH<sub>3</sub>), 2.80 (br s, 4H, COCH<sub>2</sub>CH<sub>2</sub>CO), 2.84–2.90 (m, 1H, cyclohexyl-H), 2.95 (*sym* m, 4H, CH<sub>2</sub>), 7.65 (s, 1H, Ar-H), 7.85 (s, 1H, Ar-H), 8.07 (s, 1H, Ar-H); <sup>13</sup>C NMR (100 MHz, DMSO-*d*<sub>6</sub>) δ 170.12, 168.22, 160.52, 155.44, 149.65, 149.10, 142.37, 126.20, 124.44, 120.49, 116.40, 116.12, 98.96, 33.02, 32.50, 28.68, 25.95, 25.69, 25.37, 22.58, 15.36. HRMS (*m/z*) (ESI): calcd for C<sub>25</sub>H<sub>26</sub>NO<sub>7</sub> [M + H]<sup>+</sup>: 452.1723, found: 452.1716. Elemental analysis: found: C, 66.20; H, 5.50; N, 3.41; calcd for C<sub>25</sub>H<sub>25</sub>NO<sub>7</sub>: C, 66.51; H, 5.58; N, 3.10%.

## Conflicts of interest

The authors declare no competing interests.

## Acknowledgements

We sincerely thank Assoc. Prof. Dr. Urban Švajger (Center for Development of Advanced Therapy Medicinal Products, Blood Transfusion Center of Slovenia) for the assistance in performing the cytokine secretion assay. The authors acknowledge the support from the Slovenian Research Agency: research core funding No. P1-0208, and grant number N1-0068 (B) to S.G.: identification of non-peptidic inhibitors of the immunoproteasome using fragment based drug discovery methods.

## Notes and references

- G. A. Collins and A. L. Goldberg, *Cell*, 2017, **169**, 792–806.
- L. Budenholzer, C. L. Cheng, Y. Li and M. Hochstrasser, *J. Mol. Biol.*, 2017, **429**, 3500–3524.
- C. S. Arendt and M. Hochstrasser, *Proc. Natl. Acad. Sci. U. S. A.*, 1997, **94**, 7156–7161.
- M. Groettrup, S. Khan, K. Schwarz and G. Schmidtke, *Biochimie*, 2001, **83**, 367–372.
- T. A. Griffin, D. Nandi, M. Cruz, H. J. Fehling, L. Van Kaer, J. J. Monaco and R. A. Colbert, *J. Exp. Med.*, 1998, **187**, 97–104.
- P. Śledź and W. Baumeister, *Annu. Rev. Pharmacol. Toxicol.*, 2016, **56**, 191–209.
- N. Micale, K. Scarbaci, V. Troiano, R. Ettari, S. Grasso and M. Zappalà, *Med. Res. Rev.*, 2014, **34**, 1001–1069.
- D. J. Kuhn, S. A. Hunsucker, Q. Chen, P. M. Voorhees, M. Orłowski and R. Z. Orłowski, *Blood*, 2009, **113**, 4667–4676.
- F. Parlati, S. J. Lee, M. Aujay, E. Suzuki, K. Levitsky, J. B. Lorens, D. R. Micklem, P. Ruurs, C. Sylvain, Y. Lu, K. D. Shenk and M. K. Bennett, *Blood*, 2009, **114**, 3439–3447.
- T. A. Thibaudeau and D. M. Smith, *Pharmacol. Rev.*, 2019, **71**, 170–197.
- M. Basler, M. M. Lindstrom, J. J. LaStant, J. M. Bradshaw, T. D. Owens, C. Schmidt, E. Maurits, C. Tsu, H. S. Overkleeft, C. J. Kirk, C. L. Langrish and M. Groettrup, *EMBO Rep.*, 2018, **19**, e46512.
- H. W. Johnson, E. Lowe, J. L. Anderl, A. Fan, T. Muchamuel, S. Bowers, D. C. Moebius, C. Kirk and D. L. McMinn, *J. Med. Chem.*, 2018, **61**, 11127–11143.
- G. de Bruin, E. M. Hubert, B.-T. Xin, E. van Rooden, K. Al-Ayed, K.-B. Kim, A. F. Kisselev, C. Driessen, M. van der Stelt, G. A. van der Marel, M. Groll and H. S. Overkleeft, *J. Med. Chem.*, 2014, **57**, 6197–6209.
- B.-T. Xin, E. M. Huber, G. de Bruin, W. Heinemeyer, E. Maurits, C. Espinal, Y. Du, M. Janssens, E. S. Weyburne, A. F. Kisselev, B. I. Florea, C. Driessen, G. A. van der Marel, M. Groll and H. S. Overkleeft, *J. Med. Chem.*, 2019, **62**, 1626–1642.
- C. Dubiella, R. Baur, H. Cui, E. M. Huber and M. Groll, *Angew. Chem., Int. Ed.*, 2015, **54**, 15888–15891.
- P. K. Singh, H. Fan, X. Jiang, L. Shi, C. F. Nathan and G. Lin, *ChemMedChem*, 2016, **11**, 2127–2131.
- E. S. Karreci, H. Fan, M. Uehara, A. B. Mihali, P. K. Singh, A. T. Kurdi, Z. Solhjoui, L. V. Riella, I. Ghobrial, T. Laragione, S. Routray, J. P. Assaker, R. Wang, G. Sukenick, L. Shi, F. J. Barrat, C. F. Nathan, G. Lin and J. Azzi, *Proc. Natl. Acad. Sci. U. S. A.*, 2016, **113**, E8425–E8432.
- H. W. Johnson, J. L. Anderl, E. K. Bradley, J. Bui, J. Jones, S. Arastu-Kapur, L. M. Kelly, E. Lowe, D. C. Moebius, T. Muchamuel, C. Kirk, Z. Wang and D. McMinn, *ACS Med. Chem. Lett.*, 2017, **8**, 413–417.
- R. Ettari, S. Previti, A. Bitto, S. Grasso and M. Zappala, *Curr. Med. Chem.*, 2016, **23**, 217–238.
- A. F. Kisselev and M. Groettrup, *Curr. Opin. Chem. Biol.*, 2014, **23**, 16–22.
- E. M. Huber and M. Groll, *Angew. Chem., Int. Ed.*, 2012, **51**, 8708–8720.
- I. Sosič, M. Gobec, B. Brus, D. Knez, M. Živec, J. Konc, S. Lešnik, M. Ogrizek, A. Obreza, D. Žigon, D. Janežič, I. Mlinarič-Raščan and S. Gobec, *Angew. Chem., Int. Ed.*, 2016, **55**, 5745–5748.
- V. Kasam, N.-R. Lee, K.-B. Kim and C.-G. Zhan, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 3614–3617.
- H. Cui, R. Baur, C. Le Chapelain, C. Dubiella, W. Heinemeyer, E. M. Huber and M. Groll, *ChemBioChem*, 2017, **18**, 523–526.
- H. Fan, N. G. Angelo, J. D. Warren, C. F. Nathan and G. Lin, *ACS Med. Chem. Lett.*, 2014, **5**, 405–410.
- E. Bosc, J. Nastri, V. Lefort, M. Valli, F. Contiguiba, R. Pioli, M. Furlan, V. da Silva Bolzani, C. El Amri and M. Reboud-Ravaux, *Biochem. Biophys. Res. Commun.*, 2018, **496**, 961–966.
- G. Lin, D. Li, L. de Carvalho, H. Deng, H. Tao, G. Vogt, K. Wu, J. Schneider, T. Chidawanyika, J. Warren, H. Li and C. Nathan, *Nature*, 2009, **461**(7264), 621–626.
- D. Niewerth, G. Jansen, L. F. V. Riethoff, J. van Meerloo, A. J. Kale, B. S. Moore, Y. G. Assaraf, J. L. Anderl, S. Zweegman, G. J. L. Kaspers and J. Cloos, *Mol. Pharmacol.*, 2014, **86**, 12–19.
- A. Daina, O. Michielin and V. Zoete, *Sci. Rep.*, 2017, **7**, 42717.