

Comment on “Global Access to Handwashing: Implications for COVID-19 Control in Low-Income Countries”

Matthew C. Freeman¹ and Bethany A. Caruso²

¹Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, USA

²Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, Georgia, USA

<https://doi.org/10.1289/EHP7852>

Refers to <https://doi.org/10.1289/EHP7200>

Brauer et al. (2020) estimate that over 2 billion people lack access to household handwashing facilities, amplifying critical questions for COVID-19 prevention measures in low-income countries. Despite these dire figures, the situation may be far worse, given unreliable and costly access to water and soap in households, schools, and health care facilities. The current World Health Organization COVID-19 guidance on hand hygiene (WHO 2020) will be infeasible for many and requires updating to support resource-poor settings.

Hundreds of millions in sub-Saharan Africa alone require at least 30 min per trip to fetch drinking water (WHO/UNICEF 2017). Data on non-drinking–water sources are not routinely collected and reported at scale, but evidence suggests that water-insecure households prioritize using water for drinking over hygiene practices (Stevenson et al. 2012). Both soap and water may be costly, prioritized for other uses, and tightly rationed, if available. Fetching water from public sources—predominantly performed by women and girls (Graham et al. 2016)—requires waiting at communal locations that may be crowded. Individuals must also touch widely used public water pump handles and taps, surfaces on which SARS-CoV-2 can remain viable for several days (van Doremalen et al. 2020).

Further, water-insecure communities are frequently served by resource-poor health care facilities. An estimated 20% of health care facilities globally—40% in sub-Saharan Africa—do not have an on-premises water source (WHO/UNICEF 2020b); the proportion with handwashing facilities is unknown (WHO/UNICEF 2020a). Local outbreaks will result in an amplification of the exposure risk for frontline health workers—also predominantly women—making it all the more crucial to get handwashing right in these settings.

What should be done? Clear, evidence-based guidance is needed on the efficient use of water and soap or soap alternatives in water-scarce settings. “Soapy water”—a mixture of powder, liquid, or bar soap diluted in water—may be a scalable and economical solution (Amin et al. 2014). Ash has been shown to be effective for cleaning hands when soap and water are scarce (Bloomfield and Nath 2009), but its effectiveness against SARS-CoV-2 needs to be established. Evidence on safe strategies for economizing water for handwashing, such as using non-drinking–water sources or reusing water, is needed.

Address correspondence to Matthew C. Freeman, Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University, 1518 Clifton Rd. NE, Atlanta, GA 30322 USA. Email: matthew.freeman@emory.edu

The authors declare they have no actual or potential competing financial interests.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Although infections in much of sub-Saharan Africa have been low relative to other regions (WHO Regional Office for Africa 2020), the pace of COVID-19 infections has been quickening, and the escalating cases in South Africa (IHME 2020a) may be a harbinger of a coming wave of infection. Cases of COVID-19 are surging in other countries with limited access to handwashing stations and water among the poorest populations (IHME 2020b). Clear, targeted, and actionable recommendations are needed for people without reliable water access (WHO/UNICEF 2017). Low-flow and locally fabricated handwashing stations, soap or alternatives, and evidence-based behavior change approaches should be developed and deployed. Lessons learned could support long-term habit formation among hard-to-reach populations and adjustments in development approaches.

Handwashing with soap is one of the most cost-effective public health interventions. Yet water inequity magnifies health disparity, making a simple behavior unattainable for the world’s poorest unless urgent action is taken.

References

- Amin N, Pickering AJ, Ram PK, Unicomb L, Najnin N, Homaira N, et al. 2014. Microbiological evaluation of the efficacy of soapy water to clean hands: a randomized, non-inferiority field trial. *Am J Trop Med Hyg* 91(2):415–423, PMID: 24914003, <https://doi.org/10.4269/ajtmh.13-0475>.
- Bloomfield SF, Nath KJ. 2009. Use of ash and mud for handwashing in low income communities. *Home Hygiene & Health, Reviews—For Developing Countries* section. 14 October 2009. <https://www.ifh-homehygiene.org/review-best-practice/use-ash-and-mud-handwashing-low-income-communities> [accessed 29 July 2020].
- Brauer M, Zhao JT, Bennitt FB, Stanaway JD. 2020. Global access to handwashing: implications for COVID-19 control in low-income countries. *Environ Health Perspect* 128(5):57005, PMID: 32438824, <https://doi.org/10.1289/EHP7200>.
- Graham JP, Hirai M, Kim S-S. 2016. An analysis of water collection labor among women and children in 24 sub-Saharan African countries. *PLoS One* 11(6):e0155981, PMID: 27248494, <https://doi.org/10.1371/journal.pone.0155981>.
- IHME (Institute for Health Metrics and Evaluation). 2020a. COVID-19 projections. South Africa. Last updated 22 July 2020. <https://covid19.healthdata.org/south-africa> [accessed 29 July 2020].
- IHME. 2020b. COVID-19 projections. Total deaths. Last updated 22 July 2020. <https://covid19.healthdata.org> [accessed 29 July 2020].
- Stevenson EGJ, Greene LE, Maes KC, Ambelu A, Tesfaye YA, Rheingans R, et al. 2012. Water insecurity in 3 dimensions: an anthropological perspective on water and women’s psychosocial distress in Ethiopia. *Soc Sci Med* 75(2):392–400, PMID: 22575697, <https://doi.org/10.1016/j.socscimed.2012.03.022>.
- van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. 2020. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. [Letter.] *N Engl J Med* 382(16):1564–1567, PMID: 32182409, <https://doi.org/10.1056/NEJMc2004973>.
- WHO (World Health Organization). 2020. Interim recommendations on obligatory hand hygiene against transmission of COVID-19. Geneva, Switzerland: WHO. <https://www.who.int/publications/m/item/interim-recommendations-on-obligatory-hand-hygiene-against-transmission-of-covid-19> [accessed 29 July 2020].
- WHO Regional Office for Africa. 2020. Africa records over 200 000 COVID-19 cases. [Press Release.] 11 June 2020. <https://www.afro.who.int/news/africa-records-over-200-000-covid-19-cases> [accessed 29 July 2020].

WHO/UNICEF (WHO and the United Nations Children's Fund). 2017. *Progress on Drinking Water, Sanitation and Hygiene: 2017 Update and SDG Baselines*. Geneva, Switzerland: WHO/UNICEF. <http://apps.who.int/iris/bitstream/10665/258617/1/9789241512893-eng.pdf?ua=1> [accessed 29 July 2020].

WHO/UNICEF. 2020a. Health care facility data—world—2016—service levels. Joint Monitoring Programme on Water Supply, Sanitation, and Hygiene

Dashboard. <https://washdata.org/data/healthcare#!/dashboard/3211> [accessed 29 July 2020].

WHO/UNICEF. 2020b. Water services in health care facilities, 2016. Health care facility data—SDG regions—sub-Saharan Africa—2016—service elements. Joint Monitoring Programme on Water Supply, Sanitation, and Hygiene Dashboard. <https://washdata.org/data/healthcare#!/dashboard/3210> [accessed 29 July 2020].