

Correction: Is there a link between very early changes of primary and secondary lymphoid organs in ^{18}F -FDG-PET/MRI and treatment response to checkpoint inhibitor therapy?

Seith F, Forschner A, Weide B, *et al.* Is there a link between very early changes of primary and secondary lymphoid organs in ^{18}F -FDG-PET/MRI and treatment response to checkpoint inhibitor therapy? *J Immunother Cancer* 2020;8:e000656. doi: 10.1136/jitc-2020-000656

This paper has been corrected since it was published online. The affiliation ‘Cluster of Excellence iFIT (EXC 2180) “Image-Guided and Functionally Instructed Tumor Therapies”, Eberhard Karls University, 72076 Tübingen, Germany’ was missing from the paper.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

J Immunother Cancer 2020;8:e000656corr1. doi:10.1136/jitc-2020-000656corr1

