

## Corrigendum to: Hospital Readmissions After Laboratory-Confirmed Influenza Hospitalization

David M. Dobrzynski Jr,<sup>1</sup> Danielle N. Ndi,<sup>2</sup> Yuwei Zhu,<sup>3</sup> Tiffanie Markus,<sup>2</sup> William Schaffner,<sup>2,4</sup> and H. Keipp Talbot<sup>2,4</sup>

<sup>1</sup>Division of Infectious Diseases, University of Rochester Medical Center, Rochester, New York, USA, <sup>2</sup>Department of Health Policy, Vanderbilt University Medical Center, Nashville, Tennessee, USA, <sup>3</sup>Department of Biostatistics, Vanderbilt University School of Medicine, Nashville, Tennessee, USA, <sup>4</sup>Division of Infectious Diseases, Vanderbilt University Medical Center, Nashville, Tennessee, USA

In “Hospital Readmissions After Laboratory-Confirmed Influenza Hospitalization” by Dobrzynski et al [J Infect Dis 2020; jiaa117], Figure 2B had incorrect values for the proportion readmitted from Time from Index Hospitalization. These values have been re-labeled and the solid black line was graphed

accordingly in the published version. Additionally, Figure 2B was incorrectly referenced after “Patients aged 65 years and older accounted for 1380 total readmissions, 215 of which occurred within 30 days among 188 patients (6%) (Figure 2B).”, which was removed in the published version. The authors regret these errors.

DOI: 10.1093/infdis/jiaa427

## Corrigendum to: A Randomized Clinical Trial to Compare *Plasmodium falciparum* Gametocytemia and Infectivity After Blood-Stage or Mosquito Bite–Induced Controlled Malaria Infection

Manon Alkema,<sup>1</sup> Isaie J. Reuling,<sup>1</sup> Gerdie M. de Jong,<sup>2</sup> Kjerstin Lanke,<sup>1</sup> Luc E. Coffeng,<sup>3</sup> Geert-Jan van Gemert,<sup>1</sup> Marga van de Vegte-Bolmer,<sup>1</sup> Quirijn de Mast,<sup>4</sup> Reinout van Crevel,<sup>4</sup> Karen Ivinson,<sup>5</sup> Christian F. Ockenhouse,<sup>5</sup> James S. McCarthy,<sup>6</sup> Robert Sauerwein,<sup>1</sup> Katharine A. Collins,<sup>1</sup> and Teun Bousema<sup>1</sup>

<sup>1</sup>Department of Medical Microbiology, Radboud University Medical Center, Nijmegen, the Netherlands, <sup>2</sup>Department of Medical Microbiology and Infectious Diseases, University Medical Center Rotterdam, Rotterdam, the Netherlands, <sup>3</sup>Department of Public Health, University Medical Center Rotterdam, Rotterdam, the Netherlands, <sup>4</sup>Department of Internal Medicine, Radboud University Medical Center, Nijmegen, the Netherlands, <sup>5</sup>PATH Malaria Vaccine Initiative, Washington, DC, USA, and <sup>6</sup>Clinical Tropical Medicine Laboratory, QIMR Berghofer Medical Research Institute, Brisbane, Australia

In “A Randomized Clinical Trial to Compare *Plasmodium falciparum* Gametocytemia and Infectivity After Blood-Stage or Mosquito Bite–Induced Controlled Malaria Infection” by Alkema et al [J Infect Dis 2020; jiaa157], data points for MB

Inoculations and IBSM were erroneously reversed for the variable: Individuals infectious to mosquitoes, % (no./total) in Table 2. This has been corrected in the article, and the authors regret the error.

DOI: 10.1093/infdis/jiaa428