

Cost-Effectiveness Analysis of Colorectal Cancer Screening: A Systematic Review

This article was published in the following Dove Press journal:
Risk Management and Healthcare Policy

Farhad Khalili ¹
Behzad Najafi ²
Fariborz Mansour-
Ghanaei ³
Mahmood Yousefi⁴
Hadi Abdollahzad ⁵
Ali Motlagh⁶

¹Department of Health Economics, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran; ²Iranian Center of Excellence in Health Management, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran; ³Gastrointestinal and Liver Diseases Research Center, Guilan University of Medical Sciences, Rasht, Iran; ⁴Department of Health Economics, Iranian Center of Excellence in Health Management, School of Management and Medical Informatics, Tabriz University of Medical Sciences, Tabriz, Iran; ⁵Research Center for Environmental Determinants of Health, Health Institute, Kermanshah University of Medical Sciences, Kermanshah, Iran; ⁶National Cancer Control Secretariat, Ministry of Health and Medical Education, Tehran, Iran

Introduction: Colorectal cancer (CRC) is a significant health problem with an increasing incidence worldwide. Screening is one of the ways, in which cases and deaths of CRC can be prevented. The objective of this systematic review was to evaluate the cost-effectiveness of the different CRC screening techniques and to specify the efficient technique from a cost-effectiveness perspective.

Methods: The economic studies of CRC screening in general populations (average risk), aged 50 years and above were reviewed. Two reviewers independently reviewed the titles, abstracts, and full-texts of the studies in five databases: Cochrane, Embase, Scopus, Web of Science and PubMed. The disagreements between reviewers were resolved through the authors' consensus. The main outcome measures in this systematic review were the incremental cost-effectiveness ratio (ICER) of screening versus no-screening and then in comparison with other screening techniques. The ICER is defined by the difference in cost between two possible interventions, divided by the difference in their effect.

Results: Eight studies were identified and retained for the final analysis. In this study, when screening techniques were compared to no-screening, all CRC screening techniques showed to be cost-effective. The lowest ICER calculated was \$PPP -16265/quality-adjusted life-year (QALY) (the negative ICERs were between purchasing power parity in US dollar (\$PPP) -16265/QALY to \$PPP -1988/QALY, whereas the positive ICERs were between \$PPP 1257/QALY to \$PPP 55987/QALY). For studies comparing various screening techniques, there was great heterogeneity in terms of the structures of the analyses, leading to diverse conclusions about their incremental cost-effectiveness.

Conclusion: All CRC screening techniques were cost-effective, compared with the no-screening methods. The cost-effectiveness of the various screening techniques mainly was dependent on the context-specific parameters and highly affected by the framework of the cost-effectiveness analysis. In order to make the studies comparable, it is important to adopt a reference-based methodology for economic evaluation studies.

Keywords: incremental cost-effectiveness ratio, screening techniques, economic evaluation, colorectal cancer

Introduction

Among the world mostly diagnosed cancers, colorectal cancer (CRC) takes the third place in males and the second in females, worldwide.¹

In 2018, there were an estimated 1.8 million new CRC cases and 861,663 deaths in both sexes at all ages (0–85+ years). This is already becoming a major public health problem across the world.^{1,2} The attributable cases and deaths of CRC can be avoided through early diagnosis. Screening for finding cases at the early stages of the disease is a well-known strategy of controlling disease, in fact, early detection

Correspondence: Mahmood Yousefi
Email mahmoodyousefi59@gmail.com

of cancer helps to remove precancerous lesions and prevent cancer from reaching advanced stages. While the European Union recommends screening by a fecal occult blood test (FOBT), in the US the guidelines recommend people older than 50 years, to choose among several recommended strategies.^{3,4}

Screening programs often target people with apparently healthy conditions, and active screening tends to be delivered collectively. Thus, the number of people using screening services is usually higher than the true number of patients. This in turn has implications for high costs per diagnosed patient. In an optimal situation, the return of benefits to society due to the implementation of screening should outweigh the associated costs. In essence, understanding this situation depends on some basic factors like the prevalence of the disease, accuracy of screening methods, target population, and cost of each strategy.⁵ Economic evaluation is a systematic and formal way of assessing the costs and benefits of screening interventions. There are several techniques for screening of CRC; colonoscopy, which is one of the most popular interventions, has been associated with relatively high but better accuracy performance,⁶ in contrast, the FOBT has advantages over colonoscopy in terms of being less costly and easy to perform. The fecal immunochemical test (FIT), which is also called either immunochemical FOBT or IFOBT, is usually favored because it does not need any dietary restriction, and it has higher specificity when it is compared to guaiac-based FOBT (G-FOBT).^{7,8} Furthermore, the non-invasive hybrid screening techniques, which is a stool DNA test (S-DNA test), have recently been approved by the US Food and Drug Administration (FDA).^{9,10} The characteristics of different types of screening techniques are presented in [Appendix 1](#).

We performed a systematic review to retrieve the evidence on the cost-effectiveness of the various CRC screening methods. The study was designed to focus on the most recent studies, which have been conducted worldwide, since 2012.

This review aimed to explore the cost-effectiveness of the different screening strategies, either in comparison with no-screening or other techniques, in the average-risk population.

Methods

Search Strategy

We conducted this systematic review based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines,¹¹ and following

the Cochrane Handbooks for systematic reviews.¹² The search strategy used in this review was registered in Prospero¹³ (registration number: CRD42018081676). The approach for identifying studies and extracting the data has been described in earlier studies, and performed in the economic evaluation of CRC screening techniques.^{14,15} Five databases were searched for identifying relevant studies, including Cochrane, Embase, Scopus, Web of Science, PubMed. We used a specific search strategy for each database and the following key terms were used: “colorectal cancer”, “cost-effectiveness” and “screening”. Our search was restricted to studies written in English and published between January 2012 and December 2018, and then updated until June 2020. In order to find the studies not covered by our database searches and studies in gray literature, we manually looked at the reference lists of the studies and contacted the authors of the concerned studies. The studies were chosen, according to the similar inclusion and exclusion criteria, as in the US Preventive Services Task Force (USPSTF) review.¹⁶

The inclusion criteria were: 1) Studies that focused on CRC screening, 2) Studies that aimed at average risk¹⁷ populations, aged 50 years and above, 3) Studies that reported the Incremental Cost-Effectiveness Ratio (ICER) or provided data for ICER calculation, 4) Studies that outlined cost per quality-adjusted life-year (QALY) gained or cost per life-year gained, 5) Studies that had full economic evaluations, and 6) Studies published in English.

The exclusion criteria were as follows: 1) Articles that reported only cost per cancer detected, cost per patient screened and cost per death prevented, 2) Non-original studies, 3) Opportunistic screening, and 4) Short-term decision trees.

Data Extraction and Quality Assessment

Firstly, two reviewers independently assessed the titles and abstracts of the retrieved studies and then excluded the irrelevant studies based on the developed criteria. When the articles could not be located based on the title and abstract, the full-text was evaluated. The disagreements between reviewers about inclusion and exclusion of studies were resolved through the authors' consensus or by the third reviewer. For data collection, we designed a data extraction form containing 1) Details of bibliography, 2) Study design, including aim and cases of studies, time horizon, interventions and alternatives, costs included in the study, sources of screening cost, outcome measures for effectiveness, data sources that are relevant to the

outcome, study's perspective, modeling sensitivity analysis and discount rate for both costs and outcomes, and 3) Main results and conclusions. Data extracted from each article were included in the table for each of the techniques under consideration such as FIT, Flexible Sigmoidoscopy (FS), FOBT, Multi-target Stool DNA (MT- S-DNA), colonoscopy, double-contrast barium enema (based on the age of the subjects and the time intervals of 2, 5 and 10 years). To compare different alternatives, costs were updated to purchasing power parity (PPP) in 2017, using the Consumer Price Index (CPI) for medical care and PPP conversion factor. The ICER was the primary outcome measure in this systematic review, for comparing the screening and no-screening methods, and then in comparison with other screening techniques. The ICERs calculated as either cost per life-year gained or cost per QALY.

$$ICER = \frac{\text{Total cost of new intervention} - \text{total cost of no screening or old intervention}}{\text{Outcome of new intervention} - \text{outcome of no screening or old intervention}}$$

In this systematic review, the threshold is defined as a standard threshold in the country of origin where the paper was published, while for the countries that did not report any threshold, the World Health Organization (WHO) recommended guideline was used.^{18–20} The criterion for assessing the cost-effectiveness was based on the recommended WHO threshold for cost-effectiveness analysis. According to this criterion, if the value of an ICER for a given intervention falls below the specified value of three-time the Gross Domestic Product (GDP) per capita, then it is considered as a cost-effective strategy.

We used the Drummond 10-point checklist, which is a standard procedure for the quality assessment of economic evaluation studies, for qualitative assessment of studies, in order to confirm their methodological quality.²¹ All questions had three options for the answer, including “yes”, “cannot tell” and “no”, and the corresponding value for each item was 1, 0.5 and 0, respectively. The reason for choosing a 0.5 value for the “cannot tell” response was that the information about that item was not complete. For items that were not relevant, we considered score 1.

Results

Study Selection

A flow diagram, based on the PRISMA, for choosing the relevant studies has been shown in [Figure 1](#). In primary

search, we identified 1701 relevant studies, which were published after January 2012. Removing the duplicates (841 papers) resulted in 830 articles for further analysis. After examining the title and abstracts of these studies, 92 articles were identified for full-text review and additional analysis. Finally, we found eight studies from 92 fully reviewed articles, which met our criteria. In [Table 1](#), we have introduced the characteristics of these studies. [Table 1](#) represents the details of methods and results, such as the lists of the models, study population, time horizon, screening techniques, the perspective of the study and outcome measure for each study, sources of cost information, type of costs, currency (type and year), discount rate, side effects, approach for expressing uncertainty and important variables in the sensitivity analysis. The results show that one study reported costs per life years saved (LYS) from CRC screening,²² six reported costs per QALYs only,^{23–28} and two reported both costs per LYS and costs per QALY.²⁹ All studies examined one or more of the available screening techniques, as well as a no-screening alternative. The perspectives of the analyses were societal or third-party payers.

Cost-Effectiveness of Common Alternative Techniques of Screening vs No-Screening

The results of the present study showed that the screening for CRC led to a decrease in the deaths from CRC in adults above 50 years of age, who were at average risk for CRC. The results demonstrated that screening by any technique was cost-effective. For one study, the cost of screening techniques (\$PPP 2028 per person to \$PPP 2428 per person) was less than the cost of no-screening (\$PPP 3580 per person).²³ In the remaining of the studies, the costs of no-screening were between \$PPP 240 per person and \$PPP 8422 per person, and for the alternative screening techniques, the costs were exceeding these ranges. In this systematic review, all the studies revealed that any type of screening strategies is more effective than the no-screening technique. The minimum ICER calculated in \$PPP was –16265/QALY (the negative ICERs were between \$PPP –16265/QALY to \$PPP –1988/QALY, whereas the positive ICERs were between \$PPP 1257/QALY to \$PPP 55987/QALY).²³ [Table 2](#) shows the details of extracted data for alternative techniques, in comparison to the no-screening strategy.

Overall, from the studies that examining only one method of screening, in four studies, the FIT was more effective and less costly than the no-screening strategy. The FIT every 2-year and FIT yearly screening techniques




Figure 1 The methods to identify studies based on the inclusion criteria.

Note: Adapted from Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097.³⁰

were more effective and less costly than the no-screening, among all techniques.

Studies also assessed the mixed method screening and found that annual FIT/COLOx1 (colonoscopy once) and FIT/sigmoidoscopy were more effective and less costly than the no-screening method (FIT/COLOx1 and FIT/sigmoidoscopy, gained 0.11 and 0.112 QALYs per person, and saved \$PPP 1396 and 1371 per person, respectively).²³

In two studies, sigmoidoscopy was dominant, compared to the no-screening technique.^{23–25} In fact, sigmoidoscopy once and sigmoidoscopy every 5 years were less costly and more effective than the no-screening method. Single sigmoidoscopy had an ICER below \$28,000/QALY, indicating the dominance of this technique to no-screening technique, in the given context-specific threshold.²⁷

Colonoscopy every 10-year, beginning at either 50 or 70 years of age, for both sexes also dominated the no-

Table 1 The Characteristics of the Included Studies

Characteristics	Wong ²²	Dinh ²³	Ladabaum ²⁴	Sharaf ²⁵	Sharp ²⁶	Dan ²⁷	James ²⁸	Wong ²⁹
Model type	Markov	Archimedes	Markov	Markov	Markov process	Markov	Markov	Markov
Study Population	GP (50 year over)	GP 50-75	GP 50, 65 and older	GP(65-71)	GP(55-74)	GP(50-75)	GP(50-100)	GP(50-70)
Time horizon	25 years	Lifetime	30 years	Lifetime	Lifetime	Lifetime	30 years	20 years
Screening technique	No-screening G-FOBT yearly and 2-year I-FOBT(FIT) yearly and 2-year Colonoscopy 10-year	No-screening FIT/ sigmoidoscopy FIT yearly/ COLOxI FIT yearly Sigmoidoscopy 5-year colonoscopy 10-year	No-screening FIT every 2 years FIT yearly MT-s DNA every 3 years Colonoscopy 10-year	No-screening FS once FOBT yearly FS every 5 years FIT yearly Colonoscopy every 10 years FS 5-year+ FIT yearly	No-screening biennial G FOBT at ages (55-74) biennial FIT at ages (55-74) FSIG once at age 60	No-screening, FIT early, barium enema 5-year, sigmoidoscopy 5-year, sigmoidoscopy 5-year+ FIT early, stool DNA every 5 years, single colonoscopy colonoscopy 10-year, colonography 10-year, FIT early+ colonoscopy 10-year	No-screening FIT yearly colonoscopy 10-year stool DNA3-year sigmoidoscopy5-year	No-screening FS5-year at ages 50-70 for male and female Colonoscopy 10-year at ages 50-60-70 for male and female FS5-year at ages 50-55 (female) FS5-year at ages 50-65 (female) FS5-year at ages 50-70 (female) +colonoscopy 10-year
Perspective	Third-party payer (healthcare service provider)	Societal	Third-party payer	Third-party payer	Third-party payer	Societal	Societal	Third-party payer (healthcare service provider)
Out-come Data sources	LYs or QALYs/ experimental studies	QALYs/ Systematic review	QALYs/ Observational and experimental studies	QALYs/ Systematic review	QALYs/ Systematic review	QALYs/Systematic review	QALYs/review of literature	LYs/experimental studies
Type of cost included	Direct	Direct	Direct	Direct	Direct	Indirect- Direct	Indirect- Direct	Direct

(Continued)

Table 1 (Continued).

Characteristics	Wong ²²	Dinh ²³	Ladabaum ²⁴	Sharaf ²⁵	Sharp ²⁶	Dan ²⁷	James ²⁸	Wong ²⁹
Source of screening cost	Hong Kong Government Printers (Government Gazette)	Medicare	Medicare payment	Medicare	DRG-2 article	Governments restructured hospitals in Singapore	Medicare and private insurance	Hong Kong Government Printers (Government Gazette)
FOBT \$PPP	NR	NR	NR	25.25	10	NR	NR	NR
FIT \$PPP	NR	2.6	19	25.25	15	33	22	NR
Barium enema	NR	NR	NR	NR	NR	115.3	NR	NR
Colonoscopy \$PPP	1351	570.3	1416	708.07	842	708.1	1075	1351
Sigmoidoscopy \$PPP	NR	184.4	NR	185.52	194	219.6	185.8	368
Sigmoidoscopy +FIT \$PPP	NR	NR	NR	NR	NR	NR	NR	NR
Stool DNA	NR	NR	661	NR	NR	329.3	NR	NR
Virtual Colonography \$PPP	NR	NR	NR	NR	550	866.15	NR	713
FIT+ Colonoscopy \$PPP	NR	NR	NR	NR	NR	NR	NR	NR
Polypectomy, \$PPP	267	172	NR	1112.05	NR	NR	NR	267
Treatment of perforation, \$PPP	15002	NR	23058	15652.14	13215	26346.7	NR	15002
Indirect costs, \$	Wong ²²	Dinh ²³	Ladabaum ²⁴	Sharaf ²⁵	Sharp ²⁶	Dan ²⁷	James ²⁸	Wong ²⁹
Screen program cost * \$PPP	NR	NR	NR	NR	NR	4.17	NR	NR
Transport and wage loss \$PPP	NR	NR	NR	NR	NR	1207.6	NR	NR
Localized, \$PPP	NR	NR	3599	2482.1	47129	14106.7	2831	23014
Regional, \$PPP	NR	NR	4795	3307.6	62229	25084.3	3080	37986

Distant, \$PPP	NR	NR	99381	78866.7	46380	46930.1	12317	99761
Side effects	Bleeding or perforation	NR	Perforation- major hemorrhage	Perforation- hemorrhage	Major bleeding- bowel perforation- deaths due to perforation	Major bleeding- Bowel Perforation- Deaths due to perforation	NR	Bleeding or perforation
Currency (type and year)	U.S dollar-2009	U.S dollar- 2010	U.S dollars 2015	U.S dollar- 2010	€ 2008	U.S dollar- 2009	U.S dollar 2014	U.S dollar-NR
Discount rate	3.5%	3%	3%	3%	4%	3%	3%	3%
Approach to expressing uncertainty	Deterministic and PSA	One-way sensitivity	One-way and multiway sensitivity	One-way sensitivity analyses	PSA and One-way sensitivity analyses	One-way and multi-way sensitivity	One-way and two-way sensitivity analyses, PSA	I-way sensitivity
Important variables in sensitivity analysis	Model parameters And outputs (compliance rates of screening, utilities, discount rate)	Compliance rates, interval between FIT screenings, performance characteristics of FIT and colonoscopy, costs of FIT, colonoscopy, and CRC treatments, and discount rate	All model inputs: costs- utility, sensitivity, specificity, side effect, interval screening	Costs of the tests, the costs of CRC care, the risk of CRC	Discount rate, costs of screening tests, costs of managing colorectal cancer	Incidence of colorectal cancer, secondary and tertiary adherence, and cost of colonoscopy, specificity and sensitivity of tests	All inputs model (specificity and sensitivity of tests, costs of the tests	Compliance rates of screening, specificity and sensitivity of tests, costs of the tests

Note: *The cost of screening program includes: screening center manpower staffing/training/case-tracking/registry/public education audit/quality control.
Abbreviations: GP, general population; FIT, fecal immunochemical test; MT-s DNA, multi-target stool DNA; FS once, flexible sigmoidoscopy once; FOBT, fecal occult blood test; QALYs, quality-adjusted life-year; LYS, life years saved; NR, not reported; PSA, probabilistic sensitivity analysis; CRC, colorectal cancer; NR, not reported.

Table 2 The Incremental Cost-Effectiveness Ratios of Different Techniques for CRC, in Contrast to the No-Screening Method

Screening Test	Study/Reference							
	Wong ²²	Dinh ²³	Ladabaum ²⁴	Sharaf ²⁵	Sharp ²⁶	Dan ²⁷	Kingsley ²⁸	Wong ²⁹
FIT 2-year	4328	-	-11391	-	1306	-	-	-
FIT yearly	4587	-16169	-7970	-7243	-	34580	-6005	-
FIT+COLOx1 ^a	-	-12695	-	-	-	-	-	-
FIT+sigmoidoscopy	-	-12243	-	-1988	-	47205	-	-
Sigmoidoscopy once	-	-	-	-11,712	1257	25981	-	-
Sigmoidoscopy 5-year	-	-	-	-3330	-	45448	-3109	40237
Colonoscopy 10-year	5265	-10400	15283	2959	-	38745	3316	22391
Single Colonoscopy	-	-	-	-	-	43911	-	-
Stool DNA	-	30014	-	-	-	48205	16046	-
Virtual colonography 5-y	-	-	-	-	-	49830	-	-

Notes: The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive.

Abbreviation: ^aCOLOx1, colonoscopy once.

screening.²⁹ Only one out of seven studies showed that performing colonoscopy every 10-year was more effective and less costly than no-screening.²³ Actually, this technique had an ICER lower than other techniques. In addition, only one out of seven studies reported a single colonoscopy technique. We found that the single colonoscopy was cost-effective, in comparison to the no-screening strategy, according to the given threshold.²⁷ The studies that reported the cost-effectiveness analysis for the newer technologies of stool DNA and virtual colonography every 5-year, revealed that both techniques were cost-effective, in comparison to the no-screening strategy.^{24,27,28}

Comparison of Different Screening Techniques

We identified eight studies, examined multiple screening methods (FOBT, FIT, colonoscopy, sigmoidoscopy and the combination of S-DNA test, FIT and sigmoidoscopy) and reached diverse conclusions about their ICER. In three studies, annual/biennial FIT was compared to annual/biennial FOBT, showing that the ICER for annual/biennial FIT in all studies was below the accepted threshold, and thus suggesting a cost-effective strategy. Also in two out of three studies, the annual FIT was more effective and less costly than annual FOBT. The analysis performed by Ladabaum et al²⁴ showed that yearly FIT was cost-effective when it was compared to FIT performed every 2 years (see Table 3). The comparisons made between annual FIT and sigmoidoscopy every 5-year showed that annual FIT was dominant in four out of five studies. In two of those studies, annual FIT showed less cost and more effectiveness, in comparison with sigmoidoscopy every 5-year.^{22,25} Only in

the analysis conducted by Dan and colleagues,²⁷ the annual FIT dominated by both single sigmoidoscopy and sigmoidoscopy every 5-year or had an ICER higher than \$50,000/per QALY (see Table 4). When FOBT was compared to sigmoidoscopy of every 5-year and sigmoidoscopy once, it was observed that all the studies had an ICER below \$50000/per QALY. In the study of Sharaf et al²⁵ the annual FOBT was more effective and less costly than sigmoidoscopy once (see Table 5). When sigmoidoscopy every 10-year or sigmoidoscopy every 5-year plus annual FIT was compared to FIT yearly, it was revealed that in four out of five studies the sigmoidoscopy every 10-year or sigmoidoscopy every 5-year plus annual FIT had an ICER of above \$50000/per QALY and was dominated (see Table 6).^{25,27} Colonoscopy every 10-year in one out of five studies was cost-effective, in comparison with the FIT yearly and annual FOBT. A study by Wong et al represented that colonoscopy every 10-year was more effective and less costly than annual FOBT (see Table 7).²² Also in one out of two comparisons, colonoscopy every 10-year was dominant, compared to the sigmoidoscopy every 5-year plus annual FIT.²⁷ However, in another study that sigmoidoscopy every 5-year plus annual FIT considered as a base technique, and was compared to colonoscopy every 10-year, we observed less cost and more effectiveness, in comparison to the colonoscopy every 10-year (see Table 8).²⁵

When colonoscopy every 10-year was compared with S-DNA test (3-year and 5-year), colonoscopy test had an ICER below \$50000/per QALY and was dominant with 100% of the certainty. Only one study showed that the virtual colonography every 5-year was less cost-effective when compared to the established technique (see Table 9).²⁷

Table 3 The Calculated Incremental Cost-Effectiveness Ratios When Using FIT Yearly vs FIT Every 2-Year, and FOBT Yearly

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Sharaf²⁵ FIT yearly FOBT yearly	2090 2187	18.7456 18.7352	-9327	\$US 50000	FIT yearly was dominant
Sharp²⁶ FIT 2-yearly FOBT 2-yearly	1443 1434	10.984 10.968	563	WHO Recommendation	FIT 2-yearly was cost-effective ^a
Wong²² FIT yearly FOBT yearly	7371 7845	15.5491 15.2339	-1504	\$US 50000	FIT yearly was dominant
Wong²² FIT 2-yearly FOBT 2-yearly	6606 6136	15.4203 15.0687	1328	\$US 50000	FIT 2-yearly was cost-effective
Ladabaum²⁴ FIT yearly FIT 2-year	2450 2251	18.747 18.741	33167	\$US 50000	FIT yearly was cost-effective

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, presented in the second row. The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; FOBT, fecal occult blood testing; FIT, fecal immunochemical testing.

Table 4 The Calculated Incremental Cost-Effectiveness Ratios When Using FIT Alone vs Flexible Sigmoidoscopy (FS)

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Dinh²³ FIT yearly FS 5-year	2028.2 2425.6	15.771 15.746	-15896	\$US 50000	FIT yearly was dominant
Sharaf²⁵ FIT yearly FS 5-year	2090 2419	18.7456 18.7372	-39167	\$US 50000	FIT yearly was dominant
Dan²⁷ FIT yearly Single FS	378.73 318.36	16.393 16.392	60370	\$US 50000	FIT yearly was dominated
Kingsley²⁸ FIT yearly FS 5-year	1607 1914	19.506 19.479	-11370	\$US 50000	FIT yearly was dominant
Sharaf²⁵ FIT yearly FS once	2090 2341	18.7372 18.6948	-5920	\$US 50000	FIT yearly was dominant
Dan²⁷ FIT yearly FS 5-year	378.73 467.65	16.393 16.394	88920	\$US 50000	FIT yearly was dominated

Notes: In this table, for each study the first row shows the new technique that is compared with the old technique, written in the second row. The negative ICER indicates that the new technique is less costly and more effective.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; FOBT, fecal occult blood testing; FIT, fecal immunochemical testing; FS 5-year, flexible sigmoidoscopy every 5 years; FS once flexible sigmoidoscopy once in lifetime.

Table 5 The Calculated Incremental Cost-Effectiveness Ratios When Using FOBT Alone vs Flexible Sigmoidoscopy (FS)

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Sharaf²⁵ FOBT yearly Sigmoidoscopy 5-year	2178 2419	18.7352 18.7372	120500	\$US 50000	FOBT yearly was dominated
Sharp²⁶ FOBT 2-year Sigmoidoscopy once	1434 1395	10.968 10.966	19500	WHO Recommendation	FOBT 2-year was cost-effective ^a
Sharaf²⁵ FOBT yearly (FS Once)Sigmoidoscopy once	2178 2341	18.7352 18.6948	-4035	\$US 50000	FOBT yearly was dominant

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, written in the second row. The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; FOBT, fecal occult blood testing yearly; FIT, fecal immunochemical testing yearly; FS once, flexible sigmoidoscopy once in lifetime.

Table 6 The Calculated Incremental Cost-Effectiveness Ratios When Using FIT+ Sigmoidoscopy vs FIT

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Dinh²³ FIT early+ sigmoidoscopy 5-year FIT yearly	2209.1 2028.2	15.787 15.771	11306	\$US 50000	FIT early+ FS 5-year was cost-effective ^a
Sharaf²⁵ FS 5-year + FIT yearly FIT yearly	2492 2090	18.7469 18.7456	309231	\$US 50000	FS 5-year + FIT yearly was dominated
Dan²⁷ FS 5-year + IFOBT yearly IFOBT yearly	570.85 378.73	16.396 16.393	64040	\$US 50000	FS 5-year + IFOBT yearly was dominated

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, written in the second row. The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; FOBT, fecal occult blood testing; FIT, fecal immunochemical testing.

Discussion

There are multiple methods, policies, and interventions for screening for CRC. These methods are used in combination or alone. These mean that we have numerous strategies for the screening of CRC, and evaluating of those strategies in terms of costs and effectiveness is, to some extent, a complex task.

As the results of this systematic review demonstrated, the CRC screening by any technique is cost-effective, in comparison with the no-screening method.

Three studies reported the cost-effectiveness analysis for the S-DNA test. The results showed that the S-DNA test was cost-effective, in comparison to the no-screening. However, this technique was less cost-effective when compared with other screening techniques (namely,

colonoscopy every 10-year, barium enema 5-year, sigmoidoscopy, FIT and virtual colonography every 5-year). In our systematic review, only one study reported that the virtual colonography every 5-year was cost-effective, in comparison to the no-screening method. However, the virtual colonography every 5-year was not cost-effective when it was compared to other techniques.²⁷ Virtual colonography every 5-year is a non-invasive procedure that potentially can be ideal for subjects who avoid invasive procedures, such as colonoscopy and FS. Although this technique has advantages, such as a reduction in complications of colonoscopy (bowel perforation, major bleeding, and deaths due to perforation), more resources need to be allocated to get the same effectiveness, in comparison with the colonoscopy.

Table 7 The Calculated Incremental Cost-Effectiveness Ratios When Using Colonoscopy vs FIT and FOBT

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Dinh²³ Colonoscopy 10-year FIT yearly	2384.3 2028.2	15.79 15.77	17805	\$US 50000	Colonoscopy 10-year was cost-effective ^a
Ladabaum²⁴ Colonoscopy 10-year FIT 2-year	4248 2251	18.746 18.741	443778	\$US 50000	Colonoscopy 10-year was dominated
Sharaf²⁵ Colonoscopy 10-year FOBT yearly	2871 2187	18.7443 18.7352	75165	\$US 50000	Colonoscopy 10-year was dominated
Wong²² Colonoscopy 10-year G-FOBT yearly	6911 7845	15.3586 15.2339	-7490	\$US 50000	Colonoscopy 10-year was dominant
Kingsley²⁸ Colonoscopy 10-year FIT yearly	2474 1607	19.517 19.506	78818	\$US 50000	Colonoscopy 10-year was dominated

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, written in the second row. The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; G-FOBT, guaiac fecal occult blood testing; FIT, fecal immunochemical testing.

Table 8 The Calculated Incremental Cost-Effectiveness Ratios When Using Colonoscopy vs FIT + Sigmoidoscopy

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Dinh²³ Colonoscopy 10 year FIT yearly+ sigmoidoscopy 5-year	2384.3 2209.1	15.79 15.787	58400	\$US 50000	Colonoscopy 10-year was dominated
Dan²⁷ Colonoscopy 10-year Sigmoidoscopy 5-year + IFOBT yearly	899.08 570.85	16.406 16.396	32823	\$US 50000	Colonoscopy 10-year was cost-effective ^a

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, written in the second row. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; IFOBT, fecal occult blood testing; FIT, fecal immunochemical testing.

Table 9 The Calculated Incremental Cost-Effectiveness Ratios When Using Colonoscopy vs Stool DNA

Author/Technique	C	Q	ICER	Threshold	Optimal Strategy
Ladabaum²⁴ Colonoscopy 10-year MT-s DNA 3-year	4248 5283	18.7455 18.7423	-323438	\$US 50000	Colonoscopy 10-year was dominant
Dan²⁷ Colonoscopy 10-year Stool DNA 5-year	899.08 614.76	16.406 16.396	28432	\$US 50000	Colonoscopy 10-year was cost-effective ^a
Kingsley²⁸ Colonoscopy 10-year Stool DNA 3-year	2474 3590	19.517 19.507	-111600	\$US 50000	Colonoscopy 10-year was dominant

Notes: In this table, for each study, the first row shows the new technique that is compared with the old technique, written in the second row. The negative ICER indicates that the new technique is less costly and more effective. The positive ICERs refer to those techniques that are more costly and more effective; both numerators and denominators are positive. ^aBased on the context threshold.

Abbreviations: C, costs per person (\$PPP); Q, QALYs (quality-adjusted life-year); ICER, incremental cost-effectiveness ratio; MT-S DNA 3-year, multi-target stool DNA every 3-year.

Based on our results, there was no consensus about starting and endpoint screening ages, but it seems that the age of 50 years was the most appropriate for initiation. Kingsley et al²⁸ examined the starting and stopping ages of screening in their analysis from the age 50 until age 100 or death, but in conclusion, they did not recommend any type of screening for people above the age of 80. The prevalence and incidence rates of CRC at different ages determine which screening technique is the most cost-effective.

In this study, two out of eight studies had reported the prevalence rate of CRC at age 50, and the decrease of CRC incidence in different screening techniques had been discussed. However, there has been no examination of the prevalence rate of CRC at different ages and the choice of age-appropriate screening techniques, which have an impact on the cost-effectiveness of screening techniques.

Overall, changes in the sensitivity and specificity of screening tests will have a different effect on the cost-effectiveness of CRC screening techniques. In the current review, four out of eight studies have demonstrated that the changes in sensitivity and specificity of the screening tests have a moderate impact on ICER, and for the other studies, this impact is minimal. For instance, in the study of Dan et al²⁷ it is shown that when the specificity and sensitivity of CRC screening tests varied from 50% to 99%, moderate impacts were observed on the cost-effectiveness of CRC screening techniques. Thereby, screening techniques may be dominated or may no longer be cost-effective.

Although most of the models used in the studies are the same (Markov model), due to the differences in cost and effectiveness of screening strategies such as time horizon, variety of techniques in different studies, differences in sensitivity and specificity rate, it is not feasible in practice to determine which strategy is the optimal technique. Dinh et al and Kingsley et al analyzed the colonoscopy but the cost of complications was not included in the model.^{23,28} These factors may be one of the main reasons, leading to inconsistency in the results of the different studies.

In addition, our findings showed that all studies conducted the sensitivity analysis with alternative assumptions to examine the uncertainty with parameters. Five studies used the Monte Carlo simulation to investigate the uncertainty of the model. Uncertainty among various parameters was assessed in each study by one-way, multi-way, and probabilistic sensitivity analysis (PSA). In 37.5% of cases with sensitivity analysis, there was no change in the results or the effect of change in parameters was not significant. For instance, in the study by Dinh et al²³ changes in the

variables had not had a significant effect on the ICER, when the FIT/COLOx1 was compared to FIT and COLO.

In 62.5% of the studies, it was found that changing the parameters had a significant effect on the results. A study conducted by Dan and colleagues showed that the risk of CRC and the cost of colonoscopy accompanied by uncertainty.²⁷ When the cost of colonoscopy was less than \$300 regardless of the risk of CRC, colonoscopy was the dominant technique. When the cost of colonoscopy was above the \$300, the IFOBT was considered the dominant technique in lower incidence levels of CRC, whereas, for the higher incidence level, the sigmoidoscopy revealed to be more cost-effective than the other techniques.

Limitation

One limitation when evaluating the cost-effectiveness and costing is that studies take place in different countries with different contexts, prices, and costs and at different times. Therefore, these need to be considered when decision-makers interested to use the overall results or some components of the cost-effectiveness analyses for different settings and times. It is recommended that in order to compare and value the cost differences over time, it should be adjusted by the inflation rate, and across countries should be adjusted by PPP. These adjustments are apart from the adjustment required for the differences in conducting processes. The magnitude of indirect costs in cancers is outstanding, and theoretically, the sum of the direct and indirect costs is almost a reflection of the cost of opportunity lost due to cancer.

Even though indirect costs constitute a significant part of the cancer costs and it is about a societal perspective, but only few studies included these types of costs in their analyses. This leads to an underestimation of the cancer costs and heterogeneity arises when there is an intention for comparing the results. In this review, we found that there are also other heterogeneous parameters such as time horizons, perspectives, and types of the models and included states that make it difficult to compare various outcomes one by one in different contexts.

One of the less paid subjects in a cost-effectiveness analysis is the threshold level for different countries. There is a general agreement that the threshold level for each country should be context-specific, but there is a discrepancy among the methods used for the implementation of the thresholds for different health systems. Furthermore, even for the same health systems, the threshold is kept constant for quite a long time. In some

countries, it might be necessary to adjust the threshold in relation to the inflation rate.

Since the feasibility and acceptability of using screening tests according to the context of the study, is one of the important factors, but it was discussed in just one out of eight studies. For example, colonoscopy is an invasive method for screening and many people do not prefer to use it in the first step. Therefore, future studies should examine the feasibility of using screening tests and consider patient preference in the selection of alternative tests.

Conclusion

Our review showed that all CRC screening techniques are cost-effective when compared with no-screening, but there is no agreement between the results of the various studies to determine the optimal technique. The newer technologies of virtual colonography and S-DNA were not cost-effective, compared with conventional techniques (FIT, sigmoidoscopy, and conventional colonoscopy). Although both techniques were less cost-effective than other techniques, S-DNA and virtual colonography are non-invasive procedures, making them a potentially ideal choice for subjects who like to avoid any invasive procedures. Finally, additional analyses are necessary to determine the optimal technique.

To compare and utilize the results of the different studies, there should be some observations like inflation, PPP, threshold levels and the subject preference for accepting the interventions.

Acknowledgments

We extend our sincere thanks to all authors who responded to our communications and shared with us their original data and documents. This paper was extracted from the master's thesis of FK, supervised by MY and supported by Tabriz University of Medical Sciences (Approval ID: IR.TBZMED.REC.1396.936)

Disclosure

The authors report no conflicts of interest in this work.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394–424. doi:10.3322/caac.21492
- Vekic B, Dragojevic-Simic V, Jakovljevic M, et al. A correlation study of the colorectal cancer statistics and economic indicators in selected Balkan countries. *Front Public Health*. 2020;8:29. doi:10.3389/fpubh.2020.00029
- Albrecht T, Conroy F, Dalmas M, et al. *European Guide for Quality National Cancer Control Programmes*. Ljubljana, Slovenia: National Institute of Public Health; 2015.
- European Council. Council recommendation of 2 December 2003 on cancer screening (2003/878/EC). *Off J Eur Union*. 2003;327.
- Keshvari-Shad F, Hajebrahimi S, Laguna Pes M, et al. A systematic review of screening tests for chronic kidney disease: an accuracy analysis. *Galen Medical J*. 2020;9:e1573. doi:10.31661/gmj.v9i0.1573
- Schoenfeld P. Quality in colorectal cancer screening with colonoscopy. *Gastrointest Endosc Clin N Am*. 2020;30(3):541–551. doi:10.1016/j.giec.2020.02.014
- Ouyang DL, Getzenberg RH, Schoen RE. Noninvasive testing for colorectal cancer: a review. *Am J Gastroenterol*. 2005;100(6):1393–1403. doi:10.1111/j.1572-0241.2005.41427.x
- Redwood D, Asay E, Roberts D, et al. Comparison of fecal occult blood tests for colorectal cancer screening in an Alaska Native population with high prevalence of *Helicobacter pylori* infection, 2008–2012. *Prev Chronic Dis*. 2014;11. doi:10.5888/pcd11.130281
- Imperiale TF, Itzkowitz SH, Levin TR, et al. Multitarget stool DNA testing for colorectal-cancer screening. *N Engl J Med*. 2014;370(14):1287–1297. doi:10.1056/NEJMoa1311194
- Administration USFaD. FDA approves first non-invasive DNA screening test for colorectal cancer. 2014. Available from: <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm409021.htm>.
- Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4:1. doi:10.1186/2046-4053-4-1
- Cochrane handbook for systematic reviews of interventions [Internet]. 2018. Available from: <http://handbook-5-1.cochrane.org/webcite>.
- International prospective register of systematic reviews. 11 February 2018. ID:CRD42018081676V. Available from: <https://www.crd.york.ac.uk/prospero/>.
- Smith H, Varshoei P, Boushey R, Kuziemyky C. Use of simulation modeling to inform decision making for health care systems and policy in colorectal cancer screening: protocol for a systematic review. *JMIR Res Protoc*. 2020;9(5):e16103–e16103. doi:10.2196/16103
- Heydari M, Yousefi M, Derakhshani N, Khodayari-Zarnaq R. Factors affecting the satisfaction of medical tourists: a systematic review. *Health Scope*. 2019;8(3):e80359. doi:10.5812/jhealthscope.80359
- Pignone M, Saha S, Hoerger T, Mandelblatt J. Cost-effectiveness analyses of colorectal cancer screening: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2002;137:96–104.
- National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology (NCCN guidelines). *CNS Cancer Ver*. 2011;2:19–21.
- Table W. Threshold values for intervention cost-effectiveness by region. 2014. Available from: http://www.who.int/choice/costs/CER_levels/en/.
- Ladabaum U. Cost-effectiveness of current colorectal cancer screening tests. *Gastrointest Endosc Clin N Am*. 2020;30(3):479–497. doi:10.1016/j.giec.2020.02.005
- Institute for Clinical and Economic Review. Overview of the ICER value assessment framework and update for 2017–2019. 2018.
- Mf D. *Methods for the Economic Evaluation of Health Care Programmes*. 3rd ed. New York: Oxford University Press; 2005.
- Wong CK, Lam CL, Wan YF, Fong DY. Cost-effectiveness simulation and analysis of colorectal cancer screening in Hong Kong Chinese population: comparison amongst colonoscopy, guaiac and immunologic fecal occult blood testing. *BMC Cancer*. 2015;15:705. doi:10.1186/s12885-015-1730-y
- Dinh T, Ladabaum U, Alperin P, Caldwell C, Smith R, Levin TR. Health benefits and cost-effectiveness of a hybrid screening strategy for colorectal cancer. *Clin Gastroenterol Hepatol*. 2013;11(9):1158–1166. doi:10.1016/j.cgh.2013.03.013

24. Ladabaum U, Mannalithara A. Comparative effectiveness and cost effectiveness of a multitarget stool DNA test to screen for colorectal neoplasia. *Gastroenterology*. 2016;151:427–439. doi:10.1053/j.gastro.2016.06.003
25. Sharaf RN, Ladabaum U. Comparative effectiveness and cost-effectiveness of screening colonoscopy vs. sigmoidoscopy and alternative strategies. *Am J Gastroenterol*. 2013;108:120–132. doi:10.1038/ajg.2012.380
26. Sharp L, Tilson L, Whyte S, et al. Cost-effectiveness of population-based screening for colorectal cancer: a comparison of guaiac-based faecal occult blood testing, faecal immunochemical testing and flexible sigmoidoscopy. *Br J Cancer*. 2012;106:805–816. doi:10.1038/bjc.2011.580
27. Dan YY, Chuah BY, Koh DC, Yeoh KG. Screening based on risk for colorectal cancer is the most cost-effective approach. *Clin Gastroenterol Hepatol*. 2012;10(3):266–271. doi:10.1016/j.cgh.2011.11.011
28. Kingsley J, Karanth S, Revere FL, Agrawal D. Cost effectiveness of screening colonoscopy depends on adequate bowel preparation rates - A modeling study. *PLoS One*. 2016;11(12):e0167452. doi:10.1371/journal.pone.0167452
29. Wong MC, Ching JY, Chan VC, et al. Colorectal cancer screening based on age and gender: a cost-effectiveness analysis. *Medicine*. 2016;95(10):e2739. doi:10.1097/MD.0000000000002739
30. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med* 6(7): e1000097

Risk Management and Healthcare Policy

Dovepress

Publish your work in this journal

Risk Management and Healthcare Policy is an international, peer-reviewed, open access journal focusing on all aspects of public health, policy, and preventative measures to promote good health and improve morbidity and mortality in the population. The journal welcomes submitted papers covering original research, basic science, clinical & epidemiological studies, reviews and evaluations,

guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/risk-management-and-healthcare-policy-journal>