



Published in final edited form as:

Nat Geosci. 2018 May 1; 11(5): 328–333. doi:10.1038/s41561-018-0092-x.

“Reduction of tree cover in West African woodlands and promotion in semi-arid farmlands”

Martin Brandt^{*1}, Kjeld Rasmussen¹, Pierre Hiernaux², Stefanie Herrmann³, Compton J. Tucker⁴, Xiaoye Tong¹, Feng Tian¹, Ole Mertz¹, Laurent Kergoat², Cheikh Mbow⁵, John David⁴, Katherine Melocik⁴, Morgane Dendoncker⁶, Caroline Vincke⁶, Rasmus Fensholt¹

¹Department of Geosciences and Natural Resource Management, University of Copenhagen, 1350 Copenhagen, Denmark.

²Geosciences Environnement Toulouse (GET), Observatoire Midi-Pyrénées, UMR 5563 (CNRS/UPS/IRD/CNES), 14 Avenue Edouard Belin, 31400 Toulouse, France.

³Agricultural and Biosystems Engineering, The University of Arizona, 1177 E. 4th Street, Tucson, AZ 85721, USA.

⁴NASA Goddard Space Flight Center, Mail Code 610.9, Greenbelt, MD 20771, USA.

⁵START International Inc., 2000 Florida Ave NW, Washington, DC 20009, USA.

⁶Université catholique de Louvain, Earth and Life Institute, Environmental Sciences, Croix du Sud 2 L7.05.09, 1348 Louvain-la-Neuve, Belgium.

Abstract

Woody vegetation in farmland acts as a carbon sink and provides ecosystem services for local people, but no macro-scale assessments of the impact of management and climate on woody cover exists for drylands. Here we make use of very high spatial resolution satellite imagery to derive wall-to-wall woody cover patterns in tropical West African drylands. Our study reveals a consistently high woody cover in farmlands along all semi-arid and sub-humid rainfall zones (16%), on average only 6% lower than in savannas. In semi-arid Sahel, farmland management increases woody cover to a greater level (12%) than found in neighbouring savannas (6%), whereas farmlands in sub-humid zones have a reduced woody cover (20%) as compared to savannas (30%). In the region as a whole, rainfall, terrain and soil are the most important (80%) determinants of woody cover, while management factors play a smaller (20%) role. We conclude

*Materials & Correspondence. Correspondence and material requests should be addressed to M.B. (mabr@ign.ku.dk).

Author Contributions. M.B., R.F., S.H., P.H. and K.R. designed the study. M.B., X.T. and F.T. conducted the analyses with support by LK, OM, KR, RF, SH, MD, PH. The data was provided by CT, JD, KM, MD, LK, CV and PH. KR and MB drafted the manuscript with contributions by all authors.

Competing financial interest. The authors declare no competing financial interests.

Data availability. Commercial very high resolution satellite images were acquired within the NextView license program. The copyright remains at DigitalGlobe and a redistribution is not possible. PROBA-V NDVI data is freely available at VITO (<http://proba-v.vgt.vito.be/>). Worldpop population data is freely available at the University of Southampton (<http://www.worldpop.org.uk/>). MODIS MCD45A1 burned area product is can be freely obtained at <http://modis-fire.umd.edu/pages/news.php>. The soil map is freely available at ISRIC (<http://www.isric.org/content/african-soilgrids-250m-geotiffs>). CHIRPS rainfall data is freely available at the Climate Hazard Group (<http://chg.geog.ucsb.edu/data/chirps/>). PALSAR mosaics are freely available from JAXA (http://www.eorc.jaxa.jp/ALOS/en/palsar_fnf/fnf_index.htm). The farmland mask is available from Marie-Julie Lambert upon request. The woody cover map at 100 m resolution is available from the corresponding author upon request.

that agricultural expansion cannot generally be claimed to cause woody cover losses, and that observations in Sahel contradict simplistic ideas of a high negative correlation between population density and woody cover.

INTRODUCTION

Concerns about declining woody cover in West Africa have been raised since the early 20th century^{1,2}. In the 1970s and 80s, negative trends in woody vegetation, presumably associated with the ‘Sahel drought’ and agricultural expansion, were observed and became part of the desertification/land degradation discourse, later termed the ‘Sahel syndrome’³. Rapidly growing settlements and urban markets demanded large amounts of firewood and charcoal, and concerns about an upcoming ‘fuelwood crisis’ were widespread⁴. Certain parts of the Sahel experienced an increase in export-oriented agriculture (e.g. groundnut production in Senegal and cotton production in Mali), which was understood to have contributed to a downward trend in woody cover as well⁵. All these concerns had substantial impact on natural resource policies of the Sahelian countries and the donors supporting them: New forests were planted (e.g. “shelterbelts” in northern Nigeria, village wood-lots in Mali and Burkina Faso) and new attempts were made to regulate firewood harvesting and charcoal production⁶. Grand schemes of ‘green belts’ across the Sahel, already suggested before the 2nd World War by Stebbing¹, were taken up again. However, from the 1980s and onwards, research by botanists⁷, foresters⁸, geographers^{9–12} and anthropologists¹³ painted a more complex picture on the relationship between humans and woody vegetation: Studies at village and landscape scales showed that increase and decrease in woody cover occurred simultaneously in different parts of the Sahel^{7,11,14}. The ‘case study’ character of this research, however, made it difficult to generalize findings, since the representativeness for the larger region was difficult to establish¹⁵.

The idea of a progressing land degradation in arid and semi-arid West Africa was also challenged from another side: Regional-scale analyses of time series of vegetation indices derived from different satellite systems showed that fluctuations of the Sahara desert boundary are common¹⁶ and that the Sahel was experiencing a ‘re-greening’ after the drought years of the 1970s and 1980s¹⁷. These studies did not, however, allow separation of the contributions from the herbaceous and woody vegetation components. Only recently has this been achieved^{18,19} revealing that the greening may be partly attributed to an increase in woody cover. The coarse spatial and limited temporal resolution of the satellite images used and the complexities of the methods applied imply that such assessments of vegetation change in the Sahel do not necessarily form a robust basis for estimating trends in woody cover locally, and leave considerable room for speculations regarding the nature of the woody vegetation changes. Attempts to produce global maps^{20,21} of tree cover focus mainly on forests in humid areas and yield unrealistically low canopy cover estimates in drylands, which are thus commonly ignored in woody vegetation assessments²². These obstacles have made it difficult to study linkages between woody vegetation, rainfall and humans for West African farmlands and savannas - knowledge that is essential in the face of demographic and climatic change.

The recent access to large volumes of DigitalGlobe, Inc. commercial satellite images with a spatial resolution as low as 0.3 m in the panchromatic band marks a technical tipping point in dryland research²³ and allows us to produce a reliable, fine-scaled assessment of woody cover²⁴. While the short period for which these data have been available does not allow to estimate long term trends, the high level of detail of such maps makes it possible to analyze how woody cover is spatially correlated with the above-mentioned causal factors, from which explanations for changes in woody cover over time can be inferred: if woody cover is threatened by the expansion of cultivation, we would expect woody cover to be substantially lower in farmlands than in the adjacent uncultivated savannas. If local harvesting of firewood is a cause of loss of woody cover, we would expect woody cover to be lowest close to settlements. Here we test these hypotheses in order to obtain a complete understanding of the distribution of woody cover in relation to human presence and thus provide a valuable reference for individual case studies that generate in-depth contextual knowledge but have a limited scope for generalization.

RESULTS

High resolution woody cover mapping.

The assessment of woody vegetation at hectare level requires high spatial resolution satellite data in order to highlight nuanced spatial differences (Supplementary Fig. 1). Here we derived canopy cover from multispectral DigitalGlobe QuickBird-2, GeoEye-1 and WorldView-2 satellite images at 1.7 m resolution without using the panchromatic band (Fig. 1, Supplementary Figs 2, 3) to train Synthetic Aperture Radar (SAR) and Normalized Difference Vegetation Index (NDVI) imagery and predict continuous woody cover from 0 to 100% at 100 m resolution for the arid (150–300 mm rainfall), semi-arid (300–600 mm) and sub-humid (600–1000 mm) zones of West Africa. The validation pixels are fairly in line with the prediction (Mean Absolute Error (MAE) of 3.7, $r=0.69$, slope=0.84, $n=661,708$; Supplementary Figs 4,5) which also agrees well with independent *in situ* data (Fig. 1b,c). The woody cover maps shown in Fig. 2 reveal a broad scale pattern following the biogeographical regions but also a high level of detail showing differences at hectare scale. Woody cover is on average 3% in the arid zone, increases to 9% in the semi-arid, and exceeds 20% in the sub-humid zone (Fig. 2).

Determinants of woody vegetation cover.

The coexistence of herbaceous and woody plants in savannas is governed by rainfall regime (mediated by run-off and water table), soil, human management (including cutting, clearing for cropping, crop-fallow management, fire and grazing)²⁵. These factors are interlinked and vary both spatially and temporally with available rainfall (Fig. 3a). Here we tested environmental variables in a decision tree ensemble model, which explained in total 67% of the predicted woody cover at 100 m resolution (Fig. 3b). Out of these, mean annual rainfall²⁶ is the major factor limiting woody cover (32%). It is followed by terrain (elevation, 23%) and human population density²⁷ is ranked third (13%), shortly before soil²⁸ (sand fraction, 12%) and inter-annual rainfall variability (12%). Distance to villages (6%) and fire frequency (2%) have a rather low relative weight. Taken together, climatic (44%) and edaphic (35%) factors are more important than management factors (21%) (Fig. 3b).

Elevation here is used to represent the terrain morphology including dune structures, depressions, plateaus, valleys, etc. Already a moderate topography can have significant impact on rainfall run off/on and soil texture, explaining the high percentage explained by terrain. A land use and rainfall zone grouping is conducted to further explore the relationships between humans and woody cover and to rule out a bias by the rainfall gradient (Fig. 3c).

Rural management impacts on woody cover.

We applied a new farmland mask at 100 m resolution²⁹ to separate the study area in uncultivated savannas and farmland (Fig. 4a-c). For savannas, there is a high positive correlation between woody cover and rainfall ($r=0.75$, $P<0.05$) with saturation around 30% canopy cover in the sub-humid zone, and with considerable spatial variations (Fig 4b). The pattern is strikingly different for farmlands (Fig. 4c): Although woody cover increases with increasing rainfall ($r=0.45$, $P<0.05$), the majority of the cultivated areas have a canopy cover of around 12%, independent of rainfall, and variability is much lower than in savannas (Fig. 4a-c). Average woody cover in arid and semi-arid Sahel is higher and less variable in farmlands (arid: 3%, semi-arid: 11%) than in savannas (arid: 2%, semi-arid: 9%). Sub-humid savannas on average have a higher woody cover (33%) and wider range than woody cover in farmlands (23%) (Figs 3c, 4a). More precisely, the median of farmland woody cover is higher as compared to savannas below 680 mm annual rainfall but lower from 680 to 1000 mm (Fig. 4a-c).

In the sub-humid zone, woody cover reaches high values primarily in rural areas with low population density, and decreases in urban areas with >100 persons km^{-2} (Fig. 4d). Interestingly, a different pattern is observed in the arid and semi-arid Sahel, where both woody cover and population density are increasing along the rainfall gradient up to 160 persons km^{-2} . Woody cover decreases at higher population densities in and around larger cities. On average, areas with a higher population density also have a higher woody cover than sparsely populated areas in the arid (7/2%) and semi-arid (12/10%) Sahel, but the opposite is observed in the sub-humid zone (31/21%) (Figs. 3c, 4a-c).

Woody cover in conservation areas is generally higher (29%) in comparison to surrounding areas (5 km) (21%) (Fig 3c). This difference is most pronounced in the semi-arid Sahel (conservation 16%; conservation surroundings 11%) and sub-humid zone (conservation 35%; conservation surroundings 23%). Differences between farmland (typically occupying sandy soils) and savannas (including vast areas of non-arable soils) become more comparable and exclude a bias by environmental pre-conditions when studying woody vegetation on sandy soils only²⁸. Sandy soils used for cultivation have remarkably higher woody cover than comparable sandy soils which are uncultivated (Fig. 4e). Buffer zones were drawn around 37,294 villages on sandy soils (Supplementary Fig. 6). Shade trees are responsible for a high canopy cover in the village centers (~12%), and areas surrounding villages within a distance up to 1.5 km have a moderately high woody cover (7–9%) which decreases gradually further away (<5%).

DISCUSSION

The traditional assumption that human presence has an exclusively adverse impact on West Africa's woody vegetation has been challenged by local studies showing that human presence can also have positive impacts on tree cover¹³, as in the case of agroforestry systems encouraging and maintaining high tree densities³⁰. Farmers' awareness of reforestation as a climate change adaptation measure has been shown³¹, and farmer managed natural regeneration or tree planting programs are common throughout West Africa. However, there regional assessments of their success are rare, and our study shows that farmlands indeed support significant woody vegetation densities, supporting the results of³². However, this is not the case in all landscapes and under all agricultural management regimes. The expansion of farmland leads to an initial reduction of woody vegetation, especially in higher rainfall zones with dense human population where savanna and woodland woody cover is dense¹⁰. If the rural population is dense, this expansion is ongoing, and forest reserves and savannas are being progressively reduced and converted into farmland, with no woodland vegetation left except in protected areas. It has also been proposed that the recent increase in woody vegetation, which is a global phenomenon in semi-arid lands supposedly driven by climate and altered atmospheric CO₂^{33,34}, often takes place in sparsely populated regions whereas high population growth decreases woody cover³⁵. However, our current study shows that this is not always the case, and once savannas and woodlands are transformed into farmland, management often aims at promoting and protecting valuable species (e.g. *Faidherbia albida*, *Vitellaria paradoxa*) by clearing/coppicing other species which also favours the growth of a few tall trees. Additionally, shade trees in village areas (e.g. *Azadirachta indica*) provide numerous ecosystem services which are more valuable for the local people³⁶ than those of typical savanna species (e.g. *Combretum glutinosum*, *Guiera senegalensis*) and also contribute to carbon storage at landscape scales.

The results presented allow a robust generalization concerning woody cover and the relationships between woody cover and various explanatory factors. First of all, we describe rainfall as the main determinant of woody cover. We confirm increases in woody cover in arid and semi-arid Sahel with rainfall up to ~650 mm³⁷. The median woody cover stabilizes in the sub-humid zone (650 – 1000 mm) around 30% woody cover.

Secondly, and most importantly, we show that the role of climate is modified by humans. The way management affects woody cover relates to the amount of annual rainfall and livelihood strategy: The median woody cover in arid and semi-arid zones is equal and partially higher in farmlands than in savannas up to an average annual rainfall of around 650 mm year⁻¹. In sub-humid zones, this difference is reversed, with median woody cover being lower in farmlands than in savannas. Unlike the rainfall driven gradient of woody cover found in savannas, the woody cover in farmlands is spatially homogeneous (constant median, narrow range) across all rainfall zones. Local studies are likely to show considerable differences between countries and eco-regions, but on average the claim that cultivated areas in the arid and semi-arid Sahel have a relatively high woody cover compared to savannas is robust. Two possible explanations may be suggested: (1) Farmers protect or plant trees due to a strong interest in the ecological services they provide³⁶. Harvesting of wood for fuel and

building material mostly takes place further away from the village areas in uncultivated land (and in fallows, which are here classified as farmland). (2) Farmland is generally located on the most suitable and fertile soils, whereas savannas also includes soil conditions less favorable for vegetation growth. Our results show, however, that the difference is still clear and even more evident when comparing only areas of sandy soils in both the cultivated and non-cultivated areas, so the latter explanation does not affect our conclusions.

Thirdly, analysis of the effect of proximity to villages on woody cover discloses that woody cover is, on average, densest within village areas and decreases with distance. This is based on a great number of villages that are very different in size and structure and this distance-function may differ depending on village size, rainfall level, agricultural practices and ethnicity of the population. Yet, at the regional scale it is clearly demonstrated that the idea that high local population pressure causes woody cover to decrease around villages does not hold true. Rather, the alternative notion that farmers protect or plant trees in and around villages¹³ is supported. The cause of a dense woody cover around villages is related to the above mentioned finding that farmlands have a relatively high woody cover. Fields are often located close to villages, while more distant savannas are mainly exploited for fuelwood. Our results showing a positive relationship between population density and woody cover seems to support the ‘more people, less erosion’ argument³⁸ of environmental recovery and sustainability associated with agricultural intensification. However, this only holds true in semi-arid areas and only up to a certain threshold of population agglomeration, i.e. at rural village level but not for larger urban settlements.

With an average canopy cover of $13 \pm 17\%$, we found substantially higher values (including larger variations) than other studies and data sets (e.g. $1.9 \pm 3\%$ in MODIS continuous fields²⁰). It has to be taken into consideration that our definition of canopy cover is more inclusive, since we include scattered woody vegetation, whereas the MODIS product is limited to forests with large sized trees. Studies based on these data sets²² are thus unable to provide detailed assessments of patterns and determinants of dryland woody cover.

The data and methods we used do not allow us to move beyond ‘woody cover’, which is the simple projected coverage of canopies. For many research applications additional variables would be of interest. From a botanical and ecological perspective, information on species would be desirable; from a climate change point of view, carbon stocks and transpiration may be in focus; foresters may require woody volume and quality; and from a pastoralist’s perspective, the annual production of green foliage of fodder species is most important. Finally, from a socio-economic perspective, we would profit from estimating the amount of trees available for each person. Additional work, more fully exploiting very high resolution imagery (e.g. mapping height and canopy size of individual trees), is likely to bring us further in these directions. This study was, however, able to demonstrate the potential of West African farmland and savannas to provide a range of ecosystem services. Moreover, the wall-to-wall coverage and the high number of pixels in our analysis provide a solid basis for understanding woody cover in different landscapes at the regional West Africa drylands scale and this can be applied to other dryland regions globally. Case studies will still remain extremely valuable as a means of obtaining insights into the complex processes linking environmental factors and land management decisions to woody cover across the variety of

local circumstances. By combining wall-to-wall analysis with process studies at local scale, a more robust basis for developing environmental policies may be established.

METHODS

We define woody cover as the percentage of ground surface covered by the vertical projection of woody plant crowns. The technical framework of this study adapts local-scale approaches of modeling dryland woody cover^{39,40} into reproducible regional/global scale assessments, as the unprecedented amount of very high spatial resolution (VHR) satellite images now available via the NextView license across the region allows for a new level of detail and larger geographic coverage. Most of the 2006 available images are from November/December (2008–2015) when most of the evergreen and deciduous woody species have green leaves, whereas the herbaceous vegetation is senescent. If no images from these months were available, the period was extended to February. The modified soil-adjusted vegetation index (MSAVI) was calculated with a spatial resolution of 1.7 m, and woody cover was extracted by using a texture based feature extraction method. Field measurements (2000–2015) of woody cover at selected sites served as an independent validation of the remote sensing mapping approach. To achieve a woody cover map of the entire area, the spatially detailed woody cover data derived from VHR images were used to train a gradient boost decision tree regressor to predict woody cover from PROBA-V NDVI and PALSAR-2 images at high resolution (100 m). We tested several filtering approaches and seasonal metrics derived with various methods^{41,42} and decided to apply a moving median window for filtering the time series and filtered 10 day composites as input variables for the regressor to keep the process reproducible. A farmland map²⁹, satellite based rainfall estimates²⁶ (CHIRPS), fire (MCD45A1) and population data²⁷ (Worldpop) were used for analysis of woody cover patterns in relation to climate and land management determinants (Supplementary Fig. 7).

Rainfall zones of the study area.

We used rainfall isohyets derived from CHIRPS²⁶ mean annual rainfall (1981–2016) to divide the study area in arid Sahel (150–300 mm), semi-arid Sahel (300–600 mm), and sub-humid lands (600–1000 mm) (Supplementary Fig. 8a). The zones correspond well with expected bioclimatic zones with different woody species⁴³. Whereas *Acacia ssp* and *Cappariaceae* are dominant in the arid and semi-arid, it is *Combretaceae* and *Fabaceae* in more sub-humid parts. In general, woody cover changes from sparsely scattered in the arid areas to closed canopies in the open woodland and riverine forest of the sub-humid zones.

Field data.

Field data is available from extensive field work in the Ferlo in Senegal (144 sites surveyed in 2015), from the CSE (Centre de Suivi Ecologique) campaigns in Senegal (24 sites surveyed between 2000 and 2015 every other year)¹⁸, from the Gourma region in Mali (23 sites)⁴⁴ and the Fakara in Niger (25 sites)⁴⁵. All surveys measure the projected canopy cover⁴⁴ over plots of various areas (50 m to 1 km), and the data were recalculated in m² per ha and percentage canopy cover.

Extraction of canopy cover from very high spatial resolution data.

The mapping technique was designed to be robust to the use of different sensor types, acquisition dates (i.e. different leaf density), atmospheric conditions, as well as being applicable to various situations ranging from sparse shrub population in arid zones to closed canopy cover woodland in the sub-humid zone. The robustness was assessed by independent field data (Fig. 1b) and is demonstrated in Supplementary Fig. 5. DigitalGlobe QuickBird-2, GeoEye-1 and WorldView-2 were orthorectified and the scenes were screened for clouds and other disturbances. All selected multispectral images were resampled (nearest neighbor) to 1.7 m resolution matching GeoEye-1. MSAVI was calculated and rescaled from 0 to 100⁴⁵ to produce a quantitative base for estimation of canopy cover. Only if a pixel is fully covered with a green leaved canopy, the MSAVI will reach higher values, partly covered pixels (e.g. parts of the crown area or small size shrubs and bushes) have relatively lower values. Visual screening of numerous images showed that most woody plants have MSAVI values above 50, which was robust across all rainfall zones and image acquisition dates. A texture based Haralick feature extraction (8 bins) was then run considering all pixels with values between 50 and 100⁴⁷. The advanced texture filter can be parameterized to extract objects (in our case crown canopies) from their surroundings and from larger objects. The feature termed “mean” was used - the objects have grayscale values depending on their distinctiveness - which was rescaled between 0 and 100, resulting in a quantitative estimate of the areas covered by canopies. Each image was visually screened and images dominated by obvious misestimations (strong under- or overestimation) were discarded. The final values represent the subpixel woody coverage, with 100 being fully covered and 0 free of any green leaved woody vegetation. The advantage of this weighted method over a binary tree/no tree classification is that a sub-pixel coverage (i.e. small crowns and edge pixels) receives a lower weight, thus preventing overestimation (Supplementary Figs 3,5). Moreover, using such weighting emphasizes larger canopies, which makes the product more robust against a rapidly changing (fire, field clearing, etc.) bush layer, which receives a lower weight. Burned areas were manually clipped to keep only high quality training images. In total, 219 images were used for the model (about 1% of the study area). The accuracy of the method was calibrated and tested with field data (144 plots) from Senegal. The square field plots are small (50 × 50 m) and include canopies of all size classes thereby being well suited to validate the VHR product. For the accuracy assessment, canopy cover surveyed for each field plot was compared with VHR imagery derived canopy cover averaged for polygons marking exactly the surveyed area.

Prediction of canopy cover at 100 m resolution.

Advanced Land Observing Satellite (ALOS) Phased Arrayed L-band Synthetic Aperture Radar (PALSAR)⁴⁸ and PROBA-V NDVI⁴⁹ were used for a large scale assessment of woody vegetation (wall-to-wall coverage of West African drylands). For PALSAR-2, we used 100 m cross-polarized HV mosaics converted to gamma-naught values and averaged from 2009 and 2010 over the study area⁴⁸. For PROBA-V, daily atmospherically corrected images at 100 m resolution were combined into 10 day maximum value composites to achieve full coverage in the lower latitudes, which are more frequently affected by cloud cover. Images are available from 2014 to 2016 and the maximum value for each 10 day composite over the 3 years was selected to avoid low values which can be caused by clouds

and burned areas. To further filter out noise, a 30 day running median window was applied, choosing the median value of 3 images. This procedure does not only filter out low value spikes caused by clouds, but also high value spikes which can be caused by herbaceous vegetation (also dry season rainfall events can lead to a flush of herbaceous plants). Both possibilities potentially introduce noise in our analysis dedicated to woody vegetation and this filter is a simple way of reducing noise but keeping the original seasonality.

The woody cover derived from the VHR imagery was used to train the PALSAR and 36 (10-day frequency) PROBA-V NDVI images to obtain a regional-scale woody cover map at 100 m resolution. First, the canopy cover images at 1.7 m resolution were aggregated to 100 m by summing all values (representing sub-pixel canopy coverage), multiplying each pixel with the original pixel size (1.7×1.7 m) and dividing it by 100 so the derived map shows the projected area within the pixel covered by woody plants with the unit percent woody cover. The data was then split into training and validation sets by randomly dividing all pixels in two groups, each including 50% of the original pixels. A large number of pixels ($n=1,323,416$) were available for training and for validation. The training set was then used to fit a non-parametric gradient boost regressor (GBR), which produces a prediction model by means of an ensemble of boosted decision trees⁵⁰. The input data were the PALSAR and 36 filtered 10 day NDVI composites covering an entire year. The quality of the model was assessed by comparing the independent validation set with the predicted woody cover. Predicted values above 100 were masked out and below 0 set to 0. Due to the large amount of training and validation pixels and their spread and representation of different landscapes, over-fitting is not a concern and the model output is expected to be robust. It should be noted that the woody cover is predicted continuously from 0 to 100 (but rounded to 1% steps), leading to a lower statistical fit than similar approaches binning canopy cover into classes of e.g. 10% intervals.

Even though all woody plants have a distinctively different phenological behavior than herbaceous annuals, six different forms of evergreen and deciduous leaf phenologies exist, ranging from short deciduous plants shedding their leaves early in the dry season to evergreen species keeping their leaves throughout the year⁵¹. To avoid an underestimation of the crown cover of stands dominated by deciduous species, the median NDVI ratio between November (a period where all trees have leaves) and February-March (most deciduous species are without or only little leaves at this time) was calculated. Field data from Senegal on species composition (ratio deciduous/evergreen per site) was compared with the NDVI-ratio for corresponding sites (Supplementary Fig. 4b). The output of the GBR prediction was then multiplied with this ratio, enhancing the predicted cover of stands with deciduous species but keeping evergreen stands unchanged. The impact of fire is mitigated by the multi-year maximum and median value over several images. Finally, wetlands and irrigated areas were masked out by combining Globland30⁵² and ESA LC CCI (2010) land cover maps. An independent accuracy assessment was conducted with field data from Senegal, Mali and Niger. These data are based on circular plots along 1 km transect lines (representing larger areas of homogeneous landscapes), spaced at 200 m intervals. The canopy cover of all woody plants was surveyed for these plots and averaged for each transect⁵¹. Polygons (3×3 km) covering the field sites were drawn and model-estimated

woody cover extracted and averaged for each site giving valuable information on the overall fit of the predicted canopy cover.

Environmental data.

Several data sets were used to analyze the relationship between woody cover, rainfall and management. CHIRPS rainfall was summed from 1981 to 2016 for each year and an average annual climatology was calculated (Supplementary Fig. 8a). The original CHIRPS resolution of 5 km was resampled (bilinear interpolation) to match the 100 m resolution of PROBA-V. A recently developed farmland map was used²⁹, which reflects the area under agriculture around 2014 (Supplementary Fig. 8b). The original resolution of the farmland map was 100 m and villages areas are masked out. Conservation areas were derived from the World Database on Protected Areas⁵³. It includes National Parks and protected forests of which most have been established during colonial time by the administration in charge of forest and wild life. The conservation areas are found predominantly in low populated regions characterized by poor soil fertility, but population growth and expansion of farmlands has often encroached into these areas. They are however edaphically different and the woody cover is therefore not entirely comparable to neighboring farmlands. Woody cover in the conservation areas was compared with woody cover in adjacent areas (within 5 km buffer around conservation area boundaries). We used Worldpop for the year 2010³⁰ as human population data set. The resolution of 1 km was resampled (bilinear interpolation) to 100 m for this study.

To improve the comparability between farmlands and savannas, we used the newly developed African soil map at 250 m resolution²⁸ to extract sandy soils (from rock outcrops, shallow soils with dense shrubland, clayey valleys, etc) (Supplementary Fig. 8c). We used the soil texture fraction to calculate a mask leaving only areas with >70% sand in the depth 0–1 m.

To test the impact of rural population on woody vegetation, all settlements with a size smaller than 5 km² were extracted from the Globeland30⁵² data set, resulting in 37,294 villages. The original resolution of 30 m was resampled to 100 m. We established buffer zones with 0.5, 1, 2, 5 and 20 km distance to the village areas (Supplementary Fig. 6).

A gradient boost classifier⁵⁰ was applied to test the determinants of predicted woody cover. Explanatory variables of this model based on an ensemble of decision trees were (1) mean annual rainfall, (2) fire frequency deriving the number of fires between 2000 and 2015 from MODIS burned area product MCD45A1 (Supplementary Fig. 8d), (3) rainfall variability (the coefficient of variation of annual sums between 1981 and 2016), (4) the sand fraction from the soil map, (5) the elevation derived from SRTM digital elevation model (90 m), (6) human population³⁰, and (7) distance from the villages (buffer zones). Predicted woody cover was grouped in classes (0–3%, 3–10%, 10–20% and >20%) to meet the requirements of the classifier and a random sample of 1% of the pixels was chosen ($n=2,812,563$) which was used as response variable. The model was run with 10 different random sets of pixels to ensure that no bias emerges by the selection. Due to the decision tree structure of the model, correlations between the explanatory variables can be neglected. The accuracy of the model

is calculated by setting aside 60% of the pixels, which are then used to test the predicted results.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements.

M.B. received funding from the European Union's Horizon 2020 Research and Innovation program under Marie Skłodowska-Curie grant agreement No. [656564]. R.F. acknowledges the funding from the Danish Council for Independent Research (DFF) Grant ID: DFF – 6111-00258. We thank Marie-Julie Lambert for providing the farmland mask. We thank the Centre de Suivi Ecologique for providing field data from Senegal. We thank DigitalGlobe for providing commercial satellite data within the NextView license program.

References

1. Stebbing EP The Encroaching Sahara: The Threat to the West African Colonies. *The Geographical Journal* 85, 506–519 (1935).
2. Aubréville A. *Climats, forêts et désertification de l'Afrique tropicale*. (Société d'éditions géographiques, maritimes et coloniales, 1949).
3. Lüdeke MKB, Moldenhauer O. & Petschel-Held G. Rural poverty driven soil degradation under climate change: the sensitivity of the disposition towards the Sahel Syndrome with respect to climate. *Environmental Modeling & Assessment* 4, 315–326 (1999).
4. Ribot JC A history of fear: imagining deforestation in the West African dryland forests. *Global Ecology and Biogeography* 8, 291–300 (1999).
5. Mbow C, Mertz O, Diouf A, Rasmussen K. & Reenberg A. The history of environmental change and adaptation in eastern Saloum-Senegal—Driving forces and perceptions. *Global and Planetary Change* 64, 210–221 (2008).
6. Hansfort SL & Mertz O. Challenging the Woodfuel Crisis in West African Woodlands. *Hum. Ecol* 39, 583–595 (2011).
7. Lykke AM, Kristensen MK & Ganaba S. Valuation of local use and dynamics of 56 woody species in the Sahel. *Biodiversity and Conservation* 13, 1961–1990 (2004).
8. Couteron P. & Kokou K. Woody vegetation spatial patterns in a semi-arid savanna of Burkina Faso, West Africa. *Plant Ecology* 132, 211–227 (1997).
9. Rasmussen K, Fog B. & Madsen JE Desertification in reverse? Observations from northern Burkina Faso. *Global Environmental Change* 11, 271–282 (2001).
10. Tappan G., Sall M, Wood E. & Cushing M. Ecoregions and land cover trends in Senegal. *Journal of Arid Environments* 59, 427–462 (2004).
11. Reij C, Tappan G. & Belemvire A. Changing land management practices and vegetation on the Central Plateau of Burkina Faso (1968–2002). *Journal of Arid Environments* 63, 642–659 (2005).
12. Mortimore MJ & Adams WM Farmer adaptation, change and crisis in the Sahel. *Global Environmental Change* 11, 49–57 (2001).
13. Fairhead J. & Leach M. False forest history, complicit social analysis: Rethinking some West African environmental narratives. *World Development* 23, 1023–1035 (1995).
14. Gonzalez P. Desertification and a shift of forest species in the West African Sahel. *Clim Res* 17, 217–228 (2001).
15. Rasmussen K. et al. Environmental change in the Sahel: reconciling contrasting evidence and interpretations. *Reg Environ Change* 1–8 (2015). doi:10.1007/s10113-015-0778-1
16. Tucker CJ & Nicholson SE Variations in the Size of the Sahara Desert from 1980 to 1997. *Ambio* 28, 587–591 (1999).
17. Olsson L, Eklundh L. & Ardo J. A recent greening of the Sahel—trends, patterns and potential causes. *Journal of Arid Environments* 63, 556–566 (2005).

18. Brandt M. et al. Ground- and satellite-based evidence of the biophysical mechanisms behind the greening Sahel. *Glob Change Biol* 21, 1610–1620 (2015).
19. Kaptué AT, Prihodko L. & Hanan NP On regreening and degradation in Sahelian watersheds. *PNAS* 201509645 (2015). doi:10.1073/pnas.1509645112
20. Hansen MC et al. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* 342, 850–853 (2013). [PubMed: 24233722]
21. Sexton JO et al. Global, 30-m resolution continuous fields of tree cover: Landsat-based rescaling of MODIS vegetation continuous fields with lidar-based estimates of error. *International Journal of Digital Earth* 6, 427–448 (2013).
22. Good SP & Caylor KK Climatological determinants of woody cover in Africa. *Proc Natl Acad Sci U S A* 108, 4902–4907 (2011). [PubMed: 21383125]
23. Browning DM et al. Emerging technological and cultural shifts advancing drylands research and management. *Frontiers in Ecology and the Environment* 13, 52–60 (2015).
24. Axelsson CR & Hanan NP Patterns in Woody Vegetation Structure across African Savannas. *Biogeosciences Discuss.* 2017, 1–16 (2017).
25. Hill MJ & Hanan NP *Ecosystem Function in Savannas: Measurement and Modeling at Landscape to Global Scales.* (CRC Press, 2010).
26. Funk C. et al. The climate hazards infrared precipitation with stations—a new environmental record for monitoring extremes. *Scientific Data* 2, 150066 (2015).
27. Tatem AJ WorldPop, open data for spatial demography. *Scientific Data* 4, 170004 (2017).
28. Hengl T. et al. Mapping Soil Properties of Africa at 250 m Resolution: Random Forests Significantly Improve Current Predictions. *PLOS ONE* 10, e0125814 (2015).
29. Lambert M-J, Waldner F. & Defourny P. Cropland Mapping over Sahelian and Sudanian Agrosystems: A Knowledge-Based Approach Using PROBA-V Time Series at 100-m. *Remote Sensing* 8, 232 (2016).
30. Bayala J, Sanou J, Teklehaimanot Z, Kalinganire A. & Ouédraogo S. Parklands for buffering climate risk and sustaining agricultural production in the Sahel of West Africa. *Current Opinion in Environmental Sustainability* 6, 28–34 (2014).
31. Mertz O. et al. Climate Factors Play a Limited Role for Past Adaptation Strategies in West Africa. *Ecology and Society* 15, (2010).
32. Bucini G. & Hanan NP A continental-scale analysis of tree cover in African savannas. *Global Ecology and Biogeography* 16, 593–605 (2007).
33. Devine AP, McDonald RA, Quaipe T. & Maclean IMD Determinants of woody encroachment and cover in African savannas. *Oecologia* 1–13 (2017). doi:10.1007/s00442-017-3807-6
34. Kulmatiski A. & Beard KH Woody plant encroachment facilitated by increased precipitation intensity. *Nature Clim. Change* 3, 833–837 (2013).
35. Brandt M. et al. Human population growth offsets climate driven woody vegetation increase in sub-Saharan Africa. *Nature Ecology & Evolution* 1, 0081 (2017).
36. Mertz O, Lykke A. & Reenberg A. Importance and seasonality of vegetable consumption and marketing in Burkina Faso. *Economic Botany* 55, 276–289 (2001).
37. Sankaran M. et al. Determinants of woody cover in African savannas. *Nature* 438, 846–849 (2005). [PubMed: 16341012]
38. Tiffen M, Mortimore M. & Gichuki F. *More People, Less Erosion: Environmental Recovery in Kenya.* (Wiley, 1994).
39. Bucini G, Saatchi S, Hanan N, Boone RB & Smit I. Woody cover and heterogeneity in the Savannas of the Kruger National Park, South Africa in 2009 *IEEE International Geoscience and Remote Sensing Symposium* 4, IV-334-IV-337 (2009).
40. Herrmann S, Wickhorst A. & Marsh S. Estimation of Tree Cover in an Agricultural Parkland of Senegal Using Rule-Based Regression Tree Modeling. *Remote Sensing* 5, 4900–4918 (2013).
41. Roerink GJ, Menenti M. & Verhoef W. Reconstructing cloudfree NDVI composites using Fourier analysis of time series. *International Journal of Remote Sensing* 21, 1911–1917 (2000).
42. Jonsson P. & Eklundh L. TIMESAT—a program for analyzing time-series of satellite sensor data* 1. *Computers & Geosciences* 30, 833–845 (2004).

43. Breman H. & Kessler J-J Woody Plants in Agro-Ecosystems of Semi-Arid Regions: with an Emphasis on the Sahelian Countries. (Springer-Verlag, 1995).
44. Hiernaux P. et al. Woody plant population dynamics in response to climate changes from 1984 to 2006 in Sahel (Gourma, Mali). *Journal of Hydrology* 375, 103–113 (2009).
45. Hiernaux P. & Ayantunde A. The Fakara: a semi-arid agro-ecosystem under stress. (ILRI, 2004).
46. Qi J, Chehbouni A, Huete AR, Kerr YH & Sorooshian S. A modified soil adjusted vegetation index. *Remote Sensing of Environment* 48, 119–126 (1994).
47. Haralick RM, Shanmugam K. & Dinstein I. Textural Features for Image Classification. *IEEE Transactions on Systems, Man, and Cybernetics SMC-3*, 610–621 (1973).
48. Shimada M. et al. New global forest/non-forest maps from ALOS PALSAR data (2007–2010). *Remote Sensing of Environment* 155, 13–31 (2014).
49. Dierckx W. et al. PROBA-V mission for global vegetation monitoring: standard products and image quality. *International Journal of Remote Sensing* 35, 2589–2614 (2014).
50. Breiman L. *Arcing the Edge*. (Statistics Department, University of California, Berkeley, 1997).
51. Brandt M. et al. Woody plant cover estimation in drylands from Earth Observation based seasonal metrics. *Remote Sensing of Environment* 172, 28–38 (2016).
52. Chen J. et al. Global land cover mapping at 30 m resolution: A POK-based operational approach. *ISPRS Journal of Photogrammetry and Remote Sensing* 103, 7–27 (2015).
53. World Conservation Union and UNEP-World Conservation Monitoring Centre, *World Database on Protected Areas*, Version 2007, WCMC, Cambridge, UK (2007).




Figure 1]. High resolution woody cover mapping and validation with field data.

a, Woody cover derived from MSAVI at 1.7 m resolution (Supplementary Figs 2-5). **b**, The woody cover map at 1.7 m resolution was validated with in situ data from northern Senegal (MAE of 3.2, $r=0.87$, slope=0.98, $n=144$). Woody cover >10% ($r=0.76$); woody cover <10% ($r=0.77$). **c**, The predicted woody cover map (100 m) was validated with independent in situ data from Senegal ($n=24$), Mali ($n=23$) and Niger ($n=25$) (MAE=0.8, $r=0.9$, slope=1.01) both for woody cover >10% ($r=0.75$) and woody cover <10% ($r=0.86$).




Figure 2]. Predicting woody cover.

a, Predicted woody cover at 100 m resolution with locations of the close-up views (b-e) indicated. **b**, Woody cover in farmlands at the semi-arid Nigeria/Niger border.. The presence of trees within villages makes them stand out as green clusters. Woody corridors (shelterbelts) can be identified. **c**, Farmlands in sub-humid Burkina Faso are expanding into remnants of forest reserves. **d**, The villages in the Malian Seno Plain are surrounded by a well managed woody vegetation **e**, The sandy pastoral zone of arid Senegal has locally high concentrations of woody plants on fine textured soils of inter-dunes..




Figure 3|. Determinants and patterns of woody cover.

a, Factors potentially impacting woody cover are averaged along the rainfall gradient (50 mm steps). **b**, The relative weight of variables in a decision tree model explaining predicted woody cover (150–1000 mm) with an overall explaining power of 67%. **c**, Mean woody cover grouped into savannas (n=148,286,890) and farmland (n=43,374,091), areas of dense (>50 persons km⁻²; n=23,127,786) and sparse (<50 persons km⁻²; n=167,752,160) population densities, as well as conservation areas (n=8,902,702) and their surroundings (5 km) (n=6,040,825). Standard deviations are shown as grey background bars. Total pixels: 191,660,981.




Figure 4|. Land management impacts on woody cover.

a, Woody cover grouped into farmland and savanna for each bioclimate zone. **b**, Woody cover (a random sample of 1%; $n=2,812,563$) is shown along the rainfall gradient (10 mm steps) for savannas and **c**, for farmlands. **d**, Woody cover is averaged within intervals of population density showing opposing patterns for arid/semi-arid (150–600 mm) and sub-humid (600–1000 mm) zones. **e**, Comparison between woody cover on farmland and on savannas, both on sandy soils only (entire region; $n= 73,848,805$). **f**, Woody cover as a function of distance to the village center (entire region; average for 37,294 villages on sandy soils).