



# Effectiveness of Action Observation Training on Upper Limb Motor Function in Children with Hemiplegic Cerebral Palsy: A Systematic Review of Randomized Controlled Trials

This article was published in the following Dove Press journal:  
*Pediatric Health, Medicine and Therapeutics*

Abayneh Alamer <sup>1</sup>  
Haimanot Melese <sup>1</sup>  
Belaynew Adugna<sup>2</sup>

<sup>1</sup>Department of Physiotherapy, School of Medicine, College of Health Sciences, Ayder Comprehensive Specialized Hospital, Mekelle University, Mekelle, Ethiopia; <sup>2</sup>Department of Physiotherapy, School of Medicine, College of Health Sciences, Tibebe Ghion Comprehensive Specialized Hospital, Bahirdar University, Bahirdar, Ethiopia

**Abstract:** The purpose of this review was to evaluate and examine the current best evidence for the effectiveness of action observation therapy on upper limb function rehabilitation in children with hemiplegic cerebral palsy. A comprehensive search of literature published between September 2010 and May 2020 was conducted using the following electronic databases: PubMed, Google Scholar, the Physiotherapy Evidence Database (PEDro), EMBASE, Cochrane library, and Scopus. Only randomized controlled trials evaluating the effect of action observation therapy on upper limb motor function in children with hemiplegic cerebral palsy were included. PEDro scale was used to assess the risk of bias of included trials. This study was reported according to the guideline of the PRISMA statement. The overall methodological quality of the studies was done using the PEDro scale and GRADE approach. The primary outcome measures of this review were the Melbourne Assessment Scale, Assisting Hand Assessment scale to evaluate physical function and structures. Furthermore, the ABILHAND-Kids test, and Box and Block Test primary outcome measures were used to determine the subjects' activities and participation. Nine randomized controlled trials involving 234 participants were analyzed. The overall quality of evidence was rated from moderate to high. This review suggests action observation therapy was found to be a promising intervention for upper limb rehabilitation in children with cerebral palsy.

**Keywords:** action observation training, hemiplegic CP, upper extremity motor function, systematic review

## Introduction

Cerebral palsy (CP) is a set of permanent movement and posture disorders that causes restriction of activities in the fetal or newborn brain.<sup>1</sup> It is a major leading cause of disabilities in childhood with a prevalence of 2.1 per 1000 live births,<sup>2,3</sup> and children with unilateral cerebral palsy (UCP) accounts for 39% of the general CP population.<sup>3,4</sup>

Often, movement deficits include poor coordination, muscle weakness, tremors, sensory deficits, poor perception, and severe difficulties with a concentration in children with CP.<sup>1,5</sup> These great burdens of health problems may be associated with restricted motor skills and activities of daily living, such as reaching and grasping, which have a significant impact on their upper limb motor function.<sup>1,5-7</sup>

Correspondence: Haimanot Melese  
Tel +251 927608383  
Fax +251 344416681  
Email haimanophysio@gmail.com

Nowadays, numerous goal-directed therapeutic approaches have been used to enhance upper limb function, including constraint-induced movement therapy (CIMT), hand-arm bimanual intensive therapy (HABIT), intramuscular botulin toxin combined with therapeutic training, and action observation therapy (AOT).<sup>8–10</sup> Among these, AOT has recently gained attention in improving upper limb function.<sup>11</sup>

AOT is a novel rehabilitation technique, which involves observation of purposeful actions with the intention to imitate and then performing those actions.<sup>12,13</sup> Often time, it is usually described as the patient being asked to carefully observe actions presented through a video-clip or performed by an operator, in order to imitate, try, and execute them after observation.<sup>14</sup> This may include the mirror neuron system (MNS) processes on the upper limb function resulting from the observation of actions (new motor skills) and actual execution of similar neural structures.<sup>15–17</sup> Current evidence<sup>18–20</sup> reported that AOT is becoming the latest treatment option among other interventions for children with UCP on upper limb function and their activities of daily living. Previous reviews have been limited to investigating AOT effectiveness on upper and lower limb rehabilitation in both neurological and orthopedic disorders,<sup>11</sup> patients with stroke and brain injuries,<sup>21–23</sup> limb pain,<sup>24</sup> and in Parkinson disease.<sup>25</sup>

However, despite the volume of our understanding, there is a dearth of current scientific evidence evaluating the efficacy of AOT on upper limb functions, particularly in children with UCP in a systematic way. Given these gaps, there is a need to perform a systematic evaluation to investigate the current scientific evidence on AOT clinical worthiness with high quality randomized controlled trials. Therefore, the purpose of this review was aimed to address the clinical research question about the effectiveness of AOT on upper limb function in children with unilateral CP.

## Methods

### Design

This systematic review was performed and reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.<sup>26</sup>

### Search Strategy

A literature search was performed to recognize all qualified randomized controlled trials. The literature search was conducted using the following databases: PubMed, Google

scholar, Cochrane library, Scopus, CINAHL, AMED, PEDro, and EMBASE. The terms used to search the literature were: “Action observation training” OR “Action observation treatment” OR “Action Observation Therapy” AND “upper limb functions” OR “movement execution” OR “physical training” AND “unilateral cerebral palsy” OR “hemiplegic/hemiplegia cerebral palsy” OR “cerebral palsy” AND “randomized clinical trial” ([Appendix](#)). The retrieval of the studies for the published articles was set for the period of September 2010 to May 2020. Manual searches of the reference list of included articles were also performed.

### Eligibility Criteria

The studies were included if they met the following established inclusion criteria: 1) population/participants: children with unilateral cerebral palsy; 2) intervention: intervention groups received AOT and/or combined with constraint-induced movement therapy; 3) comparison: control groups received sham AOT/repetitive physical training; 4) outcomes: physical function and structure, activities and participation. All randomized control trials (RCT) performed to evaluate the efficacy of AOT on UCP survivors were included in this review. Studies in which the addition of AOT over other conventional interventions (experimental group) compared with other interventions only (control group) were also included. Only full-length articles reported in English were included. This review excluded observational studies, quasi-experimental studies, conference abstracts, and RCTs which include healthy children, children with stroke, and traumatic brain injuries.

### Study Selection

Two reviewers (H.M, A.A) independently screened papers from the identified lists on the basis of title/abstract, based on the determined inclusion standards. The studies have been retrieved in detail by means of methodological quality and data extraction tools. Reference lists of recognized studies have been manually searched for extra RCTs. The duplicated articles were eliminated. Eligible papers have been collected in complete-textual content, independently screened by the same reviewers. The third reviewer (B.A) was there to resolve the disagreements among the two reviewers.

### Risk of Bias in Individual Studies

The risk of bias of the included studies was independently assessed by two review authors (H.M and A.A) by using the Physiotherapy Evidence Database (PEDro) scale. The

methodological quality of each included study has been assessed using the PEDro scale.<sup>27,28</sup> This tool comprises 10 items with the first item (external validity of the article) quality assessments of controlled intervention study tool.<sup>26</sup> The following topics were assessed based on critical criteria: concealed assignment, intention-to-treat analysis, and adequate treatment follow-up. These characteristics make the PEDro scale a valuable apparatus to evaluate the methodological quality of RCTs. This review considered trials with a score of 5 to 7 as moderate quality, and a score of  $\geq 8$  as high quality (Table 1). The overall quality of the evidence and strength recommendation was evaluated by using the GRADE approach.<sup>29</sup> The GRADE approach specifies four levels of quality (high, moderate, low, and very low). The overall evidence was downgraded depending on the presence of five factors: limitations (due to risk of bias); consistency of results; directness (eg, whether participants are similar to those about whom conclusions are drawn); precision (ie, sufficient data to produce narrow confidence intervals); and other (eg, publication bias).

## Data Extraction

Information extraction was done based on the data retrieval template of the Cochrane Consumer and Communication Review Group. Two reviewers (H.M & A.A) extracted the data independently and the third author checked the extracted data. Disagreements were resolved by consensus among the two authors. If the two authors did not reach a consensus, the third author (B.A) was there to decide. The following data were extracted from each trial using PICOS standards: participant characteristics (sex, mean age, sample size, duration of the disease), the objective of the study, author name and year of publication, PEDro score, follow-up time and intensity of the training, treatment outcomes (at baseline, follow-up, and end-of-treatment), type of treatment, study design, study results, and study conclusions (Table 2).

## Results

### Study Selection

A total of 2507 articles were identified from databases by the searching strategy. After adjusting for duplicates, 1882 remained. After the title and abstract screening among 644 studies, 594 studies were excluded. After full-text screening out of 50 articles, 9 randomized controlled trials were included in this review (Figure 1).

## Study Characteristics

The detailed descriptions, and characteristics of the included studies are summarized in Table 1. All nine randomized controlled trials, which focused on the effectiveness of AOT in the treatment of the upper limb function of children with hemiplegic cerebral palsy, were included in the current systematic review.

## Participants

All subjects who participated in individual studies had a clinical diagnosis of hemiplegic cerebral palsy (CP). The sample size ranged from 10<sup>30</sup> to 59 participants.<sup>31</sup> The mean age (SD) of participants ranged from 9.13 (2.36)<sup>32</sup> to 9.48 (2.12)<sup>20</sup> years in experimental groups and 9.25 (3.15)<sup>32</sup> to 9.94 (3.15)<sup>20</sup> years in controlled groups.

## Interventions

Randomized controlled trials were included that assessed the effectiveness of AOT combined or not with mCIMT, compared with a comparison/control group; assessed sham/placebo AOT (watching videos of computer games and landscape photographs with no specific motor content); and/routine conventional rehabilitation programs. The training durations were ranged from 20 minutes to 1 hour per session, 1 up to 5 times per week, for three weeks to six months.<sup>18,20,33</sup> The characteristics of the intervention expressed as mean values (range) were: 12 min of AOT for each session (9–18); 4 min for each video; 10.7 min of observed action performance (3–36); 5 sessions a week (3–7); the total duration of treatment=46.7 days (9–180 days). In all studies, the control group performed the same actions as the experimental group for the same amount of time. All eight studies administered AOT through goal-directed videos with specific hand/arm motor skill contents and afterward, the participants were asked to execute the same action.<sup>18,20,30-34</sup> Only one study involved home-based AOT with repeated practice,<sup>35</sup> and two studies conducted combining AOT with mCIMT.<sup>31,33</sup>

## Outcome Measures

The outcomes of individual studies were extracted from the following: physical function and structures, activities, and participation. The primary outcome measures of this systematic review were Melbourne Assessment Scale, ABILHAND-Kids test, Box and Block Test (BBT), and Assisting Hand Assessment scale (AHA) to evaluate the upper limb function of children with hemiplegic CP. Furthermore, the ABILHAND-Kids test and Box and Block Test (BBT) were

**Table 1** Methodological Quality Assessment of Included RCTs

| <b>PEDro Items</b>              | <b>Buccino et al, 2012<sup>18</sup></b> | <b>Sgandurra et al, 2013<sup>20</sup></b> | <b>Young Kim et al, 2014<sup>32</sup></b> | <b>Kirkpatrick et al, 2016<sup>35</sup></b> | <b>Kim et al, 2018<sup>36</sup></b> | <b>Buccino et al, 2018<sup>34</sup></b> | <b>Simon-Martinez et al, 2019<sup>33</sup></b> | <b>Simon-Martinez, et al, 2020<sup>31</sup></b> | <b>Kim, 2020<sup>30</sup></b> |
|---------------------------------|---|---|---|---|-------------------------------------|---|--|---|-------------------------------|
| Eligibility criteria            | Yes                                     | No  | Yes                                       | Yes   | Yes                                 | Yes                                     | No   | Yes   | Yes                           |
| Random allocation               | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | Yes                                     | Yes  | Yes   | Yes                           |
| Concealed allocation            | No                                      | Yes                                       | No  | Yes   | No                                  | No                                      | No   | No  | No                            |
| Baseline comparability          | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | Yes                                     | Yes  | Yes   | Yes                           |
| Blind participants              | Yes                                     | No  | No  | No  | No                                  | Yes                                     | No   | No  | No                            |
| Blind therapists                | No                                      | No  | No  | No  | No                                  | No                                      | No   | No  | No                            |
| Adequate follow-up              | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | No                                      | Yes  | Yes   | Yes                           |
| Blind assessors                 | Yes                                     | Yes                                       | No  | Yes   | Yes                                 | Yes                                     | No   | No  | No                            |
| Intention to treat analysis     | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | Yes                                     | Yes  | Yes   | Yes                           |
| Between group comparisons       | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | Yes                                     | Yes  | Yes   | Yes                           |
| Point estimates and variability | Yes                                     | Yes                                       | Yes                                       | Yes   | Yes                                 | Yes                                     | Yes  | Yes   | Yes                           |
| Total score out of 10           | 8/10                                    | 8/10                                      | 6/10                                      | 8/10  | 7/10                                | 7/10                                    | 6/10   | 6/10  | 6/10                          |
| Graded approach scale           | High                                    | High                                      | Moderate                                  | High  | Moderate                            | Moderate                                | Moderate                                       | Moderate  | Moderate                      |

**Table 2** Summary of Included Randomized Controlled Trials

| Authors, Publication Years          | Participant Characteristics   | Interventions   | Outcome Measures                        | Results/Conclusion   |
|-------------------------------------|---|---|---|--|
| Buccino et al, 2012 <sup>18</sup>   | -15 total participants: EG; (n=8), CG; (n=7)<br>-Sex: 9 males and 6 females, age range: 7 to 11 years<br>-Median age/years: EG; 7 and 6 Months, CG; 8 years. GMFCS: not stated              | -EG: asked to observe videos with motor content (every day arm/hand actions), afterwards to imitate them for 3 weeks.<br>-CG: asked to observe videos with no motor specific content and afterwards to execute the same actions as the case group for 3 weeks.<br>-Both groups received routine conventional rehabilitation program.  | -Melbourne Assessment Scale             | - After treatment, the functional score gain (D) was significantly different in the case and control groups ( $p=0.02$ ). A Bonferroni post-hoc test indicated that motor scores as measured by the Melbourne Assessment Scale at T3 (mean 88.3) improved compared with T2 (mean 85.1; $p=0.04$ ) and T1 (mean 83.9; $p=0.004$ ). A Bonferroni post-hoc test showed that an increased score on the Melbourne Assessment Scale was achieved only in the case group (T1=86.8, T2=88.0, T3=94.2; T1 vs T3, $p=0.002$ ; T2 vs T3, $p=0.01$ ), but not in the control group (T1=80.5, T2=81.8, T3=81.5; T1 vs T3, $p=1.0$ ; T2 vs T3, $p=1.0$ ).<br>-Action observation treatment can be an effective part of the rehabilitation program in children with CP.                       |
| Sgandurra et al, 2013 <sup>20</sup> | -24 total participants: EG; (n=12), CG; (n=12)<br>-Sex (M/F): 16/8<br>-Age: mean age (years) (SD): EG; 9.48 (2.12), CG; 9.94 (2.77)<br>Range: EG; 6.4–12.6, CG; 6.1–14.3. GMFCS: not stated | -EG: observed goal directed Videos each lasted 3 minutes and child was required to perform the same observed action for 3 minutes in the exact order and with the same objects as shown in the video for 60 minutes per day for 3 consecutive weeks.<br>-CG: watched computer games and then were verbally instructed to perform the same for 60 minutes per day for 3 consecutive weeks. | -AHA<br>-MUJUL<br>-ABILHAND-Kids        | -The experimental group improved more ( $p=0.008$ ) in score changes for the AHA at the primary endpoints T1 ( $p=0.008$ ), T2 ( $p=0.019$ ), and T3 ( $p=0.049$ ). No between-group significant changes were found for ABILHAND-Kids or Melbourne assessment.<br>-The Upper Limb Children Action Observation Training (UP-CAT) improved daily upper limb (UL) activities with UCP children, suggesting a new rehabilitation approach based on a neurophysiological model of motor learning.   |
| Young Kim et al, 2014 <sup>32</sup> | -16 total participants: EG; (n=8), CG; (n=8)<br>-Sex (M/F) 8/8<br>-Mean age/year (SD): EG; 9.13 (2.36) CG; 9.25 (3.15). GMFCS: EG; 2.38 ± 0.74, CG; 2.25 ± 1.16                             | EG: received AOPT & practiced repeating the actions they observed on video clips for twelve 30-min sessions, 3 days a week, for 4 weeks. For each video five minutes and 10 minutes per tasks, two videos were shown during each training session.<br>CG: PT group performed the same actions as the AOPT group did after observing landscape photographs                                 | -MAS<br>-BBT<br>-WeeFIM<br>-Grasp power | -The results of this study showed that, in comparison with the physiotherapy group, the functioning of the upper extremities in the AOT group was significantly improved in body functions and activity and participation according to the ICF framework. Comparison between the groups before and after training indicated significantly more improvement for the action observation physical training group than for the physical training group ( $p<0.05$ ); MAS in AOT (-0.6 ± 0.3) vs in PT (-0.2 ± 0.3), BBT in AOT (7.8 ± 2.9) vs in PT (2.3 ± 1.9), ABILHAND-Kids in AOT (3.7 ± 3.0) vs PT (1.6 ± 2.6) and WeeFIM in AOT (1.1 ± 0.8) vs (0.7 ± 0.7).<br>-Indicates that AOT has a positive influence on the functioning of the upper extremities in children with CP. |

(Continued)

Table 2 (Continued).

| Authors, Publication Years            | Participant Characteristics   | Interventions  | Outcome Measures               | Results/Conclusion  |
|---------------------------------------|---|--|--------------------------------|---|
| Kirkpatrick et al, 2016 <sup>35</sup> | -59 total participants: EG; (n=28), CG (n= 31) -Sex (M/F): 28/31 -Median age (IQR): EG; 5y 2mo (4y), CG; 5y 4mo (3y 4mo), age range, 3 to 10 years. GMFCS: not stated | -EG: received AO plus RP 15 minutes/day, 5 days/week for 3 months<br>CG: received 15 minutes/day, 5day/week for 3 months.  | - AHA<br>-MA2<br>ABILHAND-Kids | -There were no between-group differences in AHA, MA2, or ABILHAND-Kids at 3 or 6 months versus baseline (all $p>0.05$ ).<br>-Combined-group improvements ( $p<0.001$ ), observed in AHA and MA2 at 3 months, were maintained at 6 months.<br>ABILHAND-Kids also showed improvement at 3 months ( $p=0.003$ ), maintained at 6 months.<br>-This study did not demonstrate a significant improvement in hand function with AO plus repeated practice (RP) compared to RP alone with children age 3 to 10 years with UCP.  |
| Kim et al, 2018 <sup>36</sup>         | -12 total participants: EG; (n=6), CG (n=6)<br>-Sex (M/F): 7/5<br>-Age ranged from 8 to 11 years. GMFCS: not stated   | -EG: received live AOT for 20 treatment sessions, each 30 minutes in duration, 5 days/week for a month.<br>-CG: received video AOT for 20 treatment sessions, each 30 minutes in duration, 5 days/week for 4 weeks.  | -JTHF<br>-BBT                  | - JTHF scores were significantly lower in the live group ( $p<0.05$ ). The BBT score was significantly higher in the live than in the video group ( $p< 0.05$ ). The findings suggest that live AOT is more effective than video AOT for improving upper limb movement acceleration and function. The functional measures, JTHF scores were significantly better for the live group ( $167.6 \pm 10.6$ s) than for the video group ( $181.2 \pm 9.5$ s) at the follow-up test ( $p=0.03$ ). Scores on the BBT were significantly higher in the live group ( $59.6 \pm 8.8$ EA) than in the video group ( $47.8 \pm 2.9$ EA) at the follow-up test ( $p=0.03$ ).   |
| Buccino et al, 2018 <sup>34</sup>     | -18 total participants: EG; (n=11), CG; (n=7)<br>-Age (years): 5-11<br>-Sex (M/F): 9/9<br>GMFCS: ranged from 1 to 4.  | - EG: observed videos with motor content (every day arm/hand actions) and perform each action plus CPT for each action 9-12 minutes, 5 times/week for 3 weeks<br>-CG: observed videos with no specific motor content plus CPT for each action 9-12 minutes, 5 times/week for 3 weeks | - MUUL<br>- AHA                | - Significant interaction between time and treatment, both for MUUL (b1 interaction=2.7, $t_{36}=3.9$ , and $p=0.00$ ) and for AHA (b1 interaction=2.3, $t_{18}=3.6$ , $p=0.002$ ). Score improvements, in both scales, were higher in the treated participants than in the controls; furthermore, in the treatment group, those improvements were not only maintained but became even stronger at T3. Post hoc analysis showed that for MUUL, results at T2 were significantly different from the results at T1 only in cases ( $p<0.001$ ), but not in controls. As for AHA, results at T2 were significantly different from the results at T1 ( $p<0.001$ ). Even more interestingly, results at T3 were different from the results at T2 ( $p<0.001$ ) for both scales, but again only in cases, but not in controls. |



|  |  |  |   |   |
|--|--|--|---|---|
| <p>Simon-Martinez et al. 2019<sup>33</sup></p> | <p>-36 total participants: EG; (n=20), CG; (n=16)<br/>                 -Mean age (SD): 9 years (1 year) in both groups<br/>                 -Sex (males): MACS (n (%)): EG, I, II, III, 5 (25), 7 (35), 8 (40) and CG, 3 (19), 6 (37), 7 (44) respectively</p> | <p>-EG: received mCIMT plus AOT, 15 hours of AOT, ie video-observation and execution of unimanual tasks at 6 months follow-up.<br/>                 -CG: received mCIMT plus placebo, watched biological-motion free videos and executed the same tasks at 6 months follow-up.</p> | <p>-Motor control parameters (RU, RGV)</p>  | <p>-The combination of mCIMT plus AOT resulted in a shorter movement time during a reaching task compared to those who did not receive additional AOT. Both groups showed improvement in RU and RGV (<math>p=0.07</math>) with large effect sizes (partial <math>\eta^2=0.15</math> for both tasks. Post-hoc analyses for RU showed significant improvements between T1-T3 (<math>p=0.02</math>), and T2-T3 (<math>p=0.03</math>). Improvements during RGV were found immediately after the intervention (T1-T2, <math>p=0.04</math>) and maintained at follow-up (T1-T3, <math>p=0.02</math>). This indicated the improvement in lateral flexion during RU (<math>p=0.03</math>, 67–100% of the movement cycle), although post-hoc analyses did not survive Bonferroni correction.<br/>                 -The mCIMT plus placebo group showed no significant improvements in movement duration (<math>p&gt;0.05</math>). The added AOT training may have a long-lasting effect on movement duration during reaching</p> |
| <p>Simon-Martinez et al. 2020<sup>31</sup></p> | <p>-44 total participants: EG; (n=22), CG; (n=22). Sex(males): 27. Mean age 9 years 6 months (1 year 10 months) MACS I, 15 MACS II, and 20 MACS III], and were allocated to the CIMT + AOT group (n=22) and CIMT + placebo group (n= 22)</p>                   | <p>EG: Received the CIMT plus AOT group received 15 hours of AOT (ie, video-observation) and executed the observed tasks. CG: the CIMT plus AOT group watched videos free of biological motion and executed the same tasks.</p>  | <p>- AHA<br/>                 -MAS<br/>                 -MA2<br/>                 ABILHAND-Kids</p> | <p>- Although no differences between-groups were observed (<math>p&gt;0.05</math>; <math>\eta^2=0-16</math>), the addition of AOT resulted in higher improvements in children with initially lower bimanual performance (<math>p=0.02</math>; <math>\eta^2=0.14</math>). Both groups improved in all outcome measures after the intervention and retained the gains at follow-up (<math>p&lt;0.01</math>; <math>\eta^2=0.02-0.71</math>). Poor sensory function resulted in larger improvements in the total group (<math>p=0.03</math>; <math>\eta^2=0.25</math>) and high amounts of mirror movements tended to result in a better response to the additional AOT training (<math>p=0.06</math>; <math>\eta^2=0.18</math>). The total group improved over time in the AHA (<math>p&lt;0.001</math>).<br/>                 - Adding AOT to CIMT resulted in a better outcome for children with poor motor function and high amounts of mirror movements.</p>   |


(Continued)

**Table 2 (Continued).**

| Authors, Publication Years | Participant Characteristics   | Interventions  | Outcome Measures                                | Results/Conclusion  |
|----------------------------|---|--|---|---|
| Kim, 2020 <sup>30</sup>    | <p>~10 total participants: EG; (n=5), CG; (n=5) Sex (male: female): EG; (2: 3), CG; (2: 3)</p> <p>Mean age (years): EG; (10.2 (1.3)), CG; (9.4 (1.3)); EG; 3(14) MACS I, 7 (32) MACS II, and 12(55) MACS III, CG; 6 (27) MACS I, 8 (36.5) MACS II, and 8 (36.5) MACS III.</p> | <p>EG: observed video tasks of goal-directed action before performing tasks, for 30 minutes per day, three times per week, for 4 weeks.</p> <p>CG: observed and performed the same actions, 60 minutes per day, three times per week, for 4 weeks.</p> | <p>-Grip power ABILHAND Kids</p> <p>- QUEST</p> | <p>-Grip power; and ABILHAND-Kids test results improved significantly in both groups (<math>p&lt;0.05</math>). The ABILHAND-Kids test within-group score improved significantly, from <math>1.6 \pm 0.4</math> logits to <math>2.3 \pm 0.7</math> logits in the 30 minutes training group (<math>p=0.04</math>), and from <math>1.4 \pm 0.3</math> logits to <math>2.1 \pm 0.5</math> logits in the 60 minute group (<math>p=0.04</math>). However, there was no between-group difference in the 30 minutes (<math>0.7 \pm 0.3</math> logits) or the 60 minute group (<math>0.6 \pm 0.1</math> logits; <math>p=1.0</math>). Grip strength at the follow-up evaluation was significantly higher, from <math>3.8 \pm 1.7</math> to <math>4.8 \pm 1.7</math> in the 30 minutes training group (<math>p=0.03</math>) and from <math>4.1 \pm 1.7</math> to <math>5.1 \pm 2.0</math> in the 60 minute group (<math>p=0.04</math>). However, there was no significant difference between the 30 minutes (<math>1.0 \pm 0.3</math>) and 60 minutes groups (<math>1.0 \pm 0.5</math>) at the end of each intervention (<math>p=0.8</math>). The 30 minutes AOT was as effective as 60 minutes AOT in improving grip power and upper limb function in children with CP. QUEST scores improved significantly, from <math>68.8 \pm 7.5\%</math> (pre-test) to <math>77.6 \pm 6.4\%</math> (post-test) in the 30 minutes (<math>p=0.04</math>) and from <math>64.9 \pm 6.1\%</math> (pretest) to <math>72.3 \pm 3.5\%</math> (post-test) in the 60 minutes groups (<math>p=0.04</math>).</p> |

**Abbreviations:** AHA, Assisting Hand Assessment scale; UCP, unilateral cerebral palsy; MUJUL, Melbourne Assessment of Unilateral Upper Limb Function; MA2, Melbourne Assessment 2; MAS, Modified Ashworth Scale; BBT, Box and Block Test; WeeFIM, Wee Functional Independent Measure; AOPT, action observation physical training; JTHF, Jebsen-Taylor Hand Function; BBT, Box and Block Test; mCIMT, modified constraint-induced movement therapy; RU, reach upwards; RGW, reach-to-grasp a vertically oriented cylinder; MAS, Modified Ashworth Scale; QUEST, Quality of Upper Extremity Skills Test; SD, standard deviation; EG, experimental group; CG, control group; GMFCS, Gross Motor Function Classification System; IQR, interquartile range; MACS, Manual Ability Classification System.





**Figure 1** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) diagram. **Note:** Adapted from Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med.* 6(7): e1000097.<sup>37</sup>

used to determine the subjects' activities and participation. The secondary outcome measures of this review were WeeFIM, QUEST, and motor control parameters. Two trials utilized WeeFIM and QUEST, designed to evaluate functional independence, movement patterns, and hand function.<sup>30,32</sup> Five studies used the ABILHAND-Kids test to assess manual ability in children with CP,<sup>19,30,31,33,35</sup> and two studies used BBT.<sup>32,36</sup> Five studies used the Melbourne Assessment Scale to evaluate the physical function and structure of children with UCP.<sup>18,20,33-35</sup> Similarly, another four studies performed a measure of upper limb motor function by using AHA.<sup>20,33-35</sup>

### Risk of Bias Across Studies

Description of risk of bias within the individual studies and their rating is provided in [Table 1](#). The PEDro score for all the included studies ranged from 6 to 8, with a mean score of 7. The quality assessment was performed in all included studies and the quality of evidence ranged between moderate to high. All included studies evaluated

random allocation, baseline comparability, intention to treat analysis, point estimate, and variability, between-group comparisons. Only two studies<sup>20,35</sup> assessed concealed allocation and, except one study,<sup>31</sup> all included trials did not blind the therapist. The most prominent potential source of bias was related to blinding.

### Effects of AOT on Body Function and Structure

Detailed extracted data on physical function and structure, from each trial, has been summarized and presented in [Table 2](#). Nine studies, with 234 individuals, supplied published treatment effects on the upper limb physical function and structure of children with UCP. Out of the nine trials, eight of them (n=175) confirmed that the body function of CP children was significantly improved in AOT groups as compared to the control groups in all outcome measures (AHA, MA2).<sup>18,20,30-34,36</sup> However, only one study<sup>35</sup> (n=59) reported the action observation

therapy had no significant improvement on the upper limb motor function compared to that of the control groups. Within the experimental intervention group, considerable improvement was found in upper limb function recovery in both outcome measures (AHA and MA2) at 3 month tests.

## Effects AOT on Activities and Participation

Out of included studies, seven of them, with 201 individuals with unilateral cerebral palsy proved that AOT has beneficial effects on the activities and participation. Their activities and participation were measured by using the ABILHAND-Kids test, BBT, WeeFIM, and QUEST whereby all outcome measures indicated a significant improvement in the experimental groups compared with the control groups in the activities and participation.<sup>20,30-32,36</sup>

## Adverse Effect

The adverse effect of this intervention was not described in any included studies.

## Discussion

The primary intention of this systematic review was to synthesize and examine the effectiveness of action observation therapy on upper limb rehabilitation in children with hemiplegic CP. To this point, there had been no reviews that determine the effectiveness of the action observation therapy on upper limb function in children with hemiplegic CP in a systematic way. Interestingly, most of the included trials with moderate to high quality of evidence confirmed that AOT was effective on upper limb motor function for children with hemiplegic CP. However, all included studies did not blind the assessor, which leads to assessor bias and could affect the estimation of the treatment effect. The overall effect of AOT on children with hemiplegic CP survivors was evaluated among included studies. Its overall effects were assessed on the different durations of intervention in children with UCP. The effect of AOT on physical function and structure was evaluated in all nine included studies. Likewise, of them seven studies evaluated AOT effects on activities and participation in subjects with hemiplegic CP. Eight studies confirmed that using AOT helps to improve physical function and structure, activities, and participation in children with hemiplegic CP.<sup>18,20,30-34,36</sup> However, in one study, AOT was not effective in upper limb motor function improvement.<sup>35</sup> For instance, the study conducted by Sgandurra et al<sup>20</sup> reported that AOT plus execution

is more effective than execution alone in children with CP. Especially, the AHA scale and ABILHAND-Kids scores differed significantly between the experimental and control groups. Similarly, Buccino et al<sup>18</sup> observed the effects of AOT by comparing action practice after action observation and action practice alone in 15 children with CP (twelve hemiplegic and three with tetraplegia). They found that the mean Melbourne Assessment score increased significantly, from 86.87% to 94.25% in the AOT group. Conversely, Kirkpatrick et al<sup>35</sup> found that home-based AOT and repeated practice did not improve the function of the upper limb more than repeated practice alone in children with CP. These possible differences could be because low-intensity upper limb therapy can be delivered at home by parents, and children in the experimental group may not always have observed movements before performing them, because of poor attention or lack of parental emphasis. Nevertheless, a study done by Kim et al<sup>32</sup> indicates that significant improvement was observed in upper limb motor functions in children with cerebral palsy on MAS, BBT, and ABILHAND-Kids in response to action observation physical training compared to simple physical training. This might be due to AOT considering motor imitation characteristics of the mirror neuron system (MNS) can positively affect neural rehabilitation that is related to enormous range neural networks,<sup>38</sup> and their changes enhance the potential to perform activities.<sup>39</sup> Despite this fact, there were no observed significant differences in WeeFIM between experimental and control groups. This could be due to the troublesome effect of measurement tools in the forms of interviews and questionnaires, which leads to the sensitivity of subjects, and could not be evaluated easily.

The study done by Kim et al<sup>36</sup> confirmed that live AOT is more effective than video AOT for improving UL movement acceleration and function. In addition, Buccino et al<sup>34</sup> evidenced that AOT plus execution is more effective than execution alone in children with CP. They found improved upper limb motor function significantly in both scales (MUUL & AHA) compared to the control groups. The possible explanation is that the brain target of AOT is exactly in the hand motor area, possibly involved in executing actions as well as in their processing which helps individuals to code for the motor properties of objects and the implementation of the most appropriate actions to act upon objects.<sup>40</sup>

Simon-Martinez et al<sup>33</sup> confirmed that a combination of mCIMT and AOT resulted in a shorter movement time during a reaching task compared to those who received mCIMT alone. This proves that the additional AOT intervention for hemiplegic CP children on upper limb motor function was more

effective, as AOT may have facilitated the activation of their mirror neuron system through video observation. Similarly, Kim et al<sup>30</sup> supported the beneficial effect of AOT in improving grip strength and upper limb function in children with hemiplegic CP. A study conducted by Simon-Martinez et al<sup>31</sup> showed that the CIMT plus AOT group benefited more than the CIMT plus placebo group on the upper extremity function of children with CP and their daily living activities. However, AOT did not show an overall added effect on improving upper limb function in children with unilateral CP when combined with CIMT in an intensive training approach.

Taken as a whole, one study reported that AOT had no beneficial effect on upper limb motor function. Nevertheless, taking into account the results of eight studies, a more plausible explanation proved that AOT had a significant improvement on grip power, hand function, activities, and participation in activity of daily living in hemiplegic CP children in comparison to the control groups.

## Limitations

This review had the following limitations: it included only English language articles. Hence, there might be a chance of missing articles published in non-English languages. Most of the included studies were not blinded to the assessor, which can lead to bias estimates of treatment effect. Due to the heterogeneity of included studies, a meta-analysis was not carried out.

## Clinical Implication

The finding of this review suggested that AOT appears to result in improved outcomes on body function and structure, activities, and participation of children with CP. Medical choice-making should rely on the cognitive and perceptual ability of children concerning the age and accessibility of needed equipment for AOT in a resource-limited setting.

## Conclusion

Taken together, these findings suggest action observation therapy was found to be a better rehabilitative intervention for upper limb rehabilitation to improve physical function and structure, activities, and participation of children with hemiplegic CP than simple motor training. However, considerable attention must be paid when we use AOT for CP children, due to the severity of motor impairment, and cognitive status; further studies are needed to determine the optimal frequency, intensity, and time of AOT on these particular study subjects.

## Ethical Approval

Ethical approval or patient consent were not required, since the present study was a review of previously published literature.

## Author Contributions

All authors have contributed to the work reported, in the conception, study design, execution, acquisition of data, analysis and interpretation, have drafted, or substantially revised or critically reviewed the article, agreed on the journal to which the article is to be submitted, checked and agreed on all versions of the article before submission, during revision. Finally, any significant changes introduced at the proofing stage, agree to take responsibility and be accountable for the contents of the article.

## Funding

There is no funding to report.

## Disclosure

The authors report no conflicts of interest for this work.

## References

1. Bax M, Goldstein M, Rosenbaum P, et al. Proposed definition and classification of cerebral palsy, April 2005. *Dev Med Child Neurol.* 2005;47(8):571–576. doi:10.1017/S001216220500112X
2. Ferrari A, Cioni G. *The Spastic Forms of Cerebral Palsy: A Guide to the Assessment of Adaptive Functions.* Springer; 2009.
3. Oskoui M, Coutinho F, Dykeman J, et al. An update on the prevalence of cerebral palsy: a systematic review and meta-analysis. *Dev Med Child Neurol.* 2013;55(6):509–519. doi:10.1111/dmcn.12080
4. Rosenbaum P, Gorter JW, Palisano R, et al. The relationship of cerebral palsy subtype and functional motor impairment: a population-based study. *Dev Med Child Neurol.* 2010;52(7):682–683. doi:10.1111/j.1469-8749.2010.03652.x
5. White DA, Christ SE. Executive control of learning and memory in children with bilateral spastic cerebral palsy. *J Int Neuropsychol Soc.* 2005;11(7):920–924. doi:10.1017/S1355617705051064
6. Steenbergen B, Gordon AM. Activity limitation in hemiplegic cerebral palsy: evidence for disorders in motor planning. *Dev Med Child Neurol.* 2006;48(9):780–783. doi:10.1017/S0012162206001666
7. Uvebrant P. Hemiplegic cerebral palsy aetiology and outcome. *Acta Paediatr.* 1988;77:1–100.
8. Gordon AM, Hung Y-C, Brandao M, et al. Bimanual training and constraint-induced movement therapy in children with hemiplegic cerebral palsy: a randomized trial. *Neurorehabil Neural Repair.* 2011;25(8):692–702. doi:10.1177/1545968311402508
9. Sakzewski L, Gordon A, Eliasson A-C. The state of the evidence for intensive upper limb therapy approaches for children with unilateral cerebral palsy. *J Child Neurol.* 2014;29(8):1077–1090. doi:10.1177/0883073814533150
10. Hoare B, Imms C, Villanueva E, et al. Intensive therapy following upper limb botulinum toxin A injection in young children with unilateral cerebral palsy: a randomized trial. *Dev Med Child Neurol.* 2013;55(3):238–247. doi:10.1111/dmcn.12054

11. Sarasso E, Gemma M, Agosta F, et al. Action observation training to improve motor function recovery: a systematic review. *Arch Physiother.* 2015;5(1):14. doi:10.1186/s40945-015-0013-x
12. Zhu M-H, Wang J, Gu X-D, et al. Effect of action observation therapy on daily activities and motor recovery in stroke patients. *Int J Nurs Stud.* 2015;2(3):279–282. doi:10.1016/j.ijnss.2015.08.006
13. Buccino G, Binkofski F, Fink GR, et al. Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study. *Soc Neurosci.* 2013;133–142.
14. Dinomais M, Lignon G, Chinier E, et al. Effect of observation of simple hand movement on brain activations in patients with unilateral cerebral palsy: an fMRI study. *Res Dev Disabil.* 2013;34(6):1928–1937. doi:10.1016/j.ridd.2013.03.020
15. Rizzolatti G, Fadiga L, Gallese V, et al. Premotor cortex and the recognition of motor actions. *Cogn Brain Res.* 1996;3(2):131–141. doi:10.1016/0926-6410(95)00038-0
16. Buccino G, Binkofski F, Fink GR, et al. Action observation activates premotor and parietal areas in a somatotopic manner: an fMRI study. *Eur J Neurosci.* 2001;13(2):400–404.
17. Sgandurra G, Biagi L, Fogassi L, et al. Reorganization of the action observation network and sensory-motor system in children with unilateral cerebral palsy: an fMRI study. *Neural Plast.* 2018;2018:1–15. doi:10.1155/2018/6950547
18. Buccino G, Arisi D, Gough P, et al. Improving upper limb motor functions through action observation treatment: a pilot study in children with cerebral palsy. *Dev Med Child Neurol.* 2012;54(9):822–828. doi:10.1111/j.1469-8749.2012.04334.x
19. Sgandurra G, Ferrari A, Cossu G, et al. Upper limb children action-observation training (UP-CAT): a randomised controlled trial in hemiplegic cerebral palsy. *BMC Neurol.* 2011;11(1):80. doi:10.1186/1471-2377-11-80
20. Sgandurra G, Ferrari A, Cossu G, et al. Randomized trial of observation and execution of upper extremity actions versus action alone in children with unilateral cerebral palsy. *Neurorehabil Neural Repair.* 2013;27(9):808–815. doi:10.1177/1545968313497101
21. Borges LR, Fernandes AB, Melo LP, et al. Action observation for upper limb rehabilitation after stroke. *Cochrane Database Syst Rev.* 2018;10.
22. Buchignani B, Beani E, Pomeroy V, et al. Action observation training for rehabilitation in brain injuries: a systematic review and meta-analysis. *BMC Neurol.* 2019;19(1):344. doi:10.1186/s12883-019-1533-x
23. Peng T-H, Zhu J-D, Chen -C-C, et al. Action observation therapy for improving arm function, walking ability, and daily activity performance after stroke: a systematic review and meta-analysis. *Clin Rehabil.* 2019;33(8):1277–1285. doi:10.1177/0269215519839108
24. Thieme H, Morkisch N, Rietz C, et al. The efficacy of movement representation techniques for treatment of limb pain—a systematic review and meta-analysis. *J Pain.* 2016;17(2):167–180. doi:10.1016/j.jpain.2015.10.015
25. Caligiore D, Mustile M, Spalletta G, et al. Action observation and motor imagery for rehabilitation in Parkinson's disease: a systematic review and an integrative hypothesis. *Neurosci Biobehav Rev.* 2017;72:210–222. doi:10.1016/j.neubiorev.2016.11.005
26. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev.* 2015;4(1):1. doi:10.1186/2046-4053-4-1
27. Maher CG, Sherrington C, Herbert RD, et al. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther.* 2003;83(8):713–721. doi:10.1093/ptj/83.8.713
28. de Morton NA. The PEDro scale is a valid measure of the methodological quality of clinical trials: a demographic study. *Aust J Physiother.* 2009;55(2):129–133. doi:10.1016/S0004-9514(09)70043-1
29. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 7. Rating the quality of evidence—inconsistency. *J Clin Epidemiol.* 2011;64(12):1294–1302. doi:10.1016/j.jclinepi.2011.03.017
30. Kim DH. Comparison of short-and long-time action observation training (AOT) on upper limb function in children with cerebral palsy. *Physiother Pract Res.* (Preprint):1–6.
31. Simon-Martinez C, Mailleux L, Hoskens J, et al. Randomized controlled trial combining constraint-induced movement therapy and action-observation training in unilateral cerebral palsy: clinical effects and influencing factors of treatment response. *Ther Adv Neurol Disord.* 2020;13:1756286419898065. doi:10.1177/1756286419898065
32. Kim JY, Kim J-M, Ko EY. The effect of the action observation physical training on the upper extremity function in children with cerebral palsy. *J Exerc Rehabil.* 2014;10(3):176. doi:10.12965/jer.140114
33. Simon-Martinez C, Mailleux L, Jaspers E, et al. Effects of combining constraint-induced movement therapy and action-observation training on upper limb kinematics in children with unilateral cerebral palsy: a randomized controlled trial. *Sci Rep.* 2020;10:1–15.
34. Buccino G, Molinaro A, Ambrosi C, et al. Action observation treatment improves upper limb motor functions in children with cerebral palsy: a combined clinical and brain imaging study. *Neural Plast.* 2018;2018:1–11. doi:10.1155/2018/4843985
35. Kirkpatrick E, Pearse J, James P, et al. Effect of parent-delivered action observation therapy on upper limb function in unilateral cerebral palsy: a randomized controlled trial. *Dev Med Child Neurol.* 2016;58(10):1049–1056. doi:10.1111/dmcn.13109
36. Kim D-H, An D-H, Yoo W-G. Effects of live and video form action observation training on upper limb function in children with hemiparetic cerebral palsy. *Technol Health Care.* 2018;26(3):437–443. doi:10.3233/THC-181220
37. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 6(7):e1000097.
38. Iacoboni M, Koski LM, Brass M, et al. Reafferent copies of imitated actions in the right superior temporal cortex. *Proc Natl Acad Sci U S A.* 2001;98(24):13995–13999. doi:10.1073/pnas.241474598
39. Buccino G, Solodkin A, Small SL. Functions of the mirror neuron system: implications for neurorehabilitation. *Cogn Behav Neurol.* 2006;19(1):55–63. doi:10.1097/00146965-200603000-00007
40. Biagi L, Cioni G, Fogassi L, et al. Action observation network in childhood: a comparative fMRI study with adults. *Dev Sci.* 2016;19(6):1075–1086. doi:10.1111/desc.12353

## Pediatric Health, Medicine and Therapeutics

### Publish your work in this journal

Pediatric Health, Medicine and Therapeutics is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries. All aspects of health maintenance, preventative measures and disease treatment interventions are addressed within the journal. Practitioners from all disciplines are

invited to submit their work as well as healthcare researchers and patient support groups. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/pediatric-health-medicine-and-therapeutics-journal>

Dovepress