

ARTICLE

Open Access

# A reporter for noninvasively monitoring gene expression and plant transformation

Yubing He<sup>1,2</sup>, Tao Zhang<sup>3,4</sup>, Hui Sun<sup>2</sup>, Huadong Zhan<sup>1</sup> and Yunde Zhao<sup>3</sup>

## Abstract

Reporters have been widely used to visualize gene expression, protein localization, and other cellular activities, but the commonly used reporters require special equipment, expensive chemicals, or invasive treatments. Here, we construct a new reporter *RUBY* that converts tyrosine to vividly red betalain, which is clearly visible to naked eyes without the need of using special equipment or chemical treatments. We show that *RUBY* can be used to noninvasively monitor gene expression in plants. Furthermore, we show that *RUBY* is an effective selection marker for transformation events in both rice and Arabidopsis. The new reporter will be especially useful for monitoring cellular activities in large crop plants such as a fruit tree under field conditions and for observing transformation and gene expression in tissue culture under sterile conditions.

## Introduction

Various genetically encodable reporters have been developed to monitor gene expression, protein subcellular localization, protein stability, hormonal signaling, and impacts of environmental signals. The green fluorescent protein (GFP) and its derivatives such as RFP, mCherry, and YFP have many applications as reporters for gene expression or as fusion proteins<sup>1,2</sup>. Although GFP is easy to use, it needs light sources to visualize the fluorescence signals. The  $\beta$ -glucuronidase (GUS) reporter has been widely used in plants for monitoring gene expression patterns and as a reporter for hormonal signaling<sup>3</sup>. For example, *DR5-GUS* transgenic lines are commonly used to monitor auxin distribution and auxin signaling<sup>4</sup>. Luciferase is another broadly used reporter in both animals and plants<sup>5</sup>. Both GUS and luciferase require the addition of expensive substrates X-Gluc (5-Bromo-4-

chloro-1*H*-indol-3-yl  $\beta$ -D-glucopyranosiduronic acid) and luciferin, respectively. Whereas the traditional reporters have been very useful, they have limitations. Fluorescent proteins are often monitored under a microscope, rendering it less useful in analyzing plants in natural growing fields or analyzing large samples such as a tree. GUS-staining is invasive and often requires sacrifice of the plants. Luciferase can be used noninvasively, but it requires a special camera and spraying the expensive substrate. It is also not very practical to use them in fields. GUS and luciferase may not be optimal for sterile conditions such as tissue culture because addition of substrates increases the chance for contamination of microbes. Therefore, there is a need to develop new reporter systems that can be widely used to monitor cellular activities noninvasively, continuously, and cost-effectively. For the past few years, gene editing has been widely used in basic research and crop improvement. A visible marker for transgenes will greatly accelerate the isolation of edited plants that no longer harbor the gene editing machinery<sup>6,7</sup>.

Plants produce many colorful compounds that potentially can serve as reporters. For example, anthocyanins display bright red-blue colors and anthocyanin-producing rice plants have been used to generate interesting patterns

Correspondence: Yubing He (yubinghe@njau.edu.cn) or Yunde Zhao (yundezhao@ucsd.edu)

<sup>1</sup>State Key Laboratory of Crop Genetics and Germplasm Enhancement, Nanjing Agricultural University, Nanjing 210095, China

<sup>2</sup>National Key Laboratory of Crop Genetic Improvement and National Center of Plant Gene Research (Wuhan), Huazhong Agricultural University, Wuhan 430070, China

Full list of author information is available at the end of the article  
These authors contributed equally: Yubing He, Tao Zhang


© The Author(s) 2020



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

in rice field. However, synthesis of anthocyanins requires multiple enzymes and varies greatly among different plants<sup>8,9</sup>. It is difficult to use anthocyanin biosynthesis pathways as a universal visible reporter. Betalains are a class of plant natural products derived from the amino-acid tyrosine<sup>10,11</sup>. The bright red color seen in beets, dragon fruit, Swiss chard, and other plants is resulted from accumulation of betalains. Biosynthesis of betalains has been well studied and only needs three enzymatic reactions to convert tyrosine into betalain (Fig. 1a)<sup>12</sup>. Tyrosine is first hydroxylated on the benzene ring, resulting in L-3,4-dihydroxyphenylalanine (L-DOPA) (Fig. 1a). The reaction is catalyzed by the P450 oxygenase CYP76AD1 (Fig. 1a). L-DOPA can be further oxidized into *cyclo*-DOPA by CYP76AD1 (Fig. 1a). Alternatively, L-DOPA is catalyzed by L-DOPA 4,5-dioxygenase (DODA) into betalamic acid, which is subsequently condensed with *cyclo*-DOPA into betanidin. The condensation reaction does not require an enzyme (Fig. 1a). Finally, a sugar moiety is added to betanidin by a glucosyltransferase to generate the colorful betalain (Fig. 1a). Betalain has a very

bright red color, which potentially can serve as a reporter to track gene expression or to visualize transgenic events. Because every cell contains the amino-acid tyrosine, exogenous application of tyrosine to tissues may not be required. We hypothesized that betalain would be a more convenient reporter than the aforementioned reporters. It is visible to naked eyes without any needs for special equipment. It does not require processing samples and it allows continuously monitoring events throughout the life cycle of an organism. Moreover, it is applicable to large plants grown under normal field conditions. Herein, we synthesize an artificial open reading frame named *RUBY* that when expressed can produce all of the enzymes required for betalain biosynthesis. We show that *RUBY* is a very effective marker for noninvasively selecting transformation events in both rice and Arabidopsis. Moreover, we show that *RUBY* can be used to visualize gene expression without any chemical treatments or special equipment, providing useful tools for visualizing gene expression in large plants under natural field growth conditions.



**Fig. 1 Components required for betalain biosynthesis.** **a** Chemical reactions for converting tyrosine into betalain, which has a red color. Tyrosine is first oxidized by the P450 CYP76AD1 into L-3,4-dihydroxyphenylalanine (L-DOPA), which can be further converted into *cyclo*-DOPA by P450 CYP76AD1. In the presence of L-DOPA 4,5-dioxygenase (DODA), DOPA is oxidized and circularized into betalamic acid. *Cyclo*-DOPA condenses with betalamic acid, a non-enzymatic reaction, to produce betanidin. Glucosylation of betanidin generates the red color betalain. **b** A strategy for expressing the whole betalain biosynthetic pathway in a single cassette. The three betalain biosynthetic genes were fused into a single open reading frame, which can be expressed using a single promoter and terminator. Between the genes, sequences that encode 2A peptides were inserted. The 2A peptides undergo self-cleavage, thus releasing the individual enzymes for betalain biosynthesis. The betalain synthesis unit can be placed under the control of a promoter of interest. The terminator used here was the Arabidopsis *HSP18.2* terminator. The open reading frame of 2A-linked betalain biosynthesis genes is named *RUBY*.

## Results and discussion

### Construction of a single open reading frame for the betalain biosynthetic pathway

Heterologous expression of *CYP76AD1*, *DODA* in tobacco, and other plants demonstrated that the betalain biosynthetic pathway can be re-constituted in plant cells<sup>13,14</sup>. In order to use betalain as a visual reporter, we need to effectively co-express the entire pathway using a single promoter. We organized *CYP76AD1*, *DODA*, and *Glucosyltransferase* into a single open reading frame (Fig. 1b) (Supplemental Fig. 1). The stop codons of *CYP76AD1* and *DODA* were removed. The three genes were linked by sequences that encode 2A peptides (Fig. 1b) (Supplemental Fig. 1)<sup>15,16</sup>. Upon transcription, the single transcript, which includes the coding regions of the three enzymes, produced the three separate enzymes through either 2A-mediated self-cleavage or ribosomal “skipping” (Fig. 1b)<sup>17</sup>. The 2A system enables the expression of multiple proteins under the control of a single promoter. We name the 2A-linked unit of *CYP76AD1*, *DODA*, and *Glucosyltransferase* *RUBY* (Fig. 1b). *RUBY* can be expressed when a promoter is placed in front of it. The expression pattern and level of a particular gene may be inferred from the red color of betalain if the gene’s promoter is used to drive *RUBY* expression.

### *RUBY* is capable of synthesizing betalain in tobacco

We first placed *RUBY* under the control of Cauliflower Mosaic Virus (CaMV) 35S promoter, which is a widely used constitutively strong promoter<sup>18</sup>. To test whether *RUBY* can produce functional enzymes for betalain synthesis, we infiltrated tobacco leaves with *Agrobacterium* that contain *RUBY*-expressing plasmid (Supplemental Fig. 2). Transient expression of *RUBY* led to the production of betalain in tobacco leaves, suggesting that the synthetic open reading frame *RUBY* can produce the functional enzymes for the synthesis of betalain. Moreover, we observed that betalain was not transported from the spots of *Agrobacterium*-infiltration spots to other leaves of the plant (Supplemental Fig. 2).

### Synthesis of betalain by *RUBY* in *Arabidopsis*

We transformed the 35S:*RUBY* construct into *Arabidopsis* using *Agrobacterium*-mediated floral dipping<sup>19</sup>. Two days after floral dipping, we noticed that the transformed plants displayed patches of red color (Supplemental Fig. 3a), indicating that the *RUBY* cassette was functionally expressed and that *RUBY* may be used to monitor transient *Arabidopsis* transformation. Once the seeds from the *Agrobacterium*-dipped plants were harvested, transgenic seeds could be easily differentiated from non-transgenic seeds (Supplemental Fig. 3b). The transformed seeds had a dark red color (Supplemental Fig. 3b), demonstrating that *RUBY* can be used as a visual


selection marker for transgenic events in *Arabidopsis*. We previously used mCherry as a very effective marker to select transgenic events (Gao et al., 2016), which requires a dissecting microscope with fluorescence capability. *RUBY* is a better option because it does not require special equipment.

The 35S:*RUBY* plants produced sufficient amount of betalain to become visually evident (Fig. 2a). Consistent with previous reports that *CaMV* 35S promoter is constitutively active, we observed red color in all tissues throughout the plant life cycle (Fig. 2a) (Supplemental Fig. 3c, d). We also expressed *RUBY* reporter under the control of the Maize *UBIQUITIN* promoter, which has been widely used to overexpress genes in monocots<sup>20</sup>. Similar to 35S:*RUBY* plants, *UBQ*:*RUBY* plants were also visibly red in leaves, stem, and flowers (Fig. 2b). These results clearly demonstrated that *RUBY* could be expressed in *Arabidopsis* and that our *RUBY* reporter was able to functionally re-constitute the betalain biosynthetic pathway.

We expressed *RUBY* using the seed specific *At2S3* promoter, which we previously used to drive *mCherry* expression in *Arabidopsis* to facilitate the selection of transgenes<sup>6</sup>. As shown in Fig. 2c, the transgenic plants were indistinguishable from wild type plants. When we checked the seeds in a silique from an *At2S3*:*RUBY* T1 plant, *RUBY*-expressing seeds displayed strong red color, whereas the non-transgenic seeds were green (Fig. 2c). *RUBY* can be conveniently used to select single T-DNA insertion events by analyzing the ratio of red seeds to green seeds, which should be ~3:1 for single insertions. The *At2S3*:*RUBY* results demonstrate that *RUBY* could be an effective marker for *Arabidopsis* transformation. Furthermore, betalain was not widely transported from the sites of synthesis to other tissues as we did not see any red color in leaves (Fig. 2c). We also expressed *RUBY* under the control of the *Arabidopsis* *YUC4* promoter (Fig. 2d). *YUC4*, which encodes a key enzyme in auxin biosynthesis, was shown to express in small regions of embryos, leaves, and flowers<sup>21,22</sup>. GUS signals were observed in leaf tips and apical region of a gynoecium in *YUC4* promoter:*GUS* transgenic plants. We observed similar patterns of betalain production in *YUC4*:*RUBY* lines (Fig. 2d).

### *RUBY* synthesizes betalain in rice

Unlike *Arabidopsis*, rice and many other plants are transformed through tissue culture and the formation of calli, which are often mosaic. A visible marker for transformation at tissue culture stage will be very useful. We placed *RUBY* under the control of *DR5*, an auxin-responsive synthetic promoter<sup>4</sup>, which was used to monitor stem cell initiation during tissue culture. We also used the rice *ACTIN1* promoter, which is considered a strong promoter, to control *RUBY* expression.



**Fig. 2 RUBY serves as an effective reporter for gene expression and plant transformation.** **a** *RUBY* expression driven by the *CaMV 35S* promoter led to a red *Arabidopsis* plant (right) compared with the WT plant (left). **b** *UBIQUITIN* promoter was effective in driving *RUBY* expression throughout the plant (right). Non-transgenic WT was shown left. **c** The seed specific promoter *At2S3* did not lead to *RUBY* expression in leaves and stems. The transgenic plant (right) and non-transgenic plant (left) were indistinguishable. The siliques of *At2S3:RUBY* were similar to those of WT, however, when the silique was opened, the seeds of *At2S3:RUBY* were clearly red. Moreover, it was obvious that transgenic and no-transgenic seeds were segregating in a silique from a T1 *At2S3:RUBY* plant. **d** *YUC4:RUBY* plants displayed patches of red at the tip of leaves and apical region of a gynoecium. **e** *DR5:RUBY* was expressed in rice calli. The red color can be used to distinguish transgenic (red) and non-transgenic calli (white). *DR5:eGFP* has been used in rice calli **f**, but it was much more difficult to distinguish transgenic from non-transgenic using *DR5:eGFP* compared with *RUBY*. **g** Roots of *DR5:RUBY* and *DR5:eGFP* rice plants, which had similar patterns. The three *DR5:eGFP* pictures were generated with the same root: bright field (left), 488 nm fluorescence field (middle), and the merged (right). **h** Activation of *DR5* promoter in *DR5:RUBY* leaves was easy to observe whereas *DR5:eGFP* was much more difficult to detect: bright field (left), 488 nm fluorescence field (middle), and the merged (right).

We transformed the *DR5:RUBY* (Fig. 2e) and *OsACTIN1:RUBY* (Supplemental Fig. 4) into rice calli. *RUBY* expression rendered calli vividly red. The presence of the *RUBY* gene correlated to the observed *RUBY* expression in rice calli (Supplemental Fig. 4). We also observed a positive correlation between the brightness of the calli the expression levels of *RUBY* gene (Supplemental Fig. 4). Introduction of *RUBY* made it easier to distinguish the transformed calli from untransformed calli (Fig. 2e). *RUBY* enables selection of calli that have better expression of transgenes among the hygromycin-resistant calli in the same tissue block (Supplemental Fig. 4). In comparison, *DR5:eGFP* has been used to visualize transgenic calli (Fig. 2f). Our *RUBY* system is clearer and much more convenient to operate during tissue culture condition. The *DR5:RUBY* and *DR5:eGFP* in rice roots showed similar patterns (Fig. 2g), but *DR5:RUBY* was easier to observe without any treatments or a change in light conditions. The advantages of *DR5:RUBY* over *DR5:eGFP* was also obvious in rice leaves (Fig. 2h). Moreover, *RUBY* as a visible marker was very useful for monitoring transgenes in intact adult rice plants (Supplemental Fig. 5).

We demonstrate that our synthetic cassette of betalain biosynthetic genes was able to produce betalain in *Arabidopsis* and in rice, providing a visible color for monitoring gene expression and plant transformation. We believe that *RUBY* will be very useful in large plants such as fruit trees and in field conditions. Because *RUBY* does not require either special equipment or expensive substrates, *RUBY* provides a cost-effective reporter and *RUBY* is a convenient alternative to the existing reporters. We envision that *RUBY* can be adapted for applications in some microbes and animals because the substrate tyrosine exists in all cells. For example, *RUBY* may provide a more convenient marker than  $\beta$ -galactosidase (LacZ) in yeast two-hybrid screens. Betalain is a natural product and was shown to have health benefits. Using *RUBY* as a reporter has less environmental and health concerns compared with antibiotic and/or herbicide resistance markers.

## Materials and methods

### *Arabidopsis* constructs and transformation

The backbone of the *RUBY* constructs for *Arabidopsis* is the plasmid *pHDE*, which was described previously<sup>6</sup>. We took advantage of the P2A peptide, which has the sequence of (GSGATNFSLLKQAGDVEENPGP), to link the *CYP76AD1*, *DODA*, and *glucosyltransferase* (*GT*) coding regions. The transcriptional terminator used was the *HSP18.2* terminator from *Arabidopsis*. The 2A-linked *CYP76AD1*, *DODA*, and *GT* unit was named *RUBY*, which was cloned into *pHDE* along with the HSP terminator by Gibson assembly at the *XbaI*<sup>23</sup>. The entire sequence of *RUBY* and terminator was shown in Supplemental Fig. 1.

Various promoters can be cloned into the *PmeI* site of *pHDE-RUBY* to drive *RUBY* expression. Promoters were amplified by PCR and the primers used in this study were listed in Supplemental Table 1. We expressed *RUBY* using *CaMV 35S*, Maize *UBIQUITIN* promoter, *At2S3* promoter, and *YUC4* promoter to test whether *RUBY* can serve as an effective reporter. The constructs were transformed into *Arabidopsis* Columbia plants by *Agrobacterium*-mediated floral dipping<sup>19</sup>. Transgenic seeds for *35S:RUBY*, *UBQ:RUBY*, and *At2S3:RUBY* were easily identified by red color. Transgenic plants for *YUC4:RUBY* were selected on MS medium containing hygromycin (16.7  $\mu$ g/ml).

### Rice constructs and transformation

The auxin-responsive promoter (*DR5*) was composed of a synthetic promoter with 16 repeats of core auxin response element (AuxRE) sequence (TGTCTC) linked with *CaMV* minimal *35S* promoter<sup>4</sup>. The *DR5* promoter was synthesized and then cloned into *pDX2181*<sup>24</sup> between *BamH* I and *Pst* I sites by Gibson assembly with primer pair Oligo15 and Oligo16 (Supplemental Table 1), resulting in the plasmid *pDR5:eGFP*.

To construct the *pDR5:RUBY* plasmid, we linked the three betalain biosynthetic genes through 2A peptides. Here, we used the F2A peptide, which has the following peptide sequence: QLLNFDLLKLAGDVESNPGP. The *RUBY* cassette replaced the *eGFP* in the *pDR5:eGFP* plasmid, resulting in the *pDR5:RUBY*. The rice version of *RUBY* was also detailed in the Supplemental Fig. 1. Both the *DR5:RUBY* and *DR5:eGFP* plasmids were transformed into Xiaowei<sup>NIP25</sup> through *Agrobacterium*-mediated plant transformation following a protocol that was previously described<sup>26</sup>.

### Quantitative analysis the expression level of *RUBY*

The relative expression levels of *RUBY* in the transgenic calli of *OsACTIN1:RUBY* were determined by reverse transcription quantitative PCR using the primer pair Oligo26/Oligo27. The primer pair Oligo28/ Oligo29 was specific for the rice *UBIQUITIN* (*UBQ*) gene, which served as the endogenous reference gene<sup>27</sup>. The primers were listed in Supplemental Table 1.

### Acknowledgements

This work was partially supported by grants from the National Transgenic Science and Technology Program (2019ZX08010-003; 2019ZX08010-001) to Y.H. and H.Z. T.Z. is a TIGS postdoctoral fellow. We thank Mr. Vitor Pinoti for helping the tobacco transformation.

### Author details

<sup>1</sup>State Key Laboratory of Crop Genetics and Germplasm Enhancement, Nanjing Agricultural University, Nanjing 210095, China. <sup>2</sup>National Key Laboratory of Crop Genetic Improvement and National Center of Plant Gene Research (Wuhan), Huazhong Agricultural University, Wuhan 430070, China. <sup>3</sup>Section of Cell and Developmental Biology, University of California San Diego, 9500

Gilman Drive, La Jolla, CA 92093-0116, USA. <sup>4</sup>Tata Institute for Genetics and Society-UCSD, La Jolla, CA 92093-0335, USA

#### Author contributions

Y.H. and Y.Z. conceived the idea. Y.H. and H.S. conducted the rice experiments. T.Z. performed the Arabidopsis experiments. Y.H., T.Z., and Y.Z. wrote the manuscript.

#### Data availability

The plasmids will be available at Addgene.

#### Conflict of interest

The authors declare that they have no conflict of interest.

**Supplementary Information** accompanies this paper at (<https://doi.org/10.1038/s41438-020-00390-1>).

Received: 6 May 2020 Accepted: 18 August 2020

Published online: 19 September 2020

#### References

- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W. W. & Prasher, D. C. Green fluorescent protein as a marker for gene expression. *Science* **263**, 802–805 (1994).
- Heim, R., Cubitt, A. B. & Tsien, R. Y. Improved green fluorescence. *Nature* **373**, 663–664 (1995).
- Jefferson, R. A., Kavanagh, T. A. & Bevan, M. W. GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**, 3901–3907 (1987).
- Sabatini, S. et al. An auxin-dependent distal organizer of pattern and polarity in the Arabidopsis root. *Cell* **99**, 463–472 (1999).
- Contag, C. H. & Bachmann, M. H. Advances in in vivo bioluminescence imaging of gene expression. *Annu. Rev. Biomed. Eng.* **4**, 235–260 (2002).
- Gao, X., Chen, J., Dai, X., Zhang, D. & Zhao, Y. An effective strategy for reliably isolating heritable and Cas9-free Arabidopsis mutants generated by CRISPR/Cas9-mediated genome editing. *Plant Physiol.* **171**, 1794–1800 (2016).
- He, Y. & Zhao, Y. Technological breakthroughs in generating transgene-free and genetically stable CRISPR-edited plants. *ABIOTECH* **1**, 88–96 (2020).
- Misyura, M., Colasanti, J. & Rothstein, S. J. Physiological and genetic analysis of Arabidopsis thaliana anthocyanin biosynthesis mutants under chronic adverse environmental conditions. *J. Exp. Bot.* **64**, 229–240 (2013).
- Li, P. et al. Regulation of anthocyanin and proanthocyanidin biosynthesis by *Medicago truncatula* bHLH transcription factor MtTT8. *N. Phytol.* **210**, 905–921 (2016).
- Strack, D., Vogt, T. & Schliemann, W. Recent advances in betalain research. *Phytochemistry* **62**, 247–269 (2003).
- Xu, J.-J., Fang, X., Li, C.-Y., Yang, L. & Chen, X.-Y. General and specialized tyrosine metabolism pathways in plants. *ABIOTECH*, <https://doi.org/10.1007/s42994-019-00006-w> (2019).
- Polturak, G. & Aharoni, A. Advances and future directions in betalain metabolic engineering. *N. Phytol.* **224**, 1472–1478 (2019).
- Polturak, G. et al. Elucidation of the first committed step in betalain biosynthesis enables the heterologous engineering of betalain pigments in plants. *N. Phytol.* **210**, 269–283 (2016).
- Polturak, G. et al. Engineered gray mold resistance, antioxidant capacity, and pigmentation in betalain-producing crops and ornamentals. *Proc. Natl Acad. Sci. USA* **114**, 9062–9067 (2017).
- Liu, Z. et al. Systematic comparison of 2A peptides for cloning multi-genes in a polycistronic vector. *Sci. Rep.* **7**, 2193 (2017).
- Wang, J. & Chen, H. A novel CRISPR/Cas9 system for efficiently generating Cas9-free multiplex mutants in Arabidopsis. *ABIOTECH* **1**, 6–14 (2020).
- Sharma, P. et al. 2A peptides provide distinct solutions to driving stop-carry on translational recoding. *Nucleic Acids Res.* **40**, 3143–3151 (2012).
- Benfey, P. N. & Chua, N. H. The cauliflower mosaic virus 35S promoter: combinatorial regulation of transcription in plants. *Science* **250**, 959–966 (1990).
- Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J.* **16**, 735–743 (1998).
- Cornejo, M. J., Luth, D., Blankenship, K. M., Anderson, O. D. & Blechl, A. E. Activity of a maize ubiquitin promoter in transgenic rice. *Plant Mol. Biol.* **23**, 567–581 (1993).
- Cheng, Y., Dai, X. & Zhao, Y. Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in Arabidopsis. *Genes Dev.* **20**, 1790–1799 (2006).
- Cheng, Y., Dai, X. & Zhao, Y. Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in Arabidopsis. *Plant Cell* **19**, 2430–2439 (2007).
- Gibson, D. G. et al. Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods* **6**, 343–345 (2009).
- Ye, R., Zhou, F. & Lin, Y. Two novel positive cis-regulatory elements involved in green tissue-specific promoter activity in rice (*Oryza sativa* L. ssp.). *Plant Cell Rep.* **31**, 1159–1172 (2012).
- Hu, S. et al. Xiaowei, a new rice germplasm for large-scale indoor research. *Mol. Plant* **11**, 1418–1420 (2018).
- Hiei, Y., Ohta, S., Komari, T. & Kumashiro, T. Efficient transformation of rice (*Oryza sativa* L.) mediated by Agrobacterium and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**, 271–282 (1994).
- He, Y. et al. PINOID is required for formation of the stigma and style in rice. *Plant Physiol.* **180**, 926–936 (2019).