


Article

Metal- and UV- Catalyzed Oxidation Results in Trapped Amyloid- β Intermediates Revealing that Self-Assembly Is Required for A β -Induced Cytotoxicity

Mahmoud B. Maina, Gunasekhar Burra, Yousra K. Al-Hilaly, Kurtis Mengham, Kate Fennell, Louise C. Serpell

l.c.serpell@sussex.ac.uk

HIGHLIGHTS

Metal- (Cu^{2+} H_2O_2) or UV-catalyzed oxidation results in dityrosine (DiY) formation

Oxidation results in DiY cross-link formation in A β and halts further assembly

Non-assembling A β (trapped A β or variant A β monomer) is not cytotoxic

Maina et al., iScience 23, 101537
October 23, 2020 © 2020 The Author(s).
<https://doi.org/10.1016/j.isci.2020.101537>

Article

Metal- and UV- Catalyzed Oxidation Results in Trapped Amyloid- β Intermediates Revealing that Self-Assembly Is Required for A β -Induced Cytotoxicity

Mahmoud B. Maina,^{1,2} Gunasekhar Burra,^{1,4} Youssra K. Al-Hilaly,^{1,3} Kurtis Mengham,¹ Kate Fennell,¹ and Louise C. Serpell^{1,5,*}

SUMMARY

Dityrosine (DiY), via the cross-linking of tyrosine residues, is a marker of protein oxidation, which increases with aging. Amyloid- β (A β) forms DiY *in vitro* and DiY-cross-linked A β is found in the brains of patients with Alzheimer disease. Metal- or UV- catalyzed oxidation of A β 42 results in an increase in DiY cross-links. Using DiY as a marker of oxidation, we compare the self-assembly propensity and DiY cross-link formation for a non-assembly competent variant of A β 42 (vA β) with wild-type A β 42. Oxidation results in the formation of trapped wild-type A β assemblies with increased DiY cross-links that are unable to elongate further. Assembly-incompetent vA β and trapped A β assemblies are non-toxic to neuroblastoma cells at all stages of self-assembly, in contrast to oligomeric, non-cross-linked A β . These findings point to a mechanism of toxicity that necessitates dynamic self-assembly whereby trapped A β assemblies and assembly-incompetent variant A β are unable to result in cell death.

INTRODUCTION

Alzheimer disease (AD) is the most common form of dementia, and it is characterized by the deposition of amyloid- β (A β) and Tau in extracellular plaques and intracellular neurofibrillary tangles, respectively. The amyloid cascade hypothesis implicates the pathological accumulation of A β and its aggregation from monomers into oligomers and fibrils, as a key event in the development of AD (Hardy and Higgins, 1992). Many pieces of evidence have subsequently shown that the oligomeric form of A β is the most toxic species, resulting in a reformulated amyloid cascade hypothesis in which A β oligomers are proposed to be central to AD pathogenesis (Selkoe and Hardy, 2016). Indeed, accumulated evidence shows that A β oligomers disrupt cellular function in cultured cells and animal models (Marshall et al., 2016; Lambert et al., 1998; Lacor et al., 2007; Zhang et al., 2014; Selkoe and Hardy, 2016). Numerous studies have searched for the elusive “toxic” species and tried to characterize its structure. For example, 12mers, *56 KDa, and hexamers have all been implicated as specific structural species that interact with particular receptors (e.g., NDMA) leading to downstream cell death (Lesne et al., 2006; Reed et al., 2011).

Oxidative stress has been proposed as a key mechanism that mediates A β toxicity (Butterfield and Halliwell, 2019; Butterfield et al., 2013) and, it is potentially one of the earliest sources of damage in human AD (Nunomura et al., 2001). Furthermore, using a cellular model, we have shown that oxidative stress is one of the earliest events induced by A β oligomers (Maina et al., 2018). One of the ways that oxidative stress causes cellular damage is through protein oxidation. The most common consequences of protein oxidation include amino acid side-chain modification, protein fragmentation, and protein cross-linking (e.g., via dityrosine [DiY] bond formation) (Lund et al., 2011). DiY cross-linking is mediated via carbon-carbon bonding between two proximal tyrosines, resulting in a stable, non-reversible covalent bond (Gross and Sizer, 1959). DiY cross-linking is known to provide elasticity, strength, and stability to proteins and has been shown to form within proteins involved with neurodegenerative diseases (e.g., A β and α -synuclein) (Galeazzi et al., 1999; Souza et al., 2000). Indeed, we have previously shown the colocalization of DiY with A β in plaques and α -synuclein in Lewy bodies in human AD and Parkinson disease brain tissue, respectively (Al-Hilaly et al., 2013, 2016). In addition to DiY, other modifications such as oxidation of histidine,

¹Sussex Neuroscience, School of Life Sciences, University of Sussex, Falmer, Brighton, East Sussex BN1 9QG, UK

²College of Medical Sciences, Yobe State University, Yobe, Nigeria

³Chemistry Department, College of Science, Mustansiriyah University, Baghdad, Iraq

⁴Present Address: Analytical Research & Development, Pharma Division, Biological E. Limited, Genome Valley, IKP-Shameerpet, Hyderabad-500078, Telangana, India

⁵Lead Contact

*Correspondence:

l.c.serpell@sussex.ac.uk

<https://doi.org/10.1016/j.isci.2020.101537>



lysine, and methionine³⁵ (met³⁵) have been shown to occur in A β (Kowalik-Jankowska et al., 2004; Ali et al., 2005; Palmblad et al., 2002; Friedemann et al., 2015), whereas hydroxylated phenylalanine has been suggested to form cross-links alongside DiY (Zhang et al., 2019). This indicates the potential relevance of these modifications in disease pathogenesis or as markers of disease progression in AD.

To learn more about the importance of DiY cross-linking in AD specifically, several studies have investigated the impact of DiY bond formation on A β aggregation and toxicity, mostly using metal-catalyzed oxidation (e.g., Cu²⁺/H₂O₂) (MCO) and peroxidase-catalyzed oxidation. Some studies have shown that DiY cross-linking of both A β 40 and A β 42 results in the generation of toxic A β assemblies with reduced assembly speed (Barnham et al., 2004; Smith et al., 2006; Kok et al., 2013; O'malley et al., 2014, 2016; Sitkiewicz et al., 2014). Others have implicated DiY in the inhibition of A β 40 assembly, especially in highly oxidative environments (Gu et al., 2018) and it has been demonstrated to be associated with the formation of non-amyloidogenic A β 42 aggregates when catalyzed with a high concentration of Cu²⁺ (Smith et al., 2007). However, whether DiY cross-linking is a driver, facilitator, or inhibitor of A β self-assembly remains unclear. Moreover, given that A β self-assembly is rapid, the time point of the cross-linking during the assembly process may influence the nature of the cross-linked A β assemblies. Here, using MCO and UV-induced oxidation to induce DiY cross-linking, we show that the oxidation process results in the stabilization of A β 42 assemblies and either prevents or very significantly slows further elongation. To specifically compare the influence of DiY cross-linking on A β assembly, we compared the effect of oxidation on a non-assembly competent variant A β (vA β) (Marshall et al., 2016) and revealed that oxidation and DiY cross-linking does not induce or promote its assembly. We show that in the absence of H₂O₂, CuCl₂ at a concentration similar to the concentration found around A β plaques (~400 μ M) (Lovell et al., 1998) is sufficient to facilitate the DiY cross-linking and formation of A β 42 oligomers into a long-lived oligomer population. A cell live/dead assay revealed that, unlike the self-assembling non-cross-linked A β , oxidized DiY containing A β 42 is non-toxic to neuron-like cells. Our results suggest that under certain conditions *in vitro*, oxidation can result in trapping of intermediate species, which cannot elongate further and that are non-toxic to neuroblastoma cells. Together with the observation that non-assembly-competent variant A β is non-toxic, this reinforces the importance of a continual self-assembly process in mechanisms of A β toxicity.

RESULTS

***In Vitro* Metal-Catalyzed Oxidation Results in the Formation of Dityrosine in Wild-Type and vA β Peptides**

To investigate the influence of oxidation on A β assembly, we compared the effect of MCO using CuCl₂ and H₂O₂ on wild-type A β 42 and variant A β 42 (henceforth called A β and vA β , respectively) (see [Transparent Methods](#) in Information for Authors for more details).

DiY serves as a useful measure of the levels of oxidation and was used here to follow oxidation effects. Other side chains can be oxidized such as Met³⁵, which can be detected using matrix-assisted laser desorption ionization mass spectrometry (Friedemann et al., 2015). We have previously detected DiY cross-links in A β and α -synuclein using a combination of techniques and shown that DiY can reliably be detected using fluorescence spectroscopy (Al-Hilaly et al., 2013, 2016). Here, rapid formation of DiY was detected for A β and vA β samples that were incubated with both CuCl₂ and H₂O₂ (A β /CuCl₂/H₂O₂ and vA β /CuCl₂/H₂O₂, henceforth called MCO), indicated by the observation of a fluorescence peak at 410 nm after only 5 min. In contrast, samples incubated with CuCl₂ alone or with buffer-only showed no peak at 410 nm (Figure 1A). Detection of tyrosine with an excitation/emission of 280/305 nm (following quenching using EDTA) showed that the formation of DiY was matched by a decrease in tyrosine fluorescence in both the A β and vA β MCO reactions compared with the samples incubated with CuCl₂ alone or buffer only (Figure 1B). DiY fluorescence intensity continued to increase for the MCO reactions up to 2 h but did not increase further after 5 days (Figure 1C). However, by 5 days, small peaks could be observed for samples incubated with CuCl₂ alone (Figure 1D), although the DiY signal remained negligible for buffer-only samples after 5 days incubation in the dark. Standard curve estimation of DiY content (Figure S1) revealed that 2 h MCO of A β and vA β induced about 5 μ M and ~12 μ M DiY, respectively. As each DiY is contributed to by two molecules of A β , the percentage of A β molecules where Tyr is cross-linked is approximately 20% for A β and 48% for vA β . Incubation of A β with CuCl₂ alone for 2 h induced ~1 μ M DiY, which further increased to ~2 μ M at 5 days, which equates to approximately 4% of A β molecules participating in cross-links. A β and vA β samples incubated with CuCl₂ showed a decrease in tyrosine fluorescence after only 5 min (Figure 1B), indicating that the CuCl₂ rapidly induces conformational changes (Roberts et al., 2012) in both A β and vA β . This appears to




Figure 1. DiY Formation in A β and vA β via Metal-Catalyzed Oxidation

(A–D) Freshly prepared A β and vA β (50 μ M) were incubated at 37°C, without CuCl₂, with CuCl₂ (400 μ M), or CuCl₂ in combination with H₂O₂. Fluorescence spectra were collected 5 min post-incubation using fluorescent excitation wavelength of 320 nm and emission collected between 340 and 600 nm, with DiY peak signal observed between 400 and 420 nm (A). Fluorescence spectra were also collected at 5 min using an excitation wavelength of 280 nm and emission collected between 290 and 500, with peak tyrosine signal observed at 305 nm (B). Fluorescence spectra were collected again at 2 h (C) and then 5 days (D) to follow DiY formation over time. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings.

be independent of DiY cross-linking, which only starts appearing ~2 h post-incubation. Overall, this suggests that the incubation of both A β and vA β with CuCl₂ alone or in combination with H₂O₂ results in the formation of DiY cross-links. DiY formation in the different samples occurs after different lengths of time depending on the oxidation conditions, which may impact A β assembly properties.

Metal-Catalyzed Oxidation Influences the Assembly of Wild-Type A β , but Does Not Affect the Structure and Aggregation Propensity of Variant A β

We have previously shown that DiY forms in both A β 42 oligomers and fibrils (Al-Hilaly et al., 2013). Thioflavin T (Th-T) fluorescence assay was used to investigate whether DiY formation correlates with changes in the A β assembly (see [Transparent Methods](#)). As expected, the assembly-incompetent vA β incubated in buffer shows no increase in Th-T fluorescence with time (Marshall et al., 2016). Wild-type A β gave the expected Th-T spectra showing a sigmoidal curve (Figure 2A) and the Th-T fluorescence increased further when A β was assembled in the presence of the metal chelator EDTA, which suggests some impact of trace metals on assembly properties (Figure S2). However, the A β and vA β MCO reactions showed no Th-T fluorescence signal increase over the time frame of 50 h, indicating that either no self-assembly had taken place or that the assembly is significantly slow enough that the threshold of Th-T detection had not been reached. Wild-type A β incubated with CuCl₂ shows a small fluorescence signal for DiY at ~2 h (Figure 1C) and Th-T spectrum showed a short lag phase (approximately up to 2 h) followed by a plateau at a low fluorescence signal. This appears to suggest that the A β assemblies formed become stabilized without further elongation (Figure 2A). vA β incubated with CuCl₂ showed no increase in Th-T fluorescence, consistent with the absence of assembly under these conditions.

We have previously shown that our A β preparation method results in the generation of monomers that assemble into oligomers detected by the oligomer-specific antibody, NU-1 (Lambert et al., 2007), before forming fibrils and amyloid plaques (Marshall et al., 2016) (see [Transparent Methods](#)). As expected, dot blotting with the NU-1 antibody revealed the presence of oligomers at 2 h in buffer-incubated A β samples, which disappeared over time (Figure 2B). In contrast, the buffer-incubated vA β reactions showed no NU-1 reactivity, indicating the absence of oligomer formation as previously reported (Figure 1F) (Marshall et al., 2016). A β oligomers were only minimally detected in the A β MCO reaction at 5 min, which disappeared over time (Figures 2B and 2C). No NU-1 reactivity was observed for vA β MCO reaction at any of the time points measured. Interestingly, the A β incubated with CuCl₂ formed more oligomers early on, which persisted throughout the time studied. The data are consistent with the possibility that slower oxidation by CuCl₂ facilitates A β oligomer formation and stabilization. The formation of DiY cross-linking correlates with this time point when A β is found in a NU-1 affinity conformation (Figures 2B and 2C).

Circular dichroism (CD) and transmission electron microscopy (TEM) were used to study the secondary structure and apparent morphologies of the resulting assemblies after 5 days incubation (see [Transparent Methods](#)). Spectra from vA β under all three conditions showed a trough at 198 nm consistent with random




Figure 2. Metal-Catalyzed Oxidation Results in the Formation of Stabilized A β Assemblies

(A–E) Th-T fluorescence was monitored for the freshly prepared A β and vA β (50 μ M) incubated at 37°C, without CuCl₂, with CuCl₂ (400 μ M), or CuCl₂ in combination with H₂O₂ (A). Dot blotting using NU-1 antibody identified A β oligomers in the A β samples, but not in vA β reactions (B). Quantification of dot blotting signal over time reveals that the oligomers in the CuCl₂-oxidized A β remain stable for 5 days, unlike in the other reactions in which very low NU-1 affinity signal was observed (C). CD at 5 days showed a high β -sheet content in the unoxidized A β sample, with reduced signal for CuCl₂ A β , and CuCl₂/H₂O₂A β . All vA β samples showed spectra consistent with random coil conformation (D). TEM imaging at 5 days revealed a network of fibers in the unoxidized A β , whereas the CuCl₂ A β showed clumped assemblies with very little fiber density and the CuCl₂/H₂O₂ sample revealed amorphous-like assemblies. Unoxidized vA β showed no assemblies, whereas both oxidized vA β reactions showed amorphous-like aggregates (E). A minimum of three independent experiments was repeated to ensure the reproducibility of the findings. Scale bars, 500 nm. Error bars are expressed as \pm SEM.

coil conformation. A β in buffer showed a minimum at 218 nm consistent with an expected β -sheet conformation, typical of fibrillar A β (Figure 2D). A β CuCl₂ shows a broad but weak minimum at 218 nm, but the signal from A β MCO reaction is too weak for assignment of secondary structure. This may be consistent with loss of protein from solution, which may arise from the formation of small, amorphous oligomers.

Electron micrographs showed an extensive fibril network for buffer-incubated A β , whereas CuCl₂-oxidized A β showed clumped assemblies with a significantly reduced fiber density, and the MCO A β sample revealed amorphous structures with very few or no fibers (Figure 2E). Buffer-incubated vA β sample showed very few assemblies of any kind. The vA β MCO and CuCl₂ reactions exhibited a few amorphous-like clumped assemblies. The CD and TEM data suggest that the MCO-catalyzed oxidation of A β prevents the formation of β -sheet-rich amyloid fibrils, whereas CuCl₂-catalyzed oxidation generates oligomeric A β assemblies with some β -sheet conformation. Taken together, the results from Th-T, dot blotting, CD, and TEM revealed that the rapid DiY cross-linking of A β induced through MCO correlates with the association and trapping of A β as intermediates and inhibits their further assembly, whereas CuCl₂ facilitates the formation and DiY cross-linking of A β oligomers into a long-lived oligomer population. In contrast, none of the conditions induced any assembly of vA β into amyloid fibrils. Instead, the MCO and CuCl₂ reactions resulted in association and trapping of vA β into amorphous assemblies that show no evidence of β -sheet conformation.

Photo-oxidation Induces Dityrosine Cross-Linking in Wild-Type A β and vA β Peptides

The aforementioned results suggest that copper contributes to DiY formation. However, previous studies have suggested that metals influence the assembly of A β (Barritt and Viles, 2015; Gu et al., 2018; Smith et al., 2006, 2007), and this is supported by the increased assembly of A β in the presence of EDTA




Figure 3. DiY Formation in A β and vA β via UV Photo-oxidation

(A–C) Freshly prepared A β and vA β (50 μ M) were incubated under UV. Fluorescence spectra were collected 5 min post-incubation using fluorescent excitation wavelength of 320 nm and emission collected between 340 and 600 nm, with DiY peak signal observed between 400 and 420 nm after 5 min of incubation (A), which increased following 2 h of incubation (B). Fluorescence intensity at 410 nm against time showed that incubation of the 2 h UV-exposed A β and vA β samples in the dark resulted in further increase in DiY formation in the absence of the UV (C). The A β and vA β samples that were not exposed to UV showed no DiY signal. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings.

(Figure S2). Therefore, to avoid the complication of using metals, which may influence assembly, UV photo-oxidation was used to oxidize A β and vA β (Figure 3) (see Transparent Methods). UV oxidation of amino acids is limited to direct damage to Trp, Tyr, His, Cys (Figure 7) and cystine, but Met can be oxidized indirectly, usually in the presence of a sensitizer (Pattison et al., 2012). As before, the measurement of DiY formation was used to follow oxidation. Fluorescence spectra showed a small peak at 410 nm for both A β and vA β after only 5 min incubation in UV. However, the intensity was lower than under MCO conditions (Figure 3A). After 2 h of UV exposure, a significant increase in intensity at 410 nm was observed (Figure 3B). Calculation of DiY content using the standard curve shown in Figure S1 suggested that approximately 12% of A β and 14% of vA β molecules were involved in DiY, which is lower than following MCO treatment. After 2 h of UV exposure, samples were stored in the dark but despite this the intensity continued to increase for 120 h following incubation (18% DiY) (Figure 3C). These results show that UV photo-oxidation can induce DiY cross-linking for A β and vA β peptides. The results are compared with peptides incubated without UV exposure for reference. Oxidation of other residues by the UV irradiation can not be ruled out, but here we focus on DiY to monitor the level of oxidation.

Photo-oxidation Influences the Assembly of Wild-Type A β , but Does Not Impact on the Structure and Assembly of vA β

Th-T fluorescence was used to monitor the assembly of A β and vA β following UV photo-oxidation for 2 h. The results are compared with peptides incubated without UV exposure for reference. A small increase in Th-T fluorescence was observed for A β +UV after a lag phase of 25 h, which may represent a slow assembly of the A β assemblies that escaped the less-efficient UV oxidation, whereas vA β + UV showed no fluorescence intensity (Figure 4A). NU-1 dot blots showed no reactivity to vA β + UV (Figure 4B). The A β +UV sample showed lower level NU-1 reactivity at 2 h compared with the A β control (Figure 4B), and reactivity disappeared after 5 days post-UV incubation in the cross-linked A β +UV sample, even though a small quantity of oligomers could still be detected in the control A β sample (Figure 4B). CD showed spectra consistent with random coil conformation for both vA β and A β incubated under UV and a loss of signal that was more evident for A β than for vA β (Figure 4C). By TEM, A β +UV and vA β +UV again showed small assemblies at 2 h, which are still present alongside occasionally clumped amorphous-like assemblies after 5 days (Figures 4D and 4E). Similar to the results from MCO (Figure 2B), this suggests that the UV-induced DiY cross-linking correlates with the stabilization of A β and vA β in a trapped, assembly-incompetent species. Other oxidation reactions may also be involved.

Co-incubation with Oxidized A β Slows the Assembly of Freshly Prepared A β

Our findings thus far suggest that oxidation results in the formation of DiY cross-links and results in stabilized A β assemblies that are prevented from further elongation into amyloid fibrils. To investigate this




Figure 4. UV-Induced DiY Cross-Linking of Early A β Assemblies Correlates with Formation of Stabilized Assemblies

(A) Th-T fluorescence spectrum shows the expected increase in fluorescence for assembling A β , but A β +UV Th-T fluorescence was significantly reduced. vA β incubated in the absence or presence UV showed no Th-T fluorescence. (B) Dot blotting using NU-1 antibody shows binding suggesting fewer oligomers in the oxidized A β +UV than in the unoxidized A β sample. No binding of NU-1 was observed for A β +UV 5 days post-UV exposure, but a small signal was detected in the A β -UV sample. (C) CD at 5 days showed a high β -sheet content in the A β sample, whereas the oxidized A β +UV showed a loss of signal but indicated some random coil. Oxidized and unoxidized vA β samples showed random coil conformation. (D) TEM after 2 h and 5 days showed that the unoxidized A β at 2 h formed oligomers, which transformed into a network of fibers at 5 days. The oxidized A β +UV samples formed small assemblies at 2 h, some of which developed into amorphous-like assemblies at 5 days. (E) vA β does not assemble into amyloid fibrils, but vA β +UV forms some amorphous aggregates after 5 days. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings. Scale bars, 500 nm.

further, DiY cross-linking was induced in A β using 2 h UV exposure, and the sample was then incubated with an equal concentration of freshly prepared A β (20 μ M:20 μ M) and compared with a 40 μ M A β sample not exposed to UV (see [Transparent Methods](#) for further detail). Fluorescence spectroscopy confirms the presence of DiY in the 40 μ M A β sample exposed to UV, and the signal reduced by half when the oxidized 40 μ M A β was diluted to 20 μ M (Figure 5A); 20 μ M oxidized A β added to 20 μ M freshly prepared unoxidized A β (A β +UV:A β) revealed the presence of DiY signal (Figure 5A). Interestingly, the levels of DiY in the A β +UV:A β mixture was higher than that of 20 μ M oxidized A β , suggesting that the presence of the UV-incubated A β in the environment results in the cross-linking of the freshly prepared A β following the co-incubation. Th-T fluorescence showed that the 40 μ M A β sample showed a shorter lag phase and higher Th-T signal compared with the 20 μ M A β sample confirming the expected concentration-dependent effect. The 20 μ M and 40 μ M oxidized A β samples showed no Th-T fluorescence intensity, similar to previous observations. When the A β mixtures were incubated together (A β +UV:A β 1:1), the mixture showed a longer lag phase (+20 h) but reached a similar Th-T intensity signal to the 20 μ M unoxidized sample at 50 h.




Figure 5. Co-incubation with Photo-oxidized, DiY-Cross-Linked A β Assemblies Slows the Aggregation of Freshly Prepared A β

(A) DiY fluorescence was measured for freshly prepared A β (40 and 20 μ M) and A β +UV:A β (20 μ M: 20 μ M) mixture immediately after co-incubation of freshly prepared A β with DiY-cross-linked A β . DiY signal was not detected in unoxidized freshly prepared 40 and 20 μ M A β but was induced in 2 h UV-oxidized (A β +UV) samples and in the mixture. (B) Th-T fluorescence showed that the unoxidized 40 and 20 μ M A β assemble at different rates, which was significantly delayed for A β +UV:A β -UV mixture and completely absent in the oxidized 40 and 20 μ M A β samples. (C) TEM imaging at 4 days revealed the presence of fibrils in both unoxidized 40 and 20 μ M A β , which was significantly reduced in the A β +UV:A β -UV mixture and absent in the oxidized 40 and 20 μ M A β samples. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings. Scale bars, 500 nm.

However, this was very significantly lower than the 40 μ M unoxidized A β sample (Figure 5B). This suggests inhibition of self-assembly in the A β +UV:A β -UV mixture. At 4 days, TEM imaging revealed the presence of A β fibrils in both 20 and 40 μ M A β samples, which were not detected in the photo-oxidized samples (Figure 5C). Only scant fibrils and smaller A β assemblies could be detected in the A β +UV:A β mixture, further confirming a reduced assembly in this mixture (Figure 5C). Taken together, this suggests that the incubation of freshly prepared A β with photo-oxidized A β results in an increase in DiY cross-linking and leads to the stabilization of some A β assemblies, resulting in the slower aggregation kinetics observed and prevention of elongation. DiY-containing, oxidized A β species may bind to A β and inhibit further elongation.

Photo-oxidation of Pre-formed A β Assemblies Slows Further Assembly of A β Oligomers/ Protofibrils and Prolongs Their Half-Life

Our findings thus far strongly suggest that oxidation stabilizes and strongly slows the aggregation and further elongation of A β assemblies. However, it is not clear whether these could also alter the assembly of preformed A β assemblies. To investigate this, A β (50 μ M) was freshly prepared and allowed to assemble for 24 h (henceforth called aA β). aA β samples were exposed for 2 h to UV to induce DiY cross-linking (aA β +UV) (Figure 6) (see Transparent Methods). Unlike samples that were not exposed to UV (aA β), the aA β +UV samples showed an increasing intensity arising from DiY, which continued to increase even after incubation in the dark following UV exposure (Figure 6A). Th-T fluorescence intensity showed that aA β continues to assemble, reaching plateau after approximately 15 h (Figure 6B). However, DiY cross-linking induced by UV correlates with the inhibition of further assembly of the aA β +UV for ~40 h, suggesting that the photo-oxidation leads to the stabilization or trapping of aA β +UV (Figure 6B). A gradual increase in fluorescence is observed beyond 40 h, which may indicate delayed assembly. Dot blotting using NU-1 revealed a similar level of oligomers in the oxidized and unoxidized aA β samples at the starting point (Figure 6C). Interestingly, the oligomers in the oxidized aA β +UV sample persisted beyond 4 days, unlike the unoxidized aA β sample, which showed a very low level of oligomers at the later time point (Figure 6C). TEM imaging showed the presence of oligomeric assemblies and scant fibers in the oxidized aA β +UV sample, compared




Figure 6. Photo-oxidation of Pre-formed A β (aA β) (24 h) Assemblies Slows Further Assembly of A β Oligomers/ Protofibrils and Prolongs the Half-Life of the Oligomers

(A) DiY signal was detected in the oxidized aA β +UV sample, which continued over time. Unoxidized aA β samples showed no DiY signal.

(B) Th-T fluorescence showed that the unoxidized aA β continues to assemble, whereas the oxidized aA β +UV became significantly inhibited from further elongation up to 40 h.

(C) Dot blotting using NU-1 antibody reveals the presence of oligomers in the unoxidized and oxidized aA β at the starting time point. However, the oligomers are still strongly detected in the oxidized aA β +UV at 4 days, unlike in the unoxidized samples.

(D) TEM imaging at 4 days revealed the presence of fibrils in the unoxidized aA β sample, whereas the oxidized aA β +UV showed oligomers and a reduced number of fibrils. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings. Scale bars, 500 nm

with the unoxidized aA β , which showed extensive fibril network. Together, these data provide further evidence indicating that photo-oxidation of pre-formed A β fibrils leads to the formation of DiY cross-links and results in stabilization of aA β assemblies, which prevent or delay further elongation. This is similar to the stabilization of A β oligomers observed following the slower/milder DiY cross-linking in the A β /CuCl₂ preparation, which occurred after the formation of A β oligomers (Figures 2B and 2C).

Self-Assembly Is Important for A β Toxicity

Multiple pieces of evidence have shown the detrimental role of A β on neuronal function that eventually leads to neuronal death (Marshall et al., 2016; Lambert et al., 1998; Lacor et al., 2007; Zhang et al., 2014; Selkoe and Hardy, 2016). Our Th-T, dot blot, and TEM data showed that A β 42 monomers self-assemble to form significant level of oligomers at 2 h, and eventually protofibrils, fibrils, and a network of fibrils at 4 days (Figures 2 and 6). Hence, the rate of self-assembly is high at early time points and plateaus at later time points when A β forms fibril network and plaques (Marshall et al., 2016). To investigate the role of self-assembly in toxicity, ReadyProbes live/dead assay was conducted on terminally differentiated SH-SY5Y neuroblastoma cells incubated for 3 days with wild-type A β 42 prepared in the following manner; A β 42 was allowed to assemble for 2, 24, 48, and 96 h to form oligomers, protofibril mixture, and fibrils and then treated for 2 h with UV (A β +UV) to induce DiY, or untreated (A β) (see Transparent Methods). vA β was used as an assembly-incompetent control (Marshall et al., 2016). Our results reveal a significantly higher level of cell death in wells incubated with A β after 2 h of preparation, compared with cells treated with A β after 24, 48, and 96 h incubation consistent with previous observations (Figure 7) (Marshall et al., 2016). However, the none of the oxidized A β samples containing DiY showed any toxic effect on the cells, similar




Figure 7. Oxidized, DiY-Containing A β Assemblies Are Not Toxic to Cells

Differentiated SH-SY5Y neuroblastoma cells were incubated with UV oxidized or unoxidized vA β or A β for 3 days, following which the percentage of dead cells was quantified using ReadyProbes Live/Dead Assay. The A β and vA β samples were freshly prepared and UV oxidized for 2 h (A β +UV/vA β +UV), or UV oxidized for 2 h followed by a further incubation on bench and in the dark for 24, 48, or 96 h, before being administered to cells. vA β and A β samples not exposed to UV were used as reference. Only the unoxidized/assembling wild-type A β induced significant cell death. Experiments were repeated five times. *** $p < 0.001$. $p \leq 0.05$ (*), < 0.01 (**), < 0.0001 (****) and > 0.05 was not significant. Error bars are expressed as \pm SEM.

to the assembly-incompetent oxidized and unoxidized vA β (Figure 7). This suggests a key role for self-assembly in the toxicity of A β .

DISCUSSION

In AD, A β self-assembles to dimers, oligomers, fibrils, and eventually amyloid plaques, one of the key hallmarks of the disease. A β is generally accepted to play a key role in AD, but the mechanism behind its toxicity is still not completely understood. Numerous studies have searched for the elusive “toxic” species and attempted to characterize its structure, and this has identified supposedly toxic assemblies such as dimers, 12mers, *56 KDa and hexamers (Lesne et al., 2006; Reed et al., 2011; Benilova et al., 2012). Here, we show that oxidation results in the formation of DiY cross-links which is one of several possible oxidative modifications and this significantly slows, or halts the self-assembly of A β 42, trapping it in a specific state. We compared the self-assembly and toxicity of assembly-incompetent vA β with wild-type A β and oxidized, trapped A β .

MCO and photo-induced oxidation of vA β revealed that DiY forms very rapidly as early as 5 min post-oxidation. However, oxidation did not lead to vA β assembly even after 5 days in the oxidative environment or post-oxidation. This demonstrates that the oxidation and the formation of DiY does not induce aggregation of the vA β , which is known not to self-assemble (Marshall et al., 2016). DiY was also rapidly induced in wild-type A β using both MCO and photo-induced oxidation; however, further assembly is inhibited or significantly slowed. Co-incubation of DiY-containing A β with freshly prepared uncross-linked A β demonstrated significantly reduced assembly. This suggests that oxidation and DiY cross-linking does not induce or facilitate the aggregation of the wild-type A β . Instead, A β assemblies are trapped and further elongation is delayed. This is supported by previous reports that showed that DiY-cross-linked A β are slow to fibrilize and form long-lived soluble oligomeric aggregates (Kok et al., 2013; O’malley et al., 2014, 2016; Sitkiewicz et al., 2014). Mass spectrometry studies have revealed that DiY cross-linking leads to the stabilization of A β 40 in compact oligomeric species (Sitkiewicz et al., 2014), which is in strong support of our findings.

Previous studies have suggested that DiY cross-linking can facilitate A β assembly (Atwood et al., 2004; Yoburn et al., 2003; Barnham et al., 2003; Zhang et al., 2017) or inhibit or slow A β self-assembly (Smith et al., 2007; Gu et al., 2018) or stabilize assemblies (Vazquez et al., 2019). Importantly, it has been shown that copper influences self-assembly in different ways depending on the concentration and ratio (Matheou

et al., 2015; Sarell et al., 2010), which may go some way to explaining these inconsistencies. These discrepancies could also arise from the different methods used to induce DiY in A β . Previous studies have used varying concentration of horseradish peroxidase and H₂O₂ (Galeazzi et al., 1999; Yoburn et al., 2003; Ali et al., 2006; Sitkiewicz et al., 2014; O'malley et al., 2014), varying concentrations of copper and H₂O₂ (Yoburn et al., 2003; Atwood et al., 2004; Barnham et al., 2004; Smith et al., 2007; Davis et al., 2011; Al-Hilaly et al., 2013; Williams et al., 2016), copper in combination with ascorbate (Gu et al., 2018), or photo-oxidation (Yoburn et al., 2003; Williams et al., 2016) to induce DiY cross-links in A β . In addition, even small differences in incubation conditions (e.g., trace metals, temperature, etc.) as well as atmospheric ozone levels (Vazquez et al., 2019) would result in different levels of DiY cross-links formed, which may result in varying impact on A β assembly. Moreover, the nature of the DiY cross-links may differ from one protocol to another. For example, other amino acids, such as phenylalanine, could modulate formation of cross-links (Zhang et al., 2019). Methodologies are likely to induce different oxidative effects on other amino acid side chains such as histidine, lysine, and met35 of A β (Kowalik-Jankowska et al., 2004; Ali et al., 2005; Palmblad et al., 2002; Friedemann et al., 2015). For example, met35 oxidation has been shown to attenuate A β oligomer formation and to enhance oxidation of A β Y10 (Palmblad et al., 2002). Mapping the specific modifications induced by the oxidation methods in a time-dependent manner using mass spectrometry would help to provide clarity on the relative role for DiY on A β assembly and whether this occurs in combination with other modifications to A β . This is a part of further studies.

Furthermore, the method of A β preparation used before DiY cross-linking may be critical in the outcome of the DiY cross-links on A β . Different sources and methods are used to prepare A β , resulting in a diverse pool of A β aggregates (Benilova et al., 2012). A β exists in a pool of monomers, soluble oligomers, and insoluble fibrils. Multiple studies have reported that the soluble A β oligomers in AD are composed of dimers, trimers, tetramers, pentamers, and decamers; A β -derived diffusible ligands (ADDLs); dodecamers; and A β *56 (Benilova et al., 2012). A question therefore arises regarding how DiY cross-linking impacts these pools of assemblies: presumably by creating new cross-linked dimeric species. Previous molecular dynamics studies have revealed that induction of DiY in a pool of monomeric A β 42 results in conformationally altered dimers that expose hydrophobic residues that may be limited to forming trimers via hydrophobic rather than polar interactions (Zhang et al., 2017). Our previous data showed that within 2 h of preparation, A β exists mostly as oligomers with a random coil conformation with a small β -sheet contribution (Marshall et al., 2016). Here, our results showed that the very slow oxidation induced by CuCl₂ alone first facilitates the formation of A β oligomers followed by DiY cross-linking of the oligomers resulting in a stabilized oligomeric population. The more rapid MCO reaction results in DiY formation as early as 5 min post-oxidation whereby the A β becomes trapped in a pre-oligomeric conformation (as assessed using the antibody NU-1). Th-T fluorescence, CD, and TEM showed that photo-oxidation of early A β species traps A β in a random coil conformation and prevents or significantly delays further assembly into amyloid fibrils. UV oxidation and DiY cross-linking in preformed oligomer/protofibril assemblies similarly results in the stabilization of this state and significantly delays further elongation to fibrils. Taken together, these results suggest that the timing of oxidation of A β critically influences its assembly, leading to the stabilization or significantly reduced assembly of the A β assemblies, which correlates with the time of cross-linking.

A β self-assembly is believed to be important for its toxicity (Marshall et al., 2016), and many studies have implicated the role of oligomeric species in cytotoxic effects (Glabe and Kaye, 2006; Saura et al., 2012; Walsh et al., 1999). Here, we compared toxicity of the unoxidized A β with A β that had been photo-oxidized and DiY cross-linked *in vitro*, whereby specific species in the assembly pathway have been stabilized. A β 42 was oxidized at different time points to stabilize a series of pre-oligomer, oligomeric, protofibrillar, and fibrillar forms. We show that none of these species is able to induce cell death following 3 days of incubation with differentiated neuroblastoma cells, whereas unoxidized, oligomeric A β remained potently toxic. This finding is in conflict with previous studies that showed that DiY A β 42 assemblies are toxic to cells (Barnham et al., 2004); DiY-cross-linked A β 40 dimers induce cell viability loss (Kok et al., 2013) and that Th-T positive, DiY-cross-linked A β 40 fibrils were able to inhibit long-term potentiation (LTP) (O'malley et al., 2014). We also observed that oxidized vA β is not toxic to cells, suggesting that the presence of DiY alone is not sufficient to induce toxicity. The oxidized and DiY-cross-linked A β 42 assemblies produced here are different from DiY A β 40 reported by others (O'malley et al., 2014; Kok et al., 2013), as the DiY A β 42 produced shows little Th-T fluorescence intensity and does not proceed to form fibrils. However, we do not rule out the possibility that DiY A β preparation in our study and others also results in other oxidative modifications, which may explain the discrepancies between these studies.

It is important to note that method of peptide preparation, peptide type, peptide concentration and aggregation state, and model system used may play a huge role in determining cell toxicity (Jana et al., 2016; Kaniyappan et al., 2017; Cecchi et al., 2008; Krishtal et al., 2015). Previous studies on DiY A β toxicity have used LTP (O'malley et al., 2014), MTS proliferation assay (Barnham et al., 2004), MTT. and lactate dehydrogenase assay (Ono et al., 2009) with varying A β concentrations to determine DiY A β toxicity. These assays, although showing the presence of cell injury, do not quantify absolute cell death. Impaired spine morphology and density, accompanied by increased reactive oxygen species and intracellular calcium, without apparent cell death have been reported as a result of tau toxicity (Kaniyappan et al., 2017). We have shown that a very short, 2 h exposure to A β oligomers of differentiated SHSY5Y cells result in oxidative and nucleolar stress without DNA damage or neuronal death (Maina et al., 2018). Thus, some discrepancies may arise from the assays and the cell model used. Moreover, if DiY A β is toxic, then it may depend on the level of DiY. None of the previous studies that studied DiY A β toxicity quantified the level of DiY formed on A β . As a result, discrepancies may also arise from the differences in the quantity of DiY cross-links in the A β used in toxicity assays. Indeed, we observed different levels DiY intensity between MCO and UV treatments.

Nonetheless, here we show that oxidation of A β *in vitro* leads to formation of DiY, halts A β self-assembly, and prevents cytotoxicity in a live-dead assay. We have previously demonstrated that assembly-incompetent vA β is not toxic to cells (Marshall et al., 2016). We therefore conclude that continued self-assembly is important for A β toxicity. We believe that the timing of the oxidation may be critical. For example, formation of DiY in A β fibrils would promote its stability and formation of amyloid plaques. Indeed, we have previously shown the presence of DiY on A β plaques in AD brain tissue and demonstrated that DiY A β fibrils become highly insoluble and resistant to formic acid denaturation (Al-Hilaly et al., 2013).

In conclusion, oxidation, which results in DiY cross-linking, promotes A β stabilization and does not induce or facilitate A β assembly. Our findings strongly suggest a role for self-assembly for A β toxicity. We show that A β exerts a high level of toxicity at a stage when self-assembly potential is high, compared with when the self-assembly rate is significantly diminished or abolished, as is the case for oxidized and vA β . This is observed even for those preparations wherein oligomeric A β has been stabilized. Our work implies that the timing of DiY formation plays a key role in further assembly and stability of A β .

Limitations of the Study

Here, we have provided evidence to show that oxidative conditions can induce the formation of DiY cross-links in A β 42 using MCO and UV photo-oxidation *in vitro*. We show that oxidation under the conditions used here halts further assembly. Stabilized A β 42 following oxidation is non-toxic to differentiated neuroblastoma cells. However, our study was not able to fully characterize whether other amino acids were affected by oxidation and what impact this might have on the prevention of assembly. We confirm that DiY is a major outcome of oxidation. Our work shows that oxidation of A β *in vitro* results in formation of non-toxic A β species. However, oxidative stress is known to be an important trigger for neurodegenerative diseases and our results do not imply a protective effect of oxidative stress. Oxidation has been performed under controlled environment *in vitro* affecting only A β self-assembly. Oxidation of A β *in vivo* is likely to have made diverse effects that have not been addressed in this study.

Resource Availability

Lead Contact

Further information and requests for resources of reagents should be directed to and will be fulfilled by the Lead Contact, Louise Serpell l.c.serpell@sussex.ac.uk.

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

This study did not generate code. The published article contains all datasets generated or analyzed during this study.

METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2020.101537>.

ACKNOWLEDGMENTS

This work was supported by funding from Alzheimer's Society (345 (AS-PG-16b-010)) awarded to L.C.S. and funding M.B.M.. K.F. is supported by Biotechnology and Biological Sciences Research Council, UK funding (BB/S003657/1), and the work was supported by Medical Research Council, UK (MR/K022105/1). Y.A. is supported by WisTa Laboratories Ltd (PAR1596). We gratefully acknowledge the valuable support of Dr P Schellenberger as electron microscope facility manager.

AUTHOR CONTRIBUTIONS

M.B.M. planned and carried out the work. G.B., Y.K.A.-H., K.F., and K.M. contributed experimental work. M.B.M. and L.C.S. wrote the paper. Y.K.A.-H. edited the paper. L.C.S. managed the project.

DECLARATION OF INTERESTS

The authors declare there are no competing interests.

Received: April 15, 2020

Revised: June 5, 2020

Accepted: September 3, 2020

Published: October 23, 2020

REFERENCES

- Al-Hilaly, Y.K., Biasetti, L., Blakeman, B.J.F., Pollack, S.J., Zibae, S., Abdul-Sada, A., Thorpe, J.R., Xue, W.-F., and Serpell, L.C. (2016). The involvement of dityrosine crosslinking in α -synuclein assembly and deposition in Lewy Bodies in Parkinson's disease. *Sci. Rep.* **6**, 39171.
- Al-Hilaly, Y.K., Williams, T.L., Stewart-Parker, M., Ford, L., Skaria, E., Cole, M., Bucher, W.G., Morris, K.L., Sada, A.A., Thorpe, J.R., and Serpell, L.C. (2013). A central role for dityrosine crosslinking of Amyloid- β in Alzheimer's disease. *Acta Neuropathol. Commun.* **1**, 83.
- Ali, F.E., Leung, A., Cherny, R.A., Mavros, C., Barnham, K.J., Separovic, F., and Barrow, C.J. (2006). Dimerisation of N-acetyl-L-tyrosine ethyl ester and Abeta peptides via formation of dityrosine. *Free Radic. Res.* **40**, 1–9.
- Ali, F.E., Separovic, F., Barrow, C.J., Cherny, R.A., Fraser, F., Bush, A.I., Masters, C.L., and Barnham, K.J. (2005). Methionine regulates copper/hydrogen peroxide oxidation products of A β . *J. Pept. Sci.* **11**, 353–360.
- Atwood, C.S., Perry, G., Zeng, H., Kato, Y., Jones, W.D., Ling, K.Q., Huang, X., Moir, R.D., Wang, D., Sayre, L.M., et al. (2004). Copper mediates dityrosine cross-linking of Alzheimer's amyloid-beta. *Biochemistry* **43**, 560–568.
- Barnham, K.J., Ciccotosto, G.D., Tickler, A.K., Ali, F.E., Smith, D.G., Williamson, N.A., Lam, Y.H., Carrington, D., Tew, D., Kocak, G., et al. (2003). Neurotoxic, redox-competent Alzheimer's beta-amyloid is released from lipid membrane by methionine oxidation. *J. Biol. Chem.* **278**, 42959–42965.
- Barnham, K.J., Haeflner, F., Ciccotosto, G.D., Curtain, C.C., Tew, D., Mavros, C., Beyreuther, K., Carrington, D., Masters, C.L., Cherny, R.A., et al. (2004). Tyrosine gated electron transfer is key to the toxic mechanism of Alzheimer's disease beta-amyloid. *FASEB J.* **18**, 1427–1429.
- Barritt, J.D., and Viles, J.H. (2015). Truncated amyloid-beta(11–40/42) from alzheimer disease binds Cu²⁺ with a femtomolar affinity and influences fiber assembly. *J. Biol. Chem.* **290**, 27791–27802.
- Benilova, I., Karran, E., and De Strooper, B. (2012). The toxic Abeta oligomer and Alzheimer's disease: an emperor in need of clothes. *Nat. Neurosci.* **15**, 349–357.
- Butterfield, D.A., and Halliwell, B. (2019). Oxidative stress, dysfunctional glucose metabolism and Alzheimer disease. *Nat. Rev. Neurosci.* **20**, 148–160.
- Butterfield, D.A., Swomley, A.M., and Sultana, R. (2013). Amyloid beta-peptide (1–42)-induced oxidative stress in Alzheimer disease: importance in disease pathogenesis and progression. *Antioxid. Redox Signal.* **19**, 823–835.
- Cecchi, C., Pensalfini, A., Liguri, G., Baglioni, S., Fiorillo, C., Guadagna, S., Zampagni, M., Formigli, L., Nosi, D., and Stefani, M. (2008). Differentiation increases the resistance of neuronal cells to amyloid toxicity. *Neurochem. Res.* **33**, 2516–2531.
- Davis, R.C., Marsden, I.T., Maloney, M.T., Minamide, L.S., Podlisny, M., Selkoe, D.J., and Bamberg, J.R. (2011). Amyloid beta dimers/trimers potentially induce cofilin-actin rods that are inhibited by maintaining cofilin-phosphorylation. *Mol. neurodegener.* **6**, 10.
- Friedemann, M., Helk, E., Tiiman, A., Zovo, K., Palumaa, P., and Tõugu, V. (2015). Effect of methionine-35 oxidation on the aggregation of amyloid- β peptide. *Biochem. Biophys. Rep.* **3**, 94–99.
- Galeazzi, L., Ronchi, P., Franceschi, C., and Giunta, S. (1999). In vitro peroxidase oxidation induces stable dimers of β -amyloid (1–42) through dityrosine bridge formation. *Amyloid* **6**, 7–13.
- Glabe, C.G., and Kaye, R. (2006). Common structure and toxic function of amyloid oligomers implies a common mechanism of pathogenesis. *Neurology* **66**, S74–S78.
- Gross, A.J., and Sizer, I.W. (1959). The oxidation of tyramine, tyrosine, and related compounds by peroxidase. *J. Biol. Chem.* **234**, 1611–1614.
- Gu, M., Bode, D.C., and Viles, J.H. (2018). Copper redox cycling inhibits A β fibre formation and promotes fibre fragmentation, while generating a dityrosine A β dimer. *Sci. Rep.* **8**, 16190.
- Hardy, J., and Higgins, G. (1992). Alzheimer's disease: the amyloid cascade hypothesis. *Science* **256**, 184–185.
- Jana, M.K., Cappai, R., Pham, C.L., and Ciccotosto, G.D. (2016). Membrane-bound tetramer and trimer A β oligomeric species correlate with toxicity towards cultured neurons. *J. Neurochem.* **136**, 594–608.
- Kaniyappan, S., Chandupatla, R.R., Mandelkow, E.M., and Mandelkow, E. (2017). Extracellular low-n oligomers of tau cause selective synaptotoxicity without affecting cell viability. *Alzheimers Dement.* **13**, 1270–1291.
- Kok, W.M., Cottam, J.M., Ciccotosto, G.D., Miles, L.A., Karas, J.A., Scanlon, D.B., Roberts, B.R., Parker, M.W., Cappai, R., and Barnham, K.J. (2013). Synthetic dityrosine-linked β -amyloid dimers form stable, soluble, neurotoxic oligomers. *Chem. Sci.* **4**, 4449–4454.
- Kowalik-Jankowska, T., Ruta, M., Wiśniewska, K., Łankiewicz, L., and Dyba, M. (2004). Products of

- Cu(II)-catalyzed oxidation in the presence of hydrogen peroxide of the 1-10, 1-16 fragments of human and mouse beta-amyloid peptide. *J. Inorg. Biochem.* 98, 940–950.
- Krishtal, J., Bragina, O., Metsla, K., Palumaa, P., and Tõugu, V. (2015). Toxicity of amyloid beta 1-40 and 1-42 on SH-SY5Y cell line. *SpringerPlus* 4, P19.
- Lacor, P.N., Buniel, M.C., Furlow, P.W., Clemente, A.S., Velasco, P.T., Wood, M., Viola, K.L., and Klein, W.L. (2007). Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J. Neurosci.* 27, 796–807.
- Lambert, M.P., Barlow, A.K., Chromy, B.A., Edwards, C., Freed, R., Liosatos, M., Morgan, T.E., Rozovsky, I., Trommer, B., Viola, K.L., et al. (1998). Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc. Natl. Acad. Sci. U S A* 95, 6448–6453.
- Lambert, M.P., Velasco, P.T., Chang, L., Viola, K.L., Fernandez, S., Lacor, P.N., Khuon, D., Gong, Y., Bigio, E.H., Shaw, P., et al. (2007). Monoclonal antibodies that target pathological assemblies of Abeta. *J. Neurochem.* 100, 23–35.
- Lesne, S., Koh, M.T., Kotilinek, L., Kaye, R., Glabe, C.G., Yang, A., Gallagher, M., and Ashe, K.H. (2006). A specific amyloid- β protein assembly in the brain impairs memory. *Nature* 440, 352–357.
- Lovell, M.A., Robertson, J.D., Teesdale, W.J., Campbell, J.L., and Markesbery, W.R. (1998). Copper, iron and zinc in Alzheimer's disease senile plaques. *J. Neurol. Sci.* 158, 47–52.
- Lund, M.N., Heinonen, M., Baron, C.P., and Estevez, M. (2011). Protein oxidation in muscle foods: a review. *Mol. Nutr. Food Res.* 55, 83–95.
- Maina, M.B., Bailey, L.J., Doherty, A.J., and Serpell, L.C. (2018). The involvement of A β 42 and tau in nucleolar and protein synthesis machinery dysfunction. *Front. Cell. Neurosci.* 12, 220.
- Marshall, K.E., Vadukul, D.M., Dahal, L., Theisen, A., Fowler, M.W., Al-Hilaly, Y., Ford, L., Kemenes, G., Day, I.J., and Staras, K. (2016). A critical role for the self-assembly of Amyloid- β 1-42 in neurodegeneration. *Scientific Rep.* 6, 30182.
- Matheou, C.J., Younan, N.D., and Viles, J.H. (2015). Cu(2)(+) accentuates distinct misfolding of Abeta(1)-(4)(0) and Abeta(1)-(4)(2) peptides, and potentiates membrane disruption. *Biochem. J.* 466, 233–242.
- Nunomura, A., Perry, G., Aliev, G., Hirai, K., Takeda, A., Balraj, E.K., Jones, P.K., Ghanbari, H., Wataya, T., Shimohama, S., and Chiba, S. (2001). Oxidative damage is the earliest event in alzheimer disease. *J. Neuropathol. Exp. Neurol.* 60, 759–767.
- O'malley, T.T., Oktaviani, N.A., Zhang, D., Lomakin, A., O'nuallain, B., Linse, S., Benedek, G.B., Rowan, M.J., Mulder, F.A., and Walsh, D.M. (2014). Abeta dimers differ from monomers in structural propensity, aggregation paths and population of synaptotoxic assemblies. *Biochem. J.* 461, 413–426.
- O'malley, T.T., Witbold, W.M., 3rd, Linse, S., and Walsh, D.M. (2016). The aggregation paths and products of Abeta42 dimers are distinct from those of the Abeta42 monomer. *Biochemistry* 55, 6150–6161.
- Ono, K., Condron, M.M., and Teplow, D.B. (2009). Structure-neurotoxicity relationships of amyloid beta-protein oligomers. *Proc. Natl. Acad. Sci. U S A* 106, 14745–14750.
- Palmblad, M., Westlind-Danielsson, A., and Bergquist, J. (2002). Oxidation of methionine 35 attenuates formation of amyloid β -peptide 1–40 oligomers. *J. Biol. Chem.* 277, 19506–19510.
- Pattison, D.I., Rahmanto, A.S., and Davies, M.J. (2012). Photo-oxidation of proteins. *Photochem. Photobiol. Sci.* 11, 38–53.
- Reed, M.N., Hofmeister, J.J., Jungbauer, L., Welzel, A.T., Yu, C., Sherman, M.A., Lesné, S., Ladu, M.J., Walsh, D.M., Ashe, K.H., and Cleary, J.P. (2011). Cognitive effects of cell-derived and synthetically derived A β oligomers. *Neurobiol. Aging* 32, 1784–1794.
- Roberts, B.R., Ryan, T.M., Bush, A.I., Masters, C.L., and Duce, J.A. (2012). The role of metallobiochemistry and amyloid-beta peptides in Alzheimer's disease. *J. Neurochem.* 120, 149–166.
- Sarell, C.J., Wilkinson, S.R., and Viles, J.H. (2010). Substoichiometric levels of Cu²⁺ ions accelerate the kinetics of fiber formation and promote cell toxicity of amyloid- β from Alzheimer disease. *J. Biol. Chem.* 285, 41533–41540.
- Selkoe, D.J., and Hardy, J. (2016). The amyloid hypothesis of Alzheimer's disease at 25 years. *EMBO Mol. Med.* 8, 595–608.
- Sitkiewicz, E., Olędzki, J., Poznański, J., and Dadlez, M. (2014). Di-tyrosine cross-link decreases the collisional cross-section of A β peptide dimers and trimers in the gas phase: an ion mobility study. *PLoS One* 9, e100200.
- Smith, D.P., Ciccotosto, G.D., Tew, D.J., Fodero-Tavoletti, M.T., Johanssen, T., Masters, C.L., Barnham, K.J., and Cappai, R. (2007). Concentration dependent Cu²⁺ induced aggregation and dityrosine formation of the alzheimer's disease amyloid- β peptide. *Biochemistry* 46, 2881–2891.
- Smith, D.P., Smith, D.G., Curtain, C.C., Boas, J.F., Pilbrow, J.R., Ciccotosto, G.D., Lau, T.L., Tew, D.J., Perez, K., Wade, J.D., et al. (2006). Copper-mediated amyloid-beta toxicity is associated with an intermolecular histidine bridge. *J. Biol. Chem.* 281, 15145–15154.
- Soura, V., Stewart-Parker, M., Williams, T.L., Ratnayaka, A., Atherton, J., Gorringer, K., Tuffin, J., Darwent, E., Rambaran, R., Klein, W., et al. (2012). Visualization of co-localization in Abeta42-administered neuroblastoma cells reveals lysosome damage and autophagosome accumulation related to cell death. *Biochem. J.* 441, 579–590.
- Souza, J.M., Giasson, B.I., Chen, Q., Lee, V.M., and Ischiropoulos, H. (2000). Dityrosine cross-linking promotes formation of stable alpha-synuclein polymers. Implication of nitrate and oxidative stress in the pathogenesis of neurodegenerative synucleinopathies. *J. Biol. Chem.* 275, 18344–18349.
- Vazquez, G., Caballero, A.B., Kokinda, J., Hijano, A., Sabaté, R., and Gamez, P. (2019). Copper, dityrosine cross-links and amyloid- β aggregation. *J. Biol. Inorg. Chem.* 24, 1217–1229.
- Walsh, D.M., Hartley, D.M., Kusumoto, Y., Fezoui, Y., Condron, M.M., Lomakin, A., Benedek, G.B., Selkoe, D.J., and Teplow, D.B. (1999). Amyloid beta-protein fibrillogenesis. Structure and biological activity of protofibrillar intermediates. *J. Biol. Chem.* 274, 25945–25952.
- Williams, T.L., Serpell, L.C., and Urbanc, B. (2016). Stabilization of native amyloid β -protein oligomers by copper and hydrogen peroxide induced cross-linking of unmodified proteins (CHICUP). *Biochim. Biophys. Acta* 1864, 249–259.
- Yoburn, J.C., Tian, W., Brower, J.O., Nowick, J.S., Glabe, C.G., and Van Vranken, D.L. (2003). Dityrosine cross-linked Abeta peptides: fibrillar beta-structure in Abeta(1-40) is conducive to formation of dityrosine cross-links but a dityrosine cross-link in Abeta(8-14) does not induce beta-structure. *Chem. Res. Toxicol.* 16, 531–535.
- Zhang, S., Fox, D.M., and Urbanc, B. (2017). Insights into formation and structure of abeta oligomers cross-linked via tyrosines. *J. Phys. Chem. B* 121, 5523–5535.
- Zhang, S., Fox, D.M., and Urbanc, B. (2019). Elucidating the role of hydroxylated phenylalanine in the formation and structure of cross-linked A β oligomers. *J. Phys. Chem. B* 123, 1068–1084.
- Zhang, Y., Lu, L., Jia, J., Jia, L., Geula, C., Pei, J., Xu, Z., Qin, W., Liu, R., Li, D., and Pan, N. (2014). A lifespan observation of a novel mouse model: in vivo evidence supports abeta oligomer hypothesis. *PLoS One* 9, e85885.

iScience, Volume 23

Supplemental Information

Metal- and UV- Catalyzed Oxidation Results in Trapped Amyloid- β Intermediates Revealing that Self-Assembly Is Required for A β -Induced Cytotoxicity

Mahmoud B. Maina, Gunasekhar Burra, Youssra K. Al-Hilaly, Kurtis Mengham, Kate Fennell, and Louise C. Serpell

Supplemental information

Transparent methods

Materials and Methods

Preparation of A β

Recombinant A β 42 (A β) and variant A β 42 (vA β) were purchased in Hexafluoroisopropanol (HFIP) films from rPeptide and JPT, respectively. vA β is a variant of A β 1-42 with F19S and G37D mutations which render the peptide assembly incompetent (see (Marshall et al. 2016) for detail). The peptides were prepared using an established protocol, and all procedures were done using protein LoBind Eppendorfs and tips. 0.2 mg/mL of the peptides were solubilized in 200 μ L HFIP (Sigma-Aldrich), vortexed for 1 min and sonicated in 50/60 Hz bath sonicator for 5 min. The HFIP was removed by air drying using a low stream of nitrogen gas. The dried peptide films were dissolved in 200 μ L Dimethyl sulfoxide (DMSO) >99.9% (Sigma-Aldrich), vortex for 1 min and sonicated for 1 min. The solution was passed through a 2 mL 7K MWCO Zeba buffer-exchange column (Thermo Scientific) and stacked with 40 μ L of 10 mM phosphate buffer (pH 7.4). The concentration of the peptides was determined using a NanoDrop spectrophotometer (Thermo Scientific) at a wavelength of 280 nm (extinction coefficient of 1490 M⁻¹cm⁻¹). The peptide solutions were immediately diluted to 50 μ M with the 10 mM phosphate buffer and used as indicated. For the experiments involving preformed A β assemblies, the 50 μ M A β was left to assemble for 24 without shaking before being subjected to UV exposure.

Metal-catalysed oxidation (MCO) of A β and vA β

Freshly prepared samples of A β 1-42 and vA β 1-42 peptides (50 μ M) in 10 mM phosphate buffer, pH7.4 were incubated i) without CuCl₂, ii) in the presence of 400 μ M CuCl₂ (peptide: CuCl₂ ratio 1:8) and iii) in the presence of 400 μ M CuCl₂ and 2.5 mM H₂O₂. At this concentration, we did not observe any precipitation of copper phosphate. An additional control was performed adding ethylenediaminetetraacetic acid (EDTA) (2 mM) to the assembly mixture of wild type A β . The peptides were incubated at 37°C without agitation and at each time point collected, the oxidation reaction was quenched using EDTA at a final concentration of 2 mM. A minimum of three independent experiments were conducted to ensure the reproducibility of the findings.

Standard curve of dityrosine

Dityrosine was synthesised as described in (Al-Hilaly *et al.*, 2013). To generate a dityrosine standard curve, a set of dityrosine standard concentrations (0.1, 0.5, 1, 2, 3, 5 and 10 μ M)

were prepared in Milli-Q water. Each concentration was prepared in triplicate and dityrosine fluorescence was recorded for each concentration using excitation wavelength 280 nm and emission wavelength 410 nm. The mean values of dityrosine fluorescence intensity for each concentration were plotted against dityrosine concentration and line plot was constructed by linear regression analysis using Microsoft Excel software. The equation of this line was used to quantify the dityrosine content of A β samples oxidized with MCO or CuCl₂ alone.

Photo-oxidation of A β and vA β

Freshly prepared samples of A β 1-42 and vA β 1-42 peptides (50 μ M) in 10 mM phosphate buffer, pH 7.4 were incubated i) without UV-C in the dark, and ii) under of UV-C for 5min or 2h using a G6T5 Germicidal 9' 6W T5 UVC lamp set to 8 J/m²/sec (General Lamps Ltd). A minimum of three independent experiments were conducted to ensure the reproducibility of the findings.

Fluorescence spectroscopy

The formation of dityrosine was monitored with a fluorescence spectrophotometer (Varian Ltd., Oxford, UK), using a 1 cm path length quartz cuvette (Starna, Essex, UK). The presence of dityrosine was detected using fluorescent excitation wavelength of 320 nm and emission collected between 340 – 600 nm, with dityrosine peak signal observed between 400-420 nm. Tyrosine fluorescence signal was monitored using an excitation wavelength of 280 nm and an emission wavelength of 305 nm, with the peak tyrosine emission observed at 305 nm. For experiments involving metal-catalysed oxidation, the reaction was quenched using EDTA to a final concentration of 2 mM. For all the measurements, the excitation and emission slits were both set to 10 nm, scan rate set to 300 nm/min with 2.5 nm data intervals and an averaging time of 0.5 s. The photomultiplier tube detector voltage was set at 500 V.

Thioflavin T fluorescence assay of A β self-assembly

Samples were incubated with 100 μ M Thioflavin T (Th-T), and the rate of Th-T binding was monitored over time at 37°C using SpectraMax i3 plate reader with samples incubated in CellCarrier-96 Ultra Microplates (PerkinElmer). The readings were collected in a black 96-well plate with a clear bottom (PerkinElmer), which was sealed with an optically clear polyolefin film to avoid evaporation (StarSeal Advanced Polyolefin Film, Starlab). The number of readings per well was set to 6, PMT voltage was set to high and blank spectra of the buffer were subtracted to protein fluorescence scans. The excitation wavelength was set at 440 nm, and emission at 483 nm and the signal collected every 30 min, with 5 sec low orbital shakes before readings. The fluorescence data were plotted against time. A minimum of three independent experiments was repeated to ensure the reproducibility of the findings.

Circular Dichroism (CD)

The secondary structure of A β and vA β peptides at 50 μ M concentration in 10 mM phosphate buffer (pH 7.4) incubated under different conditions was assessed using Jasco J715 CD spectrometer (Jasco, Goh-Umstadt, Germany). 140 μ L of each sample was placed into a 1 mm path length quartz cuvette (Hellma) and scanned between 190 nm and 260 nm. The CD spectra were collected in triplicate at a maintained temperature of 21 °C.

Negative-stain transmission electron microscopy (TEM)

The morphology of control and cross-linked A β and vA β peptides was assessed by negative stain TEM. Briefly, 4 μ L of each sample was dropped onto 400-mesh carbon-coated grids (Agar Scientific, Essex, UK). After 1 min incubation, the excess sample was blotted using filter paper, and the grid was washed with 4 μ L filtered Milli-Q water and blotted. The grid was then negatively stained for 40 sec using 4 μ L of filtered 2% (w/v) uranyl acetate. The excess stain was blotted with filter paper and grids left to air-dry before storage. The grids were examined on a Jeol Jem1400-plus transmission electron microscope (Jeol, USA), operated at 80 kV fitted with a Gatan Orius SC100 camera (UK).

Dot-blotting

A total of 4 μ L was spotted onto a 0.2 μ M pore nitrocellulose membrane and allowed to dry for 10 min. The membrane was boiled with PBS for 1 min twice and then blocked with blocking buffer (5% milk in 0.05% TBS-T) for 1 hour at room temperature on a rocker. The blocking buffer was next replaced with mouse NU-1 primary antibody (1/2000) and left to bind overnight at 4°C on a rocker. The membrane was washed 6 times for 5 min with washing buffer (0.05% TBS-T), then incubated with an HRP-conjugated goat anti-mouse secondary antibody for 1 hour. The membrane was washed six times for 5 min with washing buffer, then incubated with Clarity Western ECL Substrate (Bio-Rad) for 1 min before being developed in the darkroom. The NU-1 antibody was a gift from the William Klein lab (Lambert et al.). A minimum of three independent experiments were conducted to ensure the reproducibility of the findings.

Cell death assay

Differentiated SHSY5Y neuroblastoma cells were used for the toxicity experiments. Firstly, undifferentiated SHSY5Y neuroblastoma cells (ATCC CRL-2266™), were maintained in Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12 (DMEM/F-12) (Life Technologies, United Kingdom), supplemented with 1% (v/v) L-glutamate (L-Gln) (Invitrogen), 1% (v/v) penicillin/streptomycin (Pen/Strep) (Invitrogen) and 10% (v/v) Fetal Calf Serum at 37°C and 5% CO₂. The undifferentiated SHSY5Y cells were seeded to 60% confluency in a CellCarrier-

96 Ultra Microplates (PerkinElmer). The cells were differentiated in a medium containing 1% Fetal Calf Serum supplemented with 10 μ M trans-Retinoic acid (Abcam) for 5 days. Next, the medium was replaced with a serum-free media supplemented with 2 nM brain-derived neurotrophic factor (BDNF) (Merck Millipore). After 2 days in the BDNF-containing media, the media was replaced with serum-free media and the cells were treated with UV cross-linked or uncross-linked vA β or A β for 3 days. At the end of the incubation period, the cells were incubated with ReadyProbes reagent (Life Technologies) for 15 min. The ReadyProbes kit contains NucBlue Live reagent that stains the nuclei of all live cells and Propidium iodide that stains the nuclei of dead cells with compromised plasma membrane. The cells were imaged at 37°C and 5% CO₂ using Operetta CLS high-content analysis system (PerkinElmer) using DAPI and TRITC filters. At least 5000 dead and live cells were analysed using the Harmony software automated analysis algorithm within the Operetta CLS high-content analysis system. A minimum of three independent experiments were repeated to ensure the reproducibility of the findings.

Supplemental Results

Figure S1




Figure S1. Related to Figure 1. Standard curve showing the concentration of DiY standard against intensity at 410 nm. Data was recorded as described in Methods above.




Figure S2. Related to Figure 1. Thioflavine T fluorescence assay comparing A β 1-42 self-assembly under MCO conditions, copper alone and in buffer with EDTA to chelate trace metals. A β 1-42 self-assembly at a higher rate than A β 1-42 in buffer suggesting that the trace metals present in the water used to make the buffer can influence the assembly rate.

References

- AL-HILALY, Y. K., WILLIAMS, T. L., STEWART-PARKER, M., FORD, L., SKARIA, E., COLE, M., BUCHER, W. G., MORRIS, K. L., SADA, A. A., THORPE, J. R. & SERPELL, L. C. 2013. A central role for dityrosine crosslinking of Amyloid-beta in Alzheimer's disease. *Acta Neuropathol Commun*, 1, 83.
- MARSHALL, K. E., VADUKUL, D. M., DAHAL, L., THEISEN, A., FOWLER, M. W., AL-HILALY, Y., FORD, L., KEMENES, G., DAY, I. J., STARAS, K. & SERPELL, L. C. 2016. A critical role for the self-assembly of Amyloid-beta1-42 in neurodegeneration. *Sci Rep*, 6, 30182.
- LAMBERT, M. P., VELASCO, P. T., CHANG, L., VIOLA, K. L., FERNANDEZ, S., LACOR, P. N., KHUON, D., GONG, Y., BIGIO, E. H., SHAW, P., DE FELICE, F. G., KRAFFT, G. A. & KLEIN, W. L. 2007. Monoclonal antibodies that target pathological assemblies of Abeta. *J Neurochem*, 100, 23-35.