



Published in final edited form as:

Hum Pathol. 2020 September ; 103: 72–82. doi:10.1016/j.humpath.2020.07.010.

IMMUNOHISTOCHEMICAL DISTINCTION OF PARAGANGLIOMAS FROM EPITHELIAL NEUROENDOCRINE TUMORS – GANGLIOCYTIC DUODENAL AND CAUDA EQUINA PARAGANGLIOMAS ALIGN WITH EPITHELIAL NEUROENDOCRINE TUMORS

Divya Mamilla, MD¹, Irena Manukyan, MD², Patricia A. Fetsch, HT ASCP², Karel Pacak, MD¹, Markku Miettinen, MD²

¹Section on Medical Neuroendocrinology, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD, 20892, USA.

²National Cancer Institute, Laboratory of Pathology, Bethesda, MD, 20892, USA.

Abstract

Distinction of paraganglioma (PGL) from epithelial neuroendocrine tumors (NETs) can be difficult as they can mimic each other by nested architecture and expression of neuroendocrine markers. In this study we examined differential diagnostic markers in 262 PGLs (142 adrenal pheochromocytomas and 120 extra-adrenal PGLs), 9 duodenal gangliocytic PGLs and 3 cauda equina PGLs, and 286 NETs (81 GI, 78 pancreatic, 42 thoracic, 37 medullary thyroid carcinomas, and 48 high-grade NETs including 32 small cell carcinomas of lung). While keratin expression was nearly uniform in NETs with the exception of few tumors, extensive keratin expression was seen in only one PGL (<1%), and focal expression in 5% PGLs. GATA 3 was present in >90% of PGLs but only in 2% of NETs, usually focally. Tyrosine hydroxylase (TH) was expressed in >90% of adrenal, abdominal, and thoracic PGLs but only in 37% of head and neck PGLs reflecting their variable catecholamine synthesis. Focal or occasional extensive TH-expression was detected in 10% of NETs. CDX2 was a helpful discriminator seen in 28% of pancreatic and most GI NETs but in no PGLs. SOX10 detected sustentacular cells in 85% of PGLs and 7% of NETs, while GFAP detected sustentacular cells mainly in PGLs of neck and was absent in NETs. Duodenal gangliocytic PGLs (n = 9) and all cauda equina PGLs (n = 3) expressed keratins, lacked GATA3, showed no or minimal TH expression as some NETs, and contained SOX10 and S100 protein-positive spindle cells negative for GFAP. Ganglion-like epithelioid cells were keratin-positive and negative for TH and SOX10 differing from true ganglion cells. We conclude that duodenal gangliocytic and cauda equina PGLs have a NET-like immunoprofile and differ from ordinary

Address for correspondence: Markku Miettinen, M.D., Head of Surgical Pathology, Laboratory of Pathology, NCI/NIH, 9000 Rockville Pike, Bldg. 10, Rm. 2S235C, Bethesda, Maryland 20892 Phone: 301-480-8947, miettinenmm@mail.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure Statement: The authors have no conflicts of interest to disclose

PGLs. NETs can be distinguished from PGLs by their expression of keratins and general lack of GATA3, TH, and GFAP-positive sustentacular cells, and sometimes by expression of CDX2 or TTF1.

Keywords

Pheochromocytoma; paraganglioma; neuroendocrine tumor; GATA3; keratins; tyrosine hydroxylase; gangliocytic paraganglioma; cauda equina paraganglioma

INTRODUCTION

Adrenal pheochromocytoma, the most common paraganglioma, and other paragangliomas (PGLs) are non-epithelial neuroendocrine neoplasms derived from chromaffin cells of the adrenal medulla and paraganglia of the sympathetic and parasympathetic nervous systems. They can occur in locations where paraganglia normally reside and as metastases almost at any site.^{1,2} The histologic diagnosis of PGLs is straightforward when tumor involves a common site and has a typical histology with a nested architecture. However, epithelial neuroendocrine tumors (NETs): carcinoids, pancreatic NETs, thyroid medullary carcinoma, and high-grade neuroendocrine carcinomas, can also have nested patterns and mimic PGLs. Small specimens can be problematic as architectural landmarks can be limited. In these instances, immunohistochemistry, is necessary for this clinically important distinction.

Both PGLs and NETs typically express pan-neuroendocrine markers chromogranin A, synaptophysin, and insulinoma-associated protein 1 (INSM1).^{1,3,4} Therefore, other markers are necessary for immunohistochemical distinction of PGLs from NETs.

Keratin monoclonal antibody cocktail AE1/AE3 detects nearly all soft epithelial (non-hair) keratins, except keratins 10, 17, 18.⁵ It is therefore expected to be positive in all epithelial NETs. Keratin antibody CAM5.2 detects specifically keratin 8.⁶ Keratins have been occasionally reported in conventional paragangliomas (of adrenal, retroperitoneum, urinary bladder, carotid body, and thorax).^{7,8}

GATA3 (GATA-binding protein 3 and trans-acting T-cell-specific factor), is one of the six members of dual zinc-finger transcription factors that binds to the DNA sequence GATA. It is essential in neuronal embryogenesis regulating the expression of the noradrenergic factor tyrosine hydroxylase and differentiation and survival of neurons and chromaffin cells.⁹ GATA3 has been reported in PGLs and generally absent in NETs but has not been studied in gangliocytic and cauda equina PGLs.¹⁰⁻¹²

Tyrosine hydroxylase, the rate limiting enzyme involved in conversion of L-tyrosine to L-DOPA has been detected biochemically and immunohistochemically in PGLs and in sporadic NETs, but it has not been extensively studied in neuroendocrine tumors.¹³⁻¹⁵

Organ-specific transcription factors, such as TTF1 and CDX2, and hormonal markers, can be useful in characterization of NETs, but none of them is universal for all subgroups.^{16,17}

TTF1 has been studied in PGLs of thorax but not at other sites. CDX2 has been detected in some pancreatic NETs¹⁶ but has not been studied in PGLs.

Sustentacular cells, elongated, slender spindle cells variably outlining the cell nests of PGLs, have also been detected in some NETs. They can be highlighted by immunostains for S100 protein, GFAP, and SOX10.^{18–21} However, these have not been systematically studied in large sets of NETs and PGLs of various sites.

Gangliocytic paraganglioma is a rare neuroendocrine tumor containing ganglion-like cells and typically occurring in the 2nd part of duodenum. It differs from conventional paraganglioma by common keratin expression.^{22,23} Neither it nor cauda equina paraganglioma, another rare type of keratin-expressing paraganglioma occurring in the caudal neural tube²⁴, has been studied for GATA3 and tyrosine hydroxylase.

In this study, we examined the expression 5 groups of markers comparatively in >500 PGLs and NETs of various sites in special groups of rare PGLs: gangliocytic PGL of duodenum and cauda equina PGL.

MATERIALS AND METHODS

A large group (n = 560) of neuroendocrine tumors was examined histologically to confirm the diagnosis and arranged in manually prepared multitumor blocks as described before.²⁵ A small number cases were studied using whole sections. The cases originated from 1970–2019 and were de-identified having sparse annotations for tumor site, size, patient age, sex, and a possible germline tumor syndrome. These tumors included 262 paragangliomas (PGLs): 142 adrenal and 120 extra-adrenal examples including 54 cervical, 49 retroperitoneal, 9 thoracic, and 8 urinary bladder PGLs. Approximately 40% of PGLs were germline syndrome-associated tumors (RET, VHL, SDH subunits, and NF1). There were also 9 duodenal gangliocytic PGLs and 3 cauda equina PGLs. The 286 epithelial neuroendocrine tumors (NETs) included 81 grade 1–2 intestinal and 42 pulmonary NETs/carcinoids, 78 pancreatic NETs, and 37 medullary carcinomas of thyroid. High-grade (Grade 3) NETs included 32 small cell carcinomas of lung and 16 high-grade neuroendocrine carcinomas of various sites (4 pulmonary large cell, 4 gastrointestinal, 2 endometrial, 1 sinonasal, 1 urinary bladder, and 4 metastatic tumors of unknown origin).

Immunohistochemical studies were performed by automated systems (Ventana Benchmark Ultra or Leica Bond). Antibodies and protocols are listed in Table 1. All tumors were confirmed immunopositive for at least one, usually all 3 neuroendocrine markers, chromogranin A, synaptophysin and INSM1.^{1–4} There was a slightly lower positivity for INSM1 in all tumor categories.

Immunohistochemical studies for keratins (AE1/AE3 cocktail, keratin 8: CAM5.2), transcription factors (GATA3 CDX2, and TTF1), tyrosine hydroxylase were tabulated in detail for estimated percentage of positive tumor cells. Expression of markers for sustentacular cells (S100 protein, GFAP, and SOX10) were evaluated in a 0–3 scale for the area of 40x magnification richest on these cells as follows: 0 = no positive cells, 1 = 1–10 cells, 2 = 11–49, 3 = 50 cells. As the results from S100 protein expression in sustentacular

cells were generally similar to those of SOX10 and there was variable labeling of chief cells/neuroendo-crine cells for S100 protein, results of only SOX10 were tabulated. Expression of phenyl-ethanolamine N-methyltransferase (PNMT) was also studied in >500 PGLs and NETs. While the results were partly similar to those of TH, they were more difficult to interpret as many NETs showed diffuse weak expression and PGLs showed variable and unpredictable expression. Therefore, this marker was not examined further.

RESULTS

The immunohistochemical results have been summarized in Table 2.

Conventional paragangliomas

These tumors included pheochromocytomas, retroperitoneal (usually paraaortic or aortocaval), thoracic/aorticopulmonary, urinary bladder, and neck PGLs (most from the carotid body, one from larynx, and one from the higher neck).

Most PGLs were negative for keratins AE1/AE3 and keratin 8 (CAM5.2). Exceptional, extensive AE1/AE3 and keratin 8-positivity was seen in one pheochromocytoma in approximately 80% of tumor cells, while this tumor was positive for GATA3 and tyrosine hydroxylase typical of PGLs (Fig. 1). Focal keratin expression with both antibodies was detected in 5% of tumor cells in 12 conventional PGLs, 0–7% in various subgroups, and these cells had a thin, slender morphology, consistent with sustentacular cells (Table 2).

GATA3 nuclear expression was seen in most pheochromocytomas PGLs of abdomen, thorax, and head and neck, usually extensively. Some cases showed zonal expression patterns probably associated with suboptimal antigen preservation. The negative cases included a small subset PGLs of different sites (Table 2).

Tyrosine hydroxylase was nearly always extensively expressed in the cytoplasm of adrenal, retroperitoneal, thoracic, and urinary bladder PGLs (88–98%). However, PGLs of the neck were heterogeneous showing positivity in only 20/54 cases (37%). Nerves and ganglion cells were also positive. CDX2 was expressed in none of the conventional PGLs. TTF1 was focally expressed of 2 adrenal PGLs in <5% of tumor cells.

SOX10-positive sustentacular cells were variably present in PGLs. They were numerous in most PGLs of the neck and thorax, and scant or absent in adrenal, urinary bladder, and retroperitoneal PGLs (Table 2). GFAP-positive sustentacular cells were predominantly detected in PGLs of the head and neck, and thorax. In adrenal, urinary bladder, and retroperitoneal PGLs they were usually absent or present only in small numbers (Table 2). SOX10 or GFAP did not label chief cells in any PGLs or NETs.

While S100 protein and SOX10 showed identical distribution in sustentacular cells, S100 protein also labeled chief cells in 50–70% PGLs of different sites, 65% of pancreatic NETs, 10% of pulmonary NETs, but only 2% of GI NETs.

Gangliocytic and cauda equina paragangliomas

The 9 gangliocytic PGLs occurred in 4 females and 4 males of ages 42–74 years (median, 65 years). Demographic information was unavailable in 1 case. All tumors were duodenal submucosal polypoid masses measuring 1.5 – 5 cm.

These tumors contained neuroendocrine nests containing scattered ganglion-like cells. Mitotic activity was <2/10 HPFs in all cases, equivalent to NET grade 1. The nests were surrounded by a variably prominent spindle cell component. The nests of epithelioid neuroendocrine cells, including ganglion-like cells with abundant cytoplasm and angulated outlines, were also positive for keratins AE1/AE3 and more extensively positive for keratin 8 (CAM5.2), which also highlighted delicate cell processes of the epithelial cells (Fig. 2). All tumors were negative for GATA3, CDX2, and TTF1. Only minimal tyrosine hydroxylase-positivity (<2% of tumor cells) was detected in 2 cases, while the remaining cases were negative. The ganglion-like cells were negative for tyrosine hydroxylase, GATA3, and SOX10, thus differing from ordinary ganglion cells. The spindle cells between the neuroendocrine nests were positive for S100 protein and SOX10, consistent with their Schwannian nature (Fig. 2), and were negative for GFAP.

The 3 cauda equina PGLs, were diffusely positive for keratins AE1/AE3 and negative for GATA3. Two the cases studied for keratin 8 were also positive. Focal tyrosine hydroxylase-positivity was seen in 2 cases (Fig. 3). A small number of SOX10-positive cells were present with no GFAP-positive cells in the 2 cases tested that were also negative for CDX2 and TTF1.

Epithelial neuroendocrine tumors (NETs)

All gastrointestinal, pancreatic, and pulmonary/mediastinal NETs and medullary thyroid carcinomas were extensively positive for keratins AE1/AE3 and keratin 8 (CAM5.2). Only 3% of NETs showed significantly higher labeling for keratin 8 than the AE1/AE3 cocktail. While most high-grade neuroendocrine carcinomas showed uniform keratin-positivity, 2 small cell carcinomas of lung were negative for both keratin antibodies (Table 2).

All gastrointestinal NETs were negative for GATA3, but 5 NETs of pancreas and 1 pulmonary carcinoid focally expressed GATA3 (Tables 2 and 3). Focal tyrosine hydroxylase positivity was detected in gastrointestinal (6/81), pancreatic (7/77) and thoracic nets (11/42), usually in <5% of tumor cells (Fig 4). Occasional pancreatic NETs and pulmonary carcinoids showed more extensive tyrosine hydroxylase-positivity (Fig. 4, Table 3). In some cases, pancreatic NETs with aberrant immunophenotypes, such as lack of keratins and strong tyrosine hydroxylase expression, were CDX2-positive supporting the diagnosis of a pancreatic NET (Fig. 5).

Sustentacular cells with SOX10-positivity were rarely detected in NETs, essentially limited to thoracic and pancreatic NETs. GFAP-positive sustentacular cells were absent in all NETs (Table 2).

DISCUSSION

Paragangliomas (PGLs) and epithelial neuroendocrine tumors (NETs) from various organs can mimic each other due to their shared nested architecture and expression of pan-neuroendocrine markers chromogranin A, synaptophysin, and INSM1. Small specimens can be more challenging and require immunohistochemical support for diagnosis.

In this study we examined >500 PGLs and NETs with a panel of 8 markers to evaluate their power in the distinction between these groups of NETs: 1) Keratins AE1/AE3 and CAM5.2 (keratin 8), 2) transcription factors GATA3, CDX2, and TTF1, 3) tyrosine hydroxylase, and 4) sustentacular cell markers S100 protein, SOX10 and GFAP.

Keratin-negativity of conventional PGL (excluding gangliocytic and cauda equina paragangliomas) is a useful parameter separating it from most NETs. However, 7% of PGLs did express keratins, but most of this expression was focal limited to sustentacular cells, and only 1 pheochromocytoma was extensively positive. Previous studies have also detected keratins in a small percentage of conventional PGLs.^{7,8}

Tyrosine hydroxylase, an enzyme involved in the biosynthesis of catecholamines, is expressed in neural tissue including paraganglia and has been detected especially in catecholamine synthesizing PGLs and some carcinoids.^{14,15} We confirmed nearly consistent tyrosine hydroxylase expression in pheochromocytoma and abdominal and thoracic PGLs, whereas expression was far less common in neck/carotid body PGLs (32%); in a previous study 27%.¹⁴ We also found some tyrosine hydroxylase expression in NETs, but this was usually focal and limited to scattered tumor cells. However, exceptions were noted with some NETs, especially pancreatic ones, being extensively TH-positive. TH has some value in distinguishing PGLs from NETs, but lack of its expression in most head and neck PGLs and extensive expression in some NETs are limitations.

Most PGLs including nearly all adrenal, retroperitoneal, thoracic, and neck/carotid body PGLs expressed GATA3, while NETs were negative making GATA3 a valuable marker in discrimination of PGLs from NETs as previously suggested.¹⁰⁻¹² However, our study reveals exceptions such as extensive GATA3 expression in 2% pancreatic and thoracic NETs, so that additional markers, such as tyrosine hydroxylase and transcription factors TTF1 and CDX2 could be used as further discriminators between NETs and PGLs. It should be considered in the differential diagnosis of neuroendocrine tumors of head and neck that GATA3 is also expressed in parathyroid and pituitary adenomas.^{26,27} Despite recently expressed concern about weaker GATA3-labeling of PGLs with immunohistochemistry protocols without robust epitope retrieval²⁸, we found that our 64-minute epitope retrieval with Roche CC1 reagent and 60 minute primary antibody incubation, and Ultraview detection gave a sufficient detection of GATA3 in PGLs (positivity >90% in all main subgroups).

While S100 protein-positive sustentacular cells are known in paragangliomas^{1,20,21}, they can also occur in some NETs.^{18,21} We found that SOX10 is superior to S100 protein in enumerating sustentacular cells, because it does not label the chief cells, as often seen in pulmonary and pancreatic NETs and PGLs of different sites. The common S100 protein-

positivity in both PGLs and NETs should not lead to confusion with S100 positive neural or epithelial tumors.

The limitation in sustentacular cell evaluation is that they are abundant in only PGLs of head and neck and only variably present in adrenal and retroperitoneal PGLs. However, their presence in NETs is essentially limited to small subsets of pulmonary and pancreatic NETs. GFAP-positive sustentacular cells are restricted to PGLs of neck and thorax, which confirms a previous observation in a smaller number of PGLs.²⁰ The assessment of sustentacular cell populations has a limited value in the diagnosis of PGLs, although their absence has been suggested a possible but weak marker of malignancy (reviewed in reference¹).

Aberations in the main patterns of expression of keratins AE1/AE3, GATA3 and tyrosine hydroxylase occurred mainly in pancreatic and pulmonary NETs (Table 3). The marker profile of one pancreatic NET indicated PGL-like differentiation: keratin and CDX2-negative and positive for GATA3 and tyrosine hydroxylase, supporting the diagnosis of a (peri)pancreatic PGL described in isolated reports.^{29,30} However, there were also cases that in addition to keratin expression, had strong expression of either GATA3 or tyrosine hydroxylase raising the question whether there are pancreatic NETs with hybrid features of NET and PGL. Extensive genomic comparison could be useful to definitively determine the type of these tumors.

Our findings indicate that duodenal gangliocytic paraganglioma and cauda equina PGLs align better with NETs than ordinary PGLs by their immunophenotype and should be classified as separate groups, perhaps termed as gangliocytic NET and cauda equina NET. In this study, all gangliocytic PGLs and cauda equina PGLs were extensively keratin-positive, negative for GATA3, and negative or minimally positive for tyrosine hydroxylase, thus having an immunohistochemical profile similar to NETs, rather than conventional PGLs. A caveat is that there may be true spinal paragangliomas with classic paraganglioma features different from cauda equina paragangliomas.³¹

Ganglion-like cells in the epithelial islands of gangliocytic paraganglioma seem to be keratin-positive and SOX10-negative epithelial neuroendocrine cells and not true ganglion cells. However, occurrence of entrapped normal ganglion cells in these tumors is possible.

Additional studies, such as extensive genomic sequencing and assessment of genomic methylation profiles, as now available for the diagnosis of brain tumors and sarcomas,³² might prove useful in determining the taxonomic status of gangliocytic and cauda equina PGLs among the neuroendocrine tumors.

Funding:

Supported by NIH's intramural Research Program

REFERENCES

1. Lack EE, Tumors of the adrenal glands and extraadrenal paraganglia AFIP Atlas of Tumor Pathology, Fourth Series; fasc 8. 2007: Washington, D.C : American Registry of Pathology in collaboration with the Armed Forces Institute of Pathology, 2007.

2. Tischler AS, Asa SL. *Histology for Pathologists*, ed. Mills SE. 2018: Wolters Kluwer; Philadelphia, PA.
3. Rosenbaum JN, Guo Z, Baus RM, et al. INSM1: A Novel Immunohistochemical and Molecular Marker for Neuroendocrine and Neuroepithelial Neoplasms. *Am J Clin Pathol*. 2015;144:579–91. [PubMed: 26386079]
4. Mukhopadhyay S, Dermawan JK, Lanigan CP, et al. Insulinoma-associated protein 1 (INSM1) is a sensitive and highly specific marker of neuroendocrine differentiation in primary lung neoplasms: an immunohistochemical study of 345 cases, including 292 whole-tissue sections. *Mod Pathol*. 2019;32:100–9. [PubMed: 30154579]
5. Sun TT, Tseng SC, Huang AJ, et al. Monoclonal antibody studies of mammalian epithelial keratins: a review. *Ann N Y Acad Sci* 1985;455:307–29. [PubMed: 2417518]
6. Kuruc N, Franke WW. Transient coexpression of desmin and cytokeratins 8 and 18 in developing myocardial cells of some vertebrate species. *Differentiation*. 1988;38:177–93. [PubMed: 2463949]
7. Chetty R, Pillay P, Jaichand, PV. Cytokeratin expression in adrenal pheochromocytomas and extra-adrenal paragangliomas *J Clin Pathol* 1998;51:477–8. [PubMed: 9771451]
8. Dermawan JK, Mukhopadhyay S, Shah AA. Frequency and extent of cytokeratin expression in paraganglioma: an immunohistochemical study of 60 cases from 5 anatomic sites and review of the literature *Hum Pathol* 2019;93:16–22. [PubMed: 31442521]
9. Moriguchi T, Takako N, Hamada M, et al. Gata3 participates in a complex transcriptional feedback network to regulate sympathoadrenal differentiation. *Development* 2006;133:3871–81. [PubMed: 16943277]
10. Nonaka D, Wang BY, Edmondson D, et al. A study of gata3 and phox2b expression in tumors of the autonomic nervous system. *Am J Surg Pathol*, 2013;37:1236–41. [PubMed: 23715162]
11. So JS, Epstein JI. GATA3 expression in paragangliomas: a pitfall potentially leading to misdiagnosis of urothelial carcinoma. *Mod Pathol*, 2013;26:1365–70. [PubMed: 23599157]
12. Miettinen M, McCue PA, Sarlomo-Rikala M, et al. GATA3: a multispecific but potentially useful marker in surgical pathology: a systematic analysis of 2500 epithelial and nonepithelial tumors. *Am J Surg Pathol*, 2014;38:13–22. [PubMed: 24145643]
13. Iwase K, Nagasaka A, Nagatsu I, et al. Tyrosine hydroxylase indicates cell differentiation of catecholamine biosynthesis in neuroendocrine tumors. *J Endocrinol Invest*. 1994;17:235–9. [PubMed: 7523477]
14. Osinga TE, Korpershoek E, de Krijger RR, et al. Catecholamine-Synthesizing Enzymes Are Expressed in Parasympathetic Head and Neck Paraganglioma Tissue. *Neuroendocrinology*. 2015;101:289–95. [PubMed: 25677368]
15. Meijer WG, Copray SC, Hollema H, et al. Catecholamine-synthesizing enzymes in carcinoid tumors and pheochromocytomas. *Clin Chem*. 2003;49:586–93. [PubMed: 12651811]
16. Hermann G, Konukiewicz B, Schmitt A, et al. Hormonally defined pancreatic and duodenal neuroendocrine tumors differ in their transcription factor signatures: expression of ISL1, PDX1, NGN3, and CDX2. *Virchows Arch*. 2011;459:147–54. [PubMed: 21739268]
17. Weissferdt A, Kalhor N, Liu H, et al. Thymic neuroendocrine tumors (paraganglioma and carcinoid tumors): a comparative immunohistochemical study of 46 cases. *Hum Pathol*. 2014;45:2463–70. [PubMed: 25294372]
18. El-Salhy M, Lundqvist M, Wilander E. Bronchial carcinoids and pheochromocytomas. A comparative study with special emphasis on S-100 protein, serotonin and neuron-specific enolase (NSE) immunoreactivity. *Acta Pathol Microbiol Immunol Scand A*. 1986;94:229–35. [PubMed: 3524110]
19. Johnson TL, Zarbo RJ, Lloyd RV, et al. Paragangliomas of the Head and Neck: Immunohistochemical Neuroendocrine and Intermediate Filament Typing. *Mod Pathol* 1988;1:216–23. [PubMed: 2467285]
20. Achilles E, Padberg BC, Holl K, et al. Immunocytochemistry of paragangliomas-value of staining for S-100 protein and glial fibrillary acid protein in diagnosis and prognosis. *Histopathology*. 1991;18:453–8. [PubMed: 1679411]
21. Tsuta K, Raso MG, Kalhor N et al. Sox10-positive Sustentacular Cells in Neuroendocrine Carcinoma of the Lung. *Histopathology*;2011;58:276–85. [PubMed: 21323953]

22. Burke AP, Helwig EB. Gangliocytic paraganglioma. *Am J Clin Pathol*, 1989;92:1–9. [PubMed: 2750701]
23. Okubo Y, Nemoto T, Wakayama M, et al. Gangliocytic paraganglioma: a multi-institutional retrospective study in Japan. *BMC Cancer*. 2016;15:269. 1308–8.
24. Orrell JM, Hales SA. Paragangliomas of the cauda equina have a distinctive cytokeratin immunophenotype. *Histopathology* 1992;21:479–81. [PubMed: 1280616]
25. Miettinen M A simple method for generating multitissue blocks without special equipment. *Appl Immunohistochem Mol Morphol*. 2012;20:410–2. [PubMed: 22495380]
26. Ordonez NG. Value of GATA3 Immunostaining in the Diagnosis of Parathyroid Tumors. *Appl Immunohistochem Mol Morphol* 2014;22:756–61. [PubMed: 25046229]
27. Mete O, Kefeli M, Çalı kan S, et al. GATA3 immunoreactivity expands the transcription factor profile of pituitary neuroendocrine tumors. *Mod Pathol*, 2019;32:484–89. [PubMed: 30390035]
28. Kimura N, Shiga K, Kaneko K, et al. The diagnostic dilemma of GATA3 immunohistochemistry in pheochromocytoma and paraganglioma. *Endocr Pathol* 2020;31:95–100. [PubMed: 32303954]
29. Fujino Y, Nagata Y, Ogino K, Watahiki H, Ogawa H, Saitoh Y. Nonfunctional paraganglioma of the pancreas: report of a case. *Surg Today*. 1998;28:209–12. [PubMed: 9525014]
30. Ginesu GC, Barmina M, Paliogiannis P, et al. Nonfunctional paraganglioma of the head of the pancreas: A rare case report. *Int J Surg Case Rep*. 2016;8:81–4.
31. Moran CA, Rush W, Mena H. Primary spinal paragangliomas: a clinicopathological and immunohistochemical study of 30 cases. *Histopathology*. 1997;31:167–73. [PubMed: 9279569]
32. Capper D, Jones DTW, Sill M, et al. DNA methylation-based classification of central nervous system tumours. *Nature* 2018;555:469–74. [PubMed: 29539639]

- Keratins, GATA3, CDX2, TTF1, and tyrosine hydroxylase are useful in distinguishing paraganglioma from NETs.
- Gangliocytic paraganglioma of duodenum and cauda equina paraganglioma have NET-like immunoproliferation.
- SOX10 is superior to S100 protein to evaluate sustentacular cells as it does not label the chief cells in paraganglioma and NETs.
- Occasional pancreatic NETs mimic paragangliomas by their antigenic patterns.




Fig 1.

A. Paraortic paraganglioma is composed of clear or eosinophilic epithelioid cells. B. Most tumor cells in this case show exceptional positivity for keratin cocktail AE1/AE3. C. Tumor cells have nuclear positivity for GATA3. D. Tumor is strongly positive for tyrosine hydroxylase.




Fig. 2. Gangliocytic paraganglioma of duodenum is composed of nests of epithelioid cells containing ganglion like cells and Schwannian stroma between nests. B. The epithelioid cells including the ganglion-like cells are positive for keratin 8 (CAM5.2), which also highlights thin cell processes. C. The Schwannian stromal cells are positive for SOX10, while the epithelioid cells are negative. D. Only few tumor cells are tyrosine hydroxylase positive.




Fig. 3. A. Cauda equina paraganglioma is a highly vascular tumor composed of epithelioid cells with perivascular arrangements. B. Tumor is positive for keratins AE1/AE3. C. Few scattered tumor cells are positive for tyrosine hydroxylase. D. Like most neuroendocrine tumors, this one shows nuclear positivity for INSM1.




Fig. 4. Examples of tyrosine hydroxylase expression in epithelial neuroendocrine tumors. A. Ileal well-differentiated NET (carcinoid) shows a 1% cell population positive. B. Duodenal well-differentiated NET shows a small number of cells strongly positive and a larger proportion cells weakly positive. C. Well-differentiated NET (carcinoid) of lung contains 50% of positive cells. D. All tumor cells are positive for tyrosine hydroxylase in this unusual pancreatic NET.




Fig. 5. Pancreatic NET mimicking paraganglioma A. The tumor is composed of sheets of neuroendocrine cells between ductal elements. B. Tumor cells are negative for keratins AE1/AE3 (also negative for keratin 8/CAM5.2). C. Most tumor cells are positive for tyrosine hydroxylase. D. Tumor cells show nuclear positivity for CDX2 supporting the diagnosis of NET.

Table 1.

Antibodies and immunohistochemical protocols used in the study.

Antibody	Source	Clone	Dilution	Automation	Epitope retrieval/time (min)	Primary antibody incubation time (min)	Detection
CDX2	Roche	EPR2764Y	Predilute	Roche	CCI 64	36	Ultra view with amplification + DAB
Chromo-Granin A	Roche	LK2H10	Predilute	Roche	CCI 36	20	Ultra view + DAB
GATA3	Roche	L50-823	Predilute	Roche	CCI 64	60	Ultra view + DAB
GFAP	Roche	EP672Y	Predilute	Roche	CCI 64	32	Ultra view + DAB
INSM1	Santa Cruz	A-8	1:1000	Leica	ER2 25	30	Bond Polymer Refine + DAB
Keratins AE1/3	Dako Agilent	AE1+AE3	1:100	Roche	CCI 36	32	Ultra view + DAB
Keratin 8	Roche	CAM5.2	Predilute	Roche	Protease 1, 8 min	20	Ultra view + DAB
PNNMT	Novus Biologicals	OTT1D2	1:400	Leica	ER2 25	30	Bond Polymer Refine + DAB
S100 protein	Biogenex	15E2E2	1:8000	Roche	CCI 64	32	Ultra view + DAB
SOX10	Cell Marque	EP268	1:100	Roche	CCI 64	32	Ultra view + DAB
Synaptophysin	Roche	SP11	Predilute	Roche	CCI 64	32	Ultra view with amplification + DAB
TTF1	Roche	SP141	Predilute	Roche	CCI 64	20	Ultra view with block
Tyrosine hydroxylase	Abcam	EP1532Y	1:3000	Leica	ER2 25	30	Bond Polymer Refine + DAB

Abbreviations: CDX2 caudal type homeobox 2, INSM1 insulinoma-associated protein 1, PNNMT phenylethanolamine N-methyl transferase, TTF1 thyroid transcription factor 1.

Summary of marker expression in 560 paragangliomas and epithelial neuroendocrine tumors.

Table 2.

Tumor category	n =	Keratins AE1/AE3	GATA3	Tyrosine Hydroxylase positive cells	CDX2	TTF1	SOX10** 0/1/2/3	GFAP** 0/1/2/3
Pheochromocytoma, adrenal	142	8/142 *	132/142 (93)	138/142 (97)	0/145	2/142 *	30/33/35/26	109/4/1/1
Paraganglioma, retroperitoneal	49	3/49 *	46/49 (94)	43/49 (88)	0/48	0/48	4/11/18/12	44/1/0/0
Paraganglioma, urinary bladder	8	0/8	6/8 (75)	8/8 (100)	0/6	0/6	0/2/1/3	All negative
Paraganglioma, neck	54	1/54 *	51/54 (93)	20/54 (37)	0/54	0/54	3/2/19/30	14/12/14/14
Paraganglioma, thoracic	9	0/9	9/9 (100)	9/9 (100)	0/9	0/9	1/0/0/7	3/2/1/2
Ganglionic paraganglioma	9	9/9	0/9 (0)	2/9 * (22)	0/9	0/9	9/9 **	All negative
Cauda equina paraganglioma	3	3/3	0/3 (0)	2/3 * (0)	0/2	0/2	0/2/0/0	All negative
Carcinoid/NET, GI-tract, Gr. 1-2	81	81/81	0/81 (0)	6/81 * (7)	73/81	0/81	79/2/0/0	81/0/0/0
Carcinoid/NET, Lung and thymus, Gr. 1-2	42	42/42	1/42 * (2)	11/42 * (26)	0/41	19/41	28/7/4/2	All negative
NET, pancreas, Gr. 1-2	78	76/78	5/77 * (6)	7/77 * (9)	22/78	1/78	74/4/1/0	All negative
Medullary carcinoma, thyroid	37	37/37	0/37 (0)	1/37 * (2)	0/37	37/37	37/0/0/0	All negative
Neuroendocrine tumor, Gr. 3, Various organs	16	15/16	0/16 (0)	2/16 * (13)	8/16	7/16	16/0/0/0	All negative
Small cell carcinoma, lung	32	30/32	0/31 (0)	0/32 (0)	0/30	30/32	30/0/0/0	ND
Total	560							

* Positivity was usually focal.

** Expression only in sustentacular cells/spindled Schwannian cells. Number of cases with each score value (Number of positive cells per 0.2 mm². Scored 0-3: 0 = negative, 1 = 10, 2 = 11-49, 3 = 50). Numbers for SOX10 and GFAP represent the cases with no positivity (Score 0) to >50 cells positive (Score 3).

Details of scoring for SOX10 and GFAP are shown in materials and methods.

Table 3.

Epithelial neuroendocrine tumors (NETs) and paragangliomas with aberrant results on expression of keratins, GATA3, or tyrosine hydroxylase (TH) potentially challenging the original classification.

Case	Starting diagnosis	Age/Sex	Size	Ker E1+AE3/ keratin 8	GATA3	Tyrosine Hydroxylase	CDX2	TTF1	Score of SOX10/ GFAP	Conclusion
1	NET, pancreas, Grade 1	65 F	7.5 cm	0/0	100	0	0	0	2/0	Probable (peri)pancreatic paraganglioma, negative for keratins
2	NET, pancreas, Grade 1	58 F	unk	0/0	0	60	100	0	0/0	Probable pancreatic NET, positive for TH and CDX2
3	NET, pancreas, Grade 1	45 M	7.5 cm	100/100	80	0	0	0	0/0	Pancreatic NET with extensive GATA 3-positivity
4	NET, pancreas, Grade 2	59 F	7 cm	100/100	20	0	0	0	0/0	Pancreatic NET with GATA 3-positivity
5	NET, pancreas Grade 1	67 F	5.5 cm	100/100	0	80	0	0	0/0	Pancreatic NET with TH-positivity
6	NET, pancreas Grade 1	69 F	unk	70/100	0	100	90	0	1/0	Pancreatic NET with TH-positivity
7	NET, lung, Grade 1	65 F	unk	5/100	0	30	0	0	0/0	NET of lung with low keratin AE1/AE3 and TH-positivity
8	NET, lung, Grade 2	69 M	unk	20/100	0	10	0	0	0/0	NET of lung with low keratin AE1/AE3 and focal TH-positivity, TTF1-negative
9	NET, lung, Grade 1	18 F	2.5 cm	5/100	0	60	0	0	1/0	NET of lung with low keratin and extensive TH-positivity, TTF1-negative
10	NET, lung, Grade 1	61 F	2.2 cm	80/100	0	20	0	100	0/0	NET of lung with TH-positivity
11	Pheochromocytoma	34 M	1.8 cm	80/100	80	100	0	0	3/0	Pheochromocytoma with extensive keratin expression

The number under an immunomarker is the estimated percentage of positive tumor cells. NET = epithelial neuroendocrine tumor. TH = tyrosine hydroxylase.

Unk = unknown. Scoring of SOX10 (left of /) and GFAP (right of /): 0 = negative; 1 = 1–10 cells per HPF; 2 = 11–49 cells per HPF; 3 = 50 cells per HPF.