



Local Sleep and Alzheimer's Disease Pathophysiology

Bryce A. Mander^{1,2*}

¹ Department of Psychiatry and Human Behavior, University of California, Irvine, Irvine, CA, United States, ² Center for the Neurobiology of Learning and Memory, University of California, Irvine, Irvine, CA, United States

OPEN ACCESS

Edited by:

Phyllis Zee,
Northwestern University,
United States

Reviewed by:

David Holtzman,
Washington University in St. Louis,
United States
Marilyn J. Duncan,
University of Kentucky, United States

*Correspondence:

Bryce A. Mander
bmander@uci.edu

Specialty section:

This article was submitted to
Sleep and Circadian Rhythms,
a section of the journal
Frontiers in Neuroscience

Received: 11 January 2020

Accepted: 01 September 2020

Published: 23 September 2020

Citation:

Mander BA (2020) Local Sleep
and Alzheimer's Disease
Pathophysiology.
Front. Neurosci. 14:525970.
doi: 10.3389/fnins.2020.525970

Even prior to the onset of the prodromal stages of Alzheimer's disease (AD), a constellation of sleep disturbances are apparent. A series of epidemiological studies indicate that multiple forms of these sleep disturbances are associated with increased risk for developing mild cognitive impairment (MCI) and AD, even triggering disease onset at an earlier age. Through the combination of causal manipulation studies in humans and rodents, as well as targeted examination of sleep disturbance with respect to AD biomarkers, mechanisms linking sleep disturbance to AD are beginning to emerge. In this review, we explore recent evidence linking local deficits in brain oscillatory function during sleep with local AD pathological burden and circuit-level dysfunction and degeneration. In short, three deficits in the local expression of sleep oscillations have been identified in relation to AD pathophysiology: (1) frequency-specific frontal deficits in slow wave expression during non-rapid eye movement (NREM) sleep, (2) deficits in parietal sleep spindle expression, and (3) deficits in the quality of electroencephalographic (EEG) desynchrony characteristic of REM sleep. These deficits are noteworthy since they differ from that seen in normal aging, indicating the potential presence of an abnormal aging process. How each of these are associated with β -amyloid (A β) and tau pathology, as well as neurodegeneration of circuits sensitive to AD pathophysiology, are examined in the present review, with a focus on the role of dysfunction within fronto-hippocampal and subcortical sleep-wake circuits. It is hypothesized that each of these local sleep deficits arise from distinct network-specific dysfunctions driven by regionally-specific accumulation of AD pathologies, as well as their associated neurodegeneration. Overall, the evolution of these local sleep deficits offer unique windows into the circuit-specific progression of distinct AD pathophysiological processes prior to AD onset, as well as their impact on brain function. This includes the potential erosion of sleep-dependent memory mechanisms, which may contribute to memory decline in AD. This review closes with a discussion of the remaining critical knowledge gaps and implications of this work for future mechanistic studies and studies implementing sleep-based treatment interventions.

Keywords: Alzheimer's disease, sleep, slow waves, sleep spindle, REM sleep, beta-amyloid protein, tau and phospho-tau protein, neurodegeneration

INTRODUCTION

Both macro and micro features of sleep architecture change across the adult lifespan, and the individual variability in the magnitude of change appears to be associated with the degree of cognitive decline and the severity of risk for dementias such as Alzheimer's disease (AD) (Ohayon et al., 2004; Redline et al., 2004; Yaffe et al., 2011; Lim et al., 2013a,b; Spira et al., 2013; Lo et al., 2014; Mander et al., 2015, 2016, 2017a; Osorio et al., 2015; Song et al., 2015; Sprecher et al., 2015, 2017; Chen et al., 2016; Bubu et al., 2017, 2019; Kabeshita et al., 2017; Pase et al., 2017; Shi et al., 2017; Tsapanou et al., 2017; Lutsey et al., 2018). While the mechanisms underlying these associations remain elusive, emerging evidence has elucidated bidirectional associations between biological mechanisms of aging and AD and the expression of sleep processes related to neuroplasticity (Ju et al., 2014; Mander et al., 2016). In particular, certain micro features of sleep are expressed locally within specific brain circuits supporting processes regulating systemic and synaptic neuroplasticity (McGaugh, 2000) in order to facilitate memory formation and consolidation (Buzsaki, 1998; Steriade, 2006; Walker, 2009; Diekelmann and Born, 2010; Wamsley and Stickgold, 2011; Watson and Buzsaki, 2015). New findings indicate that topographic and frequency-specific disruptions in the expression of these brain oscillations, including slow waves and sleep spindles during non-rapid eye movement (NREM) sleep and EEG desynchrony during REM sleep, occur in the context of AD pathophysiology—in some cases even prior to onset of mild cognitive impairment (MCI)—and these local sleep deficits differ from those observed in normal aging (Hassainia et al., 1997; Petit et al., 2004; Jyoti et al., 2015; Mander et al., 2015, 2016, 2017a; Brayet et al., 2016; Gorgoni et al., 2016; De Gennaro et al., 2017; Holth et al., 2017; Lucey et al., 2019). It is therefore possible that certain local sleep deficits may reflect an abnormal aging process in its initial stages, and that disruptions in sleep-dependent memory mechanisms may contribute to the biological and cognitive consequences of AD pathophysiology. In this review, we summarize the state of the science linking local sleep to core features of AD pathophysiology and AD-related hippocampus-dependent memory impairment.

LOCAL SLEEP

The brain expresses a complex and dynamic organization of oscillatory activities during sleep, cycling between NREM and REM sleep stages that each distinctly organize local brain activities across frequency, topography, and time (Landolt et al., 1996; Liscombe et al., 2002; Steriade, 2006; Diekelmann and Born, 2010; Staresina et al., 2015; Watson and Buzsaki, 2015; Sprecher et al., 2016; Helfrich et al., 2018; Scarpelli et al., 2019). The spatially circumscribed nature of the expression of these sleep-specific oscillatory activities and their frequency specificity highlight the local nature of some canonical sleep features. For example, most NREM slow wave and sleep spindle oscillations are expressed regionally and migrate across cortex

in a predictable manner (Massimini et al., 2004; Nir et al., 2011; Muller et al., 2016). Further, in cases of excessive sleep pressure, isolated neuronal ensembles have even been shown to express these sleep-specific rhythms while the majority of cortex remains in a state of wakefulness (Van Dongen et al., 2011; Krueger et al., 2019). The fact that this occurs has led to the hypothesis that sleep is not entirely global, i.e., it does not always occur equally throughout the entire brain (Van Dongen et al., 2011; Krueger et al., 2019). Furthermore, even during sleep local expression of certain oscillatory activities, such as slow waves during NREM sleep, can be influenced by prior waking experiences in a topographically-specific manner. More specifically, neuronal ensembles more active during learning will exhibit greater local intensity of slow wave activity relative to surrounding cortex (Huber et al., 2004, 2006). These phenomena demonstrate that certain sleep features are local and not global.

Two specific NREM sleep oscillations of note that are organized in this fashion are slow waves and sleep spindles. Slow waves, defined as low frequency (0.5–4.5 Hz) and high amplitude ($>75 \mu\text{V}$) waveforms, are generated in cortex, predominantly within association cortices including medial prefrontal cortex, insular cortex, and the cingulate cortex (Steriade, 2006; Murphy et al., 2009). Their expression is dependent on the coordinated synchrony of hyperpolarized down states and depolarized up states of local neuronal ensembles (Steriade et al., 1993b; Steriade, 2006). The spectral power of slow waves, termed slow wave activity (SWA), peaks over frontal electroencephalography (EEG) derivations in the first quartile of the night, dissipating across the sleep period in a manner reflective of a homeostatic “Process S” (Borbely et al., 1981; Borbely, 1982; Landolt et al., 1996; Carrier et al., 2001; Mander, 2013; Mander et al., 2017a). Recent optogenetic findings support the existence of two kinds of slow waves: lower frequency slow oscillations (SO; <1 Hz) that support memory consolidation processes and faster delta waves (1–4.5 Hz) that facilitate forgetting (Steriade, 2006; Kim et al., 2019). Whether the temporal or topographical organization of these waveforms is distinct or is influenced by aging or AD differentially remains to be seen, and should be a focus of future studies.

A second characteristic sleep oscillation that is critically tied to neuroplasticity is the sleep spindle. Sleep spindles are transient (0.5–3 s) bursts of oscillatory activity in the sigma frequency range (11–16 Hz) with waxing and waning components. They are generated within the reticular nucleus of the thalamus and expressed through cortico-thalamic loops (De Gennaro and Ferrara, 2003; Steriade, 2006). Two kinds of sleep spindles have been described as well: fast (~ 13 –16 Hz) and slow (~ 11 –13 Hz) frequency sleep spindles (Jobert et al., 1992; De Gennaro and Ferrara, 2003). Though exact frequency definitions vary by study, the distinct topographic nature of their expression does not. Fast frequency sleep spindles tend to peak over midline central and parietal EEG derivations, while slow frequency sleep spindles tend to peak in expression over frontal EEG derivations (Jobert et al., 1992; De Gennaro and Ferrara, 2003). Moreover, while SWA may dissipate across the sleep period,

spindle activity tends to peak in the morning (Landolt et al., 1996). However, these NREM sleep oscillations are not expressed in isolation, tending to couple during the rising phase of the depolarizing slow wave up state (Steriade et al., 1993a; Steriade, 2006; Staresina et al., 2015). This phase-locking is specific to the SO and is also accompanied by phase locking of another oscillation, the hippocampus ripple, within the troughs of the sleep spindle oscillation (Staresina et al., 2015; Helfrich et al., 2018). Coordinated firing of neuronal ensembles during initial encoding has been observed to spontaneously reoccur during ripples (Wilson and McNaughton, 1994; Ji and Wilson, 2007; Davidson et al., 2009), with ripples instigating information transfer from the hippocampus to the cortex when phase locked to SOs and sleep spindles (Olcese et al., 2018; Helfrich et al., 2019). The coupling of these three cortical, thalamic, and hippocampal brain oscillations, and the “replay” associated with them, is thought to support the consolidation of memories acquired during prior wakefulness (Steriade, 2006; Diekelmann and Born, 2010; Watson and Buzsaki, 2015; Helfrich et al., 2018, 2019). Indeed, recent evidence demonstrates that stimulation and suppression of sleep spindles only impacts sleep-dependent memory consolidation if it occurs during the rising phase of the depolarizing up-state and in coordination with hippocampal ripples (Latchoumane et al., 2017).

The spectral content and topographic organization of REM sleep differs from that observed in NREM sleep. The apparent wake-like nature of EEG-measured neuronal activity during REM sleep is why it is also known as “paradoxical sleep” (Peever and Fuller, 2016), though there are important distinctions that are observed when using other neuroimaging modalities (Braun et al., 1997; Maquet, 2000) or analytic approaches (Lendner et al., 2020). Of particular note, while the low voltage mixed frequency content of the “desynchronized” EEG observed during wakefulness is supported by a series of brainstem, midbrain, and hypothalamic glutamatergic, monoaminergic, and cholinergic inputs, cholinergic input plays a more dominant role in generating the desynchronized EEG in REM sleep than monoaminergic inputs (Peever and Fuller, 2016; Scammell et al., 2017). Thus, while the EEG looks the same in wake and REM sleep, the neurobiological correlates are distinct.

In terms of spectral content, both alpha and theta power are particularly prominent in REM sleep (Landolt et al., 1996), with theta peaking over fronto-central derivations, and alpha peaking over occipito-parietal derivations (Scarpelli et al., 2019). Similar to theta activity, higher frequency content, including beta and gamma power, peak over fronto-central derivations, but while low frequency power, including delta and theta power decrease across successive REM sleep periods, high frequency content remains relatively stable (Liscombe et al., 2002). Little is known about the functional relevance of the organized expression of EEG activities during REM sleep, and this should be a topic of future study.

Substantial evidence indicates that both aging and AD influence local NREM and REM sleep expression in distinct ways, and these effects are reviewed below.

LOCAL SLEEP IN AGING AND ALZHEIMER'S DISEASE (AD)

Aging

Increasing age is associated with reductions in oscillatory activities across multiple frequency bands in non-rapid eye movement (NREM) sleep (Sprecher et al., 2016). The biggest reductions occur globally across the frequency bands of slow waves (0.5–4.5 Hz) and sleep spindles (12–16 Hz), with the largest reductions occurring over frontal EEG derivations, due to the reduced incidence and amplitude of slow wave and sleep spindle oscillations (Dijk et al., 1989; Landolt et al., 1996; Carrier et al., 2001; Mander, 2013; Martin et al., 2013; Mander et al., 2014, 2017a,b; Sprecher et al., 2016). Aging also disrupts the phase-locked synchrony between slow waves and sleep spindles, with slow wave-sleep spindle coupling being more variable and occurring at an earlier phase of the slow wave, closer to the hyperpolarized slow wave down state (Helfrich et al., 2018; Muehlroth et al., 2019). All these age effects on NREM sleep oscillation expression have been associated with age effects on frontal and hippocampal gray and white matter volume and integrity (see Mander et al., 2017a, for an in depth review). Central to this work is the consideration of whether these age-related changes contribute to cognitive decline in aging, and this will be reviewed in detail later in this review.

In addition to NREM sleep, a few studies have explored age effects on REM sleep architecture. These studies have shown that older adults experience a modest reduction in REM sleep duration (0.6% per decade) that emerges much later than age effects on NREM sleep (Ehlers and Kupfer, 1997; Van Cauter et al., 2000; Carrier et al., 2001; Gori et al., 2004; Ohayon et al., 2004; Redline et al., 2004; Floyd et al., 2007; Scarpelli et al., 2019). However, qualitative differences in REM sleep are also present in older adults even if REM duration reductions are minimal, including increased awakenings from REM sleep, decreased REM latency, decreased REM density, and shorter and more disorganized REM bursts, particularly in adults over 65 years (Gillin et al., 1981; Ehlers and Kupfer, 1989, 1997; Vegni et al., 2001; Darchia et al., 2003; Conte et al., 2014). In terms of REM sleep microarchitecture, a few reports have indicated spectral power is reduced across delta, theta, and alpha frequencies in older adults, particularly over central derivations (Landolt et al., 1996; Scarpelli et al., 2019). It remains unknown why REM sleep microarchitecture is reduced in aging, and whether these changes are functionally relevant or epiphenomenal.

Alzheimer's Disease

Local sleep deficits have been observed in the context of MCI, AD, and even in healthy older adults with AD pathology (**Figure 1**). These local sleep deficits diverge from those observed in the context of normal aging both in terms of topography and frequency (Petit et al., 2004; Mander et al., 2016, 2017a). Reports have shown that slow wave sleep (SWS) is reduced and more fragmented in MCI, AD, and in rodent models of AD (Prinz et al., 1982; Gagnon et al., 2008; Hita-Yanez et al., 2012, 2013; Roh et al., 2012; Westerberg et al., 2012; Mander, 2013; Jyoti et al., 2015;

Kam et al., 2016), SWA is lower over midline frontal, central, and parietal derivations in healthy older adults with AD pathology and patients with MCI (Westerberg et al., 2012; Mander et al., 2015; Varga et al., 2016b; Lucey et al., 2019), and there are fewer frontal K-complexes in AD (De Gennaro et al., 2017). However, this effect of AD pathology and diagnosis on slow wave expression appears to depend on slow wave frequency, particular AD pathology, and, potentially, disease stage (**Figure 1A**). More specifically, in healthy older adults with cortical A β pathology, SWA deficits appear to be specific to lower frequencies (i.e., in the SO frequency range; < 1 Hz), with increases in SWA observed in the delta frequency range (1–4 Hz) (Mander et al., 2015; Kastanenka et al., 2017, 2019). This effect may change to reflect a more global loss of SWA across frequencies once tau pathology reaches the cortex in the MCI stage (De Gennaro et al., 2017; Lucey et al., 2019). Similar findings were observed in transgenic mouse models of AD, though this depended on which model was examined. Decreases in fronto-parietal SWA and increases in higher frequency power were observed during NREM sleep in Tg2576 and APP/PS1 transgenic mouse models of AD, but not in 3xTgAD mice (Zhang et al., 2005; Kent et al., 2018). This is noteworthy given that Tg2576 and APP/PS1 transgenic AD mouse models both exhibit rapid increases in A β pathology in the absence of neurofibrillary tangles, while the 3xTgAD mice express a more mild level of both A β and tau pathology. Together, these findings indicate that the influence of AD pathophysiology on SWA may not be fixed, but instead may change depending on the location and degree of burden of distinct AD pathologies.

Reduction in sleep spindle expression has also been observed in the context of AD. There are fewer sleep spindles and lower spindle activity in patients with MCI and AD (Rauchs et al., 2008; Westerberg et al., 2012; Gorgoni et al., 2016). However, this effect appears to be specific to faster frequencies (13–16 Hz) over parietal derivations (**Figure 1B**), with no discernable effect on frontal sleep spindles in slower frequencies (11–13 Hz) (Westerberg et al., 2012; Gorgoni et al., 2016). This is in stark contrast to the effect of age, which selectively disrupts frontal sleep spindles regardless of frequency, leaving parietal fast frequency sleep spindles largely spared (Martin et al., 2013; Sprecher et al., 2016; Mander et al., 2017a). The reduction of sleep spindles, and in particular fast frequency sleep spindles, is also apparent in healthy older adults with tau pathology (Kam et al., 2019). The effect of AD pathologies and MCI and AD diagnosis on slow wave-sleep spindle coupling remain unclear, though a recent study showed that tau pathological burden in the medial temporal lobe was associated with reduced slow wave-sleep spindle coupling (Winer et al., 2019). Because these parietal fast frequency sleep spindle deficits associated with MCI and AD are so topographically distinct from age-related deficits in sleep spindles, it is possible that these deficits could represent an early physiological signal distinguishing between abnormal and normal aging processes.

A robust literature demonstrates that MCI and AD diagnosis is associated with characteristic changes in REM sleep microarchitecture in a topographically-specific manner (**Figure 1C**). Specifically, a loss of EEG desynchrony during REM sleep is observed as early as MCI (Brazete et al., 2013;

Brayet et al., 2016), and can be used to differentiate individuals with and without AD with high diagnostic accuracy (Hassainia et al., 1997). This measure, quantified as the ratio of low frequency to high frequency power can even distinguish between individuals with amnesic versus non-amnesic MCI (Brayet et al., 2016), but can also predict non-AD dementia onset and cognitive decline in individuals with REM behavior disorder (Brazete et al., 2016). It is therefore possible that this signature may be more specific to dementia-related cognitive decline than AD, *per se*. The effect of MCI and AD diagnoses on EEG desynchrony is particularly prominent over central and posterior EEG derivations (Hassainia et al., 1997; Brazete et al., 2013, 2016; Brayet et al., 2016). Alongside these findings, others have found loss of central-parietal theta activity during REM sleep in patients with amnesic MCI (Westerberg et al., 2012), and both AD and tauopathy rodent models show reduced theta and alpha activity (Holth et al., 2017). Hence, the quantitative expression of REM sleep EEG is also impacted by MCI and AD, and these deficits may track cognitive decline associated with dementia.


The distinct nature of the effects of AD pathophysiology on the expression of local sleep features relative to those observed with advancing age hints at a distinction in underlying mechanisms, which may herald the emergence and progression of AD in its initial stages. Recent work highlights the mechanistic links between local sleep disturbance and AD pathophysiological features, which are examined in detail in the next section.

EMERGING LINKS BETWEEN LOCAL SLEEP AND AD PATHOPHYSIOLOGY

The Alzheimer's Disease Cascade—The Role of Amyloid (A), Tau (T), and Neurodegeneration (N)

Contemporary theories of AD pathogenesis describe the emergence of a biological cascade of pathological events which trigger a progressive degenerative process that ultimately results in the clinical and cognitive symptoms of dementia (Jack et al., 2010, 2013; Jagust, 2018; Marquez and Yassa, 2019). While there remains controversy over which pathological event emerges first (see Herrup, 2015; Musiek and Holtzman, 2015, for further details), what is clear is that the initial stages involve the build-up of cortical A β (A) and subcortical tau (T; **Figure 2**). As they converge and interact, tau begins to spread cortically and A β subcortically, triggering progressive and widespread neurodegeneration (N) which results in progressive cognitive and clinical dysfunction, and ultimately gross loss of basic functions (Jack et al., 2010, 2013, 2019).

The probability of onset and rate of progression of this neurodegenerative process is influenced by other biological factors which directly or indirectly influence this core biology of AD. For example, many of the known genetic risk factors for familial and sporadic AD, such as mutations in the amyloid precursor protein (APP), presenilin, and apolipoprotein E (APOE) genes, result in an acceleration of A β accumulation and the resulting neurodegeneration (Bagyinszky et al., 2014).




Sex also influences AD, with women being at twofold greater risk for developing AD relative to men (Podcasy and Epperson, 2016). Recent evidence indicates that one potential reason for this is that the same degree of A β burden may facilitate a greater degree of accumulation of tau pathology in women relative to men (Buckley et al., 2019).

Emerging evidence indicates that chronic inflammation plays a critical role in mediating the interactions between multiple pathophysiologic features of AD, facilitating its progression (for in depth reviews see, Heppner et al., 2015; Spangenberg and Green, 2017; Wang et al., 2017). Microglia—glial cells supporting innate immunity within the central nervous system—are activated by A β through its interaction with the triggering receptor expressed on myeloid cells 2 (TREM2), resulting in the release of proinflammatory cytokines and chemokines, including interleukin-1 β (IL-1 β), IL-6, tumor necrosis factor alpha (TNF- α), among others (Heppner et al., 2015; Spangenberg and Green, 2017). This activation, alongside activation of astrocytes, facilitates A β clearance through degradation and phagocytosis (Heppner et al., 2015; Spangenberg and Green, 2017). When A β plaques are formed, both astrocytes and microglia surround them to degrade and clear them from the central nervous system, as well as to form a barrier to minimize the toxic

effects of A β on surrounding neural tissue (Heppner et al., 2015; Spangenberg and Green, 2017).

Unfortunately, A β also impairs microglial function, suppressing its ability to degrade and clear A β . This leads to the induction of a chronic inflammatory state, which facilitates microglial burn-out, astrocytic atrophy, sustained A β production, decreased A β clearance, disrupted tau-microtubule binding, neurofibrillary tangle (NFT) formation, and propagation of tau pathology (Heppner et al., 2015; Spangenberg and Green, 2017). Chronically activated microglia also contribute to neurodegeneration and cognitive decline through dysfunctional phagocytosis, which results in pathological stripping of synapses and facilitation of neuronal loss (Spangenberg et al., 2016), a process also observed following sleep deprivation (Belleli et al., 2017). Indeed, the number of activated microglia have been shown to correlate with cognitive decline (Cagnin et al., 2001; Versijpt et al., 2003; Edison et al., 2008). These findings are supported by studies in humans showing that chronic systemic inflammatory conditions, such as rheumatoid arthritis, increase risk for developing AD through sustained increases in IL-1 β , IL-6, TNF- α , and C-reactive protein (CRP) levels, which can cross the blood brain barrier and directly impact brain A β metabolism or impact A β in the periphery which



then interacts with central A β pools (Wang et al., 2017). Further, in addition to genetic mutations in genes impacting A β accumulation, genetic mutations in numerous genes affecting microglial function, such as mutations in TREM2, also increase AD risk (Spangenberg and Green, 2017). In addition, release of APOE by astrocytes is thought to critically support the capacity of microglia to clear A β (Terwel et al., 2011), and thus mutations in the APOE gene may impact A β in part through its impact on central immunity. It is important to note, however, that chronic inflammation can precede A β plaque deposition, and facilitate AD pathogenesis (Krstic et al., 2012), indicating that chronic inflammation may not only be an outcome of A β pathophysiology but also an initiating event. Thus, onset into and progression of AD depends on the interaction between two distinct pathologies, that of A β and tau, and the influence of other biological factors that affect them, their interaction, and potentially their impact on neuronal integrity and function.

Prior to widespread degeneration, A β and tau also influence neuronal function in distinct ways. A recent study implemented multiple rodent models and examined neuronal dysfunction resulting from them (Busche et al., 2019). In a model expressing only A β , neuronal hyperexcitability was predominant (Busche et al., 2019), mirroring the dysfunctional hippocampus and default mode network hyperactivation observed in early stages of MCI and A β positive but cognitively asymptomatic older adults (Sperling et al., 2009; Yassa et al., 2010b; Leal et al., 2017). This is also consistent with other studies reporting increased incidence of interictal spikes, particularly during sleep, in rodent models overexpressing A β pathology (Kam et al., 2016). In further support of this notion, recent studies have shown that antiepileptic drugs targeting this presumably A β -related hyperactivity resolved hippocampus-dependent memory impairments observed in early stages of amnesic MCI (Bakker et al., 2012, 2015). In contrast, rodent models expressing either tau alone or both A β and tau resulted

in neuronal silence (Busche et al., 2019), meaning that, when convergent, tau trumps the effects of A β on neuronal function. This finding supports the possibility that the form of neuronal dysfunction observed in AD, and the cognitive consequences of that dysfunction, may depend on the neural network examined and the relative abundance of both A β and tau pathology within that network. This theoretical model of AD offers critical insight into the mechanistic links between local NREM and REM sleep dysfunctions and AD (Figure 1D). It also highlights the potential of these local sleep dysfunctions to offer a window into AD stage and progression, and the cognitive consequences of their relationships, all of which are reviewed below.

β -Amyloid and Local Expression of Slow Wave Activity (A)

A β levels increase during wakefulness and decrease during sleep in the interstitial fluid (ISF) in rodents and CSF in humans, and subjective and objective measures of sleep duration and quality are associated with A β burden (Kang et al., 2009; Roh et al., 2012; Ju et al., 2013; Spira et al., 2013; Mander et al., 2015; Sprecher et al., 2015, 2017; Varga et al., 2016b), indicating that A β peptides are regulated by the sleep/wake cycle. Collectively, these studies show that NREM sleep loss is associated with greater A β burden while greater A β burden further disrupts NREM sleep, potentially triggering a vicious cycle fostering AD progression (Ju et al., 2014; Mander et al., 2016). The mechanisms for these bidirectional relationships remain unclear, though there are a few candidates highlighted by recent work. First and foremost considers how NREM sleep disruption may result in increased A β burden. A β burden increases with increasing neuronal activity (Li et al., 2013), and neurometabolic activity is higher in wake than in SWS (Buchsbaum et al., 1989; Braun et al., 1997; Dworak et al., 2010). Thus, reductions in SWS duration increases the duration of wakefulness which then results in increased net A β accumulation.

A second possible mechanism regards findings linking disruptions in slow oscillation expression in APP transgenic mice to A β plaque burden (Kastanenka et al., 2017, 2019). In APP transgenic mice, SWA is reduced specifically in the slow oscillation frequency range (<1 Hz) by 3 months of age, 2 months before A β plaques are apparent (Kastanenka et al., 2017). This is accompanied by reduced GABA levels and downregulation of GABA_A and GABA_B receptor expression, which directly impacted the expression of slow oscillations by disrupting neuronal synchrony. Slow oscillation expression was restored by both increasing GABA levels and by optogenetically enhancing neuronal synchrony, and this resulted in reduced A β plaque burden and intracellular calcium overload (Kastanenka et al., 2017). In contrast, doubling the frequency of the slow oscillation to >1 Hz resulted in increased A β production, increased intracellular calcium overload, and decreased dendritic spine density (Kastanenka et al., 2019).

Yet another possibility involves the recently discovered glymphatic system, which is a biological process in the brain that actively clears toxins and waste from the ISF by flushing the brain with CSF pulses (Jessen et al., 2015; Rasmussen et al., 2018). Two studies actively link this process to SWS. First,

glymphatic flow is increased during SWS and results in increased clearance of A β proteins in rodents (Xie et al., 2013). Second, SWA peaks prior to and is phase locked to CSF flow through the brain in humans (Fultz et al., 2019). These findings suggest that it may not be NREM sleep, *per se*, but the expression of slow waves that is mechanistically linked to glymphatic clearance during sleep. If true, the efficacy of this system could depend on the integrity of astrocytic mechanisms regulating neuronal function. Astrocytes directly regulate the transition to and expression of slow oscillations, potentially through spatial K⁺ buffering through the inwardly rectifying potassium channel Kir4.1 expressed in astrocytes in tripartite synapses (Fellin et al., 2009; Poskanzer and Yuste, 2016; Haydon, 2017; Cucchiara et al., 2020). Astrocytic function is also central to the expression of the glymphatic system (Nedergaard, 2013). However, high A β burden in AD results in loss of Kir4.1 potassium channels, as well as aquaporin 4 (AQP4) channels which are also hypothesized to be critical for glymphatic function (Wilcock et al., 2009; Nedergaard, 2013; Kress et al., 2014; Zeppenfeld et al., 2017). Hence, disruptions in slow waves during NREM sleep, associated with astrocytic dysfunction, may then decrease active clearance of AD pathology, also potentially yielding a net increase in A β accumulation. These possibilities are supported by recent studies showing significant associations between slow wave incidence and power and A β burden (Zhang et al., 2005; Mander et al., 2015; Varga et al., 2016b; Kastanenka et al., 2017; Kent et al., 2018). Directly addressing this possibility, a recent study showed that active suppression of SWA using acoustic stimulation resulted in increased CSF A β levels in middle-aged adults, indicating the potential for a causal relationship (Ju et al., 2017). However, it is important to note that the role of glymphatic clearance during SWS on A β protein levels has only been shown using injected radiolabeled A β proteins, and no study to date has shown direct evidence for endogenous clearance of A β via the glymphatic system during SWS (Smith and Verkman, 2018). This limitation is a critical caveat that must be addressed in future studies.

Lastly, it has been hypothesized that another potential pathway linking sleep deficits to increased A β pathology and AD risk is through an association with chronic inflammation (Irwin and Vitiello, 2019). Aging is coincidentally associated with chronic inflammation, SWS disruption, and increased AD risk (Ingiosi et al., 2013; Besedovsky et al., 2019). Sleep, and in particular SWS, has a complex bidirectional interaction with biological processes supporting immune function (for in depth reviews see, Besedovsky et al., 2019; Irwin, 2019). A recent meta-analysis of over 70 studies indicated that evidence of sleep disturbance, short sleep duration, and excessively long sleep duration was associated with elevated plasma levels of CRP and IL-6 (Irwin et al., 2016). Further, short sleep duration, greater wake after sleep onset (WASO), worse sleep efficiency, and greater percentages of the sleep period with blood oxygen saturations below 90% are associated with increased IL-6, CRP, TNF- α , and interferon γ (INF- γ) in older adults and AD caregivers (Friedman et al., 2005; Von Kanel et al., 2006; Smagula et al., 2016). In addition, experimental sleep restriction to four hours increased nuclear factor κ -light-chain-enhancer of activated B cells (NF- κ B) in plasma, which is a critical

factor regulating the inflammatory signal cascade (Irwin et al., 2008). Moreover, chronic sleep restriction to 4 h per night for 12 days resulted in increased IL-6 plasma levels in young adults (Haack et al., 2007).

In addition to sleep disturbance triggering an inflammatory response, inflammation directly impacts sleep expression as well. However, the specifics of this effect depend on the duration and magnitude of inflammatory activation. While acute inflammation can induce SWS and enhance SWA, chronic inflammatory conditions are associated with fragmented sleep, alpha intrusions in NREM sleep, and reduced SWS, SWA, and REM sleep (Besedovsky et al., 2019; Irwin, 2019). Together these findings portray a complex interaction between inflammation and SWS in later life, with age-related increases in chronic inflammation potentially impairing SWS expression and overall sleep quality, ultimately resulting in greater systemic inflammation. Since chronic inflammation promotes A β accumulation (Krstic et al., 2012; Heppner et al., 2015; Spangenberg and Green, 2017), SWS deficits may increase A β burden through an interaction with markers of systemic and central inflammation. However, it should be noted that data linking either systemic or central inflammation markers to the local expression of any sleep oscillation is limited. Future work should examine this more closely to more fully address the hypothesis that slow wave deficits increase A β burden through interactions with inflammatory processes.

The second mechanistic question involves how A β burden may impact the expression of slow waves. NREM sleep is disrupted in rodent models over-expressing A β (Roh et al., 2012). Moreover, our recent work has shown that there is an association between SWA and cortical A β burden in cognitively asymptomatic older adults, but that this association is distinct from that due to age and age-related frontal cortical atrophy (Mander et al., 2015). A β was not associated with a global loss of slow waves or a global reduction in SWA, as is common in aging (Mander et al., 2017a), but instead showed a negative association with slow waves and SWA in the SO frequency range (0.6–1 Hz) and a positive association with slow waves and SWA in the delta frequency range (1–4 Hz), representing a shift in the slow wave distribution (**Figure 1A**) (Mander et al., 2015; Winer et al., 2019). If A β did not disrupt slow waves, but only was increased by slow wave disruption, than it would be expected that age-related changes in slow waves would predict increases in A β burden. This was not what we found. Instead, deficits associated with SWA were specific to A β and independent of age, indicating that A β may further disrupt slow wave expression independently of age-related changes. Interestingly, a similar effect on the SWA distribution was also observed in APP transgenic mice, a rodent model of AD selectively overexpressing A β pathology, with the loss of low frequency SO activity emerging as early as 3 months of age (Kastanenka et al., 2017). The mechanisms for how A β may disrupt slow wave expression is unknown, though there are some theoretical possibilities. One possibility is that A β may disrupt the generation and expression of slow waves as it accumulates in cortical slow wave generators (**Figures 1A,D**). Indeed, A β plaques initially deposit preferentially within many slow wave generating regions, including within the medial prefrontal cortex

(mPFC) and anterior and posterior cingulate gyri (Buckner et al., 2005; Murphy et al., 2009; Sepulcre et al., 2013; Jagust, 2018). Supporting this possibility, the association between A β burden and SWA is not global, but peaks locally over midline and frontal derivations (Mander et al., 2015; Varga et al., 2016b). Further, source analysis confirmed that the strongest predictor of frontal SWA was A β burden within the same mPFC region where frontal slow waves were sourced (Mander et al., 2015). It remains to be determined exactly how A β burden could disrupt slow wave generation and expression. One hypothesis is that A β directly acts on the neurobiological mechanisms controlling slow wave expression (Mander et al., 2016). There is some evidence that A β disrupts GABA levels and GABA and NMDA receptor signaling (Kurup et al., 2010; Busche et al., 2015; Kastanenka et al., 2017), which are critical for SO expression (Steriade et al., 1993b). Another potential mechanism is the effect of A β on the inwardly rectifying potassium channel Kir4.1 in astrocytes (Wilcock et al., 2009). Astrocytes have been shown to impact slow oscillation expression through an impact on NMDA receptor activity in neurons (Fellin et al., 2009). Further, a recent report has linked a gain of function mutation in Kir4.1 channels in children expressing the Autism-Epilepsy Phenotype with a lengthening of the slow oscillation period during NREM sleep (Cucchiara et al., 2020). Since the presence of A β results in a loss of Kir4.1 channels (Wilcock et al., 2009), rather than a gain of function, it is likely that A β could trigger the opposite effect, i.e., a shortening of period and thus an increase in the mean slow oscillation frequency. This hypothesis remains to be tested and should be a focus of future studies. Yet another possibility is that A β burden triggers generalized cortical hyperexcitability, which then results in a shift toward a greater abundance of delta waves and a reduction in SO events as a consequence. Indeed, optogenetically doubling the slow oscillation frequency from 0.6 to 1.2 Hz in APP transgenic mice resulted in increased A β production, supporting the frequency specificity and bidirectional nature of this effect (Kastanenka et al., 2019). Lastly, since A β increases markers of inflammation and chronic inflammation can disrupt SWA expression, inflammation may also contribute to deficits in SWA associated with A β (Besedovsky et al., 2019; Irwin, 2019). Supporting the frequency specificity of this effect, relative to healthy controls, patients with myalgic encephalomyelitis/chronic fatigue syndrome (ME/CFS) exhibit reduced SWA in the SO frequency range and increased SWA in the delta frequency range, particularly over prefrontal cortex (Le Bon et al., 2012). This is notable, because central inflammation is a core component of ME/CFS (VanElzakker et al., 2019) and this frequency effect on SWA is highly consistent with the reported effects of A β (Mander et al., 2015; Winer et al., 2019). Regardless of the mechanism or the direction of causality, what is clear is that A β burden is associated with local disruptions in frontal slow waves, even prior to MCI diagnosis both cross-sectionally (Mander et al., 2015; Varga et al., 2016b; Winer et al., 2019) and longitudinally (Winer et al., 2020 in press). Some evidence also indicates that this effect of A β burden on local sleep may be specific to slow wave expression, as it does not appear to be associated with the REM sleep and spindle deficits observed in MCI and AD (Brazete et al., 2013, 2016;

Mander et al., 2015; Brayet et al., 2016; Gorgoni et al., 2016; Winer et al., 2019).

Tau Pathology and Local Sleep (T)

Habitual poor sleep and chronic sleep deprivation are associated with increased tau burden, tau hyperphosphorylation, and the spread of tau pathology across brainstem and hippocampal circuits, resulting in degeneration and memory impairment (Rothman et al., 2013; Di Meco et al., 2014; Ju et al., 2017; Zhu et al., 2018; Holth et al., 2019). Similar to the effects of sleep deprivation on A β , one recent finding indicated that ISF and CSF tau levels in rodents and humans also increase during wakefulness and decrease during sleep (Holth et al., 2019). Sleep deprivation exaggerated this wake-related increase, similar to its effects on A β , and chronic sleep deprivation increased the spread of tau pathology from the hippocampus to the locus coeruleus in the brainstem (Holth et al., 2019), a norepinephrine-expressing brain region known to accumulate tau pathology in the earliest stages of AD (Grudzien et al., 2007; Braak et al., 2011; Ehrenberg et al., 2017).

In addition to sleep deficits impacting tau pathology, more recent work shows that tau burden may be related to a number of distinct global and local sleep deficits observed in MCI and AD, each of which likely depends on the location of tau accumulation. Alongside the initial deposition of tau in the medial temporal lobe (MTL), tau also deposits in the brainstem, midbrain, and hypothalamus, preferentially within many sleep and wake-promoting nuclei (Oh et al., 2019), and this may decrease the consolidation of sleep and wakefulness which are regulated by these circuits (Lim et al., 2013a, 2014). Within the MTL, tau burden disrupts the GABA-regulated temporal synchrony of hippocampal ripple events during sleep and quiet wakefulness in a rodent model of tauopathy (Witton et al., 2016). Another rodent model of tauopathy identified the evolution of changes in sleep physiology, such that increases in NREM delta power and REM theta power in P301S tau transgenic mice relative to wild type mice at 6 and 9 months of age was followed by a decrease in NREM delta and REM theta power at 11 months (Holth et al., 2017). This finding indicated that, similar to the effect of MCI and AD on hippocampus activation, there may be a u-shaped curve in the expression of NREM and REM sleep oscillations as tau burden increases in the cortex.

In human studies, sleep spindle density and duration, but not SWA, are negatively associated with CSF-measured total and hyperphosphorylated tau levels in cognitively asymptomatic older adults (Kam et al., 2019). A similar finding was observed with slow wave-sleep spindle coupling strength, over frontal and parietal derivations, in relation to MTL tau burden in cognitively asymptomatic older adults (Figures 1B,D) (Winer et al., 2019). Similar to (Kam et al., 2019), this study did not find an association between SWA and MTL tau burden. In contrast, other reports showed that cortical tau was associated with a global loss of SWA across frequencies and prolonged duration of slow wave hyperpolarized down states (Menkes-Caspi et al., 2015; Lucey et al., 2019). However, these studies included patients with MCI (Lucey et al., 2019), and animal models that deposit tau in the cortex (Menkes-Caspi et al., 2015),

which is not typical of early preclinical AD stages (Braak and Braak, 1996). It is possible that when tau remains confined to MTL and brainstem regions in early stages, SWA is not affected, while sleep features that depend on these structures are affected. Once tau begins to spread cortically, global deficits in SWA may begin to be apparent (Figures 1A,D), drowning out the hyperexcitability effects of A β (Busche et al., 2019). This hypothesis, which remains to be tested, would indicate that global and local sleep deficits due to AD pathophysiology are not static across disease stages, but instead evolve across disease stages as distinct AD pathophysiological features build up and spread throughout various cortical and subcortical brain circuits. If true, this would mean that a comprehensive picture of global and local sleep deficits may offer unique insight into how AD pathophysiology is progressing in a given individual, even in preclinical stages. Whether these tau effects on sleep are prior to or following tau-related neurodegeneration remain unclear, and should be distinguished in future studies.

Neurodegeneration and Local Sleep (N)

Similar to tau pathological burden, the influence of neurodegeneration on global and local sleep likely depends on the location of neurodegeneration. Moreover, the distinction between degeneration due to age-related processes and dementia is not always clear. In the context of global sleep, loss of galanin-expressing inhibitory neurons in the intermediate nucleus of the hypothalamus, a potential human homolog to the ventral lateral preoptic nucleus which regulates sleep maintenance, is associated with sleep fragmentation (Lim et al., 2014). In patients with AD, the loss of galanin neurons and sleep fragmentation were both more extreme, but the overall relationship remained unchanged. Hence, it is not clear whether galanin neuron degeneration was due to an AD-specific process that began early in preclinical stages, or whether it was primarily age-related. However, there was also a trend for neurofibrillary tau tangles to be more concentrated in those with greater sleep fragmentation, indicating that galanin neurodegeneration may be driven by tau pathological burden in the hypothalamus (Lim et al., 2014). New evidence may help begin to distinguish between the role of tau and tau-related neurodegeneration on sleep expression. A recent study conducted comprehensive histopathological analysis of tau inclusion, neurotransmitter synthesis, and neuronal loss in a series of brainstem wake-promoting nuclei in healthy controls and patients with a variety of tauopathies, including AD, corticobasal degeneration (CBD), and progressive supranuclear palsy (PSP) (Oh et al., 2019). Increased tau inclusion and a decrease in neurons synthesizing neurotransmitters was apparent in all tauopathies relative to controls. However, it was only in AD that widespread neuronal loss was observed. Given these findings, it is therefore possible that distinct sleep phenotypes observed in distinct tauopathies is driven not by tau pathology, *per se*, but by the neurodegenerative effects of that pathology. For example, an extreme form of insomnia characterized by the inability to nap despite a profoundly shortened nighttime sleep duration (2–4 h per night) is consistently seen in PSP patients (Gagnon et al., 2008; Walsh et al., 2017), while sleep/wake instability is more typical in AD patients (Gagnon et al., 2008).

This makes sense if widespread neuronal loss in wake-promoting brainstem regions is observed in AD and not in PSP. The reason why tau-related neurodegeneration in the brainstem differs by tauopathy remains unclear, though one possible mechanism regards potential differences in interactions between central inflammation and tau pathology (Leyns and Holtzman, 2017).

Less is clear about the role of neurodegenerative processes in the expression of local sleep. While it has been shown that frontal atrophy is consistently associated with slow wave deficits and slow wave-sleep spindle coupling deficits (Mander, 2013; Varga et al., 2016a; Helfrich et al., 2018), and hippocampal atrophy and frontal white matter degeneration are associated with sleep spindle deficits (Fogel et al., 2017; Mander et al., 2017b), it is unclear if these are secondary to an AD process or generally related to aging. However, recent findings have linked deficits in subjective sleep duration and quality to cross sectional and longitudinal volumetric differences in signature cortical amnesic MCI and AD regions as well as overall ventricular enlargement (Lo et al., 2014; Spira et al., 2016; Alperin et al., 2019), though the directionality of these relationships remains unclear.

In terms of loss of EEG desynchrony during REM sleep in MCI and AD, a double blind placebo controlled clinical trial administering donepezil, an anticholinesterase inhibitor, in patients with AD showed that donepezil treatment for 6 months reduced REM sleep EEG slowing and increased REM sleep duration (Dos Santos Moraes et al., 2006). This indicates that REM sleep EEG slowing may be due to progressive loss of cholinergic activity in AD (Figures 1C,D). This is consistent with the fact that EEG desynchrony during REM sleep is critically dependent on brainstem and basal forebrain cholinergic inputs (Peever and Fuller, 2016; Scammell et al., 2017), and that basal forebrain cholinergic projections are selectively degenerated early in the AD pathophysiological process (Mufson et al., 1988; Mesulam et al., 2004; Oh et al., 2019). These findings support the hypothesis that REM sleep EEG slowing may be related to tau pathological burden, but only because of the tau-dependent degeneration of the cholinergic system (Oh et al., 2019). However, these hypotheses remain to be directly tested. Further, whether global and local REM sleep deficits are a biomarker of or directly related to cognitive deficits in AD is unknown.

Hence, new emerging evidence indicates that signature pathological features of AD (i.e., A β , tau, and neurodegeneration) are all associated with distinct effects on global and local sleep expression, and that the sleep deficits observed in AD are likely dependent on the location and relative severity of each of these processes and the manner in which they synergistically interact. However, much less is known about the direct and indirect roles these local sleep deficits may play in the cognitive decline associated with AD pathophysiology.

LOCAL SLEEP AND AD-RELATED MEMORY IMPAIRMENT

Critical components of a functioning MTL memory system include a network linking the entorhinal cortex (EC) layer II neurons to the dentate gyrus (DG) and cornu ammonis 3 (CA3)

hippocampal subfields by way of the perforant path (Hyman et al., 1986; Witter, 2007). This network, which supports successful separation between similar yet distinct memories—i.e., pattern separation (Yassa et al., 2011b)—is highly sensitive to age and sleep loss, and is among the earliest and most dramatically affected in AD (Jagust, 2013; Leal and Yassa, 2013). Indeed, degeneration of this MTL circuit is a critical preclinical structural biomarker of MCI and AD (Desikan et al., 2012; Holland et al., 2012). Targeted studies have identified diminished pattern separation ability both in aging (Wilson et al., 2006; Stark et al., 2010, 2013; Yassa et al., 2011a,b) and following sleep deprivation (Saletin et al., 2016), and impaired pattern separation is even more dramatic in patients with amnesic MCI (Bakker et al., 2012, 2015). In aging and MCI, these deficits are associated with DG/CA3 region hyperactivity (Yassa et al., 2011a,b), EC thinning (Reagh et al., 2018), and perforant path degeneration (Yassa et al., 2010a, 2011b). Levetiracetam (LEV) is an anti-epileptic drug that modulates synaptic excitability by binding to synaptic vesicle protein SV2A, thus impacting action potential-dependent neurotransmitter release (Lynch et al., 2004). LEV administration in aMCI patients significantly reduces hyperactivity and enhances object pattern separation performance relative to placebo (Bakker et al., 2012, 2015), supporting the notion that this hyperactivity is dysfunctional rather than compensatory. Of note, in addition to its impact on functional hyperactivity and cognition during wakefulness, LEV has also been shown to consolidate SWS and increase its duration in healthy adults, suggesting that it may also influence SWS-related memory processing as well (Cicolin et al., 2006).

Beyond the MTL, cortical network dysfunction and impaired functional connectivity have been observed in the context of AD pathophysiology. In particular, breakdown of connectivity and disinhibition of activity within the default mode network (DMN) is associated with both hippocampus hyperactivity and cognitive impairment in individuals with AD pathologies (Pihlajamaki et al., 2008; Jacobs et al., 2013; Pereira et al., 2019). Disruptions in DMN functional connectivity are also observed following sleep deprivation (Gujar et al., 2010) and are associated with the extent of daytime sleepiness (Ward et al., 2013). This may be due, in part, to a deprivation or disruption of infraslow EEG oscillations during SWS that organize DMN expression supporting sleep-dependent memory consolidation (Picchioni et al., 2011). Together, these findings implicate widespread circuit dysfunction within frontal, parietal, and MTL networks, which are associated with memory impairments even prior to AD-related neurodegeneration. Further, dysfunction within this circuit has also been associated with sleep disturbance, including in disorders such as sleep apnea and attention deficit hyperactivity disorder (Gujar et al., 2010; Ward et al., 2013; Khazaie et al., 2017; Tashjian et al., 2017), indicating that sleep disturbance may also be related to the effects of AD pathophysiology on cortical brain function.

The role of AD pathology in this memory-related circuit dysfunction is still being investigated. However, it appears that MTL tau burden, microglial dysfunction, and related synaptic loss are more strongly associated with memory impairment than A β burden (Jack et al., 2010, 2013; Scholl et al., 2016; Spangenberg

et al., 2016; Pereira et al., 2019). This has led to the view that tau pathology is more directly linked with cognitive impairments in AD than A β , though the truth is likely more complicated (Bloom, 2014; Maass et al., 2018; Sperling et al., 2019). For example, while tau is more likely to deposit early in the MTL, the tau-related memory impairments observed may be more typical of age-related memory decline (Maass et al., 2018). Once tau spreads beyond the MTL, which may depend on the presence of A β (Bloom, 2014; Tosun et al., 2017; Sperling et al., 2019), AD-related cognitive decline may begin to emerge and progress (Tosun et al., 2017; Sperling et al., 2019). Regardless, the pathophysiology of the long-term trend in tau-related cognitive decline may be distinct from the memory impairments due to hippocampus hyperexcitability.

What role, if any, that local sleep dysfunction plays in AD-related memory impairment remains unclear, though there are multiple theoretical ways in which local sleep could interact with AD pathologies to disrupt a variety of memory functions. The state of the science supporting each of these possibilities is reviewed below.


Direct Effects

An established literature demonstrates a clear relationship between local sleep features and multiple forms of memory encoding and consolidation (Fischer et al., 2002; Walker et al., 2002; Marshall et al., 2004, 2006; Spencer et al., 2006, 2007; Mander et al., 2011, 2013, 2014, 2015, 2017a,b; Antonenko et al., 2013; Fogel et al., 2013, 2017; Westerberg et al., 2015; Lustenberger et al., 2016; Papalambros et al., 2017). Three, non-mutually exclusive core theories of sleep-dependent memory consolidation implicate specific synaptic and systemic processes that mechanistically support the role of sleep in the long-term retention of procedural and episodic memories. There is evidence that many of the neurobiological mechanisms supporting these process may be directly disrupted by AD pathophysiology.

First, one theory posits that the coordinated expression of cortical slow oscillations (0.5–1 Hz), cortico-thalamic sleep spindles (12–16 Hz), and hippocampal ripples (140–220 Hz) actively supports memory consolidation by triggering memory replay during SWS to transform memory traces to be more cortically-dependent (**Figure 3A**) (Steriade, 2006; Diekelmann and Born, 2010; Abel et al., 2013; Staresina et al., 2015; Mander et al., 2017a; Helfrich et al., 2018). This framework has been supported by studies demonstrating induction of LTP-related plasticity, coordinated replay of neuronal ensembles, and coordinated hippocampal-neocortical information transfer during this tripartite coupling of slow wave, sleep spindle, and ripple events (Rosanova and Ulrich, 2005; Wierzynski et al., 2009; Chauvette et al., 2012; Abel et al., 2013; Latchoumane et al., 2017; Helfrich et al., 2019). More specifically, it has been shown that the hyperpolarized downstate of the cortical slow oscillation induces the expression of sleep spindles in the thalamic reticular nucleus which are preferentially expressed during the rising phase toward the depolarizing upstate of the slow oscillation (Mak-McCully et al., 2017). In addition, hippocampal ripple expression is maximal during the excitatory troughs of sleep spindles (Staresina et al., 2015; Latchoumane et al., 2017). Recent

closed-loop optogenetic evidence in rodents has demonstrated that this precise phase-relationship of coupling between sleep spindles with slow oscillations is essential for successful hippocampus-dependent memory consolidation (Latchoumane et al., 2017). Induction of sleep spindle expression in thalamic reticular neurons during the rising phase of the slow oscillation up-state increases phase locking of hippocampal ripples to spindle troughs and enhances memory consolidation, while suppression of sleep spindle expression during this SO phase impairs memory consolidation. Conversely, either inducing or suppressing sleep spindle expression outside of this SO rising up phase has no effect on memory consolidation. Prior studies offer mechanistic support for this effect of SO-sleep spindle coupling on memory consolidation. Sleep spindles have a local effect on neuroplasticity, but this depends on its coupling with slow oscillations. When sleep spindles are appropriately phase locked to slow oscillations local evidence of long term potentiation (LTP) is apparent, and when sleep spindles are expressed without slow oscillations long term depression (LTD) is instead observed (Rosanova and Ulrich, 2005). Another recent optogenetic study also demonstrated a causal role for SO events (<1 Hz slow waves) and not faster frequency delta waves (1–4 Hz slow waves) in memory retention (Kim et al., 2019). Hence, optogenetic manipulation of both SO and sleep spindle events directly impacts long-term memory retention, supporting their causal role in sleep-dependent memory consolidation. This model is further supported by studies that mechanistically enhance slow waves, sleep spindles, and slow wave-sleep spindle coupling through external electrical and auditory stimulation methods in humans, resulting in improved procedural and episodic memory, even in older adults and patients with MCI (Marshall et al., 2004, 2006; Westerberg et al., 2015; Lustenberger et al., 2016; Ladenbauer et al., 2017; Papalambros et al., 2017, 2019).

Theoretical depictions of this form of sleep-dependent memory describe a process by which hippocampal-neocortical neuronal ensembles which are primed by learning during prior wakefulness are reactivated during slow wave sleep in a temporally precise manner. This is supported by classic studies showing replay of coordinated place cell activity during slow wave sleep, and specifically during hippocampal ripples (Wilson and McNaughton, 1994; Ji and Wilson, 2007; Davidson et al., 2009). It has been hypothesized that this replay provides a window for information transfer from hippocampal nodes to neocortical nodes facilitating neuroplasticity in support of memory transformation and long-term memory retention (Buzsaki, 1998). This supposition has been supported by studies in rodents and intracranial EEG studies in humans describing evidence of information transfer from hippocampus to cortex during triple coupling of SOs, sleep spindles, and ripple events (Olcese et al., 2018; Helfrich et al., 2019), with this information transfer being particularly prominent in neuronal ensembles modulated by a memory task in prior waking periods (Olcese et al., 2018). Further support for the causal role of replay in memory consolidation is demonstrated by a series of studies implementing targeted memory reactivation (TMR) methods to trigger the replay of memories encoded prior to sleep during SWS. This approach implements pairing learning with external



sensory stimulation, such as pairing odor cues with a memory task (Rasch et al., 2007) and auditory cues with individual memory trials (Rudoy et al., 2009; Antony et al., 2012; Van Dongen et al., 2012; Batterink et al., 2016; Berkers et al., 2018). These sensory cues are then replayed during sleep with memory performance being assessed in sensory cued versus un-cued trials in subsequent waking periods. These studies have shown that sensory cueing specifically during SWS enhances slow wave-sleep spindle coupling and memory performance (Rasch et al., 2007; Rudoy et al., 2009; Antony et al., 2012; Van Dongen et al., 2012; Feld and Diekelmann, 2015; Batterink et al., 2016; Berkers et al., 2018; Bar et al., 2020), with the efficacy of this effect depending on the phase of the slow wave during sensory cueing (Batterink et al., 2016; Goldi et al., 2019). Hence, these findings implicate a mechanism by which triple coupling of SO, sleep spindle, and ripple events coordinate replay of relevant neuronal ensembles supporting targeted neuroplasticity in facilitation of memory consolidation.

Cortical slow oscillations are diminished in the presence of both A β and tau pathology in the cortex (Mander et al., 2015;

Lucey et al., 2019), and this effect is independent of gray matter atrophy (Mander et al., 2015). Furthermore, this effect of A β on SWA was correlated with retrieval-related hippocampus hyperactivation, which was negatively associated with sleep-dependent memory retention (Mander et al., 2015). Further supporting the importance of the <1 Hz SO, doubling the frequency of the SO from 0.6 to 1.2 Hz in APP transgenic mice resulted in reduced dendritic spine density, likely directly impacting memory functions (Kastanenka et al., 2019). In addition, MTL tau burden has been shown to disrupt hippocampal ripple expression and slow wave-sleep spindle coupling during NREM sleep (Witton et al., 2016; Winer et al., 2019). Hence, current evidence indicates that AD pathologies disrupt the expression and coupling of all three of the core sleep oscillations underlying sleep-dependent memory consolidation (Figure 3A). Beyond their direct effects on local sleep expression, AD pathophysiology could also disrupt cortical network structure and function in a way that diminishes the efficacy of local sleep expression to support consolidation across memory systems, as observed in aging (Mander et al., 2017a).

Second, the synaptic homeostasis hypothesis posits that slow waves act to globally downscale synapses to manage the growing energy demands of the brain following continuous learning and thus wide-spread synaptic potentiation (**Figure 3B**) (Tononi and Cirelli, 2014). In theory, memory is consolidated and enhanced by this process as it acts to enhance the signal to noise of neuronal ensembles related to a given memory trace. A series of studies across model systems support key tenants of this hypothesis. Synaptic potentiation is saturated following extended wakefulness, resulting in partial occlusion of long term potentiation (LTP) induction, which is refreshed following SWS (Vyazovskiy et al., 2011). This is mirrored in some human studies, with slow wave stimulation and sleep spindle expression both predicting restoration of memory encoding ability following sleep (Mander et al., 2011; Antonenko et al., 2013). Other studies have shown that memory tasks targeting localized brain networks result in local increases in SWA in subsequent sleep periods, with the intensity of SWA predicting the degree of motor learning (Huber et al., 2004, 2006). A recent optogenetic paper has showed that while SO events support memory retention, delta waves may support forgetting (Kim et al., 2019). This finding may support the presence of both hypothesized mechanisms in SWS, with an active process enhancing synaptic strength of a small number of neuronal ensembles directly relevant to triggered replay of specific prior waking experiences, while another globally downscales synapses to maintain energy homeostasis and reduce neural noise.

This latter mechanism may also be influenced by AD pathology (**Figure 3B**). While A β decreased SO events, it also increased delta waves (Mander et al., 2015). It is possible this shift in SO-delta wave balance may shift memory systems toward greater forgetting and lower memory retention (Kim et al., 2019). Indeed, while SO events were positively associated with sleep-dependent memory consolidation, delta events were negatively associated with sleep-dependent memory consolidation in older adults with A β pathology (Mander et al., 2015). In addition to this potential “hyper downscaling” in the presence of A β , cortical tau may impoverish the downscaling process altogether, indirectly impacting the ability of the brain to encode novel experiences in subsequent waking periods due to incomplete depotentiation of synapses (**Figure 3B**). Therefore, AD pathologies may either exaggerate a downscaling process or disrupt it altogether, impairing sleep-dependent memory consolidation and/or sleep-related memory encoding ability.

Lastly, another model concerns coordinated sequential synaptic remodeling across NREM-REM cycles (Yang et al., 2014; Miyawaki and Diba, 2016; Li et al., 2017; Mizuseki and Miyawaki, 2017). This theory is supported by animal work showing that NREM sleep can trigger a limited formation of new synapses following a learning experience in a dendritic branch-specific manner (Yang et al., 2014). REM sleep then eliminates many of these, but stabilizes the synapses not eliminated (Li et al., 2017). AD could impact this process, too, by disrupting the stability of NREM and REM sleep, shortening REM duration, and/or disrupting the EEG desynchrony characteristic of REM sleep. This would likely result in fewer new synapses formed following learning, all of which would be more labile and vulnerable to

decay (**Figure 3C**). As with the other sleep-dependent memory models, this possibility has not been directly tested.

Each of the theoretical mechanisms proposed to support sleep-dependent memory encoding and consolidation are likely negatively influenced by AD pathophysiology (**Figure 3**), which would contribute to the memory impairments observed in AD. However, the impact of local sleep deficits on memory in AD is not necessarily limited to the effects of AD pathophysiology on sleep-dependent memory mechanisms.

Indirect Effects

There is also evidence that AD pathology interacts with sleep deficits to influence cognition. For example, the interaction between A β and sleep efficiency predicts cognitive deficits on cognitive tasks, such that poor sleep quality results in impaired memory to a greater extent when an individual has AD pathology in their brain (Molano et al., 2017). Moreover, the influence of APOE genotype—the strongest genetic risk factor for late onset, sporadic AD (O'Donoghue et al., 2018)—on cognition depended on the quality of sleep, with APOE genotype having minimal impact in those with high quality sleep and substantial impact on those with low quality sleep (Lim et al., 2013b). This indicates that sleep may be related to a neural reserve factor making the brain more or less vulnerable to the presence of AD pathology or risk factors. Indeed, there is a literature that indicates that deficits in sleep quantity and quality result in cortical brain atrophy (Spira et al., 2016; Alperin et al., 2019) and overall ventricular enlargement even over the course of 2 years in healthy older adults (Lo et al., 2014), which would likely diminish neural reserves to compensate for AD pathophysiology.

Another possibility is that sleep may be related to cognitive impairment independently of sleep-dependent memory processes through its direct effects on AD pathophysiology. Growing evidence indicates the bidirectional relationship between slow waves and A β and tau pathology (Zhang et al., 2005; Xie et al., 2013; Mander et al., 2015, 2016; Menkes-Caspi et al., 2015; Varga et al., 2016b; Ju et al., 2017; Kastanenka et al., 2017, 2019; Kent et al., 2018; Fultz et al., 2019; Lucey et al., 2019). More specifically, slow wave deficits and sleep/wake instability appear to foster greater production and lower clearance of both A β and tau pathology (Zhang et al., 2005; Mander et al., 2015; Varga et al., 2016b; Ju et al., 2017; Kastanenka et al., 2017, 2019; Kent et al., 2018; Musiek et al., 2018; Lucey et al., 2019), which may then exacerbate central inflammation and MTL-related neurodegeneration, thus accelerating memory decline (Jack et al., 2010, 2013, 2019; Yassa et al., 2010b, 2011b; Desikan et al., 2012; Holland et al., 2012; Jagust, 2013; Leal and Yassa, 2013; Scholl et al., 2016; Spangenberg et al., 2016; Leal et al., 2017; Spangenberg and Green, 2017; Reagh et al., 2018; Pereira et al., 2019).

Hence, in addition to the disruption of local sleep processes related to sleep-dependent memory encoding and consolidation, sleep disruption may further accelerate AD pathophysiology or make the brain more vulnerable to it. Each of these could contribute to AD-related cognitive decline, and could do so at distinct stages. For example, sleep-dependent memory deficits may emerge as soon as local sleep deficits emerge in

the initial stages of AD pathogenesis (Mander et al., 2015), while deficits associated with neural reserve or AD-related neurodegeneration may emerge later once AD pathological burden has reached a tipping point or degeneration has progressed sufficiently to result in clinical symptoms (Westerberg et al., 2012; Lim et al., 2013b; Liguori et al., 2014). This would indicate that the role of sleep in AD-related cognitive decline may also be multifaceted and evolve across AD pathophysiological stages.

CONCLUSION AND FUTURE DIRECTIONS

Evidence continues to mount in support of the relationship between local sleep processes and AD pathophysiology. Recent studies have established that A β , tau, and neurodegeneration, defining pathological components of AD, all result in specific deficits in local and global sleep expression that depend on the underlying brain regions impacted (**Figure 1**). Pathological accumulation and degeneration within brainstem and hypothalamic sleep/wake regulatory centers fosters sleep state instability, pathology and degeneration in the MTL impacts ripple and spindle expression, cortical deposition and degeneration alters slow wave expression, and AD-related cholinergic degeneration likely disrupts REM sleep expression. These deficits are likely not static, but evolve as AD pathophysiology progresses through brain networks and across disease stages. For example, the slow wave deficits due to A β may differ from those observed with tau (**Figure 1A**). In some cases, these deficits may actively contribute to the progression of the disease, especially as they relate to the facilitation of A β and tau accumulation and neurodegeneration of circuits vulnerable to these pathologies (e.g., disruption of glymphatic clearance of AD pathologies). There is some evidence that these associations between AD pathophysiology and local sleep expression may be relevant for the memory impairments observed in the context of AD, though mechanisms of these relationships remain hypothetical.

The current state of the science indicates that there are more questions that remain than answers, both at mechanistic and therapeutic levels. At the mechanistic level, a clearer understanding of how local sleep is influenced by AD pathophysiology at each stage of the disease process is warranted. This means conducting larger scale studies that can implement the ATN framework to understand how the presence and absence and relative burdens of different AD pathologies at different AD stages may impact local sleep. This will also require examining other mechanisms, including metabolic dysfunction, chronic inflammation, and HPA-axis dysregulation, all of which contribute to AD pathophysiology and are related to sleep disturbance (Heppner et al., 2015; Spangenberg and Green, 2017; Caruso et al., 2018; Carroll and Macauley, 2019; Irwin and Vitiello, 2019; Chappel-Farley et al., 2020). Chronic inflammation, in particular, is a promising target for future research, because of its known independent interactions with

sleep expression and multiple aspects of AD pathophysiology, as well as the paucity of data addressing its links with age-related deficits in local sleep expression. Further, targeted studies examining how the flow of CSF in the brain that is phase-locked to SWA (Fultz et al., 2019) may change with age and in relation to endogenously generated pathological features of AD will critically advance our understanding of the importance of the glymphatic system in AD pathogenesis. A greater understanding of the intersection between sleep disorders and local sleep, and their relationship with AD pathophysiological progression is needed, to better understand how various forms of sleep disturbance impact AD risk. Studies examining how local sleep differs across forms of dementia or kinds of AD will be informative both in terms of understanding mechanisms and the utility of local sleep to aid differential diagnosis. Longitudinal studies incorporating local sleep measures alongside AD biomarkers are of great need to determine the time course of sleep changes in relation to AD pathophysiology. Lastly, a comprehensive examination of how local sleep disruption is related to various forms of impairment in sleep-dependent and sleep-independent cognitive processes in relation to AD pathophysiology will help unpack the role(s) sleep plays in AD-related cognitive decline across disease severity. This will necessarily include examination of the influence of MCI, AD, and AD pathophysiology on ripple expression and memory replay, as well as the efficacy of targeted memory reactivation paradigms to facilitate memory transformation and consolidation.

Alongside these efforts, an emphasis on large-scale studies implementing sleep interventions must be considered. At present, it is not clear whether local sleep deficits are a biomarker of AD progression, or are active contributors to AD pathophysiology or related cognitive decline. The utility of sleep interventions, both targeting global sleep and local sleep, to address AD pathophysiological progression remains entirely unknown. While there are some data that suggest that treating sleep disorders may impact AD biomarkers (Osorio et al., 2015; Liguori et al., 2017a,b), large scale studies are lacking. Further, while a few reports show that enhancing local sleep in older adults and patients with MCI will improve overnight memory consolidation (Ladenbauer et al., 2017; Papalambros et al., 2019), the efficacy of this type of intervention, even for more than one night, is unknown. As important is an examination of the impact of AD disease stage on the efficacy of sleep interventions to impact AD pathophysiology and related cognitive impairments. Will sleep interventions help preclinical, MCI, or AD patients equally, or will sleep interventions have diminishing returns as the disease progresses, or affect distinct outcomes? This will be a critical question to answer, as it will help guide expectations related to disease management and identify the appropriate therapeutic targets based on the disease stage.

While much remains unknown, it is clear that local sleep processes are intricately intertwined with AD pathophysiology, likely in a bidirectional manner and through multiple mechanisms. These findings also open up the possibility that local sleep may offer a unique and sensitive window into

circuit level dysfunction across different forms of dementia, aiding early efforts in differential diagnosis of dementia risk. As different forms of dementia target distinct neural networks, the profile of deficits in local and global sleep appear to differ as well (Petit et al., 2004; Gagnon et al., 2008; Oh et al., 2019). Ultimately, the importance of local sleep for the optimal functioning of the brain means that local sleep disruption will likely be a crucial factor in the progression of all dementias, whether as a biomarker of a diseased brain, a contributor to disease pathogenesis, or an independent factor influencing the brain's resiliency or vulnerability to dementia. Uncovering the specific roles of local sleep in dementias such as AD will thus offer new opportunities to target interventions to offer symptomatic relief, impact disease progression and risk, and slow affiliated cognitive decline.

REFERENCES

- Abel, T., Havekes, R., Saletin, J. M., and Walker, M. P. (2013). Sleep, plasticity and memory from molecules to whole-brain networks. *Curr. Biol.* 23, R774–R788.
- Alperin, N., Wiltshire, J., Lee, S. H., Ramos, A. R., Hernandez-Cardenache, R., Rundek, T., et al. (2019). Effect of sleep quality on amnesic mild cognitive impairment vulnerable brain regions in cognitively normal elderly individuals. *Sleep* 42:zsy254.
- Antonenko, D., Diekelmann, S., Olsen, C., Born, J., and Molle, M. (2013). Napping to renew learning capacity: enhanced encoding after stimulation of sleep slow oscillations. *Eur. J. Neurosci.* 37, 1142–1151. doi: 10.1111/ejn.12118
- Antony, J. W., Gobel, E. W., O'hare, J. K., Reber, P. J., and Paller, K. A. (2012). Cued memory reactivation during sleep influences skill learning. *Nat. Neurosci.* 15, 1114–1116. doi: 10.1038/nn.3152
- Bagyinszky, E., Youn, Y. C., An, S. S., and Kim, S. (2014). The genetics of Alzheimer's disease. *Clin. Interv. Aging* 9, 535–551.
- Bakker, A., Albert, M. S., Krauss, G., Speck, C. L., and Gallagher, M. (2015). Response of the medial temporal lobe network in amnesic mild cognitive impairment to therapeutic intervention assessed by fMRI and memory task performance. *Neuroimage Clin.* 7, 688–698. doi: 10.1016/j.nicl.2015.02.009
- Bakker, A., Krauss, G. L., Albert, M. S., Speck, C. L., Jones, L. R., Stark, C. E., et al. (2012). Reduction of hippocampal hyperactivity improves cognition in amnesic mild cognitive impairment. *Neuron* 74, 467–474. doi: 10.1016/j.neuron.2012.03.023
- Bar, E., Marmelshtein, A., Arzi, A., Perl, O., Livne, E., Hizmi, E., et al. (2020). Local targeted memory reactivation in human sleep. *Curr. Biol.* 30, 1435–1446.e5.
- Batterink, L. J., Creery, J. D., and Paller, K. A. (2016). Phase of spontaneous slow oscillations during sleep influences memory-related processing of auditory cues. *J. Neurosci.* 36, 1401–1409. doi: 10.1523/jneurosci.3175-15.2016
- Bellesi, M., De Vivo, L., Chini, M., Gilli, F., Tononi, G., and Cirelli, C. (2017). Sleep Loss Promotes Astrocytic Phagocytosis and Microglial Activation in Mouse Cerebral Cortex. *J. Neurosci.* 37, 5263–5273. doi: 10.1523/jneurosci.3981-16.2017
- Berkers, R., Ekman, M., Van Dongen, E. V., Takashima, A., Barth, M., Paller, K. A., et al. (2018). Cued reactivation during slow-wave sleep induces brain connectivity changes related to memory stabilization. *Sci. Rep.* 8:16958.
- Besedovsky, L., Lange, T., and Haack, M. (2019). The sleep-immune crosstalk in health and disease. *Physiol. Rev.* 99, 1325–1380. doi: 10.1152/physrev.00010.2018
- Bloom, G. S. (2014). Amyloid-beta and Tau the trigger and bullet in alzheimer disease pathogenesis. *JAMA Neurol.* 71, 505–508. doi: 10.1001/jamaneurol.2013.5847
- Borbely, A. A. (1982). A two process model of sleep regulation. *Hum. Neurobiol.* 1, 195–204.
- Borbely, A. A., Baumann, F., Brandeis, D., Strauch, I., and Lehmann, D. (1981). Sleep deprivation: effect on sleep stages and EEG power density in man. *Electroencephalogr. Clin. Neurophysiol.* 51, 483–495. doi: 10.1016/0013-4694(81)90225-x

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and has approved it for publication.

FUNDING

This work was supported by K01AG068353.

ACKNOWLEDGMENTS

I would like to acknowledge Joseph R. Winer, Ph.D., for his feedback on this review.

- Braak, H., and Braak, E. (1991). Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol.* 82, 239–259. doi: 10.1007/bf00308809
- Braak, H., and Braak, E. (1995). Staging of Alzheimer's disease-related neurofibrillary changes. *Neurobiol. Aging* 16, 271–278. doi: 10.1016/0197-4580(95)00021-6
- Braak, H., and Braak, E. (1996). Evolution of the neuropathology of Alzheimer's disease. *Acta Neurol. Scand.* 93, 3–12.
- Braak, H., Thal, D. R., Ghebremedhin, E., and Del Tredici, K. (2011). Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. *J. Neuropathol. Exp. Neurol.* 70, 960–969. doi: 10.1097/nen.0b013e318232a379
- Braun, A. R., Balkin, T. J., Wesensten, N. J., Carson, R. E., Varga, M., Baldwin, P., et al. (1997). Regional cerebral blood flow throughout the sleep-wake cycle - An (H₂O)-O-15 PET study. *Brain* 120, 1173–1197. doi: 10.1093/brain/120.7.1173
- Brayet, P., Petit, D., Frauscher, B., Gagnon, J. F., Gosselin, N., Gagnon, K., et al. (2016). Quantitative EEG of rapid-eye-movement sleep: a marker of amnesic mild cognitive impairment. *Clin. EEG Neurosci.* 47, 134–141. doi: 10.1177/1550059415603050
- Brazete, J. R., Gagnon, J. F., Postuma, R. B., Bertrand, J. A., Petit, D., and Montplaisir, J. (2016). Electroencephalogram slowing predicts neurodegeneration in rapid eye movement sleep behavior disorder. *Neurobiol. Aging* 37, 74–81. doi: 10.1016/j.neurobiolaging.2015.10.007
- Brazete, J. R., Montplaisir, J., Petit, D., Postuma, R. B., Bertrand, J. A., Marchand, D. G., et al. (2013). Electroencephalogram slowing in rapid eye movement sleep behavior disorder is associated with mild cognitive impairment. *Sleep Med.* 14, 1059–1063. doi: 10.1016/j.sleep.2013.06.013
- Bubu, O. M., Brannick, M., Mortimer, J., Umasabor-Bubu, O., Sebastiao, Y. V., Wen, Y., et al. (2017). Sleep, cognitive impairment and Alzheimer's disease: a systematic review and meta-analysis. *Sleep* 40:zsw032.
- Bubu, O. M., Pirraglia, E., Andrade, A. G., Sharma, R. A., Gimenez-Badia, S., Umasabor-Bubu, O. Q., et al. (2019). Obstructive sleep apnea and longitudinal Alzheimer's disease biomarker changes. *Sleep* 42:zsz048.
- Buchsbaum, M. S., Gillin, J. C., Wu, J., Hazlett, E., Sicotte, N., Dupont, R. M., et al. (1989). Regional cerebral glucose metabolic-rate in human sleep assessed by positron emission tomography. *Life Sci.* 45, 1349–1356. doi: 10.1016/0024-3205(89)90021-0
- Buckley, R. F., Mormino, E. C., Rabin, J. S., Hohman, T. J., Landau, S., Hanseeuw, B. J., et al. (2019). Sex differences in the association of global amyloid and regional tau deposition measured by positron emission tomography in clinically normal older adults. *JAMA Neurol.* 76, 542–551. doi: 10.1001/jamaneurol.2018.4693
- Buckner, R. L., Snyder, A. Z., Shannon, B. J., Larossa, G., Sachs, R., Fotenos, A. F., et al. (2005). Molecular, structural, and functional characterization of Alzheimer's disease: evidence for a relationship between default activity, amyloid, and memory. *J. Neurosci.* 25, 7709–7717. doi: 10.1523/jneurosci.2177-05.2005
- Busche, M. A., Kekus, M., Adelsberger, H., Noda, T., Forstl, H., Nelken, I., et al. (2015). Rescue of long-range circuit dysfunction in Alzheimer's disease models. *Nat. Neurosci.* 18, 1623–1630. doi: 10.1038/nn.4137

- Busche, M. A., Wegmann, S., Dujardin, S., Commins, C., Schiantarelli, J., Klickstein, N., et al. (2019). Tau impairs neural circuits, dominating amyloid-beta effects, in Alzheimer models *in vivo*. *Nat. Neurosci.* 22, 57–64. doi: 10.1038/s41593-018-0289-8
- Buzsaki, G. (1998). Memory consolidation during sleep: a neurophysiological perspective. *J. Sleep Res.* 7(Suppl. 1), 17–23. doi: 10.1046/j.1365-2869.7.s1.3.x
- Cagnin, A., Brooks, D. J., Kennedy, A. M., Gunn, R. N., Myers, R., Turkheimer, F. E., et al. (2001). In-vivo measurement of activated microglia in dementia. *Lancet* 358, 461–467. doi: 10.1016/s0140-6736(01)05625-2
- Carrier, J., Land, S., Buysse, D. J., Kupfer, D. J., and Monk, T. H. (2001). The effects of age and gender on sleep EEG power spectral density in the middle years of life (ages 20–60 years old). *Psychophysiology* 38, 232–242. doi: 10.1111/1469-8986.3820232
- Carroll, C. M., and Macauley, S. L. (2019). The interaction between sleep and metabolism in Alzheimer's disease: Cause or consequence of disease? *Front. Aging Neurosci.* 11:258. doi: 10.3389/fnagi.2019.00258
- Caruso, A., Nicoletti, F., Mango, D., Saidi, A., Orlando, R., and Scaccianoce, S. (2018). Stress as risk factor for Alzheimer's disease. *Pharmacol. Res.* 132, 130–134.
- Chappel-Farley, M. G., Lui, K. K., Dave, A., Chen, I. Y., and Mander, B. A. (2020). Candidate mechanisms linking insomnia disorder to Alzheimer's disease risk. *Curr. Opin. Behav. Sci.* 33, 92–98. doi: 10.1016/j.cobeha.2020.01.010
- Chauvette, S., Seigneur, J., and Timofeev, I. (2012). Sleep oscillations in the thalamocortical system induce long-term neuronal plasticity. *Neuron* 75, 1105–1113. doi: 10.1016/j.neuron.2012.08.034
- Chen, J. C., Espeland, M. A., Brunner, R. L., Lovato, L. C., Wallace, R. B., Leng, X. Y., et al. (2016). Sleep duration, cognitive decline, and dementia risk in older women. *Alzheimers Dement.* 12, 21–33. doi: 10.1016/j.jalz.2015.03.004
- Cicolin, A., Magliola, U., Giordano, A., Terreni, A., Bucca, C., and Mutani, R. (2006). Effects of levetiracetam on nocturnal sleep and daytime vigilance in healthy volunteers. *Epilepsia* 47, 82–85. doi: 10.1111/j.1528-1167.2006.00376.x
- Conte, F., Arzilli, C., Errico, B. M., Giganti, F., Iovino, D., and Ficca, G. (2014). Sleep Measures Expressing 'Functional Uncertainty' in Elderlies'. *Sleep Gerontol.* 60, 448–457. doi: 10.1159/000358083
- Cucchiara, F., Frumento, P., Banfi, T., Sesso, G., Di Galante, M., D'ascanio, P., et al. (2020). Electrophysiological features of sleep in children with Kir4.1 channel mutations and Autism-Epilepsy phenotype: a preliminary study. *Sleep* 43:zsz255.
- Darchia, N., Campbell, I. G., and Feinberg, I. (2003). Rapid eye movement density is reduced in the normal elderly. *Sleep* 26, 973–977. doi: 10.1093/sleep/26.8.973
- Davidson, T. J., Kloosterman, F., and Wilson, M. A. (2009). Hippocampal replay of extended experience. *Neuron* 63, 497–507. doi: 10.1016/j.neuron.2009.07.027
- De Gennaro, L., and Ferrara, M. (2003). Sleep spindles: an overview. *Sleep Med. Rev.* 7, 423–440. doi: 10.1053/smr.2002.0252
- De Gennaro, L., Gorgoni, M., Reda, F., Lauri, G., Truglia, I., Cordone, S., et al. (2017). The Fall of Sleep K-Complex in Alzheimer Disease. *Sci. Rep.* 7:39688.
- Desikan, R. S., Mcevoy, L. K., Thompson, W. K., Holland, D., Brewer, J. B., Aisen, P. S., et al. (2012). Amyloid-beta-associated clinical decline occurs only in the presence of elevated P-tau. *Arch. Neurol.* 69, 709–713.
- Di Meco, A., Joshi, Y. B., and Pratico, D. (2014). Sleep deprivation impairs memory, tau metabolism, and synaptic integrity of a mouse model of Alzheimer's disease with plaques and tangles. *Neurobiol. Aging* 35, 1813–1820. doi: 10.1016/j.neurobiolaging.2014.02.011
- Diekelmann, S., and Born, J. (2010). The memory function of sleep. *Nat. Rev. Neurosci.* 11, 114–126.
- Dijk, D. J., Beersma, D. G. M., and Van Den Hoofdakker, R. H. (1989). All night spectral analysis of EEG sleep in young adult and middle-aged subjects. *Neurobiol. Aging* 10, 677–682. doi: 10.1016/0197-4580(89)90004-3
- Dos Santos Moraes, W. A., Poyares, D. R., Guilleminault, C., Ramos, L. R., Ferreira Bertolucci, P. H., and Tufik, S. (2006). The effect of donepezil on sleep and REM sleep EEG in patients with Alzheimer disease: a double-blind placebo-controlled study. *Sleep* 29, 199–205. doi: 10.1093/sleep/29.2.199
- Dworak, M., Mccarley, R. W., Kim, T., Kalinchuk, A. V., and Basheer, R. (2010). Sleep and brain energy levels: ATP changes during sleep. *J. Neurosci.* 30, 9007–9016. doi: 10.1523/jneurosci.1423-10.2010
- Edison, P., Archer, H. A., Gerhard, A., Hinz, R., Pavese, N., Turkheimer, F. E., et al. (2008). Microglia, amyloid, and cognition in Alzheimer's disease: an [11C](R)PK11195-PET and [11C]PIB-PET study. *Neurobiol. Dis.* 32, 412–419. doi: 10.1016/j.nbd.2008.08.001
- Ehlers, C. L., and Kupfer, D. J. (1989). Effects of age on delta and rem-sleep parameters. *Electroencephalogr. Clin. Neurophysiol.* 72, 118–125. doi: 10.1016/0013-4694(89)90172-7
- Ehlers, C. L., and Kupfer, D. J. (1997). Slow-wave sleep: Do young adult men and women age differently? *J. Sleep Res.* 6, 211–215. doi: 10.1046/j.1365-2869.1997.00041.x
- Ehrenberg, A. J., Nguy, A. K., Theofilas, P., Dunlop, S., Suemoto, C. K., Di Lorenzo Alho, A. T., et al. (2017). Quantifying the accretion of hyperphosphorylated tau in the locus coeruleus and dorsal raphe nucleus: the pathological building blocks of early Alzheimer's disease. *Neuropathol. Appl. Neurobiol.* 43, 393–408. doi: 10.1111/nan.12387
- Feld, G. B., and Diekelmann, S. (2015). Sleep smart-optimizing sleep for declarative learning and memory. *Front. Psychol.* 6:622. doi: 10.3389/fpsyg.2015.00622
- Fellin, T., Halassa, M. M., Terunuma, M., Succol, F., Takano, H., Frank, M., et al. (2009). Endogenous nonneuronal modulators of synaptic transmission control cortical slow oscillations *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.* 106, 15037–15042. doi: 10.1073/pnas.0906419106
- Fischer, S., Hallschmid, M., Elsner, A. L., and Born, J. (2002). Sleep forms memory for finger skills. *Proc. Natl. Acad. Sci. U.S.A.* 99, 11987–11991. doi: 10.1073/pnas.182178199
- Floyd, J. A., Janisse, J. J., Jenuwine, E. S., and Ager, J. W. (2007). Changes in REM-sleep percentage over the adult lifespan. *Sleep* 30, 829–836. doi: 10.1093/sleep/30.7.829
- Fogel, S., Vien, C., Karni, A., Benali, H., Carrier, J., and Doyon, J. (2017). Sleep spindles: a physiological marker of age-related changes in gray matter in brain regions supporting motor skill memory consolidation. *Neurobiol. Aging* 49, 154–164. doi: 10.1016/j.neurobiolaging.2016.10.009
- Fogel, S. M., Albouy, G., Vien, C., Popovicci, R., King, B. R., Hoge, R., et al. (2013). fMRI and sleep correlates of the age-related impairment in motor memory consolidation. *Hum. Brain Mapp.* 35, 3625–3645. doi: 10.1002/hbm.22426
- Friedman, E. M., Hayney, M. S., Love, G. D., Urry, H. L., Rosenkranz, M. A., Davidson, R. J., et al. (2005). Social relationships, sleep quality, and interleukin-6 in aging women. *Proc. Natl. Acad. Sci. U.S.A.* 102, 18757–18762. doi: 10.1073/pnas.0509281102
- Fultz, N. E., Bonmassar, G., Setsompop, K., Stickgold, R. A., Rosen, B. R., Polimeni, J. R., et al. (2019). Coupled electrophysiological, hemodynamic, and cerebrospinal fluid oscillations in human sleep. *Science* 366, 628–631. doi: 10.1126/science.aax5440
- Gagnon, J.-F., Petit, D., Latreille, V., and Montplaisir, J. (2008). Neurobiology of sleep disturbances in neurodegenerative disorders. *Curr. Pharm. Des.* 14, 3430–3445. doi: 10.2174/138161208786549353
- Gillin, J. C., Duncan, W. C., Murphy, D. L., Post, R. M., Wehr, T. A., Goodwin, F. K., et al. (1981). Age-related-changes in sleep in depressed and normal subjects. *Psychiatry Res.* 4, 73–78. doi: 10.1016/0165-1781(81)90010-x
- Goldi, M., Van Poppel, E. A. M., Rasch, B., and Schreiner, T. (2019). Increased neuronal signatures of targeted memory reactivation during slow-wave up states. *Sci. Rep.* 9:2715.
- Gorgoni, M., Lauri, G., Truglia, I., Cordone, S., Sarasso, S., Scarpelli, S., et al. (2016). Parietal fast sleep spindle density decrease in Alzheimer's disease and amnesic mild cognitive impairment. *Neural Plast.* 2016:8376108.
- Gori, S., Ficca, G., Giganti, F., Di Nasso, I., Murri, L., and Salzarulo, P. (2004). Body movements during night sleep in healthy elderly subjects and their relationships with sleep stages. *Brain Res. Bull.* 63, 393–397. doi: 10.1016/j.brainresbull.2003.12.012
- Grudzien, A., Shaw, P., Weintraub, S., Bigio, E., Mash, D. C., and Mesulam, M. M. (2007). Locus coeruleus neurofibrillary degeneration in aging, mild cognitive impairment and early Alzheimer's disease. *Neurobiol. Aging* 28, 327–335. doi: 10.1016/j.neurobiolaging.2006.02.007
- Gujar, N., Yoo, S. S., Hu, P., and Walker, M. P. (2010). The unrested resting brain: sleep deprivation alters activity within the default-mode network. *J. Cogn. Neurosci.* 22, 1637–1648. doi: 10.1162/jocn.2009.21331
- Haack, M., Sanchez, E., and Mullington, J. M. (2007). Elevated inflammatory markers in response to prolonged sleep restriction are associated with increased pain experience in healthy volunteers. *Sleep* 30, 1145–1152. doi: 10.1093/sleep/30.9.1145

- Hassainia, F., Petit, D., Neilsen, T., Gauthier, S., and Montplaisir, J. (1997). Quantitative EEG and statistical mapping of wakefulness and REM sleep in the evaluation of mild to moderate Alzheimer's disease. *Eur. Neurol.* 37, 219–224. doi: 10.1159/000117446
- Haydon, P. G. (2017). Astrocytes and the modulation of sleep. *Curr. Opin. Neurobiol.* 44, 28–33. doi: 10.1016/j.conb.2017.02.008
- Helfrich, R. F., Lendner, J. D., Mander, B. A., Guillen, H., Paff, M., Mnatsakanyan, L., et al. (2019). Bidirectional prefrontal-hippocampal dynamics organize information transfer during sleep in humans. *Nat. Commun.* 10:3572.
- Helfrich, R. F., Mander, B. A., Jagust, W. J., Knight, R. T., and Walker, M. P. (2018). Old brains come uncoupled in sleep: slow wave-spindle synchrony, brain atrophy, and forgetting. *Neuron* 97, 221–230.e4.
- Heppner, F. L., Ransohoff, R. M., and Becher, B. (2015). Immune attack: the role of inflammation in Alzheimer disease. *Nat. Rev. Neurosci.* 16, 358–372. doi: 10.1038/nrn3880
- Herrup, K. (2015). The case for rejecting the amyloid cascade hypothesis. *Nat. Neurosci.* 18, 794–799. doi: 10.1038/nn.4017
- Hita-Yanez, E., Atienza, M., and Cantero, J. L. (2013). Polysomnographic and subjective sleep markers of mild cognitive impairment. *Sleep* 36, 1327–1334. doi: 10.5665/sleep.2956
- Hita-Yanez, E., Atienza, M., Gil-Neciga, E., and Cantero, J. L. (2012). Disturbed sleep patterns in elders with mild cognitive impairment: the role of memory decline and ApoE epsilon4 genotype. *Curr. Alzheimer Res.* 9, 290–297. doi: 10.2174/156720512800107609
- Holland, D., Mcevoy, L. K., Dale, A. M., and Alzheimer's Disease Neuroimaging, I. (2012). Unbiased comparison of sample size estimates from longitudinal structural measures in ADNI. *Hum. Brain. Mapp.* 33, 2586–2602. doi: 10.1002/hbm.21386
- Holth, J. K., Fritsch, S. K., Wang, C., Pedersen, N. P., Cirrito, J. R., Mahan, T. E., et al. (2019). The sleep-wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans. *Science* 363, 880–884. doi: 10.1126/science.aav2546
- Holth, J. K., Mahan, T. E., Robinson, G. O., Rocha, A., and Holtzman, D. M. (2017). Altered sleep and EEG power in the P301S Tau transgenic mouse model. *Ann. Clin. Transl. Neurol.* 4, 180–190. doi: 10.1002/acn3.390
- Huber, R., Felice Ghilardi, M., Massimini, M., and Tononi, G. (2004). Local sleep and learning. *Nature* 430, 78–81. doi: 10.1038/nature02663
- Huber, R., Ghilardi, M. F., Massimini, M., Ferrarelli, F., Riedner, B. A., Peterson, M. J., et al. (2006). Arm immobilization causes cortical plastic changes and locally decreases sleep slow wave activity. *Nat. Neurosci.* 9, 1169–1176. doi: 10.1038/nn1758
- Hyman, B. T., Van Hoesen, G. W., Kromer, L. J., and Damasio, A. R. (1986). Perforant pathway changes and the memory impairment of Alzheimer's disease. *Ann. Neurol.* 20, 472–481. doi: 10.1002/ana.410200406
- Ingiosi, A. M., Opp, M. R., and Krueger, J. M. (2013). Sleep and immune function: glial contributions and consequences of aging. *Curr. Opin. Neurobiol.* 23, 806–811. doi: 10.1016/j.conb.2013.02.003
- Irwin, M. R. (2019). Sleep and inflammation: partners in sickness and in health. *Nat. Rev. Immunol.* 19, 702–715. doi: 10.1038/s41577-019-0190-z
- Irwin, M. R., Olmstead, R., and Carroll, J. E. (2016). Sleep disturbance, sleep duration, and inflammation: a systematic review and meta-analysis of cohort studies and experimental sleep deprivation. *Biol. Psychiatry* 80, 40–52. doi: 10.1016/j.biopsych.2015.05.014
- Irwin, M. R., and Vitiello, M. V. (2019). Implications of sleep disturbance and inflammation for Alzheimer's disease dementia. *Lancet Neurol.* 18, 296–306. doi: 10.1016/s1474-4422(18)30450-2
- Irwin, M. R., Wang, M., Ribeiro, D., Cho, H. J., Olmstead, R., Breen, E. C., et al. (2008). Sleep loss activates cellular inflammatory signaling. *Biol. Psychiatry* 64, 538–540. doi: 10.1016/j.biopsych.2008.05.004
- Jack, C. R. Jr., Knopman, D. S., Jagust, W. J., Shaw, L. M., Aisen, P. S., Weiner, M. W., et al. (2010). Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol.* 9, 119–128. doi: 10.1016/s1474-4422(09)70299-6
- Jack, C. R. Jr., Wiste, H. J., Therneau, T. M., Weigand, S. D., Knopman, D. S., Mielke, M. M., et al. (2019). Associations of amyloid, Tau, and neurodegeneration biomarker profiles with rates of memory decline among individuals without dementia. *JAMA* 321, 2316–2325. doi: 10.1001/jama.2019.7437
- Jack, C. R., Knopman, D. S., Jagust, W. J., Petersen, R. C., Weiner, M. W., Aisen, P. S., et al. (2013). Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. *Lancet Neurol.* 12, 207–216. doi: 10.1016/s1474-4422(12)70291-0
- Jacobs, H. I. L., Radua, J., Luckmann, H. C., and Sack, A. T. (2013). Meta-analysis of functional network alterations in Alzheimer's disease: toward a network biomarker. *Neurosci. Biobehav. Rev.* 37, 753–765. doi: 10.1016/j.neubiorev.2013.03.009
- Jagust, W. (2013). Vulnerable neural systems and the borderland of brain aging and neurodegeneration. *Neuron* 77, 219–234. doi: 10.1016/j.neuron.2013.01.002
- Jagust, W. (2018). Imaging the evolution and pathophysiology of Alzheimer disease. *Nat. Rev. Neurosci.* 19, 687–700. doi: 10.1038/s41583-018-0067-3
- Jessen, N. A., Munk, A. S. F., Lundgaard, I., and Nedergaard, M. (2015). The Glymphatic system: a Beginner's Guide. *Neurochem. Res.* 40, 2583–2599.
- Ji, D., and Wilson, M. A. (2007). Coordinated memory replay in the visual cortex and hippocampus during sleep. *Nat. Neurosci.* 10, 100–107. doi: 10.1038/nn1825
- Jobert, M., Poiseau, E., Jahng, P., Schulz, H., and Kubicki, S. (1992). Topographical analysis of sleep spindle activity. *Neuropsychobiology* 26, 210–217. doi: 10.1159/000118923
- Ju, Y. E., Mclelland, J. S., Toedebusch, C. D., Xiong, C., Fagan, A. M., Duntley, S. P., et al. (2013). Sleep quality and preclinical Alzheimer disease. *JAMA Neurol.* 70, 587–593.
- Ju, Y. E. S., Lucey, B. P., and Holtzman, D. M. (2014). Sleep and Alzheimer disease pathology—a bidirectional relationship. *Nat. Rev. Neurol.* 10, 115–119. doi: 10.1038/nrneurol.2013.269
- Ju, Y. S., Ooms, S. J., Sutphen, C., Macauley, S. L., Zangrilli, M. A., Jerome, G., et al. (2017). Slow wave sleep disruption increases cerebrospinal fluid amyloid-beta levels. *Brain* 140, 2104–2111. doi: 10.1093/brain/awx148
- Jyoti, A., Plano, A., Riedel, G., and Platt, B. (2015). Progressive age-related changes in sleep and EEG profiles in the PLB1 (Triple) mouse model of Alzheimer's disease. *Neurobiol. Aging* 36, 2768–2784. doi: 10.1016/j.neurobiolaging.2015.07.001
- Kabeshita, Y., Adachi, H., Matsushita, M., Kanemoto, H., Sato, S., Suzuki, Y., et al. (2017). Sleep disturbances are key symptoms of very early stage Alzheimer disease with behavioral and psychological symptoms: a Japan multi-center cross-sectional study (J-BIRD). *Int. J. Geriatr. Psychiatry* 32, 222–230. doi: 10.1002/gps.4470
- Kam, K., Duffy, A. M., Moretto, J., Lafrancois, J. J., and Scharfman, H. E. (2016). Interictal spikes during sleep are an early defect in the Tg2576 mouse model of beta-amyloid neuropathology. *Sci. Rep.* 6:20119.
- Kam, K., Parekh, A., Sharma, R. A., Andrade, A. G., Lewin, M., Castillo, B., et al. (2019). Sleep oscillation-specific associations with Alzheimer's disease CSF biomarkers: novel roles for sleep spindles and tau. *Mol. Neurodegener.* 14:10.
- Kang, J. E., Lim, M. M., Bateman, R. J., Lee, J. J., Smyth, L. P., Cirrito, J. R., et al. (2009). Amyloid-beta dynamics are regulated by orexin and the sleep-wake cycle. *Science* 326, 1005–1007. doi: 10.1126/science.1180962
- Kastanenka, K. V., Calvo-Rodriguez, M., Hou, S. S., Zhou, H., Takeda, S., Arbel-Ornath, M., et al. (2019). Frequency-dependent exacerbation of Alzheimer's disease neuropathophysiology. *Sci. Rep.* 9:8964.
- Kastanenka, K. V., Hou, S. S., Shakerdge, N., Logan, R., Feng, D., Wegmann, S., et al. (2017). Optogenetic restoration of disrupted slow oscillations halts amyloid deposition and restores calcium homeostasis in an animal model of Alzheimer's disease. *PLoS One* 12:e0170275. doi: 10.1371/journal.pone.0170275
- Kent, B. A., Strittmatter, S. M., and Nygaard, H. B. (2018). Sleep and EEG power spectral analysis in three transgenic mouse models of Alzheimer's Disease: APP/PS1, 3xTgAD, and Tg2576. *J. Alzheimers Dis.* 64, 1325–1336. doi: 10.3233/jad-180260
- Khazaie, H., Veronese, M., Noori, K., Emamian, F., Zarei, M., Ashkan, K., et al. (2017). Functional reorganization in obstructive sleep apnoea and insomnia: a systematic review of the resting-state fMRI. *Neurosci. Biobehav. Rev.* 77, 219–231. doi: 10.1016/j.neubiorev.2017.03.013
- Kim, J., Gulati, T., and Ganguly, K. (2019). Competing roles of slow oscillations and delta waves in memory consolidation versus forgetting. *Cell* 179:514. doi: 10.1016/j.cell.2019.08.040
- Kress, B. T., Iliff, J. J., Xia, M., Wang, M., Wei, H. S., Zeppenfeld, D., et al. (2014). Impairment of paravascular clearance pathways in the aging brain. *Ann. Neurol.* 76, 845–861. doi: 10.1002/ana.24271

- Krstic, D., Madhusudan, A., Doehner, J., Vogel, P., Notter, T., Imhof, C., et al. (2012). Systemic immune challenges trigger and drive Alzheimer-like neuropathology in mice. *J. Neuroinflammation* 9:151.
- Krueger, J. M., Nguyen, J. T., Dykstra-Aiello, C. J., and Taishi, P. (2019). Local sleep. *Sleep Med. Rev.* 43, 14–21. doi: 10.1016/j.smrv.2018.10.001
- Kurup, P., Zhang, Y. F., Xu, J., Venkataramani, D. V., Haroutunian, V., Greengard, P., et al. (2010). A beta-Mediated NMDA receptor endocytosis in Alzheimer's disease involves ubiquitination of the tyrosine phosphatase STEP61. *J. Neurosci.* 30, 5948–5957. doi: 10.1523/jneurosci.0157-10.2010
- Ladenbauer, J., Ladenbauer, J., Kulzow, N., De Boor, R., Avramova, E., Grittner, U., et al. (2017). Promoting sleep oscillations and their functional coupling by transcranial stimulation enhances memory consolidation in mild cognitive impairment. *J. Neurosci.* 37, 7111–7124. doi: 10.1523/jneurosci.0260-17.2017
- Landolt, H. P., Dijk, D. J., Achermann, P., and Borbely, A. A. (1996). Effect of age on the sleep EEG: slow-wave activity and spindle frequency activity in young and middle-aged men. *Brain Res.* 738, 205–212. doi: 10.1016/s0006-8993(96)00770-6
- Latchoumane, C. F. V., Ngo, H. V. V., Born, J., and Shin, H. S. (2017). Thalamic spindles promote memory formation during sleep through triple phase-locking of cortical, thalamic, and hippocampal rhythms. *Neuron* 95, 424–435.e6.
- Le Bon, O., Neu, D., Berquin, Y., Lanquart, J. P., Hoffmann, R., Mairesse, O., et al. (2012). Ultra-slow delta power in chronic fatigue syndrome. *Psychiatry Res.* 200, 742–747. doi: 10.1016/j.psychres.2012.06.027
- Leal, S. L., Landau, S. M., Bell, R. K., and Jagust, W. J. (2017). Hippocampal activation is associated with longitudinal amyloid accumulation and cognitive decline. *eLife* 6:e22978.
- Leal, S. L., and Yassa, M. A. (2013). Perturbations of neural circuitry in aging, mild cognitive impairment, and Alzheimer's disease. *Ageing Res. Rev.* 12, 823–831. doi: 10.1016/j.arr.2013.01.006
- Lendner, J. D., Helfrich, R. F., Mander, B. A., Romundstad, L., Lin, J. J., Walker, M. P., et al. (2020). An electrophysiological marker of arousal level in humans. *eLife* 9:e55092.
- Leyns, C. E. G., and Holtzman, D. M. (2017). Glial contributions to neurodegeneration in tauopathies. *Mol. Neurodegener.* 12:50.
- Li, W., Ma, L., Yang, G., and Gan, W. B. (2017). REM sleep selectively prunes and maintains new synapses in development and learning. *Nat. Neurosci.* 20, 427–437. doi: 10.1038/nn.4479
- Li, X. J., Uemura, K., Hashimoto, T., Nasser-Ghods, N., Arimon, M., Lill, C. M., et al. (2013). Neuronal activity and secreted amyloid beta lead to altered amyloid beta precursor protein and presenilin 1 interactions. *Neurobiol. Dis.* 50, 127–134. doi: 10.1016/j.nbd.2012.10.002
- Liguori, C., Romigi, A., Nuccetelli, M., Zannino, S., Sancesario, G., Martorana, A., et al. (2014). Orexinergic system dysregulation, sleep impairment, and cognitive decline in Alzheimer disease. *JAMA Neurol.* 71, 1498–1505. doi: 10.1001/jamaneurol.2014.2510
- Liguori, C., Chiaravalloti, A., IZZI, F., Nuccetelli, M., Bernardini, S., Schillaci, O., et al. (2017a). Sleep apnoeas may represent a reversible risk factor for amyloid-beta pathology. *Brain* 140:e75. doi: 10.1093/brain/awx281
- Liguori, C., Mercuri, N. B., IZZI, F., Romigi, A., Cordella, A., Sancesario, G., et al. (2017b). Obstructive sleep apnea is associated with early but possibly modifiable Alzheimer's disease biomarkers changes. *Sleep* 40:zsx011.
- Lim, A. S., Kowgier, M., Yu, L., Buchman, A. S., and Bennett, D. A. (2013a). Sleep fragmentation and the risk of incident Alzheimer's disease and cognitive decline in older persons. *Sleep* 36, 1027–1032. doi: 10.5665/sleep.2802
- Lim, A. S., Yu, L., Kowgier, M., Schneider, J. A., Buchman, A. S., and Bennett, D. A. (2013b). Modification of the relationship of the apolipoprotein E epsilon4 allele to the risk of Alzheimer disease and neurofibrillary tangle density by sleep. *JAMA Neurol.* 70, 1544–1551. doi: 10.1001/jamaneurol.2013.4215
- Lim, A. S. P., Ellison, B. A., Wang, J. L., Yu, L., Schneider, J. A., Buchman, A. S., et al. (2014). Sleep is related to neuron numbers in the ventrolateral preoptic/intermediate nucleus in older adults with and without Alzheimer's disease. *Brain* 137, 2847–2861. doi: 10.1093/brain/awu222
- Liscombe, M. P., Hoffmann, R. F., Trivedi, M. H., Parker, M. K., Rush, A. J., and Armitage, R. (2002). Quantitative EEG amplitude across REM sleep periods in depression: preliminary report. *J. Psychiatry Neurosci.* 27, 40–46.
- Lo, J. C., Loh, K. K., Zheng, H., Sim, S. K. Y., and Chee, M. W. L. (2014). Sleep duration and age-related changes in brain structure and cognitive performance. *Sleep* 37, 1171–1178.
- Lucey, B. P., McCullough, A., Landsness, E. C., Toedebusch, C. D., Mcleland, J. S., Zaza, A. M., et al. (2019). Reduced non-rapid eye movement sleep is associated with tau pathology in early Alzheimer's disease. *Sci. Transl. Med.* 11:eaa06550. doi: 10.1126/scitranslmed.aau6550
- Lustenberger, C., Boyle, M. R., Alagapan, S., Mellin, J. M., Vaughn, B. V., and Frohlich, F. (2016). Feedback-controlled transcranial alternating current stimulation reveals a functional role of sleep spindles in motor memory consolidation. *Curr. Biol.* 26, 2127–2136. doi: 10.1016/j.cub.2016.06.044
- Lutsey, P. L., Misialek, J. R., Mosley, T. H., Gottesman, R. F., Punjabi, N. M., Shahar, E., et al. (2018). Sleep characteristics and risk of dementia and Alzheimer's disease: the atherosclerosis risk in communities study. *Alzheimers Dement.* 14, 157–166. doi: 10.1016/j.jalz.2017.06.2269
- Lynch, B. A., Lambeng, N., Nocka, K., Kinsel-Hammes, P., Bajjalieh, S. M., Matagne, A., et al. (2004). The synaptic vesicle protein SV2A is the binding site for the antiepileptic drug levetiracetam. *Proc. Natl. Acad. Sci. U.S.A.* 101, 9861–9866. doi: 10.1073/pnas.0308208101
- Maass, A., Lockhart, S. N., Harrison, T. M., Bell, R. K., Mellinger, T., Swinerton, K., et al. (2018). Entorhinal Tau Pathology, episodic memory decline, and neurodegeneration in aging. *J. Neurosci.* 38, 530–543. doi: 10.1523/jneurosci.2028-17.2017
- Mak-McCully, R. A., Rolland, M., Sargsyan, A., Gonzalez, C., Magnin, M., Chauvel, P., et al. (2017). Coordination of cortical and thalamic activity during non-REM sleep in humans. *Nat. Commun.* 8:15499.
- Mander, B. A. (2013). Disturbed sleep in preclinical cognitive impairment: Cause and effect? *Sleep* 36, 1275–1276. doi: 10.5665/sleep.2942
- Mander, B. A., Marks, S. M., Vogel, J. W., Rao, V., Lu, B., Saletin, J. M., et al. (2015). beta-amyloid disrupts human NREM slow waves and related hippocampus-dependent memory consolidation. *Nat. Neurosci.* 18, 1051–1057. doi: 10.1038/nn.4035
- Mander, B. A., Rao, V., Lu, B., Saletin, J. M., Ancoli-Israel, S., Jagust, W. J., et al. (2014). Impaired prefrontal sleep spindle regulation of hippocampal-dependent learning in older adults. *Cereb. Cortex* 24, 3301–3309. doi: 10.1093/cercor/bht188
- Mander, B. A., Rao, V., Lu, B., Saletin, J. M., Lindquist, J. R., Ancoli-Israel, S., et al. (2013). Prefrontal atrophy, disrupted NREM slow waves and impaired hippocampal-dependent memory in aging. *Nat. Neurosci.* 16, 357–364. doi: 10.1038/nn.3324
- Mander, B. A., Santhanam, S., Saletin, J. M., and Walker, M. P. (2011). Wake deterioration and sleep restoration of human learning. *Curr. Biol.* 21, R183–R184.
- Mander, B. A., Winer, J. R., Jagust, W. J., and Walker, M. P. (2016). Sleep: A novel mechanistic pathway, biomarker, and treatment target in the pathology of Alzheimer's disease? *Trends Neurosci.* 39, 552–566. doi: 10.1016/j.tins.2016.05.002
- Mander, B. A., Winer, J. R., and Walker, M. P. (2017a). Sleep and Human Aging. *Neuron* 94, 19–36.
- Mander, B. A., Zhu, A. H., Lindquist, J. R., Villeneuve, S., Rao, V., Lu, B., et al. (2017b). White matter structure in older adults moderates the benefit of sleep spindles on motor memory consolidation. *J. Neurosci.* 37, 11675–11687. doi: 10.1523/jneurosci.3033-16.2017
- Maquet, P. (2000). Functional neuroimaging of normal human sleep by positron emission tomography. *J. Sleep Res.* 9, 207–231. doi: 10.1046/j.1365-2869.2000.00214.x
- Marquez, F., and Yassa, M. A. (2019). Neuroimaging biomarkers for Alzheimer's disease. *Mol. Neurodegener.* 14:21.
- Marshall, L., Helgadottir, H., Molle, M., and Born, J. (2006). Boosting slow oscillations during sleep potentiates memory. *Nature* 444, 610–613. doi: 10.1038/nature05278
- Marshall, L., Molle, M., Hallschmid, M., and Born, J. (2004). Transcranial direct current stimulation during sleep improves declarative memory. *J. Neurosci.* 24, 9985–9992. doi: 10.1523/jneurosci.2725-04.2004
- Martin, N., Lafortune, M., Godbout, J., Barakat, M., Robillard, R., Poirier, G., et al. (2013). Topography of age-related changes in sleep spindles. *Neurobiol. Aging* 34, 468–476. doi: 10.1016/j.neurobiolaging.2012.05.020
- Massimini, M., Huber, R., Ferrarelli, F., Hill, S., and Tononi, G. (2004). The sleep slow oscillation as a traveling wave. *J. Neurosci.* 24, 6862–6870. doi: 10.1523/jneurosci.1318-04.2004

- McGaugh, J. L. (2000). Memory—a century of consolidation. *Science* 287, 248–251. doi: 10.1126/science.287.5451.248
- Menkes-Caspi, N., Yamin, H. G., Kellner, V., Spires-Jones, T. L., Cohen, D., and Stern, E. A. (2015). Pathological Tau disrupts ongoing network activity. *Neuron* 85, 959–966. doi: 10.1016/j.neuron.2015.01.025
- Mesulam, M., Shaw, P., Mash, D., and Weintraub, S. (2004). Cholinergic nucleus basalis tauopathy emerges early in the aging-MCI-AD continuum. *Ann. Neurol.* 55, 815–828. doi: 10.1002/ana.20100
- Miyawaki, H., and Diba, K. (2016). Regulation of hippocampal firing by network oscillations during sleep. *Curr. Biol.* 26, 893–902. doi: 10.1016/j.cub.2016.02.024
- Mizuseki, K., and Miyawaki, H. (2017). Hippocampal information processing across sleep/wake cycles. *Neurosci. Res.* 118, 30–47. doi: 10.1016/j.neures.2017.04.018
- Molano, J. R. V., Roe, C. M., and Ju, Y. E. (2017). The interaction of sleep and amyloid deposition on cognitive performance. *J. Sleep Res.* 26, 288–292. doi: 10.1111/jsr.12474
- Muehlroth, B. E., Sander, M. C., Fandakova, Y., Grandy, T. H., Rasch, B., Shing, Y. L., et al. (2019). Precise slow oscillation-spindle coupling promotes memory consolidation in younger and older adults. *Sci. Rep.* 9:1940.
- Mufson, E. J., Mash, D. C., and Hersh, L. B. (1988). Neurofibrillary tangles in cholinergic pedunculopontine neurons in Alzheimers-disease. *Ann. Neurol.* 24, 623–629. doi: 10.1002/ana.410240506
- Muller, L., Piantoni, G., Koller, D., Cash, S. S., Halgren, E., and Sejnowski, T. J. (2016). Rotating waves during human sleep spindles organize global patterns of activity that repeat precisely through the night. *eLife* 5:e17267.
- Murphy, M., Riedner, B. A., Huber, R., Massimini, M., Ferrarelli, F., and Tononi, G. (2009). Source modeling sleep slow waves. *Proc. Natl. Acad. Sci. U.S.A.* 106, 1608–1613.
- Musiek, E. S., Bhimasani, M., Zangrilli, M. A., Morris, J. C., Holtzman, D. M., and Ju, Y. S. (2018). Circadian rest-activity pattern changes in aging and preclinical Alzheimer disease. *JAMA Neurol.* 75, 582–590. doi: 10.1001/jamaneurol.2017.4719
- Musiek, E. S., and Holtzman, D. M. (2015). Three dimensions of the amyloid hypothesis: time, space and 'wingmen'. *Nat. Neurosci.* 18, 800–806. doi: 10.1038/nn.4018
- Nedergaard, M. (2013). Neuroscience. Garbage truck of the brain. *Science* 340, 1529–1530. doi: 10.1126/science.1240514
- Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., et al. (2011). Regional slow waves and spindles in human sleep. *Neuron* 70, 153–169. doi: 10.1016/j.neuron.2011.02.043
- O'Donoghue, M. C., Murphy, S. E., Zamboni, G., Nobre, A. C., and Mackay, C. E. (2018). APOE genotype and cognition in healthy individuals at risk of Alzheimer's disease: a review. *Cortex* 104, 103–123. doi: 10.1016/j.cortex.2018.03.025
- Oh, J., Eser, R. A., Ehrenberg, A. J., Morales, D., Petersen, C., Kudlacek, J., et al. (2019). Profound degeneration of wake-promoting neurons in Alzheimer's disease. *Alzheimers Dement.* 15, 1253–1263. doi: 10.1016/j.jalz.2019.06.3916
- Ohayon, M. M., Carskadon, M. A., Guilleminault, C., and Vitiello, M. V. (2004). Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan. *Sleep* 27, 1255–1273. doi: 10.1093/sleep/27.7.1255
- Olcese, U., Bos, J. J., Vinck, M., and Pennartz, C. M. A. (2018). Functional determinants of enhanced and depressed interareal information flow in nonrapid eye movement sleep between neuronal ensembles in rat cortex and hippocampus. *Sleep* 41:zsy167.
- Osorio, R. S., Gumb, T., Pirraglia, E., Varga, A. W., Lu, S. E., Lim, J., et al. (2015). Sleep-disordered breathing advances cognitive decline in the elderly. *Neurology* 84, 1964–1971.
- Palmqvist, S., Scholl, M., Strandberg, O., Mattsson, N., Stomrud, E., Zetterberg, H., et al. (2017). Earliest accumulation of beta-amyloid occurs within the default-mode network and concurrently affects brain connectivity. *Nat. Commun.* 8:1214.
- Papalambros, N. A., Santostasi, G., Malkani, R. G., Braun, R., Weintraub, S., Paller, K. A., et al. (2017). Acoustic enhancement of sleep slow oscillations and concomitant memory improvement in older adults. *Front. Hum. Neurosci.* 11:109. doi: 10.3389/fnhum.2017.00109
- Papalambros, N. A., Weintraub, S., Chen, T., Grimaldi, D., Santostasi, G., Paller, K. A., et al. (2019). Acoustic enhancement of sleep slow oscillations in mild cognitive impairment. *Ann. Clin. Transl. Neurol.* 6, 1191–1201. doi: 10.1002/acn3.796
- Pase, M. P., Himali, J. J., Grima, N. A., Beiser, A. S., Satizabal, C. L., Aparicio, H. J., et al. (2017). Sleep architecture and the risk of incident dementia in the community. *Neurology* 89, 1244–1250. doi: 10.1212/wnl.0000000000004373
- Peever, J., and Fuller, P. M. (2016). The Biology of REM Sleep. *Curr. Biol.* 26, R34–R35.
- Pereira, J. B., Ossenkoppele, R., Palmqvist, S., Strandberg, T. O., Smith, R., Westman, E., et al. (2019). Amyloid and tau accumulate across distinct spatial networks and are differentially associated with brain connectivity. *eLife* 8:e50830.
- Petit, D., Gagnon, J. F., Fantini, M. L., Ferini-Strambi, L., and Montplaisir, J. (2004). Sleep and quantitative EEG in neurodegenerative disorders. *J. Psychosom. Res.* 56, 487–496. doi: 10.1016/j.jpsychores.2004.02.001
- Picchioni, D., Horovitz, S. G., Fukunaga, M., Carr, W. S., Meltzer, J. A., Balkin, T. J., et al. (2011). Infralow EEG oscillations organize large-scale cortical-subcortical interactions during sleep: a combined EEG/fMRI study. *Brain Res.* 1374, 63–72. doi: 10.1016/j.brainres.2010.12.035
- Pihlajamaki, M., Depeau, K. M., Blacker, D., and Sperling, R. A. (2008). Impaired medial temporal repetition suppression is related to failure of parietal deactivation in Alzheimer disease. *Am. J. Geriatr. Psychiatry* 16, 283–292. doi: 10.1097/jgp.0b013e318162a0a9
- Podcasy, J. L., and Epperson, C. N. (2016). Considering sex and gender in Alzheimer disease and other dementias. *Dialogues Clin. Neurosci.* 18, 437–446.
- Poskanzer, K. E., and Yuste, R. (2016). Astrocytes regulate cortical state switching in vivo. *Proc. Natl. Acad. Sci. U.S.A.* 113, E2675–E2684.
- Prinz, P. N., Vitaliano, P. P., Vitiello, M. V., Bokan, J., Raskind, M., Peskind, E., et al. (1982). Sleep, EEG and mental function changes in senile dementia of the Alzheimer's type. *Neurobiol. Aging* 3, 361–370. doi: 10.1016/0197-4580(82)90024-0
- Rasch, B., Buchel, C., Gais, S., and Born, J. (2007). Odor cues during slow-wave sleep prompt declarative memory consolidation. *Science* 315, 1426–1429. doi: 10.1126/science.1138581
- Rasmussen, M. K., Mestre, H., and Nedergaard, M. (2018). The glymphatic pathway in neurological disorders. *Lancet Neurol.* 17, 1016–1024. doi: 10.1016/s1474-4422(18)30318-1
- Rauchs, G., Schabus, M., Parapatics, S., Bertran, F., Clochon, P., Hot, P., et al. (2008). Is there a link between sleep changes and memory in Alzheimer's disease? *Neuroreport* 19, 1159–1162.
- Reagh, Z. M., Noche, J. A., Tustison, N. J., Delisle, D., Murray, E. A., and Yassa, M. A. (2018). Functional Imbalance of Anterolateral Entorhinal Cortex and Hippocampal Dentate/CA3 Underlies Age-Related Object Pattern Separation Deficits. *Neuron* <cpsjrn> 97, 1187–1198.e4.
- Redline, S., Kirchner, H. L., Quan, S. F., Gottlieb, D. J., Kapur, V., and Newman, A. (2004). The effects of age, sex, ethnicity, and sleep-disordered breathing on sleep architecture. *Arch. Intern. Med.* 164, 406–418. doi: 10.1001/archinte.164.4.406
- Roh, J. H., Huang, Y., Bero, A. W., Kasten, T., Stewart, F. R., Bateman, R. J., et al. (2012). Disruption of the sleep-wake cycle and diurnal fluctuation of β -amyloid in mice with Alzheimer's disease pathology. *Sci. Transl. Med.* 4:150ra122. doi: 10.1126/scitranslmed.3004291
- Rosanova, M., and Ulrich, D. (2005). Pattern-specific associative long-term potentiation induced by a sleep spindle-related spike train. *J. Neurosci.* 25, 9398–9405. doi: 10.1523/jneurosci.2149-05.2005
- Rothman, S. M., Herdener, N., Frankola, K. A., Mughal, M. R., and Mattson, M. P. (2013). Chronic mild sleep restriction accentuates contextual memory impairments, and accumulations of cortical A beta and pTau in a mouse model of Alzheimer's disease. *Brain Res.* 1529, 200–208. doi: 10.1016/j.brainres.2013.07.010
- Rudoy, J. D., Voss, J. L., Westerberg, C. E., and Paller, K. A. (2009). Strengthening individual memories by reactivating them during sleep. *Science* 326:1079. doi: 10.1126/science.1179013
- Saletin, J. M., Goldstein-Piekarski, A. N., Greer, S. M., Stark, S., Stark, C. E., and Walker, M. P. (2016). Human hippocampal structure: a novel biomarker predicting mnemonic vulnerability to, and recovery from, Sleep Deprivation. *J. Neurosci.* 36, 2355–2363. doi: 10.1523/jneurosci.3466-15.2016

- Scammell, T. E., Arrigoni, E., and Lipton, J. O. (2017). Neural circuitry of wakefulness and sleep. *Neuron* 93, 747–765. doi: 10.1016/j.neuron.2017.01.014
- Scarpelli, S., D'atri, A., Bartolacci, C., Mangiaruga, A., Gorgoni, M., and De Gennaro, L. (2019). Oscillatory EEG Activity During REM Sleep in elderly people predicts subsequent dream recall after awakenings. *Front. Neurol.* 10:985. doi: 10.3389/fneur.2019.00985
- Scholl, M., Lockhart, S. N., Schonhaut, D. R., O'Neil, J. P., Janabi, M., Ossenkoppele, R., et al. (2016). PET imaging of Tau deposition in the aging human brain. *Neuron* 89, 971–982. doi: 10.1016/j.neuron.2016.01.028
- Sepulcre, J., Sabuncu, M. R., Becker, A., Sperling, R., and Johnson, K. A. (2013). *In vivo* characterization of the early states of the amyloid-beta network. *Brain* 136, 2239–2252. doi: 10.1093/brain/awt146
- Shi, L., Chen, S. J., Ma, M. Y., Bao, Y. P., Han, Y., Wang, Y. M., et al. (2017). Sleep disturbances increase the risk of dementia: a systematic review and meta-analysis. *Sleep Med. Rev.* 40, 4–16. doi: 10.1016/j.smrv.2017.06.010
- Smagula, S. F., Stone, K. L., Redline, S., Ancoli-Israel, S., Barrett-Connor, E., Lane, N. E., et al. (2016). Actigraphy- and Polysomnography-measured sleep disturbances, inflammation, and mortality among older men. *Psychosom. Med.* 78, 686–696. doi: 10.1097/psy.0000000000000312
- Smith, A. J., and Verkman, A. S. (2018). The "glymphatic" mechanism for solute clearance in Alzheimer's disease: Game changer or unproven speculation? *FASEB J.* 32, 543–551. doi: 10.1096/fj.201700999
- Song, Y. S., Blackwell, T., Yaffe, K., Ancoli-Israel, S., Redline, S., Stone, K. L., et al. (2015). Relationships between sleep stages and changes in cognitive function in older men: the MrOS sleep study. *Sleep* 38, 411–421. doi: 10.5665/sleep.4500
- Spangenberg, E. E., and Green, K. N. (2017). Inflammation in Alzheimer's disease: lessons learned from microglia-depletion models. *Brain Behav. Immun.* 61, 1–11. doi: 10.1016/j.bbi.2016.07.003
- Spangenberg, E. E., Lee, R. J., Najafi, A. R., Rice, R. A., Elmore, M. R., Blurton-Jones, M., et al. (2016). Eliminating microglia in Alzheimer's mice prevents neuronal loss without modulating amyloid-beta pathology. *Brain* 139, 1265–1281. doi: 10.1093/brain/aww016
- Spencer, R. M., Gouw, A. M., and Ivry, R. B. (2007). Age-related decline of sleep-dependent consolidation. *Learn. Mem.* 14, 480–484. doi: 10.1101/lm.569407
- Spencer, R. M., Sunm, M., and Ivry, R. B. (2006). Sleep-dependent consolidation of contextual learning. *Curr. Biol.* 16, 1001–1005. doi: 10.1016/j.cub.2006.03.094
- Sperling, R. A., Laviolette, P. S., O'Keefe, K., O'Brien, J., Rentz, D. M., Pihlajamaki, M., et al. (2009). Amyloid deposition is associated with impaired default network function in older persons without dementia. *Neuron* 63, 178–188. doi: 10.1016/j.neuron.2009.07.003
- Sperling, R. A., Mormino, E. C., Schultz, A. P., Betensky, R. A., Papp, K. V., Amariglio, R. E., et al. (2019). The impact of amyloid-beta and tau on prospective cognitive decline in older individuals. *Ann. Neurol.* 85, 181–193.
- Spira, A. P., Gamaldo, A. A., An, Y., Wu, M. N., Simonsick, E. M., Bilgel, M., et al. (2013). Self-reported sleep and beta-amyloid deposition in community-dwelling older adults. *JAMA Neurol.* 70, 1537–1543.
- Spira, A. P., Gonzalez, C. E., Venkatraman, V. K., Wu, M. N., Pacheco, J., Simonsick, E. M., et al. (2016). Sleep duration and subsequent cortical thinning in cognitively normal older adults. *Sleep* 39, 1121–1128. doi: 10.5665/sleep.5768
- Sprecher, K. E., Bendlin, B. B., Racine, A. M., Okonkwo, O. C., Christian, B. T., Kosciak, R. L., et al. (2015). Amyloid burden is associated with self-reported sleep in nondemented late middle-aged adults. *Neurobiol. Aging* 36, 2568–2576. doi: 10.1016/j.neurobiolaging.2015.05.004
- Sprecher, K. E., Kosciak, R. L., Carlsson, C. M., Zetterberg, H., Blennow, K., Okonkwo, O. C., et al. (2017). Poor sleep is associated with CSF biomarkers of amyloid pathology in cognitively normal adults. *Neurology* 89, 445–453. doi: 10.1212/wnl.00000000000004171
- Sprecher, K. E., Riedner, B. A., Smith, R. F., Tononi, G., Davidson, R. J., and Benca, R. M. (2016). High resolution topography of age-related changes in non-rapid eye movement sleep electroencephalography. *PLoS One* 11:e0149770. doi: 10.1371/journal.pone.0149770
- Staresina, B. P., Bergmann, T. O., Bonfond, M., Van Der Meij, R., Jensen, O., Deuker, L., et al. (2015). Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. *Nat. Neurosci.* 18, 1679–1686. doi: 10.1038/nn.4119
- Stark, S. M., Yassa, M. A., Lacy, J. W., and Stark, C. E. (2013). A task to assess behavioral pattern separation (BPS) in humans: data from healthy aging and mild cognitive impairment. *Neuropsychologia* 51, 2442–2449. doi: 10.1016/j.neuropsychologia.2012.12.014
- Stark, S. M., Yassa, M. A., and Stark, C. E. (2010). Individual differences in spatial pattern separation performance associated with healthy aging in humans. *Learn. Mem.* 17, 284–288. doi: 10.1101/lm.1768110
- Steriade, M. (2006). Grouping of brain rhythms in corticothalamic systems. *Neuroscience* 137, 1087–1106. doi: 10.1016/j.neuroscience.2005.10.029
- Steriade, M., Nunez, A., and Amzica, F. (1993a). Intracellular analysis of relations between the slow (< 1 Hz) neocortical oscillation and other sleep rhythms of the electroencephalogram. *J. Neurosci.* 13, 3266–3283. doi: 10.1523/jneurosci.13-08-03266.1993
- Steriade, M., Nunez, A., and Amzica, F. (1993b). A Novel Slow (Less-Than-1 Hz) oscillation of neocortical neurons *in-vivo* - depolarizing and hyperpolarizing components. *J. Neurosci.* 13, 3252–3265. doi: 10.1523/jneurosci.13-08-03252.1993
- Tashjian, S. M., Goldenberg, D., and Galvan, A. (2017). Neural connectivity moderates the association between sleep and impulsivity in adolescents. *Dev. Cogn. Neurosci.* 27, 35–44. doi: 10.1016/j.dcn.2017.07.006
- Terwel, D., Steffensen, K. R., Verghese, P. B., Kummer, M. P., Gustafsson, J. A., Holtzman, D. M., et al. (2011). Critical role of astroglial apolipoprotein E and liver X receptor-alpha expression for microglial Abeta phagocytosis. *J. Neurosci.* 31, 7049–7059. doi: 10.1523/jneurosci.6546-10.2011
- Thal, D. R., Rub, U., Orantes, M., and Braak, H. (2002). Phases of A beta-deposition in the human brain and its relevance for the development of AD. *Neurology* 58, 1791–1800. doi: 10.1212/wnl.58.12.1791
- Tononi, G., and Cirelli, C. (2014). Sleep and the price of plasticity: from synaptic and cellular homeostasis to memory consolidation and integration. *Neuron* 81, 12–34. doi: 10.1016/j.neuron.2013.12.025
- Tosun, D., Landau, S., Aisen, P. S., Petersen, R. C., Mintun, M., Jagust, W., et al. (2017). Association between tau deposition and antecedent amyloid-beta accumulation rates in normal and early symptomatic individuals. *Brain* 140, 1499–1512. doi: 10.1093/brain/awx046
- Tsapanou, A., Gu, Y., O'shea, D. M., Yannakoulia, M., Kosmidis, M., Dardiotis, E., et al. (2017). Sleep quality and duration in relation to memory in the elderly: initial results from the Hellenic Longitudinal Investigation of Aging and Diet. *Neurobiol. Learn. Mem.* 141, 217–225. doi: 10.1016/j.nlm.2017.04.011
- Van Cauter, E., Leproult, R., and Plat, L. (2000). Age-related changes in slow wave sleep and REM sleep and relationship with growth hormone and cortisol levels in healthy men. *JAMA* 284, 861–868. doi: 10.1001/jama.284.7.861
- Van Dongen, E. V., Takashima, A., Barth, M., Zapp, J., Schad, L. R., Paller, K. A., et al. (2012). Memory stabilization with targeted reactivation during human slow-wave sleep. *Proc. Natl. Acad. Sci. U.S.A.* 109, 10575–10580. doi: 10.1073/pnas.1201072109
- Van Dongen, H. P., Belenky, G., and Krueger, J. M. (2011). A local, bottom-up perspective on sleep deprivation and neurobehavioral performance. *Curr. Top. Med. Chem.* 11, 2414–2422. doi: 10.2174/156802611797470286
- VanElzakker, M. B., Brumfield, S. A., and Lara Mejia, P. S. (2019). Corrigendum: neuroinflammation and Cytokines in Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS): a Critical Review of Research Methods. *Front. Neurol.* 10:316. doi: 10.3389/fneur.2019.00316
- Varga, A. W., Duca, E. L., Kishi, A., Fischer, E., Parekh, A., Koushyk, V., et al. (2016a). Effects of aging on slow-wave sleep dynamics and human spatial navigational memory consolidation. *Neurobiol. Aging* 42, 142–149. doi: 10.1016/j.neurobiolaging.2016.03.008
- Varga, A. W., Wohlleber, M. E., Gimenez, S., Romero, S., Alonso, J. F., Duca, E. L., et al. (2016b). Reduced slow-wave sleep is associated with high cerebrospinal fluid a beta 42 levels in cognitively normal elderly. *Sleep* 39, 2041–2048. doi: 10.5665/sleep.6240
- Vegni, C., Ktonas, P., Giganti, F., Ficca, G., Trammell, J., Gori, S., et al. (2001). The organization of rapid eye movement activity during rapid eye movement sleep is further impaired in very old human subjects. *Neurosci. Lett.* 297, 58–60. doi: 10.1016/s0304-3940(00)01636-0
- Versijpt, J. J., Dumont, F., Van Laere, K. J., Decoo, D., Santens, P., Audenaert, K., et al. (2003). Assessment of neuroinflammation and microglial activation in Alzheimer's disease with radiolabelled PK11195 and single photon emission computed tomography. A pilot study. *Eur. Neurol.* 50, 39–47. doi: 10.1159/000070857

- Vogel, J. W., Iturria-Medina, Y., Strandberg, O. T., Smith, R., Levitis, E., Evans, A. C., et al. (2020). Spread of pathological tau proteins through communicating neurons in human Alzheimer's disease. *Nat. Commun.* 11:2612.
- Von Kanel, R., Dimsdale, J. E., Ancoli-Israel, S., Mills, P. J., Patterson, T. L., Mckibbin, C. L., et al. (2006). Poor sleep is associated with higher plasma proinflammatory cytokine interleukin-6 and procoagulant marker fibrin D-dimer in older caregivers of people with Alzheimer's disease. *J. Am. Geriatr. Soc.* 54, 431–437. doi: 10.1111/j.1532-5415.2005.00642.x
- Vyazovskiy, V. V., Cirelli, C., and Tononi, G. (2011). Electrophysiological correlates of sleep homeostasis in freely behaving rats. *Prog Brain Res.* 193, 17–38. doi: 10.1016/b978-0-444-53839-0.00002-8
- Walker, M. P. (2009). The role of sleep in cognition and emotion. *Ann. N. Y. Acad. Sci.* 1156, 168–197. doi: 10.1111/j.1749-6632.2009.04416.x
- Walker, M. P., Brakefield, T., Morgan, A., Hobson, J. A., and Stickgold, R. (2002). Practice with sleep makes perfect: sleep-dependent motor skill learning. *Neuron* 35, 205–211. doi: 10.1016/s0896-6273(02)00746-8
- Walsh, C. M., Ruoff, L., Walker, K., Emery, A., Varbel, J., Karageorgiou, E., et al. (2017). Sleepless night and day, the plight of progressive supranuclear palsy. *Sleep* 40:zsx154.
- Wamsley, E. J., and Stickgold, R. (2011). Memory, sleep and dreaming: experiencing consolidation. *Sleep Med. Clin.* 6, 97–108. doi: 10.1016/j.jsmc.2010.12.008
- Wang, J., Gu, B. J., Masters, C. L., and Wang, Y. J. (2017). A systemic view of Alzheimer disease - insights from amyloid-beta metabolism beyond the brain. *Nat. Rev. Neurol.* 13:703. doi: 10.1038/nrneurol.2017.147
- Ward, A. M., McLaren, D. G., Schultz, A. P., Chhatwal, J., Boot, B. P., Hedden, T., et al. (2013). Daytime sleepiness is associated with decreased default mode network connectivity in both young and cognitively intact elderly subjects. *Sleep* 36, 1609–1615. doi: 10.5665/sleep.3108
- Watson, B. O., and Buzsaki, G. (2015). Sleep, Memory & Brain Rhythms. *Daedalus* 144, 67–82.
- Westerberg, C. E., Florczak, S. M., Weintraub, S., Mesulam, M. M., Marshall, L., Zee, P. C., et al. (2015). Memory improvement via slow-oscillatory stimulation during sleep in older adults. *Neurobiol. Aging* 36, 2577–2586. doi: 10.1016/j.neurobiolaging.2015.05.014
- Westerberg, C. E., Mander, B. A., Florczak, S. M., Weintraub, S., Mesulam, M. M., Zee, P. C., et al. (2012). Concurrent impairments in sleep and memory in amnesic mild cognitive impairment. *J. Int. Neuropsychol. Soc.* 18, 490–500. doi: 10.1017/s135561771200001x
- Whitwell, J. L. (2010). Progression of atrophy in Alzheimer's disease and related disorders. *Neurotox. Res.* 18, 339–346.
- Wierzynski, C. M., Lubenov, E. V., Gu, M., and Siapas, A. G. (2009). State-dependent spike-timing relationships between hippocampal and prefrontal circuits during sleep. *Neuron* 61, 587–596. doi: 10.1016/j.neuron.2009.01.011
- Wilcock, D. M., Vitek, M. P., and Colton, C. A. (2009). Vascular amyloid alters astrocytic water and potassium channels in mouse models and humans with Alzheimer's disease. *Neuroscience* 159, 1055–1069. doi: 10.1016/j.neuroscience.2009.01.023
- Wilson, I. A., Gallagher, M., Eichenbaum, H., and Tanila, H. (2006). Neurocognitive aging: prior memories hinder new hippocampal encoding. *Trends Neurosci.* 29, 662–670. doi: 10.1016/j.tins.2006.10.002
- Wilson, M. A., and McNaughton, B. L. (1994). Reactivation of hippocampal ensemble memories during sleep. *Science* 265, 676–679. doi: 10.1126/science.8036517
- Winer, J. R., Mander, B. A., Helfrich, R. F., Maass, A., Harrison, T. M., Baker, S. L., et al. (2019). Sleep as a Potential Biomarker of Tau and beta-Amyloid Burden in the Human Brain. *J. Neurosci.* 39, 6315–6324. doi: 10.1523/jneurosci.0503-19.2019
- Winer, J. R., Mander, B. A., Kumar, S., Reed, M., Baker, S. L., Jagust, W. J., et al. (2020). Sleep disturbance forecasts b-amyloid accumulation across subsequent years. *Curr. Biol.* 30, 1–8.
- Witter, M. P. (2007). The perforant path: projections from the entorhinal cortex to the dentate gyrus. *Prog. Brain Res.* 163, 43–61. doi: 10.1016/s0079-6123(07)63003-9
- Witton, J., Staniaszek, L. E., Bartsch, U., Randall, A. D., Jones, M. W., and Brown, J. T. (2016). Disrupted hippocampal sharp-wave ripple-associated spike dynamics in a transgenic mouse model of dementia. *J. Physiol.* 594, 4615–4630. doi: 10.1113/jphysiol.2014.282889
- Xie, L., Kang, H., Xu, Q., Chen, M. J., Liao, Y., Thiyagarajan, M., et al. (2013). Sleep drives metabolite clearance from the adult brain. *Science* 342, 373–377. doi: 10.1126/science.1241224
- Yaffe, K., Laffan, A. M., Harrison, S. L., Redline, S., Spira, A. P., Ensrud, K. E., et al. (2011). Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA* 306, 613–619.
- Yang, G., Lai, C. S., Cichon, J., Ma, L., Li, W., and Gan, W. B. (2014). Sleep promotes branch-specific formation of dendritic spines after learning. *Science* 344, 1173–1178. doi: 10.1126/science.1249098
- Yassa, M. A., Muftuler, L. T., and Stark, C. E. (2010a). Ultra-high-resolution microstructural diffusion tensor imaging reveals perforant path degradation in aged humans *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.* 107, 12687–12691. doi: 10.1073/pnas.1002113107
- Yassa, M. A., Stark, S. M., Bakker, A., Albert, M. S., Gallagher, M., and Stark, C. E. (2010b). High-resolution structural and functional MRI of hippocampal CA3 and dentate gyrus in patients with amnesic Mild Cognitive Impairment. *Neuroimage* 51, 1242–1252. doi: 10.1016/j.neuroimage.2010.03.040
- Yassa, M. A., Lacy, J. W., Stark, S. M., Albert, M. S., Gallagher, M., and Stark, C. E. (2011a). Pattern separation deficits associated with increased hippocampal CA3 and dentate gyrus activity in nondemented older adults. *Hippocampus* 21, 968–979.
- Yassa, M. A., Mattfeld, A. T., Stark, S. M., and Stark, C. E. (2011b). Age-related memory deficits linked to circuit-specific disruptions in the hippocampus. *Proc. Natl. Acad. Sci. U.S.A.* 108, 8873–8878. doi: 10.1073/pnas.1101567108
- Zeppenfeld, D. M., Simon, M., Haswell, J. D., D'abreo, D., Murchison, C., Quinn, J. F., et al. (2017). Association of perivascular localization of aquaporin-4 with cognition and Alzheimer disease in aging brains. *JAMA Neurol.* 74, 91–99. doi: 10.1001/jamaneurol.2016.4370
- Zhang, B., Veasey, S. C., Wood, M. A., Leng, L. Z., Kaminski, C., Leight, S., et al. (2005). Impaired rapid eye movement sleep in the Tg2576 APP murine model of Alzheimer's disease with injury to pedunculopontine cholinergic neurons. *Am. J. Pathol.* 167, 1361–1369. doi: 10.1016/s0002-9440(10)61223-0
- Zhu, Y., Zhan, G. X., Fenik, P., Brandes, M., Bell, P., Francois, N., et al. (2018). Chronic sleep disruption advances the temporal progression of tauopathy in p301s mutant mice. *J. Neurosci.* 38, 10255–10270. doi: 10.1523/jneurosci.0275-18.2018

Conflict of Interest: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Mander. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.