

Rising Temperature May Trigger Deep Soil Carbon Loss Across Forest Ecosystems

Jinquan Li, Junmin Pei, Elise Pendall, Peter B. Reich, Nam Jin Noh, Bo Li, Changming Fang, and Ming Nie*

Significantly more carbon (C) is stored in deep soil than in shallow horizons, yet how the decomposition of deep soil organic C (SOC) will respond to rising temperature remains unexplored on large scales, leading to considerable uncertainties to predictions of the magnitude and direction of C-cycle feedbacks to climate change. Herein, short-term temperature sensitivity of SOC decomposition (expressed as Q_{10}) from six depths within the top 1 m soil from 90 upland forest sites (540 soil samples) across China is reported. Results show that Q_{10} significantly increases with soil depth, suggesting that deep SOC is more vulnerable to loss with rising temperature in comparison to shallow SOC. Climate is the primary regulator of shallow soil Q_{10} but its relative influence declines with depth; in contrast, soil C quality has a minor influence on Q_{10} in shallow soil but increases its influence with depth. When considering the depth-dependent Q_{10} variations, results further show that using the thermal response of shallow soil layer for the whole soil profile, as is usually done in model predictions, would significantly underestimate soil C-climate feedbacks. The results highlight that Earth system models need to consider multilayer soil C dynamics and their controls to improve prediction accuracy.

future dynamics of soil C can substantially affect not only the climate but also soil fertility.^[3] Despite their low C concentrations, deep soil horizons contain more than half of global soil organic C (SOC) stocks,^[4] and thus may be even more important in terms of influencing atmospheric CO₂ concentrations than shallow soil C.^[5] Although shallow soil C responses to climate change have been relatively well studied,^[6–8] major questions remain unsolved regarding dynamics in deep soil C in response to climate change.^[5,9] Determining the thermal sensitivity of SOC in deep soil horizons is an important step toward predicting contributions of soil to global C cycle and potential feedbacks to climate change.^[8,10–13] A comprehensive analysis of the dynamics of deep soil C should help improve accuracy and precision in modeling feedbacks between climate and the global C cycle.^[14–16]

The thermal sensitivity of SOC in deep soil horizons has received increasing attention over recent years.^[10,17–19] A recent deep soil warming experiment in a

1. Introduction

Soils store at least three times as much carbon (C) as is found either in the atmosphere or in living plants,^[1] making them a huge potential source or sink for atmospheric C.^[2] The


temperate forest ecosystem showed that all soil depths had similar temperature sensitivities in response to warming;^[10] that study described “apparent” temperature sensitivity which is constrained by field conditions (e.g., soil moisture and root growth).^[20–23] Given that many Earth system models (ESMs),

Dr. J. Li, J. Pei, Prof. B. Li, Prof. C. Fang, Prof. M. Nie
Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering
Coastal Ecosystems Research Station of the Yangtze River Estuary
School of Life Sciences
Fudan University
Shanghai 200438, P. R. China
E-mail: mnjie@fudan.edu.cn

Prof. E. Pendall, Prof. P. B. Reich, Dr. N. J. Noh
Hawkesbury Institute for the Environment
Western Sydney University
Penrith NSW 2751, Australia

Prof. P. B. Reich
Department of Forest Resources
University of Minnesota
St. Paul, MN 55108, USA

Dr. N. J. Noh
Forest Technology and Management Research Center
National Institute of Forest Science
Pocheon 11186, Republic of Korea

 The ORCID identification number(s) for the author(s) of this article can be found under <https://doi.org/10.1002/advs.202001242>

© 2020 The Authors. Published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

DOI: 10.1002/advs.202001242




Figure 1. Geographic and climatic distribution of experimental sites. a) The 90 sampling sites across China's forests, with numbers in parenthesis indicating the number of sampling sites and the mean value of the mean annual temperature of each biome type. b) The 90 sites represent a wide range of mean annual temperature and mean annual precipitation.

such as the CanESM2,^[24] HadGEM2-ES,^[25] and INMCM4.0^[26] require the temperature sensitivity which is the inherent property of SOC decomposition,^[23] information from laboratory incubation temperature response studies is irreplaceable.^[23,27]

Until now, however, results from various laboratory incubations have been highly controversial and contradictory regarding the depth-dependence of temperature sensitivity.^[19,28–34] We synthesized data from 31 published laboratory incubation experiments of 98 soil profiles, and found no general pattern of temperature sensitivity variations with soil depth (see Supplementary Text and Figure S1, Supporting Information). The lack of consistency in individual outcomes could largely be explained by three possible reasons: 1) soil properties changed gradually and nonlinearly with depth (e.g., SOC and soil pH), yet ~66% of these soil profiles included only two depths; 2) the methods used, such as temperature range during incubation and calculation of temperature sensitivity, were different, leading to a difficult comparison across studies; and 3) these studies focused mainly on individual sites, and thus differences among studies could be due to context-dependencies rather than represent any problem or conflict per se (see Supplementary Text, Supporting Information). In addition, factors regulating soil C-temperature response with soil depth over large scales have yet to be evaluated, which also adds uncertainties in predictions of soil C-climate feedbacks.^[8]

To address these knowledge gaps, we designed a laboratory incubation study using a uniform method with soils of 1-m depth divided into six layers from 90 upland forest sites, spanning large gradients of mean annual temperature (MAT) from −2.2 to 25.0 °C and mean annual precipitation (MAP) from 98 to 1888 mm across most of the major global forest biomes in China (Figure 1). Forests, covering ~30% of the Earth's land surface,^[35] store ~47% of the terrestrial SOC,^[4] and play a vital role in the global C cycle.^[36] Most of the major types of global forest biomes occur in China, covering from tropical to boreal forests,^[37] providing an ideal system for studying general patterns and controls

of the temperature sensitivity of SOC decomposition with soil depth on large geographical scales.

In this study, we defined the temperature sensitivity of SOC decomposition (expressed as Q_{10} , proportional change in decomposition rate for a 10 °C difference in temperature) as the change in decomposition rate with temperature under otherwise constant conditions.^[23] The short-term (hours to days) Q_{10} was estimated using a dynamic temperature ramping method to minimize the substrate effects.^[30,38] In addition, we considered climate (MAT and MAP), plant productivity (normalized difference vegetation index, NDVI), soil physical and chemical properties of clay content and pH, C quantity of SOC content, and C quality indicated by the ratio of carbohydrates to aromatics. We hypothesized that 1) the temperature sensitivity of SOC decomposition increases with soil depth across large geographical scales, and 2) climate factors primarily regulate shallow soil Q_{10} , while soil C quality plays the most important role in deep soil.

2. Results and Discussion

2.1. The Temperature Sensitivity with Soil Depth

Temperature sensitivity of SOC decomposition increased with increasing soil depths across the broad geographical scale (Figure 2, $P < 0.001$). Because global temperatures are projected to increase approximately 3 °C by the end of the 21st century,^[39] we calculated Q_{10} under a 3 °C range—that was Q_{10} was calculated on the basis of the fitted decomposition rates (incubated under the same temperature range of 4–28 °C with a step of 4 °C) at MAT and MAT + 3 °C of each site (which is more ecologically relevant than a single fixed reference temperature). Results showed that Q_{10} significantly increased with increasing soil depths but with greater variability in deep soil compared to shallow soil (Figure 2a, $P < 0.001$). Specifically, Q_{10} values were 3.21 ± 0.73 (Median \pm SD), 3.34 ± 0.71 , 3.56 ± 1.01 , 3.90 ± 1.12 , $4.21 \pm$




Figure 2. Temperature sensitivity increases with soil depth across China's forests. Box plots of a) the temperature sensitivity of soil organic carbon decomposition (Q_{10}) and b) activation energy (E_a) with soil depth. Lines in boxes represent median, left and right of boxes represent first and third quartiles; dots represent single observations. Linear mixed-effects models are used to evaluate the effect of soil depth on Q_{10} and E_a excluding auto-correlations of different sampling sites and depths, showing that Q_{10} and E_a significantly increase with soil depth ($P < 0.001$). $N = 90$ for each soil depth.




Figure 3. The temperature sensitivity of soil organic carbon decomposition (Q_{10}) with soil depth among biome types. Shaded areas indicate the 95% confidence intervals for each biome type. Linear mixed-effects models are used to evaluate the effect of soil depth on Q_{10} with excluding auto-correlations of different sampling sites and depths, showing that Q_{10} for each biome type significantly increases with soil depth ($P < 0.001$).

1.27, and 4.53 ± 1.38 at soil depths of 0–10, 10–20, 20–35, 35–50, 50–70, and 70–100 cm, respectively (Figure S2, Supporting Information). Similarly, a significant effect of soil depth was found on activation energy (E_a , which can be described as a small “push” needed to begin chemical reactions^[23]) (Figure 2b, $P < 0.001$). Q_{10} values across all sites at some fixed temperatures (e.g., Q_{10} at 15 °C was calculated on the basis of fitted decomposition rates at 15 and 15 + 3 °C) also significantly increased with soil depth (Figure S3, Supporting Information). In the present study, the reported Q_{10} value hereafter was standardized to site-specific MATs of each site unless otherwise specified. Furthermore, the pattern of increasing Q_{10} with soil depth was common for all biomes (Figure 3; and Figure S4, Supporting Information), and Q_{10} increased with soil depth to a larger extent in colder biomes than in warmer

biomes. Collectively, these results clearly show that the temperature sensitivity of SOC decomposition increased with soil depth across diverse forest ecosystems, demonstrating that deep soil C is at high risk of increased loss in the face of rising temperature.

2.2. Factors Regulating the Depth-Dependent Temperature Sensitivity

Climate, plant productivity, clay content, soil pH, SOC quantity, and soil C quality are recognized as important factors to potentially control the heterogeneity of Q_{10} value.^[38,40–42] We conducted boosted regression tree (BRT) analyses^[43,44] to identify the relative contributions of all factors considered in explaining Q_{10} at each soil depth. Our comprehensive study showed that Q_{10} values were primarily regulated by climatic factors in shallow soil, while they were mainly influenced by climate and C quality in deep soil (Figure 4). This is because the effect of climate on soil properties is depth dependent, with stronger effect on topsoil than deep soil.^[4]

Across all soils, we found that Q_{10} was significantly and negatively correlated to soil C quality indicated by the ratio of carbohydrates to aromatics along soil depth (Figure S5, Supporting Information), in agreement with the C quality-temperature (CQT) hypothesis, which suggests decomposition of higher quality C has lower Q_{10} than that of lower quality C.^[23] Results from our experiment are consistent with other studies supporting the CQT hypothesis,^[42,45,46] but suggest that soil C quality plays a less important role in shallow soil than in deep soil. In addition, although the relative contributions of clay to Q_{10} was small in shallow soil, the importance increased with soil depth (Figure 4), which might be strongly related to clay associated SOC in microaggregates in deep soil compared to shallow soil.^[5] The greater variability in Q_{10} in deep soil might be attributable to the




Figure 4. Changes in relative contributions (%) of factors in explaining the temperature sensitivity of SOC decomposition (Q_{10}) with soil depth based on boosted regression tree analysis. The relative contributions of climate are the sum of relative contribution of mean annual temperature and the relative contribution of mean annual precipitation. SOC, soil organic carbon; NDVI, normalized difference vegetation index.

high variability in clay compared to shallow soil (Figure S6, Supporting Information), which could be potentially incorporated into models to better predict the spatial variation in temperature sensitivity in deep soil.

2.3. Implications for Soil C Cycling

Deep soil horizons contain large quantities of sequestered organic C^[4] and are projected to warm at roughly the same rate as shallow soils over the next century.^[10] Thus, any increase in deep soil C mineralization rates with increasing temperature could, over time, have significant effects on global C dynamics. However, published ESMS have typically used the Q_{10} value of shallow soil for the whole-soil profile^[11,47] due to the lack of data for Q_{10} in deep soil horizons. Here, we used two model scenarios (with vs. without consideration of depth-associated Q_{10} variations) to predict SOC stock across China's forests under a gradual increase of 3.0 °C by the end of this century (Figure 5): a single-layer model treating the soil profile as a homogeneous unit and using the Q_{10} value of the surface layer (0–10 cm) for the whole unit, and a six-layer model using depth-associated Q_{10} values for each soil layer. Acknowledging a wide range of potential changes in C inputs,^[20] the models were fit under three arbitrary scenarios of the temperature sensitivity of C input rate (hereafter, Q_{10} of C input) (i.e., Q_{10} value of C input 50% lower, equal to, or 50% higher than that of SOC decomposition^[48]). These three scenarios were not intended to represent predicted futures, but instead to provide a visualization of how variation in soil temperature sensitivity with depth would impact soil C cycling across the entire plausible range of inputs and outputs. Under predicted global warming, simulation results showed that ignoring the depth-associated variation in the thermal response of SOC decomposition (i.e., single-layer modeling) would significantly underestimate the release of soil C across China's forests at any rate of SOC sequestration (~2.1–




Figure 5. Modeling the stock of soil organic carbon (SOC) in the top 1-m depth subjected to a gradual increase of 3.0 °C over the period 2018–2100. Two alternatives are compared (with vs. without depth consideration): using Q_{10} value (the temperature sensitivity of SOC decomposition) of the top layer (0–10 cm) for the single-layer model or using Q_{10} values of six depths for the six-layer model. Meanwhile, these two alternatives are calculated with the asynchrony of the temperature responses of SOC sequestration rate (hereafter, Q_{10} value of C input) and output (i.e., Q_{10} value of C input is 50% lower, equal to, or 50% higher than that of SOC decomposition). All scenarios show that, compared to the six-layer model, the single-layer model would significantly underestimate the C-climate feedback intensity to temperature increase. Details of the calculations are shown in Table S1 (Supporting Information).

2.4% of the initial SOC stock depending on C input scenarios, see Figure 5). This would amount to ~15.5–17.7 Pg C emission underestimated from global forest soils, equivalent to about 2.0 times the annual C emissions from fossil fuel globally in 2017.^[49]

2.4. Uncertainty Analyses

In most incubation studies, soil samples are separately incubated at different but constant temperatures.^[50,51] In such cases, the differential depletion of C pools^[30] and microbial adaptation^[52] to different temperatures may make it difficult to derive an accurate estimation of temperature sensitivity. Our dynamic temperature ramping method is usually completed within 1 or 2 weeks to ensure soil conditions (e.g., substrates and microbial communities) relatively stable,^[19,38,40,53] and thus decomposition rates might more closely reflect the actual soil C dynamics as temperatures increase. In addition, this approach captures the temperature sensitivity of labile C decomposition on a short-term basis. Similarly, in field conditions, the short-term temperature sensitivity of soil respiration reflects the more labile C component due to continual input of new C to soil.^[20] However, understanding long-term adjustments^[54,55] and the temperature response of recalcitrant C pool decomposition^[18,30] would be necessary as a complement to the short-term responses, in order to fully quantify soil C responses to temperature increase across global forests.

Uncertainties existed in the effects of disturbances (e.g., soil sampling, sieving, and aerobic incubation) on SOC decomposition especially for deep soil horizons. Although soil sampling and sieving may damage the physical integrity of soil samples relating to the location of SOC within the soil matrix,^[56]

sieving removes autotrophic C sources (e.g., roots) and previous studies found that the effects of soil sieving on Q_{10} can be neglected if soil samples are sieved under field-moist conditions.^[57] Thus, we sieved soil samples immediately after transporting to the lab. Moreover, most laboratory incubations are conducted under aerobic conditions, i.e., sometimes at higher oxygen availability compared to in situ soil environment. High oxygen availability may accelerate SOC decomposition differentially for samples from different depths, but is unlikely to change the ranking of their Q_{10} values, except for sites where deep soils are systematically and consistently either more anaerobic or more aerobic than shallow soil horizons, which is unlikely to be a general pattern across our 90 sites of upland forest soils. All soil profiles in the present study were collected above the water table (soil gravimetric water content < 41% for all sites; Table S2, Supporting Information), thus *in situ*, aerobic decomposition rather than anaerobic decomposition would dominate across all depths.^[58,59] Therefore, changes in oxygen availability during soil incubation were unlikely to obscure the main finding of this study that the short-term Q_{10} values increased significantly with soil depth.

3. Conclusions

Our findings provide strong empirical evidence that under rising temperatures, SOC in deep soil is likely more vulnerable to loss than that in shallow soil. This vulnerability is not yet a component of ESMs, which could be an important source of uncertainty in predictions of the global terrestrial C cycle. Given the urgent need to accurately quantify and predict future soil C-climate feedbacks, our documentation of a predictable variation in the temperature sensitivity with depth across all major forest biomes represents a valuable step toward better quantifying the response of the huge but largely ignored deep soil C to temperature increase over broad geographical scales in a fashion that allows its integration into the current framework of global modeling. Although decomposition rates are likely to be very low in deep soil horizons and cold environments, our work shows that the huge deep SOC stocks are particularly vulnerable to warming, especially in cold biomes. Thus, even a slight variation in Q_{10} there can significantly influence both the atmospheric CO₂ concentration and C cycling. Moreover, Q_{10} values were primarily regulated by climate in shallow soil, while they were mainly controlled by climate and soil C quality in deep soil. The high Q_{10} value in deep soil is attributable to the low C quality, supporting the CQT hypothesis. Finally, long-term and multi-site warming experiments are required to study the apparent temperature sensitivity with soil depth in the future, especially for highly-sensitive temperate and boreal ecosystems.

4. Experimental Section

Study Area and Soil Sampling: In this study, a total of 90 sites, representing a diverse array of soil and site characteristics, were sampled across China's forests (Figure 1). These sites were all from upland forests, and site information (e.g., geographic and climatic information, dominant tree species, and some soil properties) are given in Table S2 (Supporting Information). These sites spanned large gradients of MAT from -2.2 to 25.0 °C, MAP from 98 to 1888 mm, and altitude from 6 to 3720 m above sea level. In general, the MAT and MAP decrease from south to north and from east to west across the sampling sites.

Soils were collected from three random locations (separated by more than 20 m) at each site in 2016–2017. The seasonal effects were not considered on Q_{10} but with the aim to reveal the vertical patterns and controls of Q_{10} with soil depth across China's forests. However, the phenology of C inputs derived from root exudates should be acknowledged and leached dissolved organic C might have some potential effects on the short-term Q_{10} determination. After removing the surface litter, soils of 1-m profile were collected at depths of 0–10, 10–20, 20–35, 35–50, 50–70, and 70–100 cm. Soil samples were then thoroughly homogenized into one composite sample by depth at each site (total of 540 soil samples, 90 sites × 6 depths). Immediately after transporting to the lab, the composite soil samples were passed through a 2-mm mesh under field-moist conditions. Approximately 50 g of homogenized soil was air-dried for physical and chemical properties (e.g., soil texture, SOC, and pH), and the rest was kept at 4 °C for incubation experiments.

Climatic Variables, Plant Productivity, and Soil Properties: Climate variables including MAT and MAP were derived from the WorldClim database.^[60] The NDVI was used, from the MODIS aboard NASA's Terra satellites, as a proxy of plant productivity.^[61] The mean value of NDVI for each site based on the monthly NDVI between the periods of 2016–2017 with 0.1° resolution was calculated.

Soil pH was measured in water solution with 1:2.5 soil:water ratio. SOC content was determined using a TOC analyzer (Vario TOC Cube, Germany) after the removal of carbonates with 1 M HCl. Soil texture (i.e., clay, silt, and sand content) was measured using a particle size analyzer (Laser Particle Sizer, LS-CWM(2), OMEC, China) after the removal of organic matter and carbonates.^[40] Soil water holding capacity (WHC) was gravimetrically determined.^[33]

Soil C quality was determined using Fourier-transform infrared spectroscopy (FT-IR, Nicolet iS5, Thermo Scientific, USA).^[62] Air-dried soil was sieved to 0.15 mm and was further dried at 60 °C and homogenized by grinding with an agate mortar and pestle. Reflectance spectra (400–4000 cm⁻¹) were obtained using a FT-IR and relative peak areas were calculated as the area of a distinct reflectance peak. Relative peak areas could reflect the relative abundance of different organic C functional groups such as carbohydrates (1024 cm⁻¹) and aromatic C groups (1637 cm⁻¹).^[63] and a higher ratio of carbohydrates to aromatics is considered as higher C quality.^[64]

Soil Incubation and Q_{10} Value Determination: Details of soil incubation and Q_{10} value determination could be found in Li et al.^[38] Briefly, 50 g (dry weight) fresh soils were adjusted to 60% WHC (under aerobic conditions), which is considered as optimal for microbial respiration,^[65,66] and incubated in 250 mL jars with four experimental replicates. Soils were preincubated at 20 °C for 7 days to activate microorganisms and minimize the possible disturbances (e.g., soil sieving). Soils were then incubated in a water bath to conduct the sequential incubation under 4–28 °C with a step of 4 °C. After being changed to a new temperature, the soils were kept for 3 h to obtain a new equilibrium stage. Following that, two headspace gas samples of 5 mL were collected before and after sealing for a period of time (1–48 h depending on soil depth and incubation temperature), and gas samples were analyzed using a gas chromatograph (Agilent 6890; Agilent Corp.). The rate of SOC decomposition at each temperature was calculated on the basis of soil weight, net CO₂ accumulation in the headspace, sealing time (respiration time), and the headspace volume.^[38,67]

A quadratic temperature response function could well fit the responses of decomposition rate to temperature change (Figure S7, Supporting Information). The fitting coefficient R^2 was higher than 0.98 for all soil samples. The quadratic temperature response function takes the form

$$R = \gamma_0 + \gamma_1 T + \gamma_2 T^2 \quad (1)$$

where R is SOC decomposition rate ($\mu\text{g C g}^{-1} \text{ soil h}^{-1}$), T is incubation temperature (°C), γ_0 , γ_1 , and γ_2 are fitted parameters.

In this study, Q_{10} value was estimated at site-specific MATs of each site.^[38] These values provide estimates of the change in respiration that would occur with climate warming if no long-term adjustments (e.g., plant phenology, soil moisture, substrate availability, biotic physiological acclimation, and biotic community compositional shift) occurred. Because

global temperatures are projected to increase approximately 3 °C by the end of this century,^[39] Q_{10} under a 3 °C range was calculated and then fixed to a 10 °C interval based on the definition of Q_{10} —that was Q_{10} was calculated based on the fitted decomposition rates at MAT and MAT + 3 °C of each site using the following function

$$Q_{10} = (R_2/R_1)^{10/(T_2-T_1)} \quad (2)$$

where R_1 and R_2 are decomposition rates fitted at $T_1 = \text{MAT}$ and $T_2 = \text{MAT} + 3$ °C, respectively.

In addition, the activation energy (E_a) was estimated by fitting the SOC decomposition rate to the Arrhenius Equation^[45]

$$k = Ae^{-\frac{E_a}{RT}} \quad (3)$$

where E_a is the activation energy (kJ mol⁻¹), k is SOC decomposition rate (μg C g⁻¹ soil h⁻¹), A is a fitted constant, R is the gas constant (8.314 J K⁻¹ mol⁻¹), and T is temperature in Kelvin.

Modeling Descriptions: Two model scenarios (with vs. without consideration of depth-associated Q_{10} variations) were used to predict SOC stock across China's forests under a gradual increase of 3.0 °C over the period 2018–2100. It was assumed that soil C output is equal to that of input in the first year of 2018, and both input and output are only affected by temperature (i.e., Q_{10} value of C input is 50% lower, equal to, or 50% higher than that of SOC decomposition). Soil C balance for each soil layer is calculated as

$$\Delta SS_i = C_{\text{input}_i} - C_{\text{output}_i} \quad (4)$$

$$C_{\text{output}_i} = D_{\text{SOC}_i} \times Q_{10,i}^{(\Delta T_i/10)} \quad (5)$$

$$C_{\text{input}_i} = C_{\text{output}_i} \times (n \times Q_{10,i})^{(\Delta T_i/10)} \quad (6)$$

where ΔSS_i is soil C balance for the i th ($i = 1, 2, 3, 4, 5,$ and 6) layer, C_{input_i} and C_{output_i} are soil C input and output for the i th layer, respectively, D_{SOC_i} is SOC decomposability per unit SOC for the i th layer, $Q_{10,i}$ is the temperature sensitivity of SOC decomposition for the i th layer (for the single-layer model, Q_{10} value of the top layer (0–10 cm) is used for each layer; for the six-layer model, Q_{10} values of six layers are used), and ΔT_i is temperature increasing for the j th year; n is the ratio of Q_{10} value of C input to Q_{10} value of SOC decomposition (C output).

The weighted soil C balance for the whole soil profile is calculated as

$$\Delta SS = \frac{\sum_{i=1}^n \Delta SS_i \times H_i \times B_i \times C_i}{\sum_{i=1}^n H_i \times B_i \times C_i} \quad (7)$$

where ΔSS is the weighted soil C balance for the top 1-m depth, and H_i , B_i , and C_i are the height, bulk density, and SOC density of each soil layer, respectively.

Soil C balance for each year is then calculated as

$$SS_j = SS_{j-1} + \Delta SS \quad (8)$$

where SS_j and SS_{j-1} are soil C balance for the j th and $(j-1)$ th year, respectively. Details of the calculations are presented in Table S1 (Supporting Information).

Statistical Analyses: BRT analyses^[43,44] were conducted to identify the relative contributions of all the considered predictors on Q_{10} at each soil depth. Data were normalized (log-transformed if needed) before doing BRT analyses. BRT is applicable to nonlinear relationships, and can be used to analyze different types of variables (predictors) and their interactive effects.^[44] The relative contributions of climate are the sum of relative

contribution of MAT and relative contribution of MAP. The BRT analyses were performed using the package *gbm* in R (version 3.4.2).

Linear mixed-effects models (LMEMs) were conducted to analyze the effect of soil depth on Q_{10} to exclude spatial autocorrelations of different sampling sites and soil depths. LMEMs were performed using the *nlme* package in R (version 3.4.2) with soil depth as a fixed effect and two random effects, including sampling site and the random slope between target variable (e.g., Q_{10}) and soil depth. Statistical analyses and correlation analyses were performed using SPSS Statistics 22 (IBM) or R (version 3.4.2).

Supporting Information

Supporting Information is available from the Wiley Online Library or from the author.

Acknowledgements

They thank Eric A. Davidson and Chao Liang for insightful discussions during the preparation of the manuscript; they also thank Zhen Li, Hui Guo, Xiao Xu, and Huimin Sun for the help with measurements and analyses; they also appreciate two anonymous reviewers for their valuable comments on the manuscript. This work was supported by the National Science Foundation of China (Nos. 91951112 and 31930070), the Postdoctoral Science Foundation of China (2020M670975), and the Australian Research Council (No. DP170102766).

Conflict of Interest

The authors declare no conflict of interest.

Author Contributions

J.L., C.F., and M.N. designed the study; J.L. conducted the overall experiment and measurements with the assistance from J.P.; C.F. and M.N. supervised the experiment and measurements; J.L. analyzed the data with the assistance from E.P., P.B.R., J.P., N.J.N., B.L., C.F., and M.N.; J.L. and M.N. wrote the first draft, and all authors jointly revised the manuscript.

Keywords

carbon decomposition, deep soil, forest ecosystems, global warming, temperature sensitivity

Received: April 4, 2020
Revised: June 29, 2020
Published online: August 6, 2020

- [1] R. Lal, *Science* **2004**, *304*, 1623.
- [2] B. Bond-Lamberty, V. L. Bailey, M. Chen, C. M. Gough, R. Vargas, *Nature* **2018**, *560*, 80.
- [3] H. Tiessen, E. Cuevas, P. Chacon, *Nature* **1994**, *371*, 783.
- [4] E. G. Jobbágy, R. B. Jackson, *Ecol. Appl.* **2000**, *10*, 423.
- [5] C. Rumpel, I. Kögel-Knabner, *Plant Soil* **2011**, *338*, 143.
- [6] T. W. Crowther, K. E. O. Todd-Brown, C. W. Rowe, W. R. Wieder, J. C. Carey, M. B. Machmuller, B. L. Snoek, S. Fang, G. Zhou, S. D. Allison, J. M. Blair, S. D. Bridgham, A. J. Burton, Y. Carrillo, P. B. Reich, J. S. Clark, A. T. Classen, F. A. Dijkstra, B. Elberling, B. A. Emmett, M. Estiarte, S. D. Frey, J. Guo, J. Harte, L. Jiang, B. R. Johnson, G. Kröel-

- Dulay, K. S. Larsen, H. Laudon, J. M. Lavalley, et al., *Nature* **2016**, 540, 104.
- [7] N. van Gestel, Z. Shi, K. J. van Groenigen, C. W. Osenberg, L. C. Andresen, J. S. Dukes, M. J. Hovenden, Y. Luo, A. Michelsen, E. Pendall, P. B. Reich, E. A. G. Schuur, B. A. Hungate, *Nature* **2018**, 554, E4.
- [8] M. W. I. Schmidt, M. S. Torn, S. Abiven, T. Dittmar, G. Guggenberger, I. A. Janssens, M. Kleber, I. Kögel-Knabner, J. Lehmann, D. A. C. Manning, P. Nannipieri, D. P. Rasse, S. Weiner, S. E. Trumbore, *Nature* **2011**, 478, 49.
- [9] J. A. Mathieu, C. Hatté, J. Balesdent, É. Parent, *Global Change Biol.* **2015**, 21, 4278.
- [10] C. E. Hicks Pries, C. Castanha, R. C. Porras, M. S. Torn, *Science* **2017**, 355, 1420.
- [11] K. E. O. Todd-Brown, J. T. Randerson, W. M. Post, F. M. Hoffman, C. Tarnocai, E. A. G. Schuur, S. D. Allison, *Biogeosciences* **2013**, 10, 1717.
- [12] J. Balesdent, I. Basile-Doelsch, J. Chadoeuf, S. Cornu, D. Derrien, Z. Fekiacova, C. Hatté, *Nature* **2018**, 559, 599.
- [13] C. D. Koven, D. M. Lawrence, W. J. Riley, *Proc. Natl. Acad. Sci. USA* **2015**, 112, 3752.
- [14] H. Q. Tian, C. Q. Lu, J. Yang, K. Banger, D. N. Huntzinger, C. R. Schwalm, A. M. Michalak, R. Cook, P. Ciais, D. Hayes, *Global Biogeochem. Cycles* **2015**, 29, 775.
- [15] N. Carvalhais, M. Forkel, M. Khomik, J. Bellarby, M. Jung, M. Migliavacca, M. Mu, S. Saatchi, M. Santoro, M. Thurner, *Nature* **2014**, 514, 213.
- [16] Y. Luo, A. Ahlström, S. D. Allison, N. H. Batjes, V. Brovkin, N. Carvalhais, A. Chappell, P. Ciais, E. A. Davidson, A. Finzi, K. Georgiou, B. Guenet, O. Hararuk, J. W. Harden, Y. He, F. Hopkins, L. Jiang, C. Koven, R. B. Jackson, C. D. Jones, M. J. Lara, J. Liang, A. D. McGuire, W. Parton, C. Peng, J. T. Randerson, A. Salazar, C. A. Sierra, M. J. Smith, H. Tian, et al., *Global Biogeochem. Cycles* **2016**, 30, 40.
- [17] T. Roy Chowdhury, E. M. Herndon, T. J. Phelps, D. A. Elias, B. Gu, L. Liang, S. D. Wullschlegel, D. E. Graham, *Global Change Biol.* **2015**, 21, 722.
- [18] S. Q. Qin, L. Y. Chen, K. Fang, Q. W. Zhang, J. Wang, F. T. Liu, J. C. Yu, Y. H. Yang, *Sci. Adv.* **2019**, 5, eaau1218.
- [19] J. Li, D. Yan, E. Pendall, J. Pei, N. J. Noh, J. He, B. Li, M. Nie, C. Fang, *Soil Biol. Biochem.* **2018**, 126, 82.
- [20] P. Smith, C. Fang, J. J. C. Dawson, J. B. Moncrieff, *Adv. Agron.* **2008**, 97, 1.
- [21] M. Nie, E. Pendall, C. Bell, C. K. Gasch, S. Raut, S. Tamang, M. D. Wallenstein, *Ecol. Lett.* **2013**, 16, 234.
- [22] J. Li, E. Pendall, F. A. Dijkstra, M. Nie, *Soil Tillage Res.* **2020**, 199, 104574.
- [23] E. A. Davidson, I. A. Janssens, *Nature* **2006**, 440, 165.
- [24] P. R. Gent, G. Danabasoglu, L. J. Donner, M. M. Holland, E. C. Hunke, S. R. Jayne, D. M. Lawrence, R. B. Neale, P. J. Rasch, M. Vertenstein, *J. Clim.* **2011**, 24, 4973.
- [25] C. Jones, J. Hughes, N. Bellouin, S. Hardiman, G. Jones, J. Knight, S. Liddicoat, F. O'Connor, R. J. Andres, C. Bell, *Geosci. Model. Dev.* **2011**, 4, 543.
- [26] E. Volodin, N. Dianskii, A. Gusev, *Izv. Atmos. Oceanic Phys.* **2010**, 46, 414.
- [27] K. Karhu, M. D. Auffret, J. A. J. Dungait, D. W. Hopkins, J. I. Prosser, B. K. Singh, J.-A. Subke, P. A. Wookey, G. I. Ågren, M.-T. Sebastià, F. Gouriveau, G. Bergkvist, P. Meir, A. T. Nottingham, N. Salinas, I. P. Hartley, *Nature* **2014**, 513, 81.
- [28] N. Fierer, A. S. Allen, J. P. Schimel, P. A. Holden, *Global Change Biol.* **2003**, 9, 1322.
- [29] C. Moni, T. Z. Lerch, K. Knoth de Zarruk, L. T. Strand, C. Forte, G. Certini, D. P. Rasse, *Soil Biol. Biochem.* **2015**, 88, 236.
- [30] C. M. Fang, P. Smith, J. B. Moncrieff, J. U. Smith, *Nature* **2005**, 433, 57.
- [31] P. Wordell-Dietrich, A. Don, M. Helfrich, *Geoderma* **2017**, 304, 40.
- [32] E. Hilasvuori, A. Akujärvi, H. Fritze, K. Karhu, R. Laiho, P. Mäkiranta, M. Oinonen, V. Palonen, P. Vanhala, J. Liski, *Soil Biol. Biochem.* **2013**, 67, 47.
- [33] J. Li, J. Pei, J. Cui, X. Chen, B. Li, M. Nie, C. Fang, *Agric., Ecosyst. Environ.* **2017**, 250, 44.
- [34] C. Mu, T. Zhang, X. Zhang, B. Cao, X. Peng, *Eur. J. Soil Sci.* **2016**, 67, 773.
- [35] G. B. Bonan, *Science* **2008**, 320, 1444.
- [36] Y. D. Pan, R. A. Birdsey, J. Fang, R. Houghton, P. E. Kauppi, W. A. Kurz, O. L. Phillips, A. Shvidenko, S. L. Lewis, J. G. Canadell, P. Ciais, R. B. Jackson, S. W. Pacala, A. D. McGuire, S. Piao, A. Rautiainen, S. Sitch, D. Hayes, *Science* **2011**, 333, 988.
- [37] J. Y. Fang, A. P. Chen, C. H. Peng, S. Q. Zhao, L. J. Ci, *Science* **2001**, 292, 2320.
- [38] J. Li, M. Nie, E. Pendall, P. B. Reich, J. Pei, N. J. Noh, T. Zhu, B. Li, C. Fang, *Global Change Biol.* **2020**, 26, 1873.
- [39] T. M. L. Wigley, S. C. B. Raper, *Science* **2001**, 293, 451.
- [40] J. Z. Ding, L. Y. Chen, B. B. Zhang, L. Liu, G. B. Yang, K. Fang, Y. L. Chen, F. Li, D. Kou, C. J. Ji, Y. Q. Luo, Y. H. Yang, *Global Biogeochem. Cycles* **2016**, 30, 1310.
- [41] J. Li, J. Pei, E. Pendall, C. Fang, M. Nie, *Soil Biol. Biochem.* **2020**, 141, 107675.
- [42] Q. K. Wang, S. G. Liu, P. Tian, *Global Change Biol.* **2018**, 24, 2841.
- [43] J. Elith, J. R. Leathwick, T. Hastie, *J. Anim. Ecol.* **2008**, 77, 802.
- [44] Z. K. Luo, W. T. Feng, Y. Q. Luo, J. Baldock, E. L. Wang, *Global Change Biol.* **2017**, 23, 4430.
- [45] J. M. Craine, N. Fierer, K. K. McLaughlan, *Nat. Geosci.* **2010**, 3, 854.
- [46] N. Fierer, B. P. Colman, J. P. Schimel, R. B. Jackson, *Global Biogeochem. Cycles* **2006**, 20, n/a.
- [47] C. D. Koven, B. Ringeval, P. Friedlingstein, P. Ciais, P. Cadule, D. Khorostyanov, G. Krinner, C. Tarnocai, *Proc. Natl. Acad. Sci. USA* **2011**, 108, 14769.
- [48] Z. Wu, P. Dijkstra, G. W. Koch, J. Peñuelas, B. A. Hungate, *Global Change Biol.* **2011**, 17, 927.
- [49] C. Le Quéré, R. M. Andrew, P. Friedlingstein, S. Sitch, J. Pongratz, A. C. Manning, J. I. Korsbakken, G. P. Peters, J. G. Canadell, R. B. Jackson, T. A. Boden, P. P. Tans, O. D. Andrews, V. K. Arora, D. C. E. Bakker, L. Barbero, M. Becker, R. A. Betts, L. Bopp, F. Chevallier, L. P. Chini, P. Ciais, C. E. Cosca, J. Cross, K. Currie, T. Gasser, I. Harris, J. Hauck, V. Haverd, R. A. Houghton, et al., *Earth Syst. Sci. Data* **2017**, 10, 405.
- [50] R. T. Conant, R. A. Drijber, M. L. Haddix, W. J. Parton, E. A. Paul, A. F. Plante, J. Six, J. M. Steinweg, *Global Change Biol.* **2008**, 14, 868.
- [51] S. Hamdi, F. Moyano, S. Sall, M. Bernoux, T. Chevallier, *Soil Biol. Biochem.* **2013**, 58, 115.
- [52] J. Å. M. Wetterstedt, T. Persson, G. I. Ågren, *Global Change Biol.* **2010**, 16, 1806.
- [53] Y. Liu, N. P. He, J. X. Zhu, L. Xu, G. R. Yu, S. Niu, X. M. Sun, X. F. Wen, *Global Change Biol.* **2017**, 23, 3393.
- [54] T. W. N. Walker, C. Kaiser, F. Strasser, C. W. Herbold, N. I. W. Leblans, D. Woebken, I. A. Janssens, B. D. Sigurdsson, A. Richter, *Nat. Clim. Change* **2018**, 8, 885.
- [55] J. C. Carey, J. Tang, P. H. Templer, K. D. Kroeger, T. W. Crowther, A. J. Burton, J. S. Dukes, B. Emmett, S. D. Frey, M. A. Heskell, L. Jiang, M. B. Machmuller, J. Mohan, A. M. Panetta, P. B. Reich, S. Reinsch, X. Wang, S. D. Allison, C. Bamminger, S. Bridgham, S. L. Collins, G. de Dato, W. C. Eddy, B. J. Enquist, M. Estiarte, J. Harte, A. Henderson, B. R. Johnson, K. S. Larsen, Y. Luo, et al., *Proc. Natl. Acad. Sci. USA* **2016**, 113, 13797.
- [56] J. A. Dungait, D. W. Hopkins, A. S. Gregory, A. P. Whitmore, *Global Change Biol.* **2012**, 18, 1781.
- [57] N. Meyer, G. Welp, W. Amelung, *Global Biogeochem. Cycles* **2018**, 32, 306.

- [58] J. Krista, F. Hannu, L. Jukka, L. Raija, *Global Change Biol.* **2007**, *13*, 491.
- [59] E. A. Davidson, S. Samanta, S. S. Caramori, K. Savage, *Global Change Biol.* **2012**, *18*, 371.
- [60] R. J. Hijmans, S. E. Cameron, J. L. Parra, P. G. Jones, A. Jarvis, *Int. J. Climatol.* **2005**, *25*, 1965.
- [61] M. Delgado-Baquerizo, F. T. Maestre, P. B. Reich, T. C. Jeffries, J. J. Gaitan, D. Encinar, M. Berdugo, C. D. Campbell, B. K. Singh, *Nat. Commun.* **2016**, *7*, 10541.
- [62] X. Hu, S. Liu, X. Li, C. Wang, X. Ni, X. Liu, Y. Wang, Y. Liu, C. Xu, *Sci. Rep.* **2019**, *9*, 8256.
- [63] S. B. Hodgkins, C. J. Richardson, R. Dommain, H. Wang, P. H. Glaser, B. Verbeke, B. R. Winkler, A. R. Cobb, V. I. Rich, M. Missilmani, N. Flanagan, M. Ho, A. M. Hoyt, C. F. Harvey, S. R. Vining, M. A. Hough, T. R. Moore, P. J. H. Richard, F. B. De La Cruz, J. Toufaily, R. Hamdan, W. T. Cooper, J. P. Chanton, *Nat. Commun.* **2018**, *9*, 3640.
- [64] M. M. Tfaily, W. T. Cooper, J. E. Kostka, P. R. Chanton, C. W. Schadt, P. J. Hanson, C. M. Iversen, J. P. Chanton, *J. Geophys. Res. Biogeosci.* **2014**, *119*, 661.
- [65] D. M. Howard, P. J. A. Howard, *Soil Biol. Biochem.* **1993**, *25*, 1537.
- [66] A. Rey, E. Pegoraro, P. G. Jarvis, *Eur. J. Soil Sci.* **2008**, *59*, 1049.
- [67] D. Yan, J. Li, J. Pei, J. Cui, M. Nie, C. Fang, *Sci. Rep.* **2017**, *7*, 5181.