



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

# Science of the Total Environment

journal homepage: [www.elsevier.com/locate/scitotenv](http://www.elsevier.com/locate/scitotenv)



## Observed and estimated prevalence of Covid-19 in Italy: How to estimate the total cases from medical swabs data

F. Bassi <sup>a,\*</sup>, G. Arbia <sup>b</sup>, P.D. Falorsi <sup>c</sup>

<sup>a</sup> Department of Statistical Sciences, University of Padova, Italy

<sup>b</sup> Department of Statistical Sciences, Catholic University of the Sacred Heart, Milano, Italy


<sup>c</sup> Italian National Statistical Institute, Italy

### HIGHLIGHTS

- There are evidences of a very high proportion of asymptomatic patients of Covid-19.
- An estimation of prevalence is extremely important.
- The sample survey conducted by the National Statistical Institute to estimate is encountering many difficulties.
- We estimate the prevalence of Covid-19 in Italy by reweighting the available official data on patients examined by nasopharyngeal swabs.
- We obtain an estimation of Covid-19 prevalence in Italy equal to 9%, corresponding to almost 6 million Italians.


### GRAPHICAL ABSTRACT

How many people have been affected by Covid-19 In Italy?



Estimation of prevalence from official data on nasopharyngeal swabs.

Applying post-stratification to the convenience sample



### ARTICLE INFO

**Article history:**

Received 1 August 2020  
 Received in revised form 29 September 2020  
 Accepted 29 September 2020  
 Available online 8 October 2020

Editor: Jay Gan

**Keywords:**

Covid-19  
 Prevalence  
 Lethality  
 Convenience sampling  
 Post-stratification

### ABSTRACT

During the Covid-19 pandemic in Italy, official data are collected with medical swabs following a pure convenience criterion which, at least in an early phase, has privileged the exam of patients showing evident symptoms. However, there are evidences of a very high proportion of asymptomatic patients. In this situation, in order to estimate the real number of infected (and to estimate the lethality rate), it should be necessary to run a properly designed sample survey through which it would be possible to calculate the probability of inclusion and hence draw sound probabilistic inference. Unfortunately, the survey run by the Italian Statistical Institute encountered many field difficulties. Some researchers proposed estimates of the total prevalence based on various approaches, including epidemiologic models, time series and the analysis of data collected in countries that faced the epidemic in earlier times. In this paper, we propose to estimate the prevalence of Covid-19 in Italy by reweighting the available official data published by the Istituto Superiore di Sanità so as to obtain a more representative sample of the Italian population. Reweighting is a procedure commonly used to artificially modify the sample composition so as to obtain a distribution which is more similar to the population. In this paper, we will use post-stratification of the official data, in order to derive the weights necessary for reweighting the sample results, using age and gender as post-stratification variables, thus obtaining more reliable estimation of prevalence and lethality. Specifically, for Italy, we obtain a prevalence of 9%. The proposed methodology represents a reasonable approximation while waiting for more reliable data obtained with a properly designed national sample survey and that it could be further improved if more data were made available.

© 2020 Elsevier B.V. All rights reserved.

\* Corresponding author.

E-mail address: [francesca.bassi@unipd.it](mailto:francesca.bassi@unipd.it) (F. Bassi).

### 1. Introduction and data issues

In the recent period of pandemic emergency, most statistical analyses on Covid-19 focused primarily on finding the best forecasting model to be able to anticipate the number of epidemic cases at a national and local level (see, e.g., Huang et al., 2020a and Ceylan, 2020, for a review), on the analysis of influencing factors (e.g. Lian et al., 2020) or on the evaluation of the effects of the lockdown measures (e.g. Huang et al., 2020b). Comparatively less attention has been paid in the literature to a detailed descriptive analysis of the publicly available data. In Italy, there are two major sources of the data on the diffusion of Covid-19: the Istituto Superiore di Sanità - Italian National Institute of Health (INIH) and the Dipartimento di Protezione Civile - Civil Protection Department (CPD). Both sources, at the moment, do not disclose individual data on the epidemic. In particular, CPD publishes daily the number of positive patients, of deaths and of individuals tested by swabs at national, and regional level. In addition, INIH releases a biweekly report with infections and deaths disaggregated by gender and age class at a national level. Positive cases were also disaggregated by age (but not by gender) in each of the 20 Italian regions; however, this information was available only till the end of May 2020. So both sources do not provide any information about the asymptomatic patients, thus limiting the usefulness of these data in view of calculating interesting epidemic parameters such as the *lethality rate*. However, a careful look at these data, especially if compared with the dimension and the demographic structure of the actual population of our country, might give important information to better understand the effects of the virus. The novelty of this paper lays in the fact that, without requesting different datasets collected on purpose for statistical reasons, we propose to exploit the existing available data in order to estimate the prevalence and the lethality of the virus in the total Italian population. In particular, we use the data published by the INIH and the CPD (henceforth the “official data”) with reference to the positive patients by Covid-19 and to the number of deaths at May 26, 2020 – unfortunately after this data the INIH stopped publishing the regional data of infections by age. Another key information to retrieve a representative sample is the proportion of infections among all performed nasopharyngeal swabs, that is available by region from the CPD website. The information on the demographic structure of the Italian population, refers instead, to 1 January 2019 and it is taken from the National Statistical Institute website ([www.istat.it/en](http://www.istat.it/en)). This paper proposes a strategy based on data collected on patients tested by nasopharyngeal swabs and post-stratification sampling techniques. The paper is organized as follows. Section 2 describes our proposed methodology while Section 3 is devoted describe the prevalence and the lethality rate as they emerge from the officially released data. Section 4 presents an estimation of the prevalence of Covid-19 in Italy based on the post-stratification. Section 5 concludes.

**Table 1**  
Positive patients, deaths and lethality rate by Covid-19 by gender and age, 26 May 2020, official data.  
Source: Official data by INIH.

Age	Positive patients				Deaths				Lethality rate		
	Male	Female	Total	Proportion of males	Male	Female	Total	Proportion of males	Male	Female	Total
0–9	1015	903	1918	52.92%	1	2	3	33.33%	0.10%	0.22%	0.16%
10–19	1724	1718	3442	50.09%	0	0	0	0.00%	0.00%	0.00%	
20–29	5685	7240	12,925	43.98%	6	3	9	66.67%	0.11%	0.04%	0.07%
30–39	8026	9902	17,928	44.77%	35	19	54	64.81%	0.44%	0.19%	0.30%
40–49	12,513	17,427	29,940	41.79%	184	62	246	74.80%	1.47%	0.36%	0.82%
50–59	19,210	22,221	41,431	46.37%	778	215	993	78.35%	4.05%	0.97%	2.40%
60–69	18,494	12,385	30,879	59.89%	2299	676	2975	77.28%	12.43%	5.46%	9.63%
70–79	19,033	14,107	33,140	57.43%	5566	2283	7849	70.91%	29.24%	16.18%	23.68%
80–89	16,497	24,027	40,524	40.71%	6593	4801	11,394	57.86%	39.96%	19.98%	28.12%
90+	3818	14,783	18,601	20.53%	1556	2873	4429	35.13%	40.75%	19.43%	23.81%
Total	106,015	124,713	230,728	45.95%	17,018	10,934	27,952	60.88%	16.05%	8.77%	12.11%

### 2. Method of investigation


Our estimation makes use of all available information on the number of positive patients and on swabs at national and regional level. As it is well known, these data suffer from the severe limitation of being observed without a proper sample design and so they can be seen as a convenience sample that cannot be used to draw probabilistic inference (Cochran, 1977). To try to reduce such distortion, we propose to post-stratify the convenience sample using gender and age as post-stratification variables in order to obtain a dataset which is closer to a representative sample of the Italian population, under some reasonable assumptions.

On 25 May 2020, the Italian Statistical Institute, in collaboration with the Italian Ministry for Health, started a sample survey to estimate the seroprevalence of Covid-19 in the country. The sample is composed 150,000 individuals living in 2015 Italian municipalities and it is representative of the Italian population by gender, age and economic activity sector. Selected people were contacted via telephone by the Italian Red Cross’s regional centers to arrange an appointment for a blood sample to be taken at one of the authorized laboratories and to answer some questions on health conditions, eventual previous infection by the virus, contacts with Coronavirus positive patients. The regional government should inform each participant residing in their territory of the result of the test. If the test was positive, the person in question should be put temporarily into isolation at home and contacted by the regional health service or the local health authority to do a nasopharyngeal swab to verify contagiousness.

Unlikely in other countries, where similar surveys were successful (see, as an example, the Spanish experience as in Pollán et al., 2020), in Italy some difficulties were encountered especially in obtaining respondents cooperation. The period of information collection was extended till 15 July. At the present moment, the National Statistical Institute released preliminary results that refer to the subsample 64,660 (less than half of the planned sample size) individuals who were tested by the serological exam before July 27 (<https://www.istat.it/it/files/2020/08/ReportPrimiRisultatiIndagineSiero.pdf>). This result suggests that, at least for Italy, some other statistical methods might be useful in order to estimate the diffusion of the Covid-19 infection at country level.

### 3. A descriptive analysis of the prevalence and deaths in Italy

Table 1 reports the absolute frequencies of infected people and of deaths for Covid-19 distinguishing by gender and age (10-year classes). The gender ratio by age is also reported. Table 1 as well contains the implied lethality rate measured as the ratio between the number of deaths and the number of infected people. This information, published twice a week by the INIH, shows the way in which positive cases are distributed in the various demographic groups. Fig. 1 reports the observed



**Fig. 1.** Prevalence in the Italian population by gender and age, 26 May 2020. Source: Official data by INIH.

prevalence rates, i.e., the proportion of infected referring to the consistency of each class in the Italian population (as of 1 January 2019), measured in each stratum as the ratio between the number of infected and the total susceptible population.

Fig. 1 clearly shows that prevalence in the sample increases with age for both genders; more specifically, this is true for people younger than 59 years and for those older than 90, while for the other age classes we observe an opposite trend. Overall, the prevalence is higher for women than for men. This is a rather new evidence. On April 92,020, for example, the prevalence in Italy was higher for males (0.24% vs. 0.21%) while on April 16, it was equal for the two genders (0.29%). An explanation of this result could be that in the last days observed in our reference period, even patients with light symptoms were tested and resulted positive to the infection and women showing less severe symptoms were examined in a larger percentage. This hypothesis is in accordance with the fact that the preliminary results from the sample survey conducted by Italian Statistical Institute did not find a statistically significant difference in the prevalence of Covid-19 among genders; a similar evidence was obtained in Spain. In the absence of conclusive results, it is worthwhile to monitor accurately this phenomenon over time to verify if this tendency is confirmed.

Another interesting result emerging from the official data, is that women with an age between 60 and 80 show a lower risk of infection than women at least 10 years younger; this is another interesting evidence that deserves further investigation. Two explanations could be valid for this evidence. Between 50 and 60 years old women experience menopause, maybe there are some effects due to hormonal change; this is a hypothesis that we, as statisticians, offer to medical doctors. A second reason of lower infections in these ages could be that women are less exposed to the virus either because usually retire from work earlier than men or because of differences in behavior, i.e., they try to avoid risky situations.

The composition indices reported in Table 2 compare the distribution of patients by gender and age (represented in Fig. 2a) with the distribution of the same variables in the Italian population (Fig. 2b). The indices are calculated by the ratio of the relative frequency in the sample of positive patients and the relative frequency in the Italian population.<sup>1</sup> A composition index lower than 1 indicates that individuals in that class are present among positive patients in a lower proportion with respect to the whole population. In contrast, a value greater than 1 provides an opposite indication. If we compare the composition indexes in the same age group between the two gender we find a confirmation of the evidences shown in Fig. 1.

<sup>1</sup> For example, the composition index for the subgroup of males between 0 and 9 years is obtained dividing the relative frequency of this class among positive patients, 0.01 (1015/106,015), by the relative frequency in the Italian population, 0.09 (2,627,094/29,384,766), obtaining the index equal to 0.1075.


**Table 2** Composition indices for patients and deaths with reference to the Italian population, 26 May 2020. Source: Official data by INIH.

Age	Patients		Deaths	
	Male	Female	Male	Female
0-9	0.1075	0.1012	0.1044	0.0007
10-19	0.1603	0.1708	0.1654	0.0000
20-29	0.4905	0.6714	0.5777	0.0032
30-39	0.6250	0.7808	0.7024	0.0170
40-49	0.7550	1.0390	0.8979	0.0692
50-59	1.1629	1.2902	1.2279	0.2934
60-69	1.4600	0.8972	1.1665	1.1306
70-79	1.9345	1.2085	1.5406	3.5243
80-89	3.2756	3.0838	3.1591	8.1550
90+	5.0555	7.2496	6.6566	12.8350


Finally, Fig. 3 reports the observed mortality rates referring to the consistency of each class in the Italian population (as of 1 January 2019) and it clearly shows that the proportion of deaths increases with age. In this case, the situation is much worse for males than for females in all age classes (see also composition indices in Table 2).

**4. Post-stratified estimation of prevalence lethality of Covid-19 in Italy**

In the reference period of our analyses, various guidelines have been followed to test individuals with swab to ascertain the presence of Covid-19 infection. In the early phase of the epidemic, the World Health Organization (WHO) recommended to test only patients with at least



**Fig. 2.** a. Age and gender distribution of Covid-19 positive patients on 26 May 2020. Source: Official data by INIH b. Age and gender distribution in the Italian population, 1 January 2019. Source: Official data by NSI.



**Fig. 3.** Proportion of deaths in the Italian population by gender and age, 26 May 2020. Source: Official data by INIH.

three specific symptoms and reporting contact with the infection. In a second moment of time, since a large proportion of positive patients did not report any symptom, these guidelines were relaxed and even people with light symptoms (or only with contact with other infected patients), were examined. Moreover, the procedures to access to swab could be different in the 20 Italian regions. These considerations, together with the evidences found in the literature of a high proportion of asymptomatic patients (e.g. Aguilar et al., 2020; Chughtai and Malik, 2020; Li et al., 2020; Mizumoto et al., 2020a, 2020b; Yelin et al., 2020), suggest that prevalence in the Italian population could be much higher than what appears from the official data.

In order to estimate the total number of infected by Covid-19, it should be necessary to run a properly designed sample survey through which it is possible to calculate the probability of inclusion and hence draw sound probabilistic inference. Such a survey has high costs and it needs time to be realized. While waiting for the results of such a desirable survey, some researchers proposed various approaches to estimate the prevalence. They are based on epidemiologic models, time series models and the analysis of data collected in countries that faced the epidemic before like for Italy and Europe countries, for example China and Korea (see, e.g., Brogi et al., 2020).

In this paper, we propose to approximate the true prevalence of Covid-19 in Italy by exploiting the official data published by the INIH (Istituto Superiore di Sanità, 2020a, 2020b) and the CPD, but reweighting them so as to obtain something closer to a representative sample of the Italian population. The procedure of reweighting (Valliant et al., 2018) is used when, in most cases due to measurement error, a sample is not representative of the reference population. More specifically, reweighting is a procedure to artificially modify the sample composition, in the phase of data analysis, so as to obtain a distribution which is closer to the population. In its simplest form, reweighting assigns appropriate weights to each sample unit where weights can be defined on the basis of the inclusion probability (if known) or on the basis of available information on the population. In this case, we call it post-stratification (Holt and Smith, 1979; Little, 1993). In this last case, after choosing one or more stratification variables, whose distribution is known in the population, the sample units are weighted with the ratio between the theoretical proportion in the population and the observed proportion in the sample. In this paper, we will use post-stratification to analyze the official data sample, using age and gender proportion in the Italian population as post-stratification variables.

In our analysis, in order to correct the observed sample, we need to introduce some working assumptions which are required because all disaggregated information is not available. Indeed, as already said, data by gender and age are available only for positive patients, and not for those that are negative at the swab. As a consequence, in order to simulate the characteristics of all patients tested by nasopharyngeal swab till 26 May 2020, we proceeded as follows.

First of all, we considered the positive patients disaggregated into age classes in each of the 20 Italian regions, an information provided twice a week by the INIH (Istituto Superiore di Sanità, 2020b). Furthermore, the total number of people subjected to pharyngeal swab was derived using the information provided by the percentage of positive tests in each region supplied by the CPD. Lacking the appropriate age disaggregation, we assumed that this percentage is constant in all age classes. The simulated sample of patients subjected to swabs is then distributed into the two genders by using the sex ratios observed among positive patients (see Table 1, column 5).

Secondly, we calculated the post-stratification weights referring to the distribution by age and gender of the Italian population at the last available date which was January 1st, 2019.

Finally, we re-estimated the prevalence after post-stratification.

The aggregate out-coming value from this operation is prevalence estimated equal to 9.22%, a number that, reported to the total population of Italy, reveals that 5,985,537 people could have been affected by Covid-19 in the country as of 26 May 2020. Furthermore, lethality rate is also re-estimated on the post-stratified sample leading to a rate of 0.47% and the median age of positive patients of 52 years.

Table 3 reports the estimated prevalence by age in the Italian population and the consequent estimated lethality rate. Even if it is known that the vast majority of swabs were obtained from symptomatic, this estimate of prevalence partly corrects for all those patients who have not been subjected to pharyngeal swab for various reasons, mainly for presenting light symptoms (pauci-symptomatic) or for being asymptomatic.

By comparing our estimation with the current estimates based on unweighted (official) data, we observe that prevalence in age classes is much higher than that calculated with official data. This result shows that there might be a much higher proportion of people infected than those measured with test, possibly due to the fact that many positive patients do not even know of their condition because they have very light symptoms or no symptoms at all (Lavezzo et al., 2020). Our estimates of prevalence have the obvious effect to produce a much lower lethality rate in all age classes with respect to that calculated with the uncorrected official data. For what concerns the lethality rate, we calculated it reporting the number of deaths in each age class to the number of estimated infections. We assume that deaths caused by Coronavirus are all correctly reported.

### 5. Summary of results and concluding remarks

In this report, we analyze data on Covid-19 infections in Italy with reference to the consistency and demographic structure of the Italian population. Looking at the official data, published by the INIH and by the CPD, in general for male patients the risk of infection increases with age. In contrast, the dynamics of infection is peculiar for female patients: it increases until the age of 59, then it decreases until 79 and it

**Table 3**  
Estimated prevalence and lethality in the Italian population by age after post-stratification.

Age	Prevalence			Lethality over 100,000		
	Male	Female	Total	Male	Female	Total
0–9	8.54%	6.85%	7.72%	0.45	1–18	0.76
10–19	7.59%	7.04%	7.32%	0.00	0.00	0.00
20–29	8.17%	9.94%	9.03%	2.29	1.01	1.61
30–39	9.08%	10.45%	9.76%	10.83	5.17	7.82
40–49	8.86%	11.49%	10.18%	45.23	11.61	26.15
50–59	10.03%	10.68%	10.36%	169.45	42.18	102.49
60–69	13.30%	8.26%	10.67%	492.17	213.97	379.93
70–79	14.29%	9.53%	11.71%	1428.68	740.04	1124.36
80–89	10.30%	13.44%	12.21%	4586.75	1654.17	2625.49
90+	5.24%	14.90%	12.29%	14,180.61	3411.68	4653.13
Total	9.91%	9.92%	9.92%	584.25	355.84	466.99



increases again for older women. Furthermore, the observed prevalence is higher for females than for males until the age of 59. After that age women show a lower prevalence and the distance from males increases with age. Finally, until the beginning of April 2020, men represented the majority among positive patients while afterwards the proportion of women was continuously increasing. The dynamic of the infection by Covid-19 in female patients deserves further study, both in terms of observation over time and with epidemiologic and statistical analyses.

Secondly, starting from official data on Covid-19, we estimate the prevalence of the infection in the Italian population assuming that only a small proportion of patients could access to pharyngeal swab test. Lacking adequate data disaggregation (for example the number of patients subjected to swab by gender and age) we had to make some working assumptions to which the results obtained are strongly depending. In particular, we are well aware of the fact that the sampling of throat swabs is not random and that tested people were mostly those who present symptoms.

However, even with the current limitations, we believe that our methodology represents a reasonable approximation while waiting for more reliable data obtained with a properly designed national sample survey and that it could be further improved if more data were made available. In particular, it would be important to have the availability of disaggregated data, at least for gender and age classes, so as to avoid to impose our restrictive hypotheses of uniformity and to obtain better estimates of prevalence and lethality in each specific group of the population. This would enable us to identify the categories that are more exposed to the risk of infection and to support a system of active surveillance also in the period of recession of the pandemic.

#### CRediT authorship contribution statement

The paper is the product of the joint work of the three authors.

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### References

Aguilar, J. B., Faust, J. S., Westafer, L. M. and Gutierrez, J. B. (2020) Investigating the Impact of Asymptomatic Carriers on COVID-19, medRxiv, doi: <https://doi.org/10.1101/2020.03.18.20037994>.

- Brogi, F., Guardabascio, B., Barcaroli, G. (2020) Covid-19 in Italy: Actual Infected Population, Testing Strategy and Imperfect Compliance, DOI: [10.13140/RG.2.2.18275.50729](https://doi.org/10.13140/RG.2.2.18275.50729).
- Ceylan, G., 2020. Estimation of COVID-19 prevalence in Italy, Spain, and France. *Science of the Total Environment* <https://doi.org/10.1016/j.scitotenv.2020.138817>.
- Chughtai, A.A., Malik, A.A., 2020. Is coronavirus disease (COVID-19) case fatality ratio underestimated? *Global Biosecurity* 1 (3).
- Cochran, W.G., 1977. *Sampling Techniques*. Wiley, New York.
- Holt, D., Smith, T.M.F., 1979. Post stratification. *Series A* 142 (1), 33–46.
- Huang, Z., Huang, J., Gu, Q., et al., 2020a. Optimal temperature zone for the dispersal of COVID-19. *Sci. Total Environ.* <https://doi.org/10.1016/j.scitotenv.2020.139487>.
- Huang, J., Zhang, L., Liu, X., et al., 2020b. Global prediction system for COVID-19 pandemic. *Science Bulletin.* <https://doi.org/10.1016/j.scib.2020.08.002>.
- Istituto Superiore di Sanità (2020a) COVID-19 Epidemic. 26 May 2020, National Update. Istituto Superiore di Sanità (2020b) COVID-19 Epidemic. 26 May 2020, National Update (Appendix).
- Lavezzo, E., Franchin, E., Ciavarella, C., Cuomo-Dannenburg, G., Barzon, L., Del Vecchio, C., Rossi, L., Manganelli, R., Lorigian, A., Navarin, N., Abate, D., Sciro, M., Merigliano, S., Decanale, E., Vanuzzo, M.C., Saluzzo, F., Onelia, F., Pacenti, M., Parisi, S., Carretta, G., Donato, D., Flor, L., Cocchio, S., Masi, G., Sperduti, A., Cattarino, L., Salvador, R., Gaythorpe, K.a.M., Brazzale, A.R., Toppo, S., Trevisan, M., Baldo, V., Donnelly, C.A., Ferguson, N.M., Dorigatti, I., Crisanti A., 2020. Suppression of COVID-19 outbreak in the municipality of Vo, Italy. *Nature* <https://doi.org/10.1038/s41586-020-2488-1>.
- Li, R., Pei, Sen, Chen, Bin, Song, Yimeng, Zhang, Tao, Wan, Yang, Shaman, Jeffrey, 2020. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science*, eabb3221 <https://doi.org/10.1126/science.abb3221> (16 Mar 2020).
- Lian, X., Huang, J., Huang, R., et al., 2020. Impact of city lockdown on the air quality of COVID-19-hit of Wuhan city. *Sci. Total Environ.* <https://doi.org/10.1016/j.scitotenv.2020.140556>.
- Little, R.J.A., 1993. Post-stratification: a modeler's perspective. *J. Am. Stat. Assoc.* 88 (423), 1001–1012. <https://doi.org/10.1080/01621459.1993.10476368>.
- Mizumoto, K., Kagaya, K., Zarebski, A., Chowell, G., 2020a. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro surveillance: bulletin European sur les maladies transmissibles = European communicable disease bulletin* 25 (10), 2000180. <https://doi.org/10.2807/1560-7917.ES.2020.25.10.2000180>.
- Mizumoto, K., Katsushi, K., Zarebski, A., Gerardo, 2020b. Estimating the Asymptomatic Proportion of 2019 Novel Coronavirus Onboard the Princess Cruises Ship, 2020. medRxiv (doi:10.1101/2020.02.20.20025866doi).
- Pollán, M., Pérez-Gómez, B., Pastor-Barriuso, R., Oteo, J., Hernán, M.A., Pérez-Olmeda, M., Sanmartín, J.L., Fernández-García, A., Cruz, I., Fernández de Larrea, N., Molina, M., Rodríguez-Cabrera, F., Martín, M., Merino-Amador, P., Paniagua, J.L., Muñoz-Montalvo, J.F., Blanco, F., Yotti, R., 2020. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *Lancet* [https://doi.org/10.1016/S0140-6736\(20\)31483-5](https://doi.org/10.1016/S0140-6736(20)31483-5).
- Valliant, R., Dever, J.A., Kreuter, F., 2018. *Practical Tools for Designing and Weighting Survey Samples*. Springer, New York.
- Yelin, I., Noga Aharon, Einat Shaer-Tamar, Amir Argoetti, Esther Messer, Dina Berenbaum, Einat Shafran, Areen Kuzli, Nagam Gandali, Tamar Hashimshony, Yael Mandel-Gutfreund, Michael Halberthal, Yuval Geffen, Moran Swarcwort-Cohen, Roy Kishony (2020) Evaluation of COVID-19 RT-qPCR Test in Multi-sample Pools, medRxiv, 27 March, 2020. doi:<https://doi.org/10.1101/2020.03.26.20039438>.