


RESEARCH PAPER

 OPEN ACCESS 

CRISPR-*cas* system in the acquisition of virulence genes in dental-root canal and hospital-acquired isolates of *Enterococcus faecalis*

Pourya Gholizadeh^a, Mohammad Aghazadeh^b, Reza Ghotaslou^b, Mohammad Ahangarzadeh Rezaee^b, Tahereh Pirzadeh^b, Şükran Köse^c, Khudaverdi Ganbarov^d, Mehdi Yousefi^{e,f}, and Hossein Samadi Kafil ^a

^aDrug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, I.R. Iran; ^bImmunology Research Center, Tabriz University of Medical Sciences, Tabriz, I.R. Iran; ^cDepartment of Infectious Diseases and Clinical Microbiology, University of Health Sciences, Tepecik Training and Research Hospital, Izmir, Turkey; ^dDepartment of Microbiology, Baku State University, Baku, Azerbaijan; ^eBiotechnology Research Center, Tabriz University of Medical Sciences, Tabriz, I.R. Iran; ^fStem Cell Research Center, Tabriz University of Medical Sciences, Tabriz, I.R. Iran

ABSTRACT

Enterococcus faecalis is one of the important causative agents of nosocomial and life-threatening infections in human. Several studies have demonstrated that the presence of CRISPR-*cas* is associated with antibiotic susceptibility and lack of virulence traits. In this study, we aimed to assess the phenotypic and genotypic virulence determinants in relation to CRISPR elements from the dental-root canals and hospital-acquired isolates of *E. faecalis*. Eighty-eight hospital-acquired and 73 dental-root canal isolates of *E. faecalis* were assessed in this study. Phenotypic screening of the isolates included biofilm formation, and gelatinase and hemolysis activities. Genotypical screening using PCR was further used to evaluate the presence of CRISPR elements and different virulence-associated genes such as *efaA*, *esp*, *cylA*, *hyl*, *gelE*, *ace*, *ebpR*, and *asa1*. Biofilm formation, gelatinase, and hemolysis activities were detected in 93.8%, 29.2%, and 19.2% of the isolates, respectively. The most prevalent virulence-associated gene was *ace*, which was followed by *efaA*, whereas *cylA* was the least identified. The presence of CRISPR1-*cas*, orphan CRISPR2, and CRISPR3-*cas* was determined in 13%, 55.3%, and 17.4% of the isolates, respectively. CRISPR elements were significantly more prevalent in the dental-root canal isolates. An inverse significant correlation was found between CRISPR-*cas* loci, *esp*, and *gelE*, while direct correlations were observed in the case of *cylA*, *hyl*, *gelE* (among CRISPR-loci 1 and 3), *asa1*, *ace*, biofilm formation, and hemolysis activity. Findings, therefore, indicate that CRISPR-*cas* might prevent the acquisition of some respective pathogenicity factors in some isolates, though not all; so selective forces could not influence pathogenic traits.

Abbreviations: BHI: brain-heart infusion agar; CRISPRs: Clustered regularly interspaced short palindromic repeats; Esp: Cell wall-associated protein; ENT: ear-nose-throat; ICU: intensive care units; OD: optical densities; PCR: polymerase chain reaction; SDS: sodium dodecyl sulfate; UTI: urinary tract infection

ARTICLE HISTORY

Received 6 June 2020
Revised 23 July 2020
Accepted 8 August 2020

KEYWORDS

Enterococcus faecalis;
CRISPR-Cas system; virulence genes; phenotypic characteristics; hospital-acquired bacteria; dental-root canal bacteria

Introduction

Enterococcus faecalis is a Gram-positive natural inhabitant of the mammalian digestive tract, including those of humans. It is also found in soil, plants, and dairy food products [1]. *E. faecalis* also behaves as an opportunistic pathogen causing life-threatening infections in humans, such as endocarditis, meningitis, septicemia, urinary tract infections, and others [2,3]. *E. faecalis* is one of the frequent isolates of the endodontic pathogens ranging in terms of prevalence from 30% to 90% of the cases [4,5]. The restriction system of *E. faecalis* enables the bacterium to acquire, accumulate, and further transfer genetic elements potentially encoding antibiotic resistance genes and virulence factors. These virulence

factors include exoenzymes and adhesins. Cytolysin is encoded by *cyl* operon, which is carried by a plasmid or integrated into the chromosome, with both hemolysin and bacteriocin activity [6,7]. Gelatinase is encoded by the chromosomal *gelE* gene, which is a zinc metalloprotease; it can hydrolyze gelatin, fibrinogen, collagen, casein, and insulin [8]. Another secreted factor is hyaluronidase, which is encoded by the *hyl* gene [7]. *E. faecalis* endocarditis antigen is encoded by the *efaA* gene that affects pathogenicity [9]. Cell wall-associated protein (Esp), encoded by pheromone-responsive plasmids or the chromosomal *esp* gene, is involved in biofilm formation and immune evasion [9]. Aggregation substance, encoded by the *thea1* gene on the

CONTACT Hossein Samadi Kafil  Kafilhs@tbzmed.ac.ir

sex pheromone-plasmid pAD1, is a surface-bound glycoprotein which mediates the conjugative transfer of plasmids through the clumping of one *E. faecalis* to another and induces the formation of the cell-cell contact [10]. In addition, *ebpR* encodes an endocarditis- and biofilm-associated pilus regulator, which activates the *ebpABC* operon [11]. Another adhesion factor is a collagen-binding protein encoded by the *ace* gene, which mediates binding to collagen type I, collagen type IV, and laminin [12].

Clustered regularly interspaced short palindromic repeats (CRISPRs) loci and CRISPR-associated (Cas) protein-encoding genes are present in approximately 45% of eubacterial genomes sequenced [13–15]. There are three types of CRISPR loci in *E. faecalis* genome: CRISPR1-*cas*, orphan CRISPR2, and CRISPR3-*cas* [16–18]. CRISPR1-*cas* and orphan CRISPR2 were first found in the *E. faecalis* OG1RF strain: CRISPR1 is located between the OG1RF homolog of EF0672 and EF0673, which has the associated *cas* genes. CRISPR2 is located between the OG1RF homolog of EF2062 and EF2063, which is an orphan consisting only of spacers and palindromes, without any *cas* genes [16]. CRISPR3 was found in two genomes of the strains Fly1, as a fruit fly *E. faecalis*, and T11, as a urine *E. faecalis* isolate. CRISPR3 is located between the homologs of the *E. faecalis* V583 open reading frames EF1760 and EF1759 [18]. CRISPR1 possesses Nmeni subtype-specific genes *csn1* and *csn2* [16,18], while CRISPR3 only possesses *csn1*, not *csn2* [18]. Both CRISPR1 and CRISPR2 contain seven repeats of a 37 bp palindromic sequence with no homology to any sequences of the 29 bp spacer [16]. Nevertheless, due to small spacer sequences, it is likely that they are derived from the pheromone-responsive type plasmids, plasmids integrated within the *E. faecalis* V583 genome, and Enterococcal prophage and phage [18]. Recent studies have demonstrated that the CRISPR/Cas system has applications for genome engineering and exerts a strong selective pressure for the acquisition of virulence factors and antibiotic resistance in pathogenic bacteria [18–21]. Mojica et al., for instance, have suggested that the pathogenicity of bacteria is largely controlled by conjugative plasmids and bacteriophages on an evolutionary timescale. As well, those CRISPR spacers that target these mobile elements might affect bacterial pathogenicity and virulence traits [22].

In this study, we aimed to assess the phenotypic and genotypic virulence determinants in relation to CRISPR elements from the dental-root canals and hospital-acquired isolates of *E. faecalis*.

Methods and materials

Bacterial strains

This study was approved by the Regional Ethics Committee of Tabriz (Tabriz University of Medical Sciences, Tabriz, Iran, No. IR.TBZMED.REC.1397.188). A total of 88 isolates of *E. faecalis* were collected from EmamReza Teaching and Treatment Hospital and pediatric hospitals of Tabriz, Iran. The specimen sources of hospital-acquired isolates included urinary tract infection (UTI) (78, 88.6%), wound (7, 7.9%), and blood (3, 3.4%). The specimens were obtained from different wards including outpatients (35, 39.8%), intensive (23, 26.1%), intensive care units (ICU) (12, 13.6%), infectious ward (13, 14.8%), emergency ward (3, 3.4%), ear-nose-throat (ENT) (1, 1.1%), urology, and nephrology (1, 1.1%). Forty-two (47.7%) isolates were from male and 46 (52.3%) were from female cases. The age range of patients was from 2 months to 86 y, with a mean of 39.04 y. At the same time, in order to collect 73 dental-root canal isolates of *E. faecalis*, patients in need of endodontic treatment were referred to the clinic of the Faculty of Dentistry at Tabriz University of Medical Sciences, Tabriz, Iran. Forty-nine (67.1%) of the isolates were obtained from the males and 24 (32.9%) from the females. The age range of endodontic treatment patients was 12–66 y, with a mean of 32.41 y. Briefly to collect the isolates, after stages of access cavity preparation by the dentist, tooth, and its surroundings were washed by sterile saline solutions and disinfected with 30% hydrogen peroxide followed by 2.5% sodium hypochlorite. Root canal of teeth with no prior endodontic treatment and teeth with previous root canal treatment that showed secondary infection was removed by drill and endodontic K-files without using any chemical solvents. After sampling the single root canal and multi-root canal of the teeth, paper points were transferred to a tube containing Enterococcal broth (Becton Dickenson microbiology systems, Cockeysville, MD) and cultured on a bile esculin azide agar (Himedia, India) and incubated at 37°C for 24–48 h [4]. Suspected colony was identified by the standard procedures of microbiology [23,24] and genotype detection was performed by *ddlE* primer [25,26], as shown in Table 1. Both clinical and tooth identified isolates for further studies were stored in a trypticase soy broth containing 10% glycerol at –70°C.

Biofilm formation

Assessment of biofilm formation was done by quantitative biofilm formation in 96-well flat-bottom polystyrene

Table 1. Primers used for the detection of virulence genes and CRISPR-associated genes.

Gene	Primer	Sequence (5'–3')	PCR product length (bp)	References
<i>esp</i>	espF	GGAACGCCTTGGTATGCTAAC	95	[46]
	espR	GCCACTTTATCAGCCTGAACC		
<i>cylA</i>	cylF	ACTCGGGGATTGATAGGC	688	[47]
	cylR	GCTGCTAAAGCTGCGCTT		
<i>hyl</i>	hylF	ACAGAAGAGCTGCAGGAAATG	276	[11]
	hylR	GACTGACGTCCAAGTTTCCAA		
<i>efaA</i>	efaF	TGGGACAGACCTCACGAATA	101	[48]
	efaR	CGCCTGTTTCTAAGTTCAAGCC		
<i>gelE</i>	gelF	TATGACAATGCTTTTGGGAT	213	[47]
	gelR	AGATGCACCCGAAATAATATA		
<i>ace</i>	aceF	GGAGAGTCAAATCAAGTACGTTGGTT	101	[49]
	aceR	TGTTGACCACTTCCTTGTGAT		
<i>ebpR</i>	ebpF	AAAAATGATTCGGCTCCAGAA	101	[11]
	ebpR	TGCCAGATTGCTCTCAAAG		
<i>asa1</i>	asaF	GCACGCTATTACGAACTATGA	375	[47]
	asaR	TAAGAAAGAACATCACACGA		
CRISPR1- <i>cas csn1</i>	For	CAGAAGACTATCAGTTGGTG	783	[18]
	Rev	CCTTCTAAATCTTCTCATAG		
CRISPR1- <i>cas</i> loci	For	GCGATGTTAGCTGATACAAC	315	[18]
	Rev	CGAATATGCCTGTGGTAAA		
CRISPR2 loci	For	CTGGCTCGCTGTACAGT	variable	[18]
	Rev	GCCAATGTTACAATACAAACA		
CRISPR3- <i>cas csn1</i>	For	GCTGAATCTGTGAAGTACTC	258	[18]
	Rev	CTGTTTTGTTACCGTTGGAT		
CRISPR3- <i>cas</i> loci	For	GATCACTAGGTTCAAGTTATTC	224	[18]
	Rev	CATCGATTATTCTCCAA		

microplates under static conditions for 48 h, as previously described [27,28]. Briefly, for each isolate, a fresh colony cultured on a Muller-Hinton agar (Merck, Germany) containing 1% glucose was suspended in sterile saline and adjusted to 0.5 McFarland. Twenty microliters of the adjusted isolates was cultured in a 180- μ l trypticase soy broth containing 1% glucose. After incubation for 48 h at 37°C, each well was washed by the 1X phosphate buffer saline (PBS; pH 7.4), fixed by methanol, and stained by 200 μ l 0.1% crystal violet for 30 min at room temperature. The excess crystal violet was discarded and washed by water flow. Biofilm formation was measured by the absorbance of the supernatant after being solubilized in 33% acetic acid at 570 nm by using a microtiter plate reader (BioTeck, Winooski, USA). The biofilm formation of each isolate was tested in three independent 96-well microplates and the average of three optical densities (OD) was used as the final biofilm formation value. The cutoff absorbance for biofilm formation was considered higher than OD = 0.524, which was the absorbance of the biofilm produced by *E. faecalis* ATCC® 29,212™. The mean of the biofilm formation of each isolate was grouped based on their level of distribution (OD_{570nm} values) and categorized in quartiles higher than the cutoff absorbance and lower than the highest absorbance. Isolates whose absorbance of OD_{570nm} fell below 0.524 were classed as non-biofilm formation, while those with 0.525–1.087 and 1.088–1.650 were grouped as low and moderate biofilm formation, respectively. Isolates with a biofilm formation greater than 1.651 were also considered with high biofilm formation.

Gelatinase production and hemolysis test

Hemolysis activity was assessed by blood agar plates prepared by brain–heart infusion agar (BHI, biomérieux, Poland, Ltd) containing 5% of the group ORh⁺ human blood. Cleared or green zone around the colonies was defined as hemolysis following incubation for 24 h at 37°C [29].

Production of gelatinase was assessed by the degradation of gelatin on the X-ray radiographic film, as described by Pickett et al. [30]. The heavy inoculum of individual isolates was cultured in the tubes containing 3 ml MHB and a strip of the X-ray radiographic film which had been cut into small strips (approximately 6 by 30 mm). The tubes were incubated for 24 h at 37°C and the cleared strip was defined as the production of gelatinase.

Genotype detection of virulence and *cas* genes

Total DNA for each isolate was extracted by the tissue buffer boiling method. Briefly, 20 μ l tissue buffer (0.25% sodium dodecyl sulfate (SDS) and 0.05 M NaOH) were mixed with one colony of bacterial isolate and incubated at 95°C for 10 min. The suspension was centrifuged at 13,000 g for 1 min, and 180 μ l DNase free water was added. Genotype analysis for each isolate was accomplished based on the multiplex polymerase chain reaction (PCR) of virulence determinants encoding the cytolysin activator *cylA*, *hyl*, *esp*, *gelE*, *efaA*, *asa1*, *ace*, *ebpR*, CRISPR1-*cas*, CRISPR1-*cascsn1*,

CRISPR2, CRISPR3-*cas*, and CRISPR3-*cascsn1*. Each of the primer sequences and the amplified size are shown in Table 1. Two microliters of total DNA was used for the multiplex PCR in a 25 µl reaction mixture. The mix for the detection of *esp*, *cyl*, *hyl* genes contained 12.5 µl of the PCR master mix (Yekta Tajhiz Azma, Iran), with 0.5 µM of each primer. The mix for *ebp*, *asa1*, and *efaA* had the same condition. The mix for the detection of *gelE* and *ace* contained 12.5 µl of the PCR master mix (Yekta Tajhiz Azma, Iran), 1.5 mM-additional MgCl₂ and 0.5 µM of each primer. The mix for CRISPR1-*cascsn1*, CRISPR3-*cascsn1*, CRISPR1-*cas*, CRISPR3-*cas*, and CRISPR2 contained 12.5 µl of the PCR master mix (Yekta Tajhiz Azma, Iran), 1 mM additional MgCl₂, and 10 mM of each primer. The amplification condition was carried out with the following thermal cycling conditions: initial denaturation at 95°C for 10 min, 34 cycles of amplification consisting of 95°C for 30 s, 30 s at 58°C for *esp*, *cylA*, *hyl*, 58°C for *efaA*, 56°C for *gel*, *ace*, 52°C for *ebpR*, *asa1*, 60°C for all *cas* genes, and 72°C for 45 s, with 72°C for 5 min in the final polymerization. PCR products were analyzed by electrophoresis in a 1% agarose gel at 100 V for 1 h in a 1X TBE buffer containing the DNA safe stain. The size of the PCR product was correlated with a 100 based-pair DNA ladder (YektaTajhizAzma, Iran) to confirm the conjunction with their expected PCR amplicon size. In addition, the PCR procedure for each isolate was carried out twice in the case of each primer in order to check the consistency and reproducibility.

Statistical analysis

SPSS software, version 17.0 (Chicago, IL, USA) was used for statistical analysis. One-tailed Fisher's exact test was used to compare the occurrence of CRISPR-*cas* loci in hospital-acquired and dental-root canal

isolates and to evaluate the distribution of biofilm formation, gelatinase and hemolysin activities, and virulence genes among strains with CRISPR-*cas*. Student's *t*-test was used to compare OD values among hospital-acquired and dental-root canal isolates. In addition, Spearman's rank correlation was calculated between the presence of different virulence genes and CRISPR-*cas* loci among isolates. Significance was set at $P \leq 0.05$.

Results

All isolates were investigated for the biofilm formation, in which the minimum, maximum, and average of biofilm formation (OD570 nm) were 0.054, 2.325, and 1.611, respectively. Most isolates showed strong biofilm formation (94, 58.4%), while 10 (6.2%) displayed no biofilm formation. Biofilm formation of hospital-acquired isolates was significantly higher than the dental-root canal isolates ($P = 0.023$). The biofilm formation absorbance according to the presence of virulence factors and CRISPR loci among *E. faecalis* isolates shown in Figure 1. Most of the isolates showed no gelatinase activity (70.8%), while hospital-acquired isolates significantly displayed the most gelatinase activity ($P = 0.001$). In addition, most isolates showed no hemolysis activity (80.7%), and all hemolysis activity was found in hospital-acquired isolates (19.2%). The most presence of the virulence genes among isolates were *ace* and *efaA* genes (88.8% and 85.1%, respectively), and the lowest one belonged to *cylA* and *asa1* (7.5% and 14.9%, respectively). The presence of *gelE* (contributing to gelatinase activity) and *cylA* (contributing to hemolysis activity) was significantly associated with phenotype gelatinase and hemolysis activity, respectively ($P < 0.001$, $P = 0.013$). In addition, the presence of *efaA*, *cylA*, and *gelE* was significantly more in hospital-acquired isolates, as compared to dental-root canal




Figure 1. Biofilm formation absorbance by *E. faecalis* isolates according to the presence of virulence factors and CRISPR loci. (Error bars illustrate the minimum and maximum of three replicates of absorbance of the biofilm formation; * P -value was significant (P -value < 0.05).)

($P = 0.002$, $P < 0.001$ and $P = 0.008$, respectively). Genotypic, and phenotypic determinants of hospital-acquired and dental-root canal isolates are shown in Table 2. The *efaA* and *gelE* harboring isolates had a higher biofilm formation than negative isolates in all isolates ($P = 0.017$ and $P = 0.042$, respectively). The biofilm formation absorbance association to virulence genes and CRISPR loci among *E. faecalis* isolates is shown in Figure 2. By comparing the presence of virulence genes among isolates, it was found that hospital-acquired isolates had higher virulence genes than dental-root canal isolates ($P = 0.007$), such that all isolates had at least one virulence gene. The distribution of virulence gene counts among *E. faecalis* isolates is presented in Figure 3. The number of virulence genes was 1–7 among hospital-acquired isolates and 16 in the case of dental-root canal isolates. Among hospital-acquired isolates, the presence of five and four virulence genes was the highest (36.4% and 30.7%, respectively); also, the presence of 4 and 3 virulence genes was the highest among isolates of the dental-root canal (39.7% and 31.5%, respectively).

The occurrence of CRISPR-*cas* is shown in Table 3. Overall, the presence of CRISPR1-*cas* loci in dental-root canal isolates (4 of 73) was lower than that of hospital-acquired isolates (17 of 88) ($P = 0.008$), whereas the presence of CRISPR3-*cas* in dental-root canal isolates (26 of 73) was higher than that of hospital-acquired isolates (2 of 88) ($P < 0.001$); also, orphan CRISPR2 made no difference between hospital-acquired and dental-root canal isolates. None of the isolates had, however, both of CRISPR1-*cas* and CRISPR3-*cas*, as well as CRISPR1-*cas*, orphan CRISPR2, and CRISPR3-*cas*, at the same time. The isolates were more likely to harbor orphan CRISPR2 than CRISPR1-*cas* and CRISPR3-*cas*. In addition, the presence of orphan CRISPR2 was significantly correlated with CRISPR1-*cas* ($P = 0.031$, correlation coefficient = 0.163), whereas it was not significant with CRISPR3-*cas*. At least one CRISPR-*cas* locus was found in 106 (65.8%) of all isolates. The results, therefore, showed the isolates containing high virulence genes tended to have more frequently investigated *cas* genes. The presence of CRISPR1 and CRISPR 2 was significantly correlated with high distribution of virulence gene numbers ($P = 0.010$ and $P = 0.011$, respectively). The virulence gene counts association to CRISPR loci among *E. faecalis* isolates is shown in Figure 4. Overall, the absence of CRISPR1-*cas* and one of CRISPR1 or CRISPR3 were significantly correlated with the absence of the *esp* gene ($P = 0.005$, correlation coefficient = 0.204 and $P = 0.033$, correlation coefficient = 0.157, respectively). In addition, the presence

Table 2. Genotypic and phenotypic determinants of hospital-acquired and dental-root canal isolates.

Source	<i>esp</i>		<i>cylA</i>		<i>hyl</i>		<i>efaA</i>		<i>gelE</i>		<i>ace</i>		<i>ebpR</i>		<i>asaI</i>		Gelatinase activity			Hemolysis			Biofilm formation							
	P	(n)	P	(n)	P	(n)	P	(n)	P	(n)	P	(n)	P	(n)	P	(n)	Positive	(n)	α	β	γ	N	+	++	+++					
Hospital-acquired (88)	73.9%	(65)	13.6%	(12)	15.9%	(14)	93.2%	(82)	37.5%	(33)	88.6%	(78)	79.5%	(70)	15.9%	(14)	39.8%	(35)	29.5%	5.7%	64.8%	(57)	4.5%	(4)	6.8%	(6)	20.5%	(18)	68.2%	(60)
UTI's (78)	78.2%	(61)	15.4%	(12)	14.1%	(11)	93.6%	(73)	34.6%	(27)	89.7%	(70)	78.2%	(61)	16.7%	(13)	39.7%	(31)	29.5%	6.4%	64.1%	(50)	5.1%	(4)	3.8%	(3)	21.8%	(17)	69.2%	(54)
Non-UTI's (10)	40%	(4)	0	(0)	30%	(3)	90%	(9)	60%	(6)	80%	(8)	90%	(9)	10%	(1)	40%	(4)	30%	0	70%	(7)	0	30%	(3)	10%	(1)	60%	(6)	
Dental root (73)	68.5%	(55)	0	(0)	24.7%	(18)	75.3%	(55)	19.2%	(14)	89%	(65)	84.9%	(62)	13.7%	(10)	16.4%	(12)	0	0	100%	(73)	8.2%	(6)	19.2%	(14)	26%	(19)	46.6%	(34)
P-value ^s	0.282	<0.001	0.118	0.002	0.008	0.570	0.002	0.008	0.008	0.008	0.570	0.249	0.435	0.249	0.435	0.001	0.001	0.001	<0.001	<0.001	0.023	0.023	0.023	0.023	0.023	0.023	0.023	0.023	0.023	0.023
Total (161)	71.4%	(115)	7.5%	(12)	19.9%	(32)	85.1%	(137)	29.2%	(47)	88.8%	(143)	82%	(132)	14.9%	(24)	29.2%	(47)	16.1%	3.1%	80.7%	(130)	6.2%	(10)	12.4%	(20)	23%	(37)	58.4%	(94)

UTIs, urinary tract infections; non-UTIs: other isolates containing wound and blood; P: presence; N: negative; α: alpha hemolysin; β: beta hemolysin; γ: none hemolysin; +: low; ++: moderate; +++: strong
^s One-tailed Fisher's exact test was used for comparison of hospital-acquired and dental-root canal groups.




Figure 2. Biofilm formation absorbance association to virulence genes and CRISPR loci among *E. faecalis* isolates. (Error bars illustrate the minimum and maximum of three replicates of absorbance of the biofilm formation; **P*-value was significant (*P*-value<0.05.)




Figure 3. Distribution of virulence gene counts among *E. faecalis* isolates.

of either CRISPR1-cas or orphan CRISPR2 and either CRISPR3-cas or orphan CRISPR2 was significantly correlated with the presence of *ace* and the absence of *gelE*, respectively (*P* = 0.019, correlation coefficient = 0.185 and *P* = 0.014, correlation coefficient = 0.184,

respectively). In addition, presence of CRISPR1-cas was significantly correlated with the absence of *hyl* (*P* = 0.048, correlation coefficient = -0.147). Other significant correlations were found between the absence of CRISPR1 and the absence of *cyaA* and *asaI* (*P* < 0.05,

Table 3. The presence of CRISPR-cas type in hospital-acquired and dental-root isolates of *E. faecalis*.

CRISPR	CRISPR1-cas	CRISPR2	CRISPR3-cas	CRISPR1-cas or CRISPR2	CRISPR1-cas or CRISPR3-cas	CRISPR2 or CRISPR3-cas	CRISPR1-cas and CRISPR2	CRISPR1-cas and CRISPR3-cas	CRISPR2 and CRISPR3-cas	CRISPR1-cas and CRISPR2 and CRISPR3-cas
Hospital-acquired isolates (88)	19.3% (17)	53.4% (47)	2.3% (2)	59.1% (52)	21.6% (19)	54.5% (48)	13.6% (12)	0	1.1% (1)	0
Dental root isolates (73)	5.5% (4)	57.5% (42)	35.6% (26)	58.9% (43)	42.5% (31)	72.6% (53)	5.5% (4)	0	20.5% (15)	0
<i>P</i> -value [§]	0.008	0.358	<0.001	0.554	0.004	0.014	0.070	NS	<0.001	NS
Total (161)	13% (21)	55.3% (89)	17.4% (28)	59% (95)	31.1% (50)	62.7% (101)	9.9% (16)	0	9.9% (16)	0

NS, not significant.

§ One-tailed Fisher's exact test was used for comparison of hospital-acquired and dental-root canal groups.




Figure 4. Virulence gene counts association to CRISPR loci among *E. faecalis* isolates. (Error bars illustrate the minimum and maximum of virulence gene counts; **P*-value was significant (*P*-value <0.05; H: Hospital-acquired; D: Dental-acquired; T: Total).

correlation coefficient = 0.171 and 0.149, respectively), and between the absence of CRISPR2 and the absence of *gelE* ($P = 0.001$, correlation coefficient = 0.248). In hospital-acquired isolates, a significant correlation was found between the absence of CRISPR loci and the absence of *gelE*, *asa1*, gelatinase and hemolysis activity ($P < 0.05$); in dental-root canal isolates, a significant correlation was found between the absence of CRISPR3-*cas* and the absence of gelatinase ($P = 0.003$, correlation coefficient = 0.365), between the absence of either CRISPR1-*cas* or CRISPR2-*cas* and the absence of *gelE* ($P = 0.021$, correlation coefficient = 0.265), and between the presence of orphan CRISPR2 and either orphan CRISPR2 or CRISPR3-*cas* and biofilm production ($P = 0.046$, correlation coefficient = 0.247 and $P = 0.044$, correlation coefficient = 0.263, respectively) (see Table 4).

Discussion

In this study, we determined the occurrence of CRISPR loci and the content of virulence factors in *E. faecalis* strains isolated from different infectious sources as a pathogenic organism and the dental-root canal of patients. We found that the presence of CRISPR1 and CRISPR3 loci was varied among *E. faecalis* strains. The abundance of CRISPR1 among the dental-root canal isolates was significantly lower than that of hospital-acquired ones, whereas the reverse was significantly true for CRISPR3. These results were consistent with those obtained by Burley et al. study [31], who found the presence of CRISPR3-*cas* was significantly more in endodontic strains, as compared to hospital-acquired strains, and the majority of strains had CRISPR3. While

these results were interesting, the reasons were not clear. In addition, we found that the presence of orphan CRISPR2 was more among *E. faecalis* strains in comparison to CRISPR1-*cas* and CRISPR3-*cas*, while CRISPR2 lacks of *cas* genes. Palmer et al. [18] and Hullahalli et al. [32] suggested that CRISPR2 is functional for sequence interference and is functionally linked to CRISPR1-Cas or CRISPR3-Cas.

The results revealed that the presence of CRISPR loci was not significantly associated with a less number of virulence factors. There are several virulence factors in *E. faecalis* which play such roles as antiphagocytosis, adherence, biofilm formation, exoenzyme, toxin, and quorum sensing system. Although several studies have reported that there is no clear relation between origin isolation or a single gene and pathogenicity, and perhaps the surface proteins of *E. faecalis* cannot be considered as virulence factors [9,33,34], we found a correlation between the absence of CRISPR1-*cas* and the absence of the *esp* gene (P -value = 0.009, coefficient correlation = 0.204) and a correlation between the absence of CRISPR1-*cas* and the absence of *cylA* (P -value = 0.03, coefficient correlation = 0.171) and *asa1* (P -value = 0.06, coefficient correlation = 0.149) genes. In addition, there was a correlation between the absence of single or multi-CRISPR loci and the absence of some virulence factors. The cytolysin operon, *cob* and *esp* genes reside in the same pathogenicity island, which are located on either the chromosome or on large pheromone-responsive plasmids such as pAD1 [35,36]. The *esp* gene encodes a large surface protein with a variable number of highly conserved 82 amino acids repeats, contributing to the promotion of primary attachment, colonization and biofilm formation of *E. faecalis* [36].

Table 4. Association between genotypic and phenotypic characteristics and the occurrence of CRISPR-cas in *E. faecalis*.

Gene	CRISPR1- present		CRISPR1- absent		P-value	CRISPR2- present		CRISPR2- absent		P-value	CRISPR3- present		CRISPR3- absent		P-value	CRISPR1 or CRISPR2- present		CRISPR1 or CRISPR2- absent		P-value	CRISPR2 or CRISPR3- present		CRISPR2 or CRISPR3- absent		P-value		
	n	%	n	%		n	%	n	%		n	%	n	%		n	%	n	%		n	%	n	%		n	%
<i>esp</i> -present	20	95	0.005	64	51	0.509	21	94	0.417	68	47	0.549	41	74	0.033	72	43	0.554	72	43	0.033	72	43	0.554	72	43	0.554
<i>esp</i> -absent	1	45		25	21		7	39		27	19		9	37		29	17		29	17		29	17		29	17	
<i>cyfA</i> -present	4	8	0.053	6	6	0.464	0	12	0.092	7	5	0.594	4	8	0.544	6	6	0.258	6	6	0.544	6	6	0.258	6	6	0.258
<i>cyfA</i> -absent	17	132		83	66		28	121		88	61		46	103		95	54		95	54		95	54		95	54	
<i>hyl</i> -present	1	31	0.048	20	12	0.237	6	26	0.500	21	11	0.260	7	25	0.149	23	9	0.161	23	9	0.149	23	9	0.161	23	9	0.161
<i>hyl</i> -absent	20	109		69	60		22	107		74	55		43	86		78	51		78	51		78	51		78	51	
<i>efaA</i> -present	20	117	0.139	75	62	0.461	22	115	0.214	81	56	0.557	43	94	0.518	86	51	0.575	86	51	0.518	86	51	0.575	86	51	0.575
<i>efaA</i> -absent	1	23		14	10		6	18		14	10		7	17		15	9		15	9		15	9		15	9	
<i>gelE</i> -present	7	40	0.415	35	12	0.001	5	42	0.108	36	11	0.003	12	35	0.217	36	11	0.014	36	11	0.217	36	11	0.014	36	11	0.014
<i>gelE</i> -absent	14	100		54	60		23	91		59	55		38	76		65	49		65	49		65	49		65	49	
<i>ace</i> -present	20	123	0.282	84	59	0.012	26	117	0.358	89	54	0.019	46	97	0.285	94	49	0.027	94	49	0.285	94	49	0.027	94	49	0.027
<i>ace</i> -absent	1	17		5	13		2	16		6	12		4	14		7	11		7	11		7	11		7	11	
<i>ebpR</i> -present	18	114	0.451	75	57	0.263	24	108	0.399	80	52	0.250	42	90	0.418	85	47	0.235	85	47	0.418	85	47	0.235	85	47	0.235
<i>ebpR</i> -absent	3	26		14	15		4	25		15	14		8	21		16	13		16	13		16	13		16	13	
<i>asaI</i> -present	6	18	0.067	15	9	0.293	3	21	0.363	17	7	0.146	9	15	0.303	16	8	0.425	16	8	0.303	16	8	0.425	16	8	0.425
<i>asaI</i> -absent	15	122		74	63		25	112		78	59		41	96		85	52		85	52		85	52		85	52	
Biofilm- positive	19	132	0.384	86	65	0.092	27	124	0.453	91	60	0.176	46	105	0.377	97	54	0.117	97	54	0.377	97	54	0.117	97	54	0.117
Biofilm- negative	2	8		3	7		1	9		4	6		4	6		4	6		4	6		4	6		4	6	
Gelatinase- positive	4	43	0.203	29	18	0.190	10	37	0.268	30	17	0.268	14	33	0.490	32	15	0.236	32	15	0.490	32	15	0.236	32	15	0.236
Gelatinase- negative	17	97		60	54		18	96		65	49		36	78		69	45		69	45		69	45		69	45	
Hemolysis- positive	4	27	0.623	19	12	0.293	0	31	0.001	19	12	0.469	4	27	0.010	19	12	0.504	19	12	0.469	19	12	0.504	19	12	0.504
Hemolysis- negative	17	113		70	60		28	102		76	54		46	84		82	48		82	48		82	48		82	48	

Our results, therefore, showed that the presence of *efaA*, *esp*, *gelE*, *ace*, and *ebpR* genes were significantly associated with biofilm formation among the hospital-acquired isolates and *efaA* and *gelE* genes were significantly associated with biofilm formation in all *E. faecalis* isolates. Conflict outcomes have been, however, published regarding the role of the genes of biofilm formation. Duggan et al., for example, suggested that *asa1*, *cylA*, *esp* and *gelE* were not associated with biofilm formation in the oral and endodontic isolates of *E. faecalis* [37], which is compatible with our results. In addition, the results revealed that 13.6% of hospital-acquired isolates carried the *cylA* gene, but only 35.2% of the isolates expressed hemolysin activity (both alpha and beta hemolysis). Several studies such as Sun et al. [38], Sedgley et al. [39] and Lindenstrauß et al. [40] have also determined 38%, 36% and 33.3% of the chronic periodontitis, endodontic, and clinical and food isolates of *E. faecalis* to be capable of producing hemolysis, respectively. These differences may be due to the differences in the types of blood used for the determination of the hemolysis activity, while we used human blood, others have employed horse and sheep blood. In addition, Sun et al. [38] and Sedgley et al. [39] reported the distribution of the *cylA* gene was detected only in 17% and 18.18% of the isolates, respectively; this was compatible with our results. These results may be due to such environmental factors as *in vitro* and *in vivo* conditions used to test for phenotypic characters, which could strongly influence gene expression [41] and can be the cause of the differences between our results and those obtained by others in the case of hemolysis activity. In addition, hemolysin activity was encoded by *cyl* operon in *E. faecalis*, where *cylA* is the only reading frame required for the expression of component A, a serine protease. As well, there is no association between CRISPR1-*cas*, biofilm-formation, and hemolysis activity. Several studies have reported that CRISPR loci play an inverse role in some virulence factors and acquisition of antibiotic resistance [18,31,40], such as Palmer and Gilmore's study [42] and Burley et al.'s study [31], reporting that CRISPR loci were inversely associated with antibiotic resistance and some virulence factors in *E. faecalis* strains. In addition, similar to our results, Toro et al. [43] and Touchon et al. [44] reported that there was no significant association with CRISPR-*cas* and acquisition of integrons, plasmids, antibiotic resistance and virulence genes in *Escherichia coli*. However, an analysis of 370 other Archaeal and Eubacteria genomes showed that there was potential evidence for the propagation of CRISPR-*cas* genes to occur via horizontal gene transfer [45]. These findings, therefore, suggested that CRISPR loci could potentially

inhibit or prevent some or part of the virulence factors and Pathogenicity Island could not serve as the selective forces to influence the pathogenic traits of *E. faecalis*.

Conclusion

The findings of this study indicated that CRISPR-*cas* might prevent the acquisition of some respective pathogenicity factors in some isolates, though not all; significant inverse correlations were found between CRISPR-*cas* loci, *esp* and *gelE*, while direct ones were found in *cylA*, *hyl*, *gelE* (between some CRISPR-loci), *asa1*, *ace*, biofilm formation, gelatinase, and hemolysis activities. However, other studies demonstrated that CRISPR-*cas* could prevent the acquisition of antibiotic resistance genes in *E. faecalis* and other bacteria. Further studies can determine the exact role of CRISPR-*cas* in the pathogenesis of Enterococcal infections.

Acknowledgments

We thank Antwerp University, Belgium, for providing us with the reference strains for study.

Disclosure statement

The authors declare that they have no conflict of interest.

Funding

This study was supported by Iran National Science Foundation (INSF) with the grant number 97015174 and Faculty of Medicine, Tabriz University of Medical Sciences.

Ethics approval and consent to participate

This study was approved by the Regional Ethics Committee of Tabriz (Tabriz University of Medical Sciences, Tabriz, Iran, No. IR.TBZMED.REC.1397.188).

Consent to publish

For this study, the formal consent was filled by all participants before any procedure according to the ethics committee approved procedure.

Authors' contributions

HSK, MY, MB and KG designed this study and discussed related methods; PG did experimental analysis, performed the statistical analysis, and was a major contributor in writing the manuscript. TP, MAR collected the data; PL reviewed the manuscript and put forward the comments; RQ, MA suggested and revised the manuscript. All authors read and approved the final manuscript.

ORCID

Hossein Samadi Kafil  <http://orcid.org/0000-0001-6026-8795>

Availability of data and materials

The datasets used in this study are available from the corresponding author on reasonable request.

References

- [1] Tannock GW, Cook G. Enterococci as members of the intestinal microflora of humans. In: Gilmore MS, Clewell DB, Courvalin P, et al., editors. *The enterococci*.
- [2] Baldassarri L, Creti R, Recchia S, et al. Virulence factors in enterococcal infections of orthopedic devices. *Int J Artif Organs*. 2006;29:402–406.
- [3] Mohamed JA, Huang W, Nallapareddy SR, et al. Influence of origin of isolates, especially endocarditis isolates, and various genes on biofilm formation by *Enterococcus faecalis*. *Infect Immun*. 2004;72:3658–3663.
- [4] Gomes BPFA, Pinheiro ET, Sousa ELR, et al. *Enterococcus faecalis* in dental root canals detected by culture and by polymerase chain reaction analysis. *Oral Sur Oral Med Oral Pathol Oral Radiol Endodontol*. 2006;102:247–253.
- [5] Hidron AI, Edwards JR, Patel J, et al. NHSN annual update: antimicrobial-resistant pathogens associated with healthcare-associated infections: annual summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2006–2007. *Infect Control Hosp Epidemiol*. 2008;29:996–1011.
- [6] Gilmore MS, Segarra RA, Booth MC, et al. Genetic structure of the *Enterococcus faecalis* plasmid pAD1-encoded cytolytic toxin system and its relationship to lantibiotic determinants. *J Bacteriol*. 1994;176:7335–7344.
- [7] Jett BD, Huycke MM, Gilmore MS. Virulence of enterococci. *Clin Microbiol Rev*. 1994;7:462–478.
- [8] Su Y, Sulavik M, He P, et al. Nucleotide sequence of the gelatinase gene (gelE) from *Enterococcus faecalis* subsp. *liquefaciens*. *Infect Immun*. 1991;59:415–420.
- [9] Eaton TJ, Gasson MJ. Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates. *Appl Environ Microbiol*. 2001;67:1628–1635.
- [10] Galli D, Lottspeich F, Wirth R. Sequence analysis of *Enterococcus faecalis* aggregation substance encoded by the sex pheromone plasmid pAD1. *Mol Microbiol*. 1990;4:895–904.
- [11] Bourgogne A, Singh KV, Fox KA, et al. EbpR is important for biofilm formation by activating expression of the endocarditis and biofilm-associated pilus operon (ebpABC) of *Enterococcus faecalis* OG1RF. *J Bacteriol*. 2007;189:6490–6493.
- [12] Nallapareddy SR, Qin X, Weinstock GM, et al. *Enterococcus faecalis* adhesin, ace, mediates attachment to extracellular matrix proteins collagen type IV and laminin as well as collagen type I. *Infect Immun*. 2000;68:5218–5224.
- [13] Brouns SJ, Jore MM, Lundgren M, et al. Small CRISPR RNAs guide antiviral defense in prokaryotes. *Science*. 2008;321:960–964.
- [14] Grissa I, Vergnaud G, Pourcel C. The CRISPRdb database and tools to display CRISPRs and to generate dictionaries of spacers and repeats. *BMC Bioinformatics*. 2007;8:172.
- [15] Makarova KS, Haft DH, Barrangou R, et al. Evolution and classification of the CRISPR–Cas systems. *Nature Rev Microbiol*. 2011;9:467–477.
- [16] Bourgogne A, Garsin DA, Qin X, et al. Large scale variation in *Enterococcus faecalis* illustrated by the genome analysis of strain OG1RF. *Genome Biol*. 2008;9:R110.
- [17] Haft DH, Selengut J, Mongodin EF, et al. A guild of 45 CRISPR-associated (Cas) protein families and multiple CRISPR/Cas subtypes exist in prokaryotic genomes. *PLoS Comput Biol*. 2005;1:e60.
- [18] Palmer KL, Gilmore MS. Multidrug-resistant enterococci lack CRISPR-cas. *MBio*. 2010;1:e00227–10.
- [19] Gholizadeh P, Aghazadeh M, Asgharzadeh M, et al. Suppressing the CRISPR/Cas adaptive immune system in bacterial infections. *Eur J Clin Microbiol Infect Dis*. 2017;36:2043–2051.
- [20] Marraffini LA, Sontheimer EJ. CRISPR Interference Limits Horizontal Gene Transfer in Staphylococci by Targeting DNA. *Science*. 2008;322:1843–1845.
- [21] Pourcel C, Salvignol G, Vergnaud G. CRISPR elements in *Yersinia pestis* acquire new repeats by preferential uptake of bacteriophage DNA, and provide additional tools for evolutionary studies. *Microbiology*. 2005;151:653–663.
- [22] Mojica FJM, Díez-Villaseñor C, García-Martínez J, et al. Intervening sequences of regularly spaced prokaryotic repeats derive from foreign genetic Elements. *J Mol Evol*. 2005;60:174–182.
- [23] Facklam R, Collins M. Identification of *Enterococcus* species isolated from human infections by a conventional test scheme. *J Clin Microbiol*. 1989;27:731–734.
- [24] Kafil HS, Mobarez AM, Moghadam MF, et al. Gentamicin induces efaA expression and biofilm formation in *Enterococcus faecalis*. *Microb Pathog*. 2016;92:30–35.
- [25] Kafil HS, Mobarez AM, Moghadam MF. Adhesion and virulence factor properties of *Enterococci* isolated from clinical samples in Iran. *Indian J Pathol Microbiol*. 2013;56:238.
- [26] Kariyama R, Mitsuhashi R, Chow JW, et al. Simple and reliable multiplex PCR assay for surveillance isolates of vancomycin-resistant enterococci. *J Clin Microbiol*. 2000;38:3092–3095.
- [27] Aghazadeh M, Zahedi Bialvaei A, Kabiri F, et al. Survey of the antibiofilm and antimicrobial effects of zingiber officinale (in Vitro Study). *Jundishapur J Microbiol*. 2016;9:e30167.
- [28] Kafil HS, Mobarez AM. Assessment of biofilm formation by enterococci isolates from urinary tract infections with different virulence profiles. *J King Saud Univ Sci*. 2015;27:312–317.
- [29] Gaspar F, Crespo M, Lopes MS. Proposal for a reliable enterococcal cytolysin production assay avoiding apparent incongruence between phenotype and genotype. *J Med Microbiol*. 2009;58:1122–1124.

- [30] Pickett M, Greenwood J, Harvey S. Tests for detecting degradation of gelatin: comparison of five methods. *J Clin Microbiol.* **1991**;29:2322–2325.
- [31] Burley KM, Sedgley CM. CRISPR-Cas, a prokaryotic adaptive immune system, in endodontic, oral, and multidrug-resistant hospital-acquired enterococcus faecalis. *J Endod.* **2012**;38:1511–1515.
- [32] Hullahalli K, Rodrigues M, Palmer KL. Exploiting CRISPR-Cas to manipulate *Enterococcus faecalis* populations. *eLife.* **2017**;6:e26664.
- [33] Hew CM, Korakli M, Vogel RF. Expression of virulence-related genes by *Enterococcus faecalis* in response to different environments. *Syst Appl Microbiol.* **2007**;30:257–267.
- [34] Martin B, Garriga M, Hugas M, et al. Genetic diversity and safety aspects of enterococci from slightly fermented sausages. *J Appl Microbiol.* **2005**;98:1177–1190.
- [35] Shankar N, Coburn P, Pillar C, et al. Enterococcal cytolysin: activities and association with other virulence traits in a pathogenicity island. *Int J Med Microbiol.* **2004**;293:609–618.
- [36] Shankar N, Lockett CV, Baghdayan AS, et al. Role of *Enterococcus faecalis* surface protein ESP in the pathogenesis of ascending urinary tract infection. *Infect Immun.* **2001**;69:4366–4372.
- [37] Duggan JM, Sedgley CM. Biofilm formation of oral and endodontic enterococcus faecalis. *J Endod.* **2007**;33:815–818.
- [38] Sun J, Sundsfjord A, Song X. *Enterococcus faecalis* from patients with chronic periodontitis: virulence and antimicrobial resistance traits and determinants. *Eur J Clin Microbiol Infect Dis.* **2012**;31:267–272.
- [39] Sedgley CM, Molander A, Flannagan SE, et al. Virulence, phenotype and genotype characteristics of endodontic *Enterococcus* spp. *Oral Microbiol Immunol.* **2005**;20:10–19.
- [40] Lindenstrauß AG, Pavlovic M, Bringmann A, et al. Comparison of genotypic and phenotypic cluster analyses of virulence determinants and possible role of CRISPR elements towards their incidence in *Enterococcus faecalis* and *Enterococcus faecium*. *Syst Appl Microbiol.* **2011**;34:553–560.
- [41] Finlay BB, Falkow S. Common themes in microbial pathogenicity revisited. *Microbiol Mol Biol Rev.* **1997**;61:136–169.
- [42] Palmer KL, Carniol K, Manson JM, et al. High-quality draft genome sequences of 28 *Enterococcus* sp. isolates. *J Bacteriol.* **2010**;192:2469–2470.
- [43] Toro M, Cao G, Ju W, et al. Association of clustered regularly interspaced short palindromic repeat (CRISPR) elements with specific serotypes and virulence potential of Shiga toxin-producing *Escherichia coli*. *Appl Environ Microbiol.* **2014**;80:1411–1420.
- [44] Touchon M, Charpentier S, Pognard D, et al. Antibiotic resistance plasmids spread among natural isolates of *Escherichia coli* in spite of CRISPR elements. *Microbiology.* **2012**;158:2997–3004.
- [45] Godde JS, Bickerton A. The repetitive DNA elements called CRISPRs and their associated genes: evidence of horizontal transfer among prokaryotes. *J Mol Evol.* **2006**;62:718–729.
- [46] Shankar V, Baghdayan AS, Huycke MM, et al. Infection-derived *Enterococcus faecalis* strains are enriched in *esp*, a gene encoding a novel surface protein. *Infect Immun.* **1999**;67:193–200.
- [47] Vankerckhoven V, Van Autgaerden T, Vael C, et al. Development of a multiplex PCR for the detection of *asa1*, *gelE*, *cylA*, *esp*, and *hyl* genes in enterococci and survey for virulence determinants among European hospital isolates of *Enterococcus faecium*. *J Clin Microbiol.* **2004**;42:4473–4479.
- [48] Lowe A, Lambert P, Smith A. Cloning of an *Enterococcus faecalis* endocarditis antigen: homology with adhesins from some oral streptococci. *Infect Immun.* **1995**;63:703–706.
- [49] Nallapareddy SR, Murray BE. Ligand-signaled upregulation of *Enterococcus faecalis* *ace* transcription, a mechanism for modulating host-*E. faecalis* interaction. *Infect Immun.* **2006**;74:4982–4989.