Open access Correction

Correction: Efficacy of prehospital National Early Warning Score to predict outpatient disposition at an emergency department of a Japanese tertiary hospital: a retrospective study

Endo T, Yoshida T, Shinozaki T, *et al.* Efficacy of prehospital National Early Warning Score to predict outpatient disposition at an emergency department of a Japanese tertiary hospital: a retrospective study. *BMJ Open* 2020;10:e034602.

This article was previously published with an error. Below typographical errors in the Results section of the Abstract have been corrected: $(n=1263, 60.3\pm3.8)$ changed to $(n=1263, 6.3\pm3.8)$ $(n=22, 110.7\pm2.9)$ changed to $(n=22, 11.7\pm2.9)$.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

BMJ Open 2020;10:e034602corr1. doi:10.1136/bmjopen-2019-034602corr1

