



Published in final edited form as:

Drugs. 2020 November ; 80(16): 1649–1676. doi:10.1007/s40265-020-01379-9.

HIV-1 Integrase Inhibitors: A Comparative Review of Efficacy and Safety

Kimberly K. Scarsi^{1,2}, Joshua P. Havens^{1,2}, Anthony T. Podany¹, Sean N. Avedissian¹, Courtney V. Fletcher^{1,2}

¹Antiviral Pharmacology Laboratory, College of Pharmacy, University of Nebraska Medical Center, Omaha, NE, USA.

²Division of Infectious Diseases, Department of Medicine, University of Nebraska Medical Center, Omaha, NE, USA.

Abstract

The newest class of antiretrovirals for all persons living with HIV are the integrase strand transfer inhibitors (INSTIs). Since 2007, five INSTIs have been introduced: raltegravir, elvitegravir, dolutegravir, bictegravir, and cabotegravir. The INSTIs have favorable pharmacokinetic and pharmacodynamic properties, which contribute to both their effectiveness and ease of use. With the exception of cabotegravir, each INSTI is US Food and Drug Administration (FDA) approved for treatment-naïve individuals initiating antiretroviral therapy. All of the INSTIs, except raltegravir, are approved for antiretroviral treatment simplification for virologically suppressed patients without INSTI resistance. Data also support the use of dolutegravir and raltegravir in individuals with antiretroviral resistance as part of an optimized antiretroviral regimen. INSTIs are generally well tolerated by people living with HIV compared with older classes of antiretrovirals, but emerging data suggest that some INSTIs contribute to weight gain. Due to their efficacy, safety, and ease of use, HIV treatment guidelines recommend oral INSTIs as preferred components of antiretroviral therapy for individuals initiating therapy. The newest INSTI, cabotegravir, represents an alternative to oral administration of life-long antiretroviral therapy with the availability of a long acting injectable formulation. This review summarizes the current use of INSTIs in adults living with HIV, highlighting the similarities and differences within the class related to pharmacodynamics, pharmacokinetics, safety, dosing and administration that contribute to their role in modern antiretroviral therapy.

1. Introduction

For the nearly 40 million persons living with HIV (PLWH), effective antiretroviral therapy (ART) reduces morbidity and mortality, leading to a life expectancy expected to be similar to age-matched peers [1, 2]. Randomized, clinical trials have demonstrated health benefits associated with early initiation of ART beyond the reduction in HIV-associated morbidity

Corresponding Author Contact Information: Kimberly K. Scarsi, PharmD, MS, Antiviral Pharmacology Laboratory, University of Nebraska Medical Center, 986145 Nebraska Medical Center, Omaha, NE 68198-6145, kim.scarsi@unmc.edu.

Conflicts of Interest: KKS, ATP, SNA, CVF declare no conflict of interest related to this content. JPH reports research grants paid to his institution from Gilead Sciences.

and mortality [3, 4]. Furthermore, effective ART offers a significant public health benefit by preventing transmission of HIV [5–7]. Based on these combined benefits, together with improvements in available antiretroviral options, ART is now recommended for all PLWH [8–10].

The integrase strand transfer inhibitor (INSTI) class of antiretrovirals contributes to the enhanced safety and efficacy of modern ART regimens. Based on data demonstrating efficacy, safety and ease of administration, INSTIs are now part of preferred or recommended ART regimens in HIV treatment guidelines throughout the world [8–10]. The efficacy of INSTIs has been well established in comparison with both protease inhibitor (PI) and non-nucleoside reverse transcriptase inhibitor (NNRTI)-based ART regimens (see Section 5.0) [8]. INSTI effectiveness is related to their antiviral activity and improved tolerability, which results in fewer treatment discontinuations [11].

Raltegravir was the first drug in the INSTI class and was approved for twice-daily administration by the United States (US) Food and Drug Administration (FDA) in 2007 [12]. A once-daily formulation was approved in 2017, but raltegravir remains the only INSTI not available as a fixed dose combination (FDC). Elvitegravir was the second INSTI approved by the FDA in 2012, and was the first once-daily, FDC single tablet regimen containing an INSTI [13]. The two most commonly used INSTIs are dolutegravir, approved in 2013, and a bicittegravir-containing FDC approved in 2018. Cabotegravir is an INSTI formulated as a long-acting product given intramuscularly in combination with rilpivirine. The combination was recently approved in Canada, and is undergoing regulatory review in the US and Europe [14].

This review will summarize key information regarding the dosing and administration, pharmacodynamics, pharmacokinetics, efficacy, and safety of INSTIs with an emphasis on treatment of adults with HIV. A detailed review on the use of INSTIs in pediatric patients is addressed by Dehority and colleagues [15]. Similarities and differences between the individual agents will be highlighted to inform the current place in therapy for each INSTI.

2. Dosage and Administration

Table 1 describes the available formulations, dosing, and indications for each INSTI. The companion antiretrovirals included in the INSTI single tablet regimens vary by agent, as described in Table 1.

3. Pharmacodynamic Properties

3.1 Antiviral Activity

INSTIs inhibit HIV by blocking the strand transfer step of viral DNA integration into the host genome. INSTIs are potent and selective antiretrovirals with sub- to low nanomolar *in vitro* activity. Against clinical isolates of HIV-1, *in vitro* 50% inhibitory concentrations (IC₅₀) reported in approved product labels are approximately, 0.1 ng/mL for cabotegravir, 0.2 ng/mL for both bicittegravir and dolutegravir; and range from 0.04–0.6 ng/mL for elvitegravir, and 2.2–5.3 ng/mL for raltegravir [12, 14, 16–18]. However, all INSTIs are

highly protein bound, which substantially influences clinical IC_{50} values. For example, in an *in vitro* assay in peripheral blood mononuclear cells, the IC_{50} of dolutegravir was 0.21 ng/mL, but was increased 75-fold in the presence of human serum albumin to a protein-adjusted IC_{50} of 16 ng/mL [19]. The protein-adjusted 90% or 95% inhibitory concentration (IC_{90} or IC_{95}) values (ng/mL) for the current INSTIs are: raltegravir, 15; elvitegravir, 45; dolutegravir, 64; bictegravir, 162; cabotegravir, 166 [20].

3.2 Resistance

To date, resistance to all antiretrovirals has been documented [21]. With regard to INSTIs, the first-generation agents, raltegravir and elvitegravir, share similar clinical resistance profiles, including cross-resistance. The second-generation INSTIs, bictegravir and dolutegravir, are characterized by similar resistance profiles and retention of potency against resistant mutants selected by first-generation INSTIs. There is evidence the resistance barrier of bictegravir and dolutegravir are higher than raltegravir and elvitegravir [22, 23].

Resistance has emerged during therapy with long-acting, intramuscular cabotegravir plus rilpivirine [24, 25]. In the combined results of two phase 3 trials, six of 586 participants (1.02%) had confirmed virologic failure with INSTI-resistance mutations. Characteristics common among all six were cabotegravir and rilpivirine plasma concentrations in the lowest quartile, despite receipt of all intramuscular injections, and HIV-1 subtypes A/A1 or AG; resistance emerged before or at week 28 of therapy in five of six. Whether clinical resistance emerges to an INSTI (or any antiretroviral) is dependent upon a variety of factors including the drug's inherent genetic barrier to resistance, the drug's structure, inhibitory quotient, therapeutic index, and pharmacokinetic forgiveness/adherence [26]. The consequences of resistance are virologic failure and reduced options for future ART regimens. Current treatment guidelines therefore contain specific recommendations for resistance testing in both naïve- and treatment-experienced PLWH, and recommendations for the use of ART regimens to maximally and durably suppress plasma HIV-RNA to minimize the emergence of resistance [8].

3.3 Clinical Pharmacodynamic Characteristics

Explicit pharmacodynamic relationships have been described between the decline in plasma HIV-RNA and trough concentrations (C_{trough}) of cabotegravir, dolutegravir, and elvitegravir [27–29]. For cabotegravir, pharmacodynamic analyses found the HIV-RNA change from baseline was associated with cabotegravir C_{trough} as a maximum effect (E_{max}) relationship: E_{max} was 2.56 \log_{10} and 50% effective concentration (EC_{50}) was 82 ng/mL [27]. The reduction in plasma HIV-RNA with dolutegravir monotherapy of 2, 10 and 50mg once-daily for 10 days was associated with dolutegravir C_{trough} in an E_{max} relationship with an estimated EC_{50} of 36 ng/mL [28]. The C_{trough} of elvitegravir in antiretroviral-naïve and experienced persons not currently on therapy, were strongly associated with the \log_{10} change in plasma HIV-RNA in an E_{max} relationship, with an EC_{50} of 14 ng/mL and 90% effective concentration (EC_{90}) of 126 ng/mL [29]. While less quantitative, exposure-response relationships have clearly been demonstrated for bictegravir and raltegravir [30, 31]. A 10-day monotherapy study of bictegravir found the mean change in HIV-RNA, from baseline to day 11 was -1.45 , -2.06 , -2.08 and -2.43 \log_{10} copies/mL for 5, 25, 50 and 100mg doses, respectively, clearly demonstrating that higher doses, and therefore higher concentrations,

were associated with a greater anti-HIV response [30]. A trial of raltegravir 800mg once-daily compared with 400mg twice-daily found C_{trough} raltegravir with once-daily dosing correlated with virologic response [31]. Participants who had raltegravir C_{trough} in the lowest quartile (median C_{trough} 12.5 ng/mL) had a clear fall off in virologic response, with <80% achieving HIV-RNA <50 copies/mL compared to 90% achieving HIV-RNA <50 copies/mL if the C_{trough} was >44 ng/mL. The recently completed phase 3 trials of intramuscular cabotegravir also provide qualitative pharmacodynamic information. The geometric mean cabotegravir concentrations at weeks 8 and 48 were approximately 1500 ng/mL and 3000 ng/mL, respectively, which are 9-fold and 18-fold greater than the protein binding-adjusted IC_{90} of 166 ng/mL [24, 25]. At weeks 4–8 after the start of intramuscular cabotegravir therapy, which is where the lowest concentrations are found, the 5th percentile cabotegravir concentration is approximately 450 ng/mL, which is 2.7-fold greater than the protein binding-adjusted IC_{90} . As noted above, INSTI-resistance mutations emerged before or at week 28 of therapy in the phase 3 trials of injectable cabotegravir. Collectively, for the INSTI class with regard to the anti-HIV response, data support relationships with dose/plasma concentrations.

The inhibitory quotient (IQ) is an intuitive concept for predicting clinical drug activity. The IQ is the ratio of drug concentration in any biologic fluid (e.g., plasma, cerebral spinal fluid) divided by an *in vitro* inhibitory concentration, (i.e. how much drug you have to how much drug you need). The IQ has utility in antiretroviral drug development, as discussed by the FDA [32], because a high IQ indicates sufficient drug concentrations can be achieved that may minimize the emergence of viral resistance and inform the selection of doses for phase 3 and 4 studies, as well as for select patient populations. For the available INSTIs, the hierarchy of IQ values, where IQ is the ratio of typical trough plasma concentration achieved with approved oral dose divided by the protein-binding adjusted IC_{90} or 95 is cabotegravir > dolutegravir > bictegravir > elvitegravir > raltegravir [20].

4. Pharmacokinetic Properties

The pharmacokinetics of INSTIs have recently been reviewed by Podany and colleagues, and readers can find detailed pharmacokinetic parameters there [20]. All oral INSTIs may be given with or without food (Table 1), except elvitegravir, where one study found that a low versus high fat meal increased the area under the concentration time curve (AUC) by 34% and 87%, respectively, compared with fasting [18]. The dolutegravir/rilpivirine FDC must also be administered with food due to the rilpivirine component. As a class, oral absorption can be impaired if taken with divalent or trivalent cations, discussed further in Section 4.2.

INSTIs are not extensively renally cleared and primarily undergo hepatic metabolism [20]. Raltegravir is primarily conjugated via uridine 5'-diphosphate glucuronosyltransferase (UGT) 1A1 [12]. Elvitegravir is primarily metabolized via cytochrome P450 (CYP) 3A4 enzyme and is the only INSTI that must be coadministered with a pharmacokinetic-enhancer, cobicistat [8]. Dolutegravir is primarily conjugated by UGT1A1, with secondary metabolism by CYP3A4 (10–15%) [17], while bictegravir is metabolized by both CYP3A4 and UGT1A [16]. Finally, cabotegravir is metabolized by both UGT1A1 (primarily) and UGT1A9, with minimal CYP involvement [33]. The metabolic pathway of each INSTI

informs the likelihood of metabolism-related drug-drug interactions, as discussed in Section 4.2.

The oral INSTIs have half-lives ranging from 9 to 38.8 hours [12, 16–18, 34]. Oral cabotegravir has the longest half-life of the INSTIs, 38.8 hours, and intramuscularly administered cabotegravir exhibits an exceptionally long and variable half-life, estimated as 2.3 to 14.7 weeks [35, 36]. Considering the prolonged time to maximum concentration (C_{max}), and the observed elimination half-life, cabotegravir is expected to accumulate over the first several months of administration, and is detectable for several months after discontinuation [37]. Landovitz and colleagues recently described that cabotegravir remained detectable in 23% of males and 63% of females one year after discontinuing cabotegravir for pre-exposure prophylaxis [37].

4.1 Special Populations

Overall, population pharmacokinetic studies have shown that oral INSTIs do not have any clinically-relevant pharmacokinetic differences based upon race or sex in adults [12, 16, 17, 38, 39].

Ontogenic enzymatic changes that occur in pediatric patients inform INSTI dosing in infants and children. For example, activity of the enzyme responsible for glucuronidation (i.e., UGT) is low at birth and increases dramatically during the first four to six-weeks of life in full-term neonates [40]. This can lead to increased clearance of INSTIs that primarily undergo UGT enzyme metabolism (i.e., dolutegravir, raltegravir, cabotegravir). Other potential differences in enzymatic activity of specific age groups in pediatric patients should be considered when selecting and dosing agents [41]. Recommendations on specific INSTIs in pediatric patients vary by agent (Table 1), with raltegravir and dolutegravir allowing dosing with formulations suitable for the smallest children. Some INSTIs may be used to treat pediatric patients weighing 20 kg or more using standard, adult doses, while many FDCs and newer agents do not yet have information for use in patients less than 18 years of age.

For many currently available INSTIs, the manufacturers recommend caution in geriatric use (> 65 years) given the greater frequency of decreased hepatic, renal or cardiac function in this population [12, 13, 17]. However, a population pharmacokinetic analysis of participants in phase 3 trials of bicitegravir showed age did not have a clinically-relevant effect on bicitegravir exposures up to 74-years of age [16]. Similarly, clinical trials of elvitegravir included participants over 65-years and found no differences in safety or efficacy compared with participants age 18- to 65-years [18]. A pharmacokinetic analysis of dolutegravir in PLWH who were > 60 years of age did find that the C_{max} for dolutegravir was significantly higher in this age group compared with a historical control population aged < 50 years (geometric mean 4246 ng/mL versus 3402 ng/mL, $p=0.005$) [42]. Despite the increased C_{max} , there was no excess risk of adverse events associated with dolutegravir. No recommendations are available for cabotegravir in geriatric populations. Despite warnings in product labeling, available data suggest INSTIs may be used in geriatric patients. Given the aging PLWH, more pharmacokinetic and pharmacodynamic studies in geriatric persons at conventional FDA-approved doses are needed.

4.1.1 Pregnancy and Lactation—The US Department of Health and Human Services (DHHS) Perinatal HIV Treatment Guidelines recommend ART therapy for all pregnant PLWH [43]. However, pregnancy can cause physiological changes that have the potential to impact ART pharmacokinetics, which may require dose considerations. For example, induction of UGT1A1 and CPY3A4 during pregnancy may influence INSTI metabolism [44, 45]. Detailed reviews of INSTI pharmacokinetics during pregnancy was recently described by van der Galien et al. and Podany et al [20, 46]; generally, INSTI exposure is lower during pregnancy, resulting in variable clinical recommendations (Table 1).

Though the exposure of dolutegravir during pregnancy was decreased between 10–50% compared to post-partum concentrations in several studies [47, 48], it still may be given without dose adjustment during pregnancy as long as it is taken with food to increase absorption [43]. This is based on the high rate of virologic suppression observed, and because the median AUC during pregnancy was similar to nonpregnant adults in clinical trials. Raltegravir concentrations were 30–50% lower during pregnancy, though highly variable, and the effectiveness during pregnancy was not affected by the reduced exposure [49, 50]. Therefore, twice-daily raltegravir is recommended; the once-daily raltegravir HD formulation should not be used in pregnant women due to the lower C_{trough} observed with this formulation compared with twice-daily [43].

Three INSTIs have insufficient evidence to support their use or are not recommended during pregnancy. Elvitegravir is not recommended due to significantly lower plasma concentrations during pregnancy with associated cases of virologic failure [43, 44, 51]. For example, Momper and colleagues performed a study in 30 pregnant women taking elvitegravir/cobicistat once-daily and found that compared to postpartum data, elvitegravir AUCs were 24% lower in the second trimester (n=14, geometric mean ratio [GMR] =0.76, 90% CI: 0.57, 1.0) and 44% lower in the third trimester (n=24, GMR=0.56, 90% CI 0.42, 0.73) [51]. The reduced elvitegravir exposure during pregnancy is related to a decrease in cobicistat exposure during pregnancy; cobicistat was 44% lower during the second trimester (GMR 0.56, 90% CI 0.37, 0.85) and 59% lower in the third trimester (GMR 0.41, 90% CI 0.30, 0.85) of pregnancy, leading to less pharmacokinetic enhancement of elvitegravir [51]. No pharmacokinetic studies are available for bictegravir during pregnancy, but studies in pregnant women are ongoing. Cabotegravir pharmacokinetic data during pregnancy are limited to three participants who became pregnant during clinical trials [52]. All three individuals had adequate cabotegravir concentrations prior to pregnancy, throughout pregnancy, and post-partum.

It is also important to consider fetal exposure throughout the antepartum period to ensure prevention of HIV transmission from mother to fetus. Pharmacokinetic studies of dolutegravir [47, 48, 53], elvitegravir [51, 54] and raltegravir [50] have shown placental transfer with mean/median cord-to-maternal plasma ratios ranging from 0.09 to 1.5. A comprehensive review of antiretroviral placental transfer is available [55].

The DHHS Perinatal HIV Treatment Guidelines recommend that women with HIV refrain from breastfeeding due to the risk of HIV transmission [43], but this recommendation is not consistent with international guidelines [56]. Data from a recent study found dolutegravir

passes to milk in pregnant women (milk-to-plasma ratio of 0.03) [48]. Dolutegravir was detectable in the plasma of the breastfed infants 10 days of age (range: 7–18) with mean C_{max} of 66.7 (range: 21–654) ng/mL, representing an infant to maternal plasma ratio of 0.03 and a mean minimum concentration of 60.9 (range: 16.3–479) ng/mL (infant:maternal ratio 0.08).

Based on pharmacokinetic results, dolutegravir and raltegravir are appropriate for use during pregnancy for treatment of maternal HIV disease and for prevention of HIV transmission to the infant [43]. There were no adverse effects observed due to infant exposure to dolutegravir during breastfeeding in one clinical trial [48].

4.1.2 Renal/Hepatic Impairment—As a class, no clinically relevant differences in INSTI pharmacokinetics have been observed between patients with renal impairment and those with normal renal function. As such, no dose adjustments are required for the INSTI component of regimens (Table 1) [57, 58]. However, plasma concentrations of dolutegravir have been found to be decreased in persons with severe renal impairment [17]; therefore, caution is advised in certain populations (e.g. those with INSTI-associated resistance). INSTIs are highly bound to plasma proteins (albumin and alpha-1-acid glycoprotein, 83–99%) and are not significantly removed by dialysis (hemodialysis or peritoneal dialysis); therefore, no dose adjustments are required in patients who are receiving hemodialysis or peritoneal dialysis [12, 16–18, 33]. Because most nucleoside reverse transcriptase inhibitors (NRTIs) are renally excreted, the use of INSTI containing FDCs in patients with renal insufficiency may be limited by the concurrent NRTIs (Table 1).

Liver disease has the potential to alter pharmacokinetics of drugs due to changes in hepatic blood flow, altered plasma protein levels, and changes to CYP enzymes and/or glucuronidation [59–61]. However, no significant differences have been seen in the exposures of INSTIs in persons with mild to moderate hepatic impairment (Child-Pugh Class A or B) and no dose adjustment are recommended (Table 1) [57, 62–66]. Given the lack of data available in persons with severe hepatic impairment (Child-Pugh Class C), INSTIs are not recommended [12, 16–18, 33].

4.2 Drug interactions

Most INSTIs are not potent inducers or inhibitors of drug metabolizing enzymes, and so infrequently cause metabolism related causing metabolism related drug-drug interactions [8]. The exception is elvitegravir, which is an inducer of CYP2C9, and must be coadministered with cobicistat, a potent CYP3A4 and a weak CYP2D6 inhibitor [13, 18]. Therefore, elvitegravir/cobicistat is commonly associated with metabolism related drug-drug interactions that effect the exposure of co-administered CYP substrates. Any drug that is a strong inducer or inhibitor of CYP3A and/or UGT1A1 may substantially influence the plasma concentrations of INSTIs [8]. Of the INSTIs, elvitegravir is the most susceptible to adverse drug-drug interactions because it is primarily metabolized by CYP3A4 [13, 18]. Bictegravir and dolutegravir are metabolized by both CYP3A4 and UGT, and comparatively, bictegravir is more susceptible to drug-drug interactions via CYP3A4 compared to

dolutegravir [16, 17]. Raltegravir and cabotegravir have a lower risks of metabolism related drug-drug interactions as they do not undergo CYP3A4 metabolism [12, 14].

Co-administration of rifampin with each INSTI illustrates the potential for metabolism related drug-drug interactions. A phase 1 drug interaction study evaluated the exposure of dolutegravir 50mg once- or twice-daily when co-administered with rifampin [67]. Twice-daily dolutegravir exposures were 54–72% lower in participants receiving rifampin compared with twice-daily dolutegravir without rifampin. However, participants receiving dolutegravir twice-daily with rifampin had higher dolutegravir exposure than those receiving standard dose dolutegravir once-daily without rifampin [GMR (90% CI): AUC_{0-24h} : 1.33 (1.15, 1.5); C_{trough} : 1.22 (1.01, 1.48)]. Clinical data in patients being treated for both HIV and tuberculosis support the efficacy of dolutegravir given twice-daily when combined with rifampin or other enzyme inducing agents [68]. Some guidelines recommend an increase in the adult dose of raltegravir from 400mg to 800mg twice-daily based on reduced raltegravir exposure of 40–61% when combined with rifampin; once-daily raltegravir HD is not recommended [8]. Alternatively, some guidelines do not recommend the increased dose based on one study that demonstrated similar virologic response among participants receiving either raltegravir 400mg or 800mg twice-daily in combination with rifampin (n=51 per group) [virologic suppression: 76% (95% CI 65%, 88%) vs. 78% (95% CI, 67%, 90%), respectively] [69]. In contrast to dolutegravir and raltegravir, coadministration of rifampin and bictegravir, cabotegravir, or elvitegravir is contraindicated [13, 14, 16, 18]. When rifampin was combined with bictegravir 50mg daily, the bictegravir exposure decreased 46–61% and could not be overcome when bictegravir was increased to twice-daily [8]. Oral cabotegravir has been investigated in one healthy-volunteer study that found that co-administration with rifampin reduced cabotegravir AUC by nearly 60% [70]. Finally, a physiologically-based pharmacokinetic model predicted that oral rifampin will decrease the exposure of intramuscular cabotegravir 41–46% [71].

A common drug-drug interaction encountered with INSTI therapy is coadministration of polyvalent cations, which may decrease INSTI absorption due to chelation. The specific management of these interactions is dependent upon the polyvalent cation and its dose, the INSTI, and how the combination was evaluated in pharmacokinetic studies [8]. For example, bictegravir may be coadministered with antacids containing calcium when taken together with food, but not on an empty stomach because food increases the exposure of bictegravir [16]. In contrast, any antacids containing aluminum or magnesium should be given at least two hours after, or six hours before, bictegravir administration. Once-daily raltegravir should not be coadministered with calcium containing antacids, but the 400mg twice-daily dose may be coadministered with calcium antacids irrespective of timing because the twice-daily formulation achieves higher C_{trough} compared with the once-daily formulation [12]. Raltegravir should not be combined with aluminum or magnesium containing antacids. Interactions with oral cabotegravir, dolutegravir, and elvitegravir may be managed by dose separation of the calcium, aluminum, or magnesium-containing antacid and the INSTI [13, 14, 17, 18]. Beyond antacid interactions, consideration of INSTI dosing is required with supplements or laxatives containing polyvalent cations. Specific recommendations exist for calcium and iron supplements, and are often extrapolated to other polyvalent cation containing products [8].

Drug transporters are a cause of some drug-drug interactions involving some INSTIs. *In vitro* studies have shown that bictegravir and dolutegravir inhibit the organic cation transporter 2 (OCT2) and multidrug and toxin extrusion transporter 1 (MATE1) [16, 17, 72]. Coadministration of bictegravir or dolutegravir with other drugs that are substrates of OCT2 and MATE1, such as metformin or dofetilide, may increase concentrations of the coadministered medications. Due to the narrow therapeutic index of dofetilide, coadministration with either dolutegravir or bictegravir is contraindicated [16, 17]. Dolutegravir and bictegravir increase metformin exposure 66–79% and 39%, respectively [17]. Case reports of dolutegravir plus metformin report the occurrence of lactic acidosis when metformin was combined with dolutegravir, as well as case reports of loss of glycemic control when the dose of metformin was empirically reduced in combination with dolutegravir [73]. Therefore, the risk/benefit of combining metformin with either bictegravir or dolutegravir should be evaluated on a case-by-case basis.

INSTIs are involved in bidirectional drug-drug interactions to a varying degree based on their unique pharmacology. An up-to-date drug-drug interaction resource is recommended when evaluating INSTI-related interactions, including the DHHS Adult and Adolescent Treatment Guideline drug-drug interaction tables [8] or the University of Liverpool's database (<http://hiv-druginteractions.org>).

5. Clinical Efficacy

The INSTIs are a key component of modern ART regimens for treatment-naïve and treatment-experienced PLWH. One unique attribute of the INSTIs is rapid virologic suppression compared with other antiretroviral classes. All of the oral INSTIs demonstrate a rapid virologic decay and potent suppression as early as 4 weeks after initiation in treatment-naïve individuals [8, 43, 74–76]. For example, in a study comparing dolutegravir- to bictegravir-containing regimens, 76% to 80% of participants demonstrated virologic suppression within four weeks of treatment initiation [74]. Table 2 summarizes the difference (95% CI) in virologic suppression for each clinical trial discussed in this section.

5.1 Raltegravir

Raltegravir efficacy was compared against an NNRTI regimen containing efavirenz in the STARTMRK study [77]. With raltegravir twice-daily plus two NRTIs, virologic suppression rates at week 48 were 86.1% compared with 81.9% in the efavirenz arm. Raltegravir efficacy remained largely durable, with 71% of raltegravir participants and 61.3% of efavirenz participants maintaining virologic suppression through week 240 [78]. The long-term success of raltegravir compared with efavirenz was largely driven by efavirenz treatment discontinuations.

Two switch studies, SWITCHMRK-1 and 2, investigated raltegravir efficacy in treatment-experienced, virologically-suppressed adults receiving a lopinavir/ritonavir-based ART regimen [79]. Participants were randomized to continue the PI-based regimen or to switch to a raltegravir twice-daily regimen. The study was terminated at week 24 due to lower than expected virologic efficacy of raltegravir: 84% of raltegravir recipients and 91% of

lopinavir/ritonavir achieved virologic suppression. These results failed to establish non-inferiority of raltegravir as a switch strategy [79].

The BENCHMRK trials explored efficacy of raltegravir in treatment-experienced participants with triple drug class resistance [80]. Both studies were randomized studies, comparing raltegravir twice-daily versus placebo in addition to optimized background therapy. Week 48 results demonstrated the efficacy of raltegravir as part of a salvage regimen (virologic suppression: raltegravir, 62.1%; placebo, 32.9%). Week 96 and 156 results largely mimic week 48 outcomes, with 51% of raltegravir recipients and 22% of placebo recipients sustaining virologic suppression through week 156 [81, 82].

Given the necessity of twice-daily dosing of original raltegravir formulation, the QDMRK study investigated the possibility of once-daily dosing in treatment-naïve individuals [83]. Comparing raltegravir once-daily (two 400mg tablets taken together every 24h) versus twice-daily (one 400mg tablet every 12 hours), both in combination with emtricitabine/tenofovir DF, the once-daily regimen had lower virologic suppression at week 48. Subsequently, the raltegravir HD 600mg tablet was developed and ONCEMRK again investigated the possibility of once-daily raltegravir dosing with the new formulation [84]. The ONCEMRK study found the once-daily raltegravir 1200mg was non-inferior to standard twice-daily raltegravir in virologic suppression at week 48 [84]: 89% of those receiving once-daily raltegravir versus 88% of the twice-daily group. These results continued through week 96, where 81.5% of the once-daily and 80.1% of the twice-daily participants with HIV-RNA <40 copies/mL [85].

In summary, raltegravir was the first-in-class INSTI with demonstrated efficacy, given as twice-daily dosing, for treatment-naïve and -experienced individuals. The raltegravir HD formulation offers an alternative, once-daily dosing regimen for treatment-naïve individuals. Raltegravir is the only INSTI not approved for antiretroviral switch in virologically-suppressed individuals.

5.2 Elvitegravir

Elvitegravir efficacy has been demonstrated in a number of studies, beginning with the FDC containing tenofovir DF in GS102 and GS103 in treatment-naïve individuals. When compared against efavirenz/tenofovir DF/emtricitabine in treatment-naïve individuals, 87.6% of participants receiving elvitegravir had virologic suppression compared with 84.1% for efavirenz at week 48 [86]. Similar rates of virologic suppression were observed through week 144 (elvitegravir, 80.2% vs. efavirenz, 75.3%) [87, 88]. Similar non-inferior performance was observed when the elvitegravir FDC was compared against boosted atazanavir-based regimens in treatment-naïve individuals [89–91]. At week 48, 89.5% of elvitegravir recipients and 86.8% of PI recipients were virologically suppressed [89]; and 77.6% and 74.6%, respectively, remained suppressed through week 144 [91].

The elvitegravir FDC containing tenofovir DF were evaluated in three switch studies for treatment-experienced, virologically suppressed individuals. In participants randomized to continue on NNRTI-based ART or switch to the elvitegravir FDC, no difference was found in virologic suppression at 48 or 96 weeks [92, 93]. Similarly, in individuals randomized to

continue a PI-containing ART regimen or switch to the elvitegravir FDC found no difference in virologic suppression at week 48; at week 96, however, the elvitegravir FDC was superior (87% vs 70%) [94, 95], driven by virologic failures and discontinuations for non-virologic reasons [94]. Finally, in individuals on twice-daily raltegravir-based ART, 100% of 48 study participants remained virologically-suppressed at week 48 after switching to the elvitegravir FDC [96].

Two studies, GS104 and GS111, compared the efficacy of the newer elvitegravir FDC containing tenofovir AF with the elvitegravir FDC containing tenofovir DF in treatment-naïve individuals [97–99]. In GS104, 93% versus 92% of participants in each of the arms met the study-defined endpoint of virologic suppression at week 48, while GS111 reported 92% (tenofovir AF arm) versus 89% (tenofovir DF arm) of participants meeting virologic suppression at week 48 [97]. No differences were found between regimens with respect to baseline CD4+, age or race. Combined long term efficacy data from both studies found similar rates of virologic suppression between groups at week 96 and 144 [98, 99].

The tenofovir AF-containing elvitegravir FDC was also studied in treatment-experienced patients [100]. In GS109, virologically-suppressed adults receiving a tenofovir DF-containing regimen (combined with either efavirenz, elvitegravir/cobicistat, or boosted-atazanavir) for 96 weeks were switched to the elvitegravir FDC containing tenofovir AF, or continued their tenofovir DF-containing ART. At week 48, switching to the tenofovir AF regimen was non-inferior to remaining on the tenofovir DF-containing regimens with virologic suppression rates of 97% vs. 93%, respectively; at week 96, the tenofovir AF arm was statistically favored (93% vs. 89%) [100, 101].

In summary, both elvitegravir FDCs have demonstrated efficacy in clinical trials of antiretroviral-naïve individuals or treatment-experienced, virologically-suppressed individuals who have not previously received INSTIs. The available data suggest the newer FDC formulation of elvitegravir containing tenofovir AF results in similar treatment efficacy compared to the original FDC containing tenofovir DF. Therefore, if elvitegravir is used, clinicians may choose between the formulations based upon clinical considerations related to pill size, cost, or adverse effects.

5.3 Dolutegravir

Dolutegravir efficacy has been demonstrated in studies of treatment-naïve individuals as well as in ART-experienced individuals. In the SINGLE trial, a once-daily regimen of dolutegravir/abacavir/lamivudine was compared against a once-daily regimen of efavirenz/tenofovir DF/emtricitabine in treatment-naïve individuals. At week 48, superiority of the dolutegravir-based regimen was realized with 88% of participants in the dolutegravir arm and 81% in the efavirenz arm achieving virologic suppression [102]. The superiority of dolutegravir continued to week 144, with 71% of dolutegravir versus 63% ($p < 0.01$) of efavirenz recipients maintaining virologic suppression [103]. Interestingly, the dolutegravir arm had lower study drug discontinuation rates at weeks 48 and 144 (2% vs. 10% at week 48 and 3% vs. 11% at week 144), likely due to central nervous system adverse events associated with efavirenz [103, 102]. Dolutegravir was also compared with ritonavir-boosted darunavir in the FLAMINGO trial. In this study of ART naïve-individuals, dolutegravir-

based ART was superior to darunavir-containing regimens at both week 48 and 96 [104, 105].

Dolutegravir was also compared with a first generation INSTI, raltegravir, for treatment-naïve individuals [106]. SPRING-2 randomized participants to either dolutegravir once-daily or raltegravir twice-daily, both with two NRTIs. At week 48, 88% of dolutegravir and 85% of raltegravir recipients achieved virologic suppression, meeting the prespecified non-inferiority margin of 10%. Non-inferiority was confirmed at the week 96 analysis with 81% of the dolutegravir group and 76% of the raltegravir group maintaining virologic suppression [107]. In the SAILING study, dolutegravir was again compared against a raltegravir-based regimen, only this time in ART-experienced participants who were INSTI-naïve but with at least two drug class resistance [108]. Participants were randomly assigned to either dolutegravir once-daily or raltegravir twice-daily in addition to optimized background therapy. At week 48, 71% of dolutegravir and 64% of raltegravir participants were virologically suppressed, demonstrating superiority of dolutegravir in this group of individuals with drug resistance.

The VIKING phase 2b study was the first study to evaluate dolutegravir in participants with prior INSTI resistance, dosed either once- (cohort 1) or twice-daily (cohort 2) [109]. Participants failing a raltegravir-containing regimen substituted dolutegravir for raltegravir for 10 days, then received an optimized background regimen containing dolutegravir through week 24. More participants receiving dolutegravir twice-daily achieved the primary endpoint of at least a 0.7 log₁₀ decline in the HIV-RNA, or HIV-RNA < 400 copies/mL, on day 11 (23/24; 96%) compared with dolutegravir once-daily (21 of 27; 78%). The advantage of twice-daily dosing persisted through 24 weeks (virologic suppression: twice-daily, 75% vs. once-daily group, 41%). The VIKING-3 study investigated dolutegravir twice-daily in participants failing either a raltegravir or elvitegravir based regimen with INSTI resistance [110]. Dolutegravir twice-daily replaced raltegravir or elvitegravir in the failing regimen during the first 7 days of the study, followed by an optimized background regimen according to resistance data. The primary study endpoints were the change in HIV-RNA from baseline to day 8 and the proportion of participants with virologic suppression at week 24. In the intention to treat population of 183 participants, the mean change in plasma HIV-RNA from baseline to day 8 was -1.43 log₁₀ and 69% of the participants were suppressed at week 24.

There are currently two FDA-approved FDC products containing dolutegravir plus a second antiretroviral, representing two-drug combination ART. Efficacy of dolutegravir/rilpivirine was demonstrated in two switch studies in participants receiving three-drug ART [111]. These trials, SWORD-1 and SWORD-2, participants were on a combination of two NRTIs plus either a PI, NNRTI, or INSTI for at least 6 months with virologic suppression. Participants were randomized to continue their current three-drug therapy or switch to the dolutegravir/rilpivirine FDC. In a pooled data analysis, 95% of participants in each treatment arm reached the primary endpoint of virologic suppression at week 48. There were no significant treatment differences by baseline CD4+ count, age, gender, race or baseline ART regimen. At week 52, 477 of the 511 participants originally randomized to the three-drug arm were eligible to switch to dolutegravir/rilpivirine [112]. An analysis of all participants

receiving dolutegravir/rilpivirine at week 100 found 93% of these participants maintained virologic suppression.

Efficacy of dolutegravir/lamivudine was demonstrated in two randomized trials of treatment-naïve adults with baseline HIV-RNA <500,000 copies/mL [113]. In the GEMINI-1 and GEMINI-2, participants were randomized to receive dolutegravir plus emtricitabine/tenofovir DF or dolutegravir plus lamivudine, each given as two separate tablets once-daily. In a pooled analysis, there were no significant differences in virologic suppression at week 48 (91% in the two-drug arm and 93% in the three-drug arm). In each of the trials, lower response rates were observed for individuals with baseline CD4+ counts ≤ 200 cells/mm³ in the two-drug group vs. the three-drug group (81 vs. 90%, respectively, GEMINI-1; 78 vs. 96%, respectively, GEMINI-2), although it must be noted only 8% (n=118) of the total study population had CD4+ ≤ 200 cells/mm³ [114]. Dolutegravir plus lamivudine remained non-inferior to the three-drug regimen at week 96, with 86% and 89.5% of participants in the two and three drug arms maintaining virologic suppression [115].

Clinical trials involving dolutegravir have demonstrated its efficacy in treatment-naïve individuals as part of both traditional, three-drug ART regimens, and as a novel, two-drug ART regimen containing lamivudine as a partner drug. Notably, the three drug regimens containing dolutegravir were superior to efavirenz-based ART and non-inferior to raltegravir-based ART, while dolutegravir/lamivudine was non-inferior to dolutegravir-containing, three-drug ART regimens for antiretroviral-naïve individuals. In addition, dolutegravir is an alternative INSTI for individuals who have failed raltegravir or elvitegravir-based regimens with INSTI resistance. Finally, the dolutegravir/rilpivirine FDC is a switch-strategy for individuals suppressed on another ART combination.

5.4 Bictegravir

Bictegravir efficacy has been demonstrated in two phase 3 randomized trials in HIV treatment-naïve individuals, each comparing the bictegravir FDC with a dolutegravir-based regimen. In study GS1489, the bictegravir FDC was non-inferior to the dolutegravir/abacavir/lamivudine FDC [116, 117]. At week 48, 92.4% of participants in the bictegravir arm and 93% in the dolutegravir arm achieved virologic suppression [116]; non-inferiority was maintained at week 96 (bictegravir: 88% vs. dolutegravir: 90%) [117]. Similar virologic suppression was observed in study GS1490 where the bictegravir FDC was compared with dolutegravir in combination with emtricitabine/tenofovir AF [118, 119]. At week 48, 89% of bictegravir recipients and 93% of dolutegravir were virologically suppressed [118]; non-inferiority was maintained through week 96 (bictegravir: 84% vs. dolutegravir: 86%) [119]. There were no differences in efficacy in participants with lower (<100,000 copies/mL) versus higher (>100,000cpm) HIV-RNA. Nor were there differences in treatment outcomes in subgroup analysis of those with varying baseline CD4+ cell counts.

In switch studies, the bictegravir FDC was non-inferior to remaining on either a dolutegravir-based regimen or a PI-based regimen. In study GS1844, the bictegravir FDC was compared against dolutegravir/abacavir/lamivudine in persons who were virologically suppressed for at least 3 months. Participants were randomly assigned to continue the dolutegravir regimen or switch to the bictegravir FDC regimen. At week 48, 3 of 282 (1%)

participants in the bictegravir arm, and 1 of 281 (<1%) in the dolutegravir arm had HIV-RNA of ≤ 50 copies/mL, demonstrating non-inferiority [120]. In a similar study design with virologically-suppressed participants receiving a boosted atazanavir or darunavir containing ART regimen, switching to the bictegravir FDC was again non-inferior to remaining on the boosted PI regimen. At week 48, each arm had 2% of participants with HIV-RNA ≤ 50 copies/mL [121]. Finally, a switch study looking at participants with pre-existing resistance to NRTIs (eg. M184V, K65R, thymidine analog mutations (TAMs)) found no significant differences in virologic suppression at week 48 between switching to the bictegravir FDC versus a dolutegravir plus emtricitabine/tenofovir AF [122].

In summary, the bictegravir FDC has demonstrated non-inferiority to dolutegravir FDC for individuals who are ART-naïve and those virologically suppressed on another regimen. Recent data suggest bictegravir may be effective in participants with a history of NRTI resistance [122], but data are needed to confirm the expectation that this second-generation INSTI will also offer an effective alternative for individuals with resistance to first-generation INSTIs.

5.5 Cabotegravir

The cabotegravir/rilpivirine combination has been studied in phase 3 clinical trials as once-monthly intramuscular injections [24, 25]. In the FLAIR trial, treatment-naïve adults were given 20 weeks of oral therapy with dolutegravir/abacavir/lamivudine [24]. Participants who reached virologic suppression by week 16 were randomized 1:1 to either continue dolutegravir-based therapy or receive one month of oral cabotegravir and rilpivirine followed by long-acting injectable cabotegravir/rilpivirine. At week 48, 2.1% of the participants in the long-acting arm and 2.5% in the oral therapy arm were virologically suppressed, demonstrating non-inferiority.

A second phase 3 study investigated monthly intramuscular cabotegravir and rilpivirine as maintenance therapy [25]. The ATLAS study enrolled PLWH who were virologically suppressed for at least six months while taking a NNRTI, PI or INSTI based regimen. Dolutegravir/abacavir/lamivudine was excluded in an attempt to broaden the generalizability of the ATLAS study, as dolutegravir-based regimens were the focus of the FLAIR study. Participants were randomized to continue their oral ART or to switch to intramuscular cabotegravir/rilpivirine. Similar to the FLAIR study, participants randomized to the cabotegravir arm of ATLAS were first given a four week lead in of daily oral cabotegravir and rilpivirine, followed by monthly injections of the long-acting cabotegravir/rilpivirine formulations. At week 48, 1.6% of participants in the long acting ART arm and 1.0% in the continued oral therapy arm were virologically suppressed, meeting the predefined criteria for non-inferiority.

The ideal dosing interval for intramuscular cabotegravir/rilpivirine is being explored in ATLAS-2M. Participants who were virologically suppressed on either oral therapy or in the ATLAS study receiving cabotegravir/rilpivirine intramuscularly were randomized to receive either cabotegravir/rilpivirine every four weeks or every eight weeks. Preliminary data show the two dosing strategies met the non-inferiority criteria, with 1.0% of those receiving every

four-week injections and 1.7% of those receiving every eight-week injections maintaining virologic suppression at week 48 [123].

Efficacy data support injectable cabotegravir/rilpivirine for individuals who attain virologic suppression on oral ART. The optimal dosing frequency of injectable cabotegravir/rilpivirine remains an area investigation, with the most data to date supporting once-monthly injections. However, emerging data suggest the potential for future dosing every eight weeks. Cabotegravir is also being investigated as a single agent for injectable pre-exposure prophylaxis, which is beyond the scope of this review [124, 125].

6. Safety and Tolerability

INSTIs are generally associated with lower rates of adverse effects than comparator classes in clinical trials. A summary of safety and tolerability data from phase 3 clinical trials is given in Table 3. Additionally, cohort studies have identified low rates of discontinuation due to adverse effects among INSTIs [11, 126]. Specific concerns related to INSTIs, including neuropsychiatric adverse events (NPAE), fetal toxicity, as well as weight and metabolic complications, have been identified and are discussed in **Sections 6.1-6.3**.

In clinical trials of raltegravir-based ART, mild to moderate rashes were most commonly observed through 96 weeks [81, 127]. Rash was more frequent with raltegravir versus placebo (11.3% and 6.3%) but lower in comparison with efavirenz (9.6% vs. 20.9%) [128]. While Stevens-Johnson Syndrome has been reported with raltegravir, no cases were observed in clinical trials [128]. Further, significantly lower rates of dyslipidemia were observed when compared with efavirenz- [77, 129] or PI-based regimens [130, 131]. Creatinine phosphokinase (CPK) elevations occurred in clinical trials with raltegravir, but none of these cases lead to treatment discontinuation [128, 132]. One cross-sectional, cohort study found higher rates of myalgias (19% vs. 3%, $p < 0.001$) and myopathies (4% vs. 0%, $p < 0.001$) in persons taking raltegravir compared to a control group [133]. There were no differences in CPK elevations (14% vs. 16%, $p < 0.001$) nor any cases of rhabdomyolysis observed, but post-marketing reports of rhabdomyolysis have been reported [134–138].

Overall, a higher rate of adverse effects are reported with elvitegravir compared with other INSTIs, which may be related to the coformulation with cobicistat. Gastrointestinal adverse events (nausea and diarrhea), abnormal dreams, CPK elevations, and headache have been reported in >5% of individuals receiving elvitegravir-based regimens (Table 3) [86, 89, 96, 98]. Compared to dolutegravir or raltegravir, the elvitegravir FDC resulted in higher rates of fatigue, malaise, and gastrointestinal adverse effects [139]. Despite cobicistat-boosting of elvitegravir, lower rates of dyslipidemias, diarrhea, and hyperbilirubinemias were observed with elvitegravir FDC compared with atazanavir/ritonavir-based regimens [89]. Greater increases in serum creatinine and reductions in glomerular filtration rates were observed with the elvitegravir FDC containing tenofovir DF compared with both NNRTI- and PI-based regimens [86, 89]. This adverse effect is secondary to a well described non-pathologic inhibition of renal tubular secretion of serum creatinine by cobicistat [140]. In contrast, the FDC containing tenofovir AF demonstrated less effect on glomerular filtration rate, as well as lower proteinuria, albuminuria, and tubular proteinuria [100]. Additionally, safe use of the

elvitegravir FDC containing tenofovir AF was demonstrated among persons with moderate to severe renal dysfunction (creatinine clearance 30–59 ml/min) and also those on hemodialysis [141].

In clinical trials, dolutegravir has been associated with nausea, headache, and elevation in aspartate aminotransferase and CPK in 5% of participants (Table 3) [102, 106, 111, 113, 116, 142]. Asymptomatic elevations in CPK across trials with dolutegravir intervention arms occurred in 5% (naïve) and 2% (experienced) of patients [132]. Insomnia was reported more often among naïve participants on dolutegravir compared with other antiretrovirals in a recent meta-analysis (6.1% vs. 4.5%; $p=0.02$) [143].

The most common adverse events associated with the bicitegravir FDC in clinical trials were gastrointestinal (diarrhea, nausea), headache, and CPK elevations (Table 3) [116, 118]. In comparison with dolutegravir/abacavir/lamivudine, lower rates of nausea were observed with bicitegravir/emtricitabine/tenofovir AF among treatment-naïve participants at week 96 (11% vs. 24%; $p<0.001$) [117]. Additionally, statistically lower rates of nausea were observed in a trial of participants randomized to switch to bicitegravir/emtricitabine/tenofovir AF or remain on dolutegravir/abacavir/lamivudine (0% vs. 2%, respectively; $p=0.03$) [120]. The difference in nausea may be related to the NRTIs (abacavir vs. tenofovir AF), rather than the INSTI component of the FDC, but because bicitegravir is not available outside of the FDC, it is difficult to separate related adverse events. One cohort study observed low rates of adverse events (8.9%; $n=18/201$) and discontinuations due to adverse events (4%; $n=9/201$) in persons treated with bicitegravir [144]. Rash was the most commonly reported adverse event ($n=10/18$; 56%) and cause for discontinuation ($n=7/9$; 78%).

Due to the high-volume intramuscular injection of cabotegravir/rilpivirine, high rates of injection site reactions (ISR), including pain, nodule and swelling at the injection site were observed among trial participants. The majority receiving injectable cabotegravir/rilpivirine experienced at least one ISR in FLAIR and ATLAS (83% and 82%, respectively) [24, 25]. ISR were typically mild to moderate in severity with most (88%) resolving within 7 days (median, 3 days). A higher incidence of ISRs was reported at week 4 after the loading dose, decreasing over time to week 48 (FLAIR: 69% to 11%; ATLAS: 71% to 20%). Similar ISR outcomes were seen in ATLAS-2M, despite the larger dosing volume in the 8-week dosing arm, as illustrated by the proportion with ISR at weeks 4 and 48, respectively (four-week arm: 53% and 19%; eight-week arm: 70% and 20%) [123]. Reassuringly, low discontinuation rates due to ISRs (range: <1–2%) were observed [24, 25, 123]. Aside from ISR, the most common drug-related adverse events were headache and pyrexia (Table 3) [24, 25].

6.2 Neuropsychiatric Adverse Events

NPAEs have commonly been reported in clinical trials and post-marketing studies of INSTIs. The reported NPAEs primarily include dizziness, depression, anxiety, headache, insomnia and other psychiatric disorders [132]. Higher rates of psychiatric disorders were observed when comparing dolutegravir with darunavir/ritonavir-based ART (6/242 [2.5%] vs. 1/242 [0.4%]) [104]. Yet, most NPAEs secondary to INSTI-based ART have been compared with NNRTI-based regimens and between INSTI-based regimens.

In comparative clinical trials, dolutegravir and elvitegravir-based ART were associated with fewer NPAEs compared with efavirenz-based ART. In STARTMRK, NPAEs were higher for participants receiving efavirenz compared with those receiving raltegravir at weeks 8 (18% vs. 10%, $p=0.0149$) and 48 (23% vs. 14%, $p=0.0044$) [77]. Dizziness (24% vs. 7%), abnormal dreams (27% vs. 15%), and insomnia (14% vs. 9%) were significantly more common (all $p<0.01$) with efavirenz in comparison with elvitegravir [86]. Further, treatment-naïve participants receiving dolutegravir-based ART experienced less NPAEs (dizziness, abnormal dreams, anxiety, somnolence) compared with efavirenz-based ART in the SINGLE trial; however, insomnia was more common in the dolutegravir arm (15% vs. 10%) [102]. Discontinuations across these trials were low and rarely associated with NPAEs

Similar rates of NPAEs (range, 4–6%) were seen within the SPRING trials comparing dolutegravir and raltegravir [106]. This was consistent with randomized clinical trials investigating bicitegravir- versus dolutegravir-based ART, which found similar, but rare, rates of NPAE or discontinuation between treatment arms [116, 120, 142]. Further, NPAEs were rarely observed in the randomized clinical trials of long-acting cabotegravir/rilpivirine in both treatment-naïve and –experienced persons [25, 123]. In contrast, the SWORD studies found NPAEs more common (5% vs. <1%) in the dolutegravir/rilpivirine treatment arm compared with the standard of care arm (INSTI-, PI-, or NNRTI-based ART) and more participants in the dolutegravir/rilpivirine arm discontinued due to NPAE ($n=5$ vs. 1) [111].

Numerous observational cohort studies have evaluated the rates NPAEs and associated ART discontinuation among patients initiated on, or switched to, INSTI-based regimens. In a cohort study of 323 participants, higher rates of NPAEs (depression, vertigo, and sleep disturbances) were observed for dolutegravir compared to elvitegravir or raltegravir (10% vs. 1–2%, respectively), while a larger cohort of 1344 participants on the same regimens found a similar trend for higher, but not statistically different, rate of NPAEs in those receiving dolutegravir (3.5%) compared to raltegravir (2.8%) or elvitegravir (1.6%) [139, 145]. Several cohort studies found higher rates of discontinuation associated with NPAEs in patients receiving dolutegravir compared to either elvitegravir or raltegravir [11, 146–148]. Female sex at birth, elderly persons (>60 years), lower CD4+ cell counts, and use of dolutegravir with or without abacavir have all been identified as predictive factors for therapy discontinuation related to NPAEs in European cohorts [11, 147–149]. However, these predictive factors have not been consistently identified across studies [126, 146]. Few cohort data are available with regard to NPAE incidence among patients treated with bicitegravir or cabotegravir; however, similar rates of NPAE were observed between bicitegravir/emtricitabine/tenofovir AF and dolutegravir/abacavir/lamivudine in one switch study [120].

Overall, though NPAEs are associated with INSTIs, the occurrence of INSTI discontinuation appears to be low based on the evidence to date. Therefore, providers should educate and monitor patients for NPAEs, and modify therapy as needed. Some NPAEs, such as sleep disturbances, may be managed by revising the dose time from evening to morning, while severe NPAEs may require therapy modification.

6.3 Fetal Toxicity

Raltegravir use among pregnant women has demonstrated an acceptable safety profile and low rates of fetal toxicity [49, 50]. The prevalence of birth defects among infants exposed to raltegravir during the first trimester of pregnancy was 3.09% (95% CI: 1.42–5.79) comparable with the US population average rate of 2.72% [150, 151]. One retrospective cohort of pregnant individuals (n=497) receiving raltegravir-based ART found similar rates of birth defects among infants exposed to raltegravir during the first versus the second/third trimesters (5.7% vs. 3.5%, respectively; p=0.29) [152].

Zash et al. reported higher rates of birth defects among infants exposed to dolutegravir at the time of conception that raised concern around the use of dolutegravir at the time of conception or during early pregnancy [153, 154]. In the initial report, rates of birth defects for those who started dolutegravir at conception, started on dolutegravir during pregnancy, started on a non-dolutegravir regimen at conception, or were HIV-negative were 0.94% (4/426), 0.00% (0/2,812), 0.12% (14/11,300), and 0.09% (61/66,057), respectively [153]. After further observations of the same study cohort, the prevalence of neural tube defects was 0.30% (5/1,683) among infants exposed to dolutegravir regimens at conception compared with 0.10% (15/14,792) among infants exposed to non-dolutegravir regimens at conception (difference, 0.20%; 95% CI: 0.01–0.59) [154]. According to the Antiretroviral Pregnancy Registry, the rate of birth defects was 3.6% (11/302) for infants exposed to dolutegravir at any time during pregnancy, while the rate of neural tube defects was 0.004% (1/248) for infants with dolutegravir exposure at periconception [150].

No increased risk of birth defects have been observed in infants exposed to elvitegravir-based ART during pregnancy [150]. The prevalence of birth defects in infants with elvitegravir exposure during the first-trimester was 2.50% (6/240) compared with the U.S. overall prevalence of 2.72% [150, 151]. However, as described in Section 4.1.1 elvitegravir exposure is decreased during pregnancy, suggesting it may not be the optimal regimen for pregnant women [43].

The DHHS Perinatal HIV Treatment Guidelines currently recommend raltegravir- and dolutegravir-based regimens for pregnant PLWH, while dolutegravir-based regimens are an alternative regimen during periconception [43]. The World Health Organization recommends dolutegravir as preferred therapy for all PLWH, irrespective of pregnancy or childbearing potential [9]. There are no data to support the use of bictegravir or cabotegravir among pregnant women, as the prevalence of birth defects among infants exposed to either is unknown.

6.4 Weight Gain and Metabolic Concerns

Recent reports of weight gain associated with the use of INSTI-based ART is a rising concern [155, 156]. A pooled analysis of eight randomized controlled trials among ART-naïve persons with 96 weeks of follow-up found INSTIs, specifically dolutegravir and bictegravir, associated with more weight gain compared with NNRTIs and PIs [157]. The use of tenofovir AF in combination with the INSTI was associated with more weight gain than other NRTI partner drugs. Predictive clinical factors associated with weight gain were

lower CD4+ cell count, higher HIV-RNA, no injection drug use, female sex at birth, and black race.

The NAMSAL trial was a randomized, non-inferiority trial in Africa evaluating three once-daily treatments consisting of dolutegravir in combination with emtricitabine/tenofovir AF, dolutegravir plus emtricitabine/tenofovir DF, or efavirenz 400mg plus emtricitabine/tenofovir DF in treatment-naïve participants; a majority of participants (65.9%) were female [158]. Analogous with Sax et al. [157], larger weight gain and obesity incidence was observed at 48 weeks in participants in the dolutegravir compared with efavirenz arms (median weight gain, 5.0 kg vs. 3.0 kg; incident obesity, 12.3% vs. 5.4%). The ADVANCE trial was a randomized trial in Africa among treatment-naïve participants with the same treatment arms as NAMSAL, except efavirenz was dosed as 600mg daily [159]. The population was nearly all black race and 59% were female. At 96 weeks, weight gain was most common in the dolutegravir groups, particularly those receiving tenofovir AF, and higher among females. Incidence in new obesity was higher overall for females, but a similar trend was observed among treatment arms and sex (incident obesity male vs. female: tenofovir AF group, 20% vs. 7%; tenofovir DF group, 11% vs. 3%; efavirenz group, 9% vs. 3%). Further, increases in mean changes of limb and truncal lean and fat mass were observed.

A retrospective, observational cohort study evaluated differences in weight gain among treatment-naïve persons 18 months after starting ART in the US. Of the 1,152 patients, significantly more weight gain was observed with dolutegravir compared with NNRTIs or elvitegravir (6.0 kg vs. 2.6 kg vs. 0.5kg respectively; $p<0.05$) [160]. Similarly, another observational cohort study of virologically-suppressed persons found larger increases in weight over an 18-month period after switching from efavirenz to either an INSTI- or PI-based regimen compared with continuing efavirenz-based ART [161]. Weight gain was greatest with INSTI-based regimen [efavirenz-based regimen: 0.9 kg vs. INSTI: 2.9 kg, $p=0.003$; or vs. PI: 0.7 kg, $p=0.81$].

Minimal data exist from clinical trials to inform our understanding of weight gain among persons treated with bicitegravir or cabotegravir. Wohl et al. found median weight gain of 3.6 kg (IQR, 0.0–8.5) in persons treated with bicitegravir/emtricitabine/tenofovir AF compared with 2.5 kg (IQR –0.4–5.8) over 96 weeks in those randomized to dolutegravir/abacavir/lamivudine [117]. Small weight increases were observed in persons receiving injectable cabotegravir/rilpivirine in FLAIR (injectable: 1.80 kg [IQR: –0.30–4.90]; oral: 0.30 kg [IQR: –1.60–2.50]) and ATLAS trials (injectable: 1.30 kg [IQR: –1.0–5.0]; oral: 1.50 kg [IQR: –1.0–3.9]) [24, 25]. No changes in weight or fasting metabolic parameters were observed in the phase 2 study evaluating cabotegravir versus placebo in HIV-negative participants using cabotegravir for HIV prevention over 41 weeks [162]. However, the phase 3 trial observed a median 1.3 kg (95% CI 0.99 to 1.6) increase per year in HIV-negative men receiving cabotegravir compared to emtricitabine/tenofovir DF [+0.31 (95% CI: –0.12 to –0.48) kg per year] [163].

Treatment-naïve persons often experience weight gain after ART initiation, which has been described as a return to health [164]. Yet, several observational cohorts have associated the

risk of onset diabetes mellitus and cardiovascular risk with weight gain after ART initiation [165, 166]. At week 96, the ADVANCE trial found the dolutegravir/emtricitabine/tenofovir AF arm had significantly higher risk of metabolic syndrome (International Diabetes Foundation; $p=0.031$), myocardial infarction or coronary death (QRISK; $p=0.027$), and 10-year risk of emergent diabetes (QDIABETES, change from baseline; $p=0.004$) [167]. However, no significant differences in cardiovascular risk (Framingham Risk Equation) were observed among arms. Several case reports highlight incident diabetes mellitus and hyperglycemia in patients taking dolutegravir [168, 169]. A single-center, retrospective cohort study among Ugandan patients identified a higher rate of hyperglycemia after initiation of dolutegravir-based ART compared to ART without dolutegravir (0.47% [16/3417] vs. 0.03% [1/3230]; $p=0.0004$) [170]. The median time to hyperglycemia after initiation of dolutegravir-based ART initiation was 4 months (IQR 2.5–4.5) and weight loss preceding hyperglycemia was observed in 80% (12/15).

Weight gain is a clinical challenge with INSTI-based regimens, particularly second generation INSTI regimens in combination with tenofovir AF [117, 119, 157, 159, 160, 171]. Further, data suggest a higher incidence of weight gain among specific populations including women, those of African descent, and Hispanic ethnicity [157, 159, 172]. Understanding the intersection of weight gain and other resultant metabolic and cardiovascular comorbidities is critical, and further data are needed to support clinical decisions on the selection of ART regimens to minimize weight gain and any associated metabolic comorbidities.

7. INSTIs in the Management of HIV-1 Infection

Clinical trials have demonstrated the INSTIs produce a rapid decline in, and maintain durable suppression of, plasma HIV-RNA, and are generally safe and well-tolerated. The rapid decline in plasma HIV-RNA associated with INSTIs is a unique characteristic compared with older antiretroviral classes, and is particularly relevant to prevent HIV transmission through rapid virologic suppression [8, 43, 74–76]. All INSTIs, except cabotegravir, are indicated for antiretroviral-naïve persons [12, 13, 16, 18, 114, 173]. With the exception of raltegravir, all INSTIs are also indicated for individuals currently suppressed on an ART regimen, who don't have INSTI-related resistance mutations, and desire treatment switch [13, 16, 18, 114, 173]. Intramuscular cabotegravir plus rilpivirine is currently approved in Canada for a non-oral ART option for virologically-suppressed adults [14]. Dolutegravir and raltegravir may be used in treatment-experienced persons with a detectable viral load in the presence of viral resistance [12, 17]. For pediatric patients, the agents have different indications for use, with raltegravir approved for use immediately after birth, dolutegravir for children at least four-weeks of age, and bictegravir and elvitegravir for adolescents weighing at least 25kg [40].

Within the INSTI class, raltegravir has the longest period of use and has minimal drug-drug interactions that cannot be managed by dose adjustment. Based on these characteristics, both the twice-daily and once-daily formulations remain a recommended antiretroviral for most individuals initiating ART in the US DHHS Adult and Adolescent Treatment Guidelines [8] but not the International AIDS Society guidelines [10]. In addition, the twice-daily

formulation is a preferred INSTI for use during pregnancy [43]. Disadvantages related to raltegravir include a higher pill burden compared with other INSTI-based regimens, and a lower barrier to resistance compared with second generation INSTIs, dolutegravir and bictegravir [22, 23]. It offers flexibility for combination with preferred NRTIs based on an individual's characteristics and may be combined with other antiretrovirals to form an ART regimen for persons with drug resistant virus.

Elvitegravir requires pharmaco-enhancement to achieve adequate concentrations, therefore, its coformulation with cobicistat results in a significantly higher risk of drug-drug interactions. In addition, a higher rate of gastrointestinal adverse events, fatigue, and malaise are reported with elvitegravir FDCs [139]. These disadvantages, along with a requirement that the medication be co-administered with food and a lower barrier to antiretroviral resistance, highlight why elvitegravir is no longer preferred for most PLWH [8, 10]. Elvitegravir is not recommended during pregnancy due to suboptimal concentrations and associated risk for virologic failure [43]. One benefit of elvitegravir is a lower risk of weight gain compared with dolutegravir in retrospective evaluations [160]. Elvitegravir is co-formulated with either tenofovir DF or tenofovir AF plus emtricitabine, allowing use of the elvitegravir FDCs in patients with chronic hepatitis B coinfection.

Dolutegravir and bictegravir are recommended options for antiretroviral-naïve individuals due to their minimal drug-drug interactions, ease of use as a FDC, and higher barrier to antiretroviral resistance compared with other INSTIs and some other antiretroviral classes [8, 10]. These characteristics also support the use of both for ART rapid start programs after diagnosis of HIV, and before resistance testing is available [174]. Dolutegravir is also the World Health Organization's only preferred option for ART in all adults, including pregnant women [56]. Dolutegravir is a preferred regimen for pregnant women according to the US DHHS Perinatal HIV Treatment Guidelines, while bictegravir does not yet have data to support its use during pregnancy [43]. Dolutegravir was non-inferior to raltegravir in ART-naïve persons [107], and superior to raltegravir in those with some drug resistance [108]. Though bictegravir has not been evaluated in a phase 3 trial compared with raltegravir, dolutegravir and bictegravir were non-inferior initial therapies for ART-naïve individuals [118, 142]. The choice between the two agents as triple therapy in adults or adolescents is often based upon the desired NRTI companion drugs present in the FDC. Though a complete discussion regarding the benefits of emtricitabine/tenofovir AF versus abacavir/lamivudine is outside the scope of this review, there are concerns related to cardiovascular risks associated with abacavir use and with weight gain related to tenofovir AF, while the emtricitabine/tenofovir AF combination is preferable for individuals with chronic hepatitis B virus coinfection [8]. Concern regarding the contribution of dolutegravir to weight gain after ART initiation is growing; however, it is not known if there is a difference in the associated risk between the two agents due to a shorter duration of clinical use of bictegravir. Finally, the bictegravir FDC is more susceptible to drug-drug interactions that cannot be overcome with dose adjustment and it is not indicated in combination with any other antiretroviral [16]. In contrast, dolutegravir may be given twice-daily, to overcome drug-drug interactions and as a treatment option for individuals with drug resistance [17].

The two-drug regimens of dolutegravir/lamivudine or dolutegravir/rilpivirine may be useful for individuals at risk for adverse events related to the companion drugs in INSTI FDCs, such as abacavir or the tenofovir prodrugs. The dolutegravir/lamivudine FDC is the only two-drug combination recommended as an initial regimen for most PLWH (with an HIV-RNA level $\leq 500,000$ copies/mL) according to the DHHS Adult and Adolescent Treatment Guidelines [8], while either two-drug regimen may be useful for treatment switch in those already suppressed on another ART regimen [175]. The dolutegravir/rilpivirine combination has added food restrictions and drug-drug interaction considerations, due to the rilpivirine component. Neither dual-therapy agent has been evaluated in adolescents, nor can they be used for patients coinfecting with chronic hepatitis B virus.

Cabotegravir is not currently included in HIV treatment guidelines, but this long-acting agent, in combination with rilpivirine, offers an alternative to PLWH who do not want to take daily oral medications. Although clinical trials have been conducted in treatment-naïve individuals initiating therapy and virologically suppressed individuals desiring treatment simplification [24, 25], the recent Canadian approval was only for treatment simplification [14]. In clinical trials, participants preferred long-acting injectable ART over oral therapy and eight-week dosing over every four-week dosing [24, 25, 123]. These results, however, may be skewed given all subjects volunteered to participate in long-acting therapy trials. The acceptability and demand for intramuscular injections outside of a clinical trial setting is unknown. Intramuscular injection will avoid some absorption related drug-drug interactions with divalent cations and food requirements for rilpivirine, but it is not yet clear how coadministration of UGT1A1 inducers or inhibitors will be managed in combination with long-acting cabotegravir.

INSTIs are part of the standard of care for most PLWH. Future ART strategies will need to demonstrate non-inferiority to the second generation INSTIs to have a role in first-line ART therapy. As long-acting ART emerges as a treatment option, clinicians will need to confront challenges managing the prolonged pharmacokinetic tail after long-acting ART discontinuation [37], how monthly or bimonthly injections will be administered in clinical settings, and non-adherence. Virologic failure has occurred in clinical trial participants receiving long-acting therapy who had perfect adherence (i.e., did not miss any injections) [24, 25]. These findings indicate the need for dosing regimen optimization, particularly as this strategy moves to settings of less than perfect adherence, and investigations of individual characteristics, such as sex and high body-mass-index that may be associated with an increased risk for virologic failure. The contribution of individual INSTIs to weight gain remains an area of clinical investigation and may influence the choice of ART. INSTIs now represent an essential component of most ART regimens and are likely to remain an important class of antiretrovirals for the foreseeable future due to their anti-HIV potency, efficacy, safety and tolerability.

Funding:

We acknowledge support from the following grants from the National Institutes of Health: 1R01HD085887-01A1 (to KS), 1K23AI134307 (to ATP), RO1 AI124965-01 and UM1AI06701 (to CVF). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References:

1. UNAIDS. Global HIV & AIDS statistics - 2019 fact sheets. <https://www.unaids.org/en/resources/fact-sheet> Accessed April 2, 2020.
2. Samji H, Cescon A, Hogg RS, Modur SP, Althoff KN, Buchacz K et al. Closing the gap: increases in life expectancy among treated HIV-positive individuals in the United States and Canada. *PLoS One*. 2013;8(12):e81355. doi:10.1371/journal.pone.0081355. [PubMed: 24367482]
3. Group ISS, Lundgren JD, Babiker AG, Gordin F, Emery S, Grund B et al. Initiation of Antiretroviral Therapy in Early Asymptomatic HIV Infection. *N Engl J Med*. 2015;373(9):795–807. doi:10.1056/NEJMoa1506816. [PubMed: 26192873]
4. Group TAS, Danel C, Moh R, Gabillard D, Badje A, Le Carrou J et al. A Trial of Early Antiretrovirals and Isoniazid Preventive Therapy in Africa. *N Engl J Med*. 2015;373(9):808–22. doi:10.1056/NEJMoa1507198. [PubMed: 26193126]
5. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, Kumarasamy N et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. *N Engl J Med*. 2016;375(9):830–9. doi:10.1056/NEJMoa1600693. [PubMed: 27424812]
6. Rodger AJ, Cambiano V, Bruun T, Vernazza P, Collins S, van Lunzen J et al. Sexual Activity Without Condoms and Risk of HIV Transmission in Serodifferent Couples When the HIV-Positive Partner Is Using Suppressive Antiretroviral Therapy. *JAMA*. 2016;316(2):171–81. doi:10.1001/jama.2016.5148. [PubMed: 27404185]
7. Townsend CL, Cortina-Borja M, Peckham CS, de Ruiter A, Lyall H, Tookey PA. Low rates of mother-to-child transmission of HIV following effective pregnancy interventions in the United Kingdom and Ireland, 2000–2006. *AIDS*. 2008;22(8):973–81. doi:10.1097/QAD.0b013e3282f9b67a. [PubMed: 18453857]
8. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV. Department of Health and Human Services Available at <http://aidsinfo.nih.gov/contentfiles/lvguidelines/AdultandAdolescentGL.pdf>.
9. Update of recommendations on first- and second-line antiretroviral regimens. Geneva, Switzerland: World Health Organization; 2019 (WHO/CDS/HIV/19.15). Licence: CC BY-NC-SA 3.0 IGO.
10. Saag MS, Benson CA, Gandhi RT, Hoy JF, Landovitz RJ, Mugavero MJ et al. Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2018 Recommendations of the International Antiviral Society-USA Panel. *JAMA*. 2018;320(4):379–96. doi:10.1001/jama.2018.8431. [PubMed: 30043070]
11. Peñafiel J, De LE, Padilla M, Rojas J, Gonzalez-Cordon A, Blanco JL et al. Tolerability of integrase inhibitors in a real-life setting. *Journal of Antimicrobial Chemotherapy*. 2017;72(6):1752–9. doi:10.1093/jac/dkx053. [PubMed: 28333231]
12. Isentress® [package insert]. Whitehouse Station, NJ Merck & Co., Inc. 1 2019.
13. Stribild® [package insert]. Foster City, CA Gilead Sciences, Inc 1 2019.
14. Cabenuva® and Vocabria® [Product Monograph]. Laval, Quebec Viiv Healthcare ULC 3 2020.
15. Dehority W, Abadi J, Wiznia A, Viani RM. Use of Integrase Inhibitors in HIV-Infected Children and Adolescents. *Drugs*. 2015;75(13):1483–97. doi:10.1007/s40265-015-0446-2. [PubMed: 26242765]
16. Biktarvy® [package insert]. Foster City, CA Gilead Sciences, Inc 2 2018.
17. Tivicay® [package insert]. Research Triangle Park, NC Viiv Healthcare 6 2020.
18. Genvoya® [package insert]. Foster City, CA Gilead Sciences, Inc 2 2019.
19. Kobayashi M, Yoshinaga T, Seki T, Wakasa-Morimoto C, Brown KW, Ferris R et al. In vitro antiretroviral properties of S/GSK1349572, a next-generation HIV integrase inhibitor; Glaxo SmithKline(United States); Sequoia(United Kingdom); Shionogi(Japan). *Antimicrobial Agents and Chemotherapy*. 2011;55(2):813–21. doi:10.1128/AAC.01209-10. [PubMed: 21115794]
20. Podany AT, Scarsi KK, Pham MM, Fletcher CV. Comparative Clinical Pharmacokinetics and Pharmacodynamics of HIV-1 Integrase Strand Transfer Inhibitors: An Updated Review. *Clin Pharmacokinet*. 2020. doi:10.1007/s40262-020-00898-8.

21. Wainberg MA, Zaharatos GJ, Brenner BG. Development of antiretroviral drug resistance. *N Engl J Med.* 2011;365(7):637–46. doi:10.1056/NEJMra1004180. [PubMed: 21848464]
22. Anstett K, Brenner B, Mesplede T, Wainberg MA. HIV drug resistance against strand transfer integrase inhibitors. *Retrovirology.* 2017;14(1):36. doi:10.1186/s12977-017-0360-7. [PubMed: 28583191]
23. Tsiang M, Jones GS, Goldsmith J, Mulato A, Hansen D, Kan E et al. Antiviral activity of bictegravir (GS-9883), a novel potent HIV-1 integrase strand transfer inhibitor with an improved resistance profile; Gilead; Toronto Research Chemicals(Canada); Sigma(United States). *Antimicrobial Agents and Chemotherapy.* 2016;60(12):7086–97. doi:10.1128/AAC.01474-16. [PubMed: 27645238]
24. Orkin C, Arasteh K, Gorgolas Hernandez-Mora M, Pokrovsky V, Overton ET, Girard PM et al. Long-Acting Cabotegravir and Rilpivirine after Oral Induction for HIV-1 Infection. *N Engl J Med.* 2020;382(12):1124–35. doi:10.1056/NEJMoa1909512. [PubMed: 32130806]
25. Swindells S, Andrade-Villanueva JF, Richmond GJ, Rizzardini G, Baumgarten A, Masia M et al. Long-Acting Cabotegravir and Rilpivirine for Maintenance of HIV-1 Suppression. *N Engl J Med.* 2020;382(12):1112–23. doi:10.1056/NEJMoa1904398. [PubMed: 32130809]
26. Boffito M, Waters L, Cahn P, Paredes R, Koteff J, van WJ et al. Perspectives on the Barrier to Resistance for Dolutegravir + Lamivudine, a 2-Drug Antiretroviral Therapy for HIV-1 Infection. *AIDS Research and Human Retroviruses.* 2019. doi:10.1089/AID.2019.0171.
27. Spreen W, Min S, Ford S, Chen S, Lou Y, Bomar M et al. Pharmacokinetics, safety, and monotherapy antiviral activity of GSK1265744, an HIV integrase strand transfer inhibitor; Glaxo SmithKline(United States). *HIV Clinical Trials.* 2013;14(5):192–203. doi:10.1310/hct1405-192. [PubMed: 24144896]
28. Min S, Sloan L, DeJesus E, Hawkins T, McCurdy L, Song I et al. Antiviral activity, safety, and pharmacokinetics/pharmacodynamics of dolutegravir as 10-day monotherapy in HIV-1-infected adults. *AIDS.* 2011;25(14):1737–45. doi:10.1097/QAD.0b013e32834a1dd9. [PubMed: 21716073]
29. DeJesus E, Berger D, Markowitz M, Cohen C, Hawkins T, Ruane P et al. Antiviral activity, pharmacokinetics, and dose response of the HIV-1 integrase inhibitor GS-9137 (JTK-303) in treatment-naïve and treatment-experienced patients; Gilead. *Journal of acquired immune deficiency syndromes.* 2006;43(1):1–5. doi:10.1097/01.qai.0000233308.82860.2f. [PubMed: 16936557]
30. Gallant JE, Thompson M, DeJesus E, Voskuhl GW, Wei X, Zhang H et al. Antiviral activity, safety, and pharmacokinetics of bictegravir as 10-day monotherapy in HIV-1-infected adults. *Journal of acquired immune deficiency syndromes.* 2017;75(1):61–6. doi:10.1097/QAI.0000000000001306. [PubMed: 28196003]
31. Rizk ML, Hang Y, Luo WL, Su J, Zhao J, Campbell H et al. Pharmacokinetics and pharmacodynamics of once-daily versus twice-daily raltegravir in treatment-naïve HIV-infected patients. *Antimicrobial Agents and Chemotherapy.* 2012;56(6):3101–6. doi:10.1128/AAC.06417-11. [PubMed: 22430964]
32. Food and Drug Administration (FDA). Guidance for Industry, Antiviral Product Development, Conducting and Submitting Virology Studies to the Agency. 2006 <https://www.fda.gov/media/71223/download> Accessed March 8, 2020.
33. Cattaneo D, Gervasoni C. Pharmacokinetics and Pharmacodynamics of Cabotegravir, a Long-Acting HIV Integrase Strand Transfer Inhibitor. *European journal of drug metabolism and pharmacokinetics.* 2019;44(3):319–27. doi:10.1007/s13318-018-0526-2. [PubMed: 30387005]
34. Bowers GD, Culp A, Reese MJ, Tabolt G, Moss L, Piscitelli S et al. Disposition and metabolism of cabotegravir: A comparison of biotransformation and excretion between different species and routes of administration in humans; Glaxo SmithKline. *Xenobiotica.* 2016;46(2):147–62. doi:10.3109/00498254.2015.1060372. [PubMed: 26134155]
35. Ford S, Crauwels H, Han K, Rossenu S, Zhang F, Huang JO, Margolis DA, Sutton K, Hudson KJ, Williams PE, Spreen W, Patel P. Cabotegravir and rilpivirine PK following long-acting HIV treatment discontinuation Conference on Retroviruses and Opportunistic Infections. 3 8–11, 2020 Boston, MA Abstract #466.
36. Spreen W, Ford SL, Chen S, Wilfret D, Margolis D, Gould E et al. GSK1265744 pharmacokinetics in plasma and tissue following single-dose long-acting (la) injectable administration in healthy

- subjects. *Journal of acquired immune deficiency syndromes*. 2014. doi:10.1097/QAI.0000000000000301.
37. Landovitz RJ, Li S, Eron JJ Jr., Grinsztejn B, Dawood H, Liu AY et al. Tail-phase safety, tolerability, and pharmacokinetics of long-acting injectable cabotegravir in HIV-uninfected adults: a secondary analysis of the HPTN 077 trial. *Lancet HIV*. 2020;7(7):e472–e81. doi:10.1016/S2352-3018(20)30106-5. [PubMed: 32497491]
 38. Vitekta® [package insert]. Foster City, CA Gilead Sciences, Inc 9 2014.
 39. Wohl DA, Dumond JB, Blevins S, Pittard D, Ragan D, Wang R et al. Raltegravir pharmacokinetics in treatment-naive patients is not influenced by race: Results from the raltegravir early therapy in african-americans living with HIV (REAL) Study. *Antimicrobial Agents and Chemotherapy*. 2013;57(2):784–8. doi:10.1128/AAC.01826-12. [PubMed: 23183440]
 40. Panel on Antiretroviral Therapy and Medical Management of Children Living with HIV. Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection. Available at <http://aidsinfo.nih.gov/contentfiles/lvguidelines/pediatricguidelines.pdf>.
 41. Lu H, Rosenbaum S. Developmental pharmacokinetics in pediatric populations. *J Pediatr Pharmacol Ther*. 2014;19(4):262–76. doi:10.5863/1551-6776-19.4.262. [PubMed: 25762871]
 42. Elliot ER, Wang X, Singh S, Simmons B, Vera JH, Miller RF et al. Increased Dolutegravir Peak Concentrations in People Living With Human Immunodeficiency Virus Aged 60 and Over, and Analysis of Sleep Quality and Cognition. *Clin Infect Dis*. 2019;68(1):87–95. doi:10.1093/cid/ciy426. [PubMed: 29771285]
 43. Panel on Treatment of Pregnant Women with HIV Infection and Prevention of Perinatal Transmission. Recommendations for the Use of Antiretroviral Drugs in Pregnant Women with HIV Infection and Interventions to Reduce Perinatal HIV Transmission in the United States. Available at <http://aidsinfo.nih.gov/contentfiles/lvguidelines/PerinatalGL.pdf>. Accessed (insert date).
 44. Colbers A, De HM, Van CR, Kruijssen M, Duisenberg-Van EM, Abbink E et al. Pharmacokinetics of crushed elvitegravir combination tablet given with drip feed. *Topics in Antiviral Medicine*. 2016;24(-1):166.
 45. Jeong H, Choi S, Song JW, Chen H, Fischer JH. Regulation of UDP-glucuronosyltransferase (UGT) 1A1 by progesterone and its impact on labetalol elimination. *Xenobiotica*. 2008;38(1):62–75. doi:10.1080/00498250701744633. [PubMed: 18098064]
 46. van der Galien R, Ter Heine R, Greupink R, Schalkwijk SJ, van Herwaarden AE, Colbers A et al. Pharmacokinetics of HIV-Integrase Inhibitors During Pregnancy: Mechanisms, Clinical Implications and Knowledge Gaps. *Clin Pharmacokinet*. 2019;58(3):309–23. doi:10.1007/s40262-018-0684-z. [PubMed: 29915921]
 47. Mulligan N, Best BM, Wang J, Capparelli EV, Stek A, Barr E et al. Dolutegravir pharmacokinetics in pregnant and postpartum women living with HIV. *AIDS*. 2018;32(6):729–37. doi:10.1097/QAD.0000000000001755. [PubMed: 29369162]
 48. Waitt C, Orrell C, Walimbwa S, Singh Y, Kintu K, Simmons B et al. Safety and pharmacokinetics of dolutegravir in pregnant mothers with HIV infection and their neonates: A randomised trial (DolPHIN-1 study). *PLoS medicine*. 2019;16(9):e1002895. doi:10.1371/journal.pmed.1002895. [PubMed: 31539371]
 49. Blonk MI, Colbers APH, Hidalgo-Tenorio C, Kabeya K, Weizsäcker K, Haberl AE et al. Raltegravir in HIV1-infected pregnant women: Pharmacokinetics, safety, and efficacy. *Clinical Infectious Diseases*. 2015;61(5):809–16. doi:10.1093/cid/civ366. [PubMed: 25944344]
 50. Watts DH, Stek A, Best BM, Wang J, Capparelli EV, Cressey TR et al. Raltegravir pharmacokinetics during pregnancy. *Journal of acquired immune deficiency syndromes*. 2014. doi:10.1097/QAI.0000000000000318.
 51. Momper JD, Best BM, Wang J, Capparelli EV, Stek A, Barr E et al. Elvitegravir/cobicistat pharmacokinetics in pregnant and postpartum women with HIV. *AIDS*. 2018;32(17):2507–16. doi:10.1097/QAD.0000000000001992. [PubMed: 30134297]
 52. Patel P, Thiagarajah S, Ford S, Margolis DA, Romach BH, Baker M, Sutton K, Harrinton CM, Shaefer MS, Spreen W, Smith K, Vannappagari V. Cabotegravir pharmacokinetic tail in pregnancy

and neonatal outcomes Conference on Retroviruses and Opportunistic Infections. 3 8–11, 2020 Boston, MA Abstract #775.

53. Schalkwijk S, Greupink R, Colbers AP, Wouterse AC, Verweij VG, van Drongelen J et al. Placental transfer of the HIV integrase inhibitor dolutegravir in an ex vivo human cotyledon perfusion model. *J Antimicrob Chemother.* 2016;71(2):480–3. doi:10.1093/jac/dkv358. [PubMed: 26538508]
54. Schalkwijk S, Colbers A, Konopnicki D, Greupink R, Russel FGM, Burger D. First reported use of elvitegravir and cobicistat during pregnancy. *AIDS.* 2016;30(5):807–8. doi:10.1097/QAD.0000000000000976. [PubMed: 26913711]
55. McCormack SA, Best BM. Protecting the fetus against HIV infection: a systematic review of placental transfer of antiretrovirals. *Clin Pharmacokinet.* 2014;53(11):989–1004. doi:10.1007/s40262-014-0185-7. [PubMed: 25223699]
56. World Health Organization, United Nations Children’s Fund. Guideline: updates on HIV and infant feeding: the duration of breastfeeding, and support from health services to improve feeding practices among mothers living with HIV. Geneva: World Health Organization; 2016.
57. Iwamoto M, Hanley WD, Petry AS, Friedman EJ, Kost JT, Breidinger SA et al. Lack of a clinically important effect of moderate hepatic insufficiency and severe renal insufficiency on raltegravir pharmacokinetics; Merck and Co(United States). *Antimicrobial Agents and Chemotherapy.* 2009;53(5):1747–52. doi:10.1128/AAC.01194-08. [PubMed: 19223645]
58. Weller S, Borland J, Chen S, Johnson M, Savina P, Wynne B et al. Pharmacokinetics of dolutegravir in HIV-seronegative subjects with severe renal impairment. *European journal of clinical pharmacology.* 2014;70(1):29–35. doi:10.1007/s00228-013-1590-9. [PubMed: 24096683]
59. Debinski HS, Lee CS, Danks JA, Mackenzie PI, Desmond PV. Localization of uridine 5'-diphosphateglucuronosyltransferase in human liver injury. *Gastroenterology.* 1995;108(5):1464–9. doi:10.1016/0016-5085(95)90695-9. [PubMed: 7729639]
60. Furlan V, Demirdjian S, Bourdon O, Magdalou J, Taburet AM. Glucuronidation of drugs by hepatic microsomes derived from healthy and cirrhotic human livers. *J Pharmacol Exp Ther.* 1999;289(2):1169–75. [PubMed: 10215701]
61. George J, Murray M, Byth K, Farrell GC. Differential alterations of cytochrome P450 proteins in livers from patients with severe chronic liver disease. *Hepatology.* 1995;21(1):120–8. [PubMed: 7806144]
62. Calza L, Danese I, Colangeli V, Manfredi R, Magistrelli E, Verucchi G et al. Plasma concentrations of efavirenz, darunavir/ritonavir and raltegravir in HIV-HCV-coinfected patients without liver cirrhosis in comparison with HIV-monoinfected patients. *Infectious Diseases.* 2015;47(9):625–36. doi:10.3109/23744235.2015.1034169. [PubMed: 25875396]
63. Custodio JM, Rhee M, Shen G, Ling KHJ, Kearney BP, Ramanathan S. Pharmacokinetics and safety of boosted elvitegravir in subjects with hepatic impairment. *Antimicrobial Agents and Chemotherapy.* 2014;58(5):2564–9. doi:10.1128/AAC.02180-13. [PubMed: 24550332]
64. Hernández-Novoa B, Moreno A, Pérez-Elías MJ, Quereda C, Drona F, Casado JL et al. Raltegravir pharmacokinetics in HIV/HCV-coinfected patients with advanced liver cirrhosis (Child-Pugh C); Merck Sharp and Dohme(United States). *Journal of Antimicrobial Chemotherapy.* 2014;69(2):471–5. doi:10.1093/jac/dkt386. [PubMed: 24097843]
65. Song IH, Borland J, Savina PM, Chen S, Patel P, Wajima T et al. Pharmacokinetics of Single-Dose Dolutegravir in HIV-Seronegative Subjects With Moderate Hepatic Impairment Compared to Healthy Matched Controls. *Clinical Pharmacology in Drug Development.* 2013;2(4):342–8. doi:10.1002/cpdd.55. [PubMed: 26097786]
66. Shaik JSB, Ford SL, Lou Y, Zhang Z, Bakshi KK, Tenorio AR et al. A Phase 1 Study to Evaluate the Pharmacokinetics and Safety of Cabotegravir in Patients With Hepatic Impairment and Healthy Matched Controls. *Clinical Pharmacology in Drug Development.* 2019;8(5):664–73. doi:10.1002/cpdd.655. [PubMed: 30811880]
67. Dooley KE, Sayre P, Borland J, Purdy E, Chen S, Song I et al. Safety, tolerability, and pharmacokinetics of the HIV integrase inhibitor dolutegravir given twice daily with rifampin or once daily with rifabutin: Results of a phase 1 study among healthy subjects. *Journal of acquired immune deficiency syndromes.* 2013;62(1):21–7. doi:10.1097/QAI.0b013e318276cda9. [PubMed: 23075918]

68. Dooley KE, Kaplan R, Mwelase N, Grinsztejn B, Ticona E, Lacerda M et al. Dolutegravir-Based Antiretroviral Therapy for Patients Co-Infected with Tuberculosis and Hiv: A Multicenter, Noncomparative, Open-Label, Randomized Trial. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2019. doi:10.1093/cid/ciz256.
69. Grinsztejn B, De CN, Arnold V, Veloso VG, Morgado M, Pilotto JH et al. Raltegravir for the treatment of patients co-infected with HIV and tuberculosis (ANRS 12 180 Reflate TB): A multicentre, phase 2, noncomparative, open-label, randomised trial; Bristol Myers Squibb(France); ViiV(France); Gilead(United States); Merck(United States). *The Lancet Infectious Diseases*. 2014;14(6):459–67. doi:10.1016/S1473-3099(14)70711-X. [PubMed: 24726095]
70. Ford SL, Sutton K, Lou Y, Zhang Z, Tenorio A, Trezza C et al. Effect of rifampin on the single-dose pharmacokinetics of oral cabotegravir in healthy subjects. *Antimicrobial Agents and Chemotherapy*. 2017;61(10). doi:10.1128/AAC.00487-17.
71. Rajoli RKR, Curley P, Chiong J, Back D, Flexner C, Owen A et al. Predicting drug-drug interactions between rifampicin and long-acting cabotegravir and rilpivirine using physiologically based pharmacokinetic modeling. *Journal of Infectious Diseases*. 2019;219(11):1735–42. doi:10.1093/infdis/jiy726. [PubMed: 30566691]
72. Reese MJ, Savina PM, Generaux GT, Tracey H, Humphreys JE, Kanaoka E et al. In vitro investigations into the roles of drug transporters and metabolizing enzymes in the disposition and drug interactions of dolutegravir, a hiv integrase inhibitor; Amersham Biosciences(United States); GE Healthcare(United States); Glaxo SmithKline(United Kingdom). *Drug Metabolism and Disposition*. 2013;41(2):353–61. doi:10.1124/dmd.112.048918. [PubMed: 23132334]
73. Cattaneo D, Resnati C, Rizzardini G, Gervasoni C. Dolutegravir and metformin: a clinically relevant or just a pharmacokinetic interaction? *AIDS*. 2018;32(4):532–3. doi:10.1097/QAD.0000000000001720. [PubMed: 29381562]
74. Acosta RK, Willkom M, Martin R, Chang S, Wei X, Garner W et al. Resistance analysis of bictegravir-emtricitabine-tenofovir alafenamide in HIV-1 treatment-naïve patients through 48 weeks. *Antimicrobial Agents and Chemotherapy*. 2019;63(5). doi:10.1128/AAC.02533-18.
75. Kintu K, Malaba TR, Nakibuka J, Papamichael C, Colbers A, Byrne K et al. Dolutegravir versus efavirenz in women starting HIV therapy in late pregnancy (DolPHIN-2): an open-label, randomised controlled trial. *Lancet HIV*. 2020;7(5):e332–e9. doi:10.1016/S2352-3018(20)30050-3. [PubMed: 32386721]
76. Jacobson K, Ogbuagu O. Integrase inhibitor-based regimens result in more rapid virologic suppression rates among treatment-naïve human immunodeficiency virus-infected patients compared to non-nucleoside and protease inhibitor-based regimens in a real-world clinical setting: A retrospective cohort study. *Medicine (Baltimore)*. 2018;97(43):e13016. doi:10.1097/MD.00000000000013016. [PubMed: 30412140]
77. Lennox JL, DeJesus E, Lazzarin A, Pollard RB, Madruga JVR, Berger DS et al. Safety and efficacy of raltegravir-based versus efavirenz-based combination therapy in treatment-naïve patients with HIV-1 infection: a multicentre, double-blind randomised controlled trial. *Lancet*. 2009;374(9692):796–806. doi:10.1016/S0140-6736(09)60918-1. [PubMed: 19647866]
78. Rockstroh JK, DeJesus E, Lennox JL, Yazdanpanah Y, Saag MS, Wan H et al. Durable efficacy and safety of raltegravir versus efavirenz when combined with tenofovir/emtricitabine in treatment-naïve HIV-1-infected patients: final 5-year results from STARTMRK. *J Acquir Immune Defic Syndr*. 2013;63(1):77–85. doi:10.1097/QAI.0b013e31828ace69. [PubMed: 23412015]
79. Eron JJ, Young B, Cooper DA, Youle M, DeJesus E, Andrade-Villanueva J et al. Switch to a raltegravir-based regimen versus continuation of a lopinavir-ritonavir-based regimen in stable HIV-infected patients with suppressed viraemia (SWITCHMRK 1 and 2): two multicentre, double-blind, randomised controlled trials; Abbott(United States); Merck(United States). *Lancet*. 2010;375(9712):396–407. doi:10.1016/S01406736(09)62041-9. [PubMed: 20074791]
80. Steigbigel RT, Cooper DA, Kumar PN, Eron JE, Schechter M, Markowitz M et al. Raltegravir with optimized background therapy for resistant HIV-1 infection; Merck(United States). *New England Journal of Medicine*. 2008;359(4):339–54. doi:10.1056/NEJMoa0708975. [PubMed: 18650512]
81. Steigbigel RT, Cooper DA, Teppler H, Eron JJ, Gatell JM, Kumar PN et al. Long-term efficacy and safety of raltegravir combined with optimized background therapy in treatment-experienced

- patients with drugresistant hiv infection: Week 96 results of the benchmrk 1 and 2 phase III trials. *Clinical Infectious Diseases*. 2010;50(4):605–12. doi:10.1086/650002. [PubMed: 20085491]
82. Eron JJ, Cooper DA, Steigbigel RT, Clotet B, Gatell JM, Kumar PN et al. Efficacy and safety of raltegravir for treatment of HIV for 5 years in the BENCHMRK studies: final results of two randomised, placebo-controlled trials. *Lancet Infect Dis*. 2013;13(7):587–96. doi:10.1016/S1473-3099(13)70093-8. [PubMed: 23664333]
 83. Eron JJ, Rockstroh JK, Reynes J, Andrade-Villanueva J, Ramalho-Madruga JV, Bekker LG et al. Raltegravir once daily or twice daily in previously untreated patients with HIV-1: A randomised, active-controlled, phase 3 non-inferiority trial; Merck. *The Lancet Infectious Diseases*. 2011;11(12):907–15. doi:10.1016/S1473-3099(11)70196-7. [PubMed: 21933752]
 84. Cahn P, Kaplan R, Sax PE, Squires K, Molina JM, Avihingsanon A et al. Raltegravir 1200 mg once daily versus raltegravir 400 mg twice daily, with tenofovir disoproxil fumarate and emtricitabine, for previously untreated HIV-1 infection: a randomised, double-blind, parallel-group, phase 3, non-inferiority trial; Gilead(Ireland). *Lancet HIV*. 2017;4(11):e486–e94. doi:10.1016/S2352-3018(17)30128-5. [PubMed: 28918877]
 85. Cahn P, Sax PE, Squires K, Molina JM, Ratanasuwan W, Rassool M et al. Raltegravir 1200 mg Once Daily vs 400 mg Twice Daily, With Emtricitabine and Tenofovir Disoproxil Fumarate, for Previously Untreated HIV-1 Infection: Week 96 Results From ONCEMRK, a Randomized, Double-Blind, Noninferiority Trial. *J Acquir Immune Defic Syndr*. 2018;78(5):589–98. doi:10.1097/QAI.0000000000001723. [PubMed: 29771789]
 86. Sax PE, DeJesus E, Mills A, Zolopa A, Cohen C, Wohl D et al. Co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus co-formulated efavirenz, emtricitabine, and tenofovir for initial treatment of HIV-1 infection: A randomised, double-blind, phase 3 trial, analysis of results after 48 weeks. *Lancet*. 2012;379(9835):2439–48. doi:10.1016/S0140-6736(12)60917-9. [PubMed: 22748591]
 87. Wohl DA, Cohen C, Gallant JE, Mills A, Sax PE, DeJesus E et al. A randomized, double-blind comparison of single tablet regimen elvitegravir/cobicistat/emtricitabine/tenofovir DF versus single tablet regimen efavirenz/emtricitabine/tenofovir DF for initial treatment of HIV-1 infection: Analysis of week 144 results. *Journal of acquired immune deficiency syndromes*. 2013. doi:10.1097/QAI.0000000000000057.
 88. Zolopa A, Sax PE, Dejesus E, Mills A, Cohen C, Wohl D et al. A randomized double-blind comparison of coformulated elvitegravir/ cobicistat/emtricitabine/tenofovir disoproxil fumarate versus efavirenz/emtricitabine/tenofovir disoproxil fumarate for initial treatment of HIV-1 infection: Analysis of week 96 results. *Journal of acquired immune deficiency syndromes*. 2013;63(1):96–100. doi:10.1097/QAI.0b013e318289545c. [PubMed: 23392460]
 89. DeJesus E, Rockstroh JK, Henry K, Molina JM, Gathe J, Ramanathan S et al. Co-formulated elvitegravir, cobicistat, emtricitabine, and tenofovir disoproxil fumarate versus ritonavir-boosted atazanavir plus co-formulated emtricitabine and tenofovir disoproxil fumarate for initial treatment of HIV-1 infection: A randomised, double-blind, phase 3, non-inferiority trial. *Lancet*. 2012;379(9835):2429–38. doi:10.1016/S0140-6736(12)60918-0. [PubMed: 22748590]
 90. Rockstroh JK, DeJesus E, Henry K, Molina JM, Gathe J, Ramanathan S et al. A randomized, double-blind comparison of coformulated elvitegravir/cobicistat/emtricitabine/tenofovir DF vs ritonavir-boosted atazanavir plus coformulated emtricitabine and tenofovir DF for initial treatment of HIV-1 infection: analysis of week 96 results. *J Acquir Immune Defic Syndr*. 2013;62(5):483–6. doi:10.1097/QAI.0b013e318286415c. [PubMed: 23337366]
 91. Clumeck N, Molina JM, Henry K, Gathe J, Rockstroh JK, DeJesus E et al. A randomized, double-blind comparison of single-tablet regimen elvitegravir/cobicistat/emtricitabine/tenofovir DF vs ritonavir-boosted atazanavir plus emtricitabine/tenofovir DF for initial treatment of HIV-1 infection: analysis of week 144 results. *J Acquir Immune Defic Syndr*. 2014;65(3):e121–4. doi:10.1097/QAI.0000000000000089. [PubMed: 24346640]
 92. Pozniak A, Markowitz M, Mills A, Stellbrink HJ, Antela A, Domingo P et al. Switching to coformulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus continuation of non-nucleoside reverse transcriptase inhibitor with emtricitabine and tenofovir in virologically suppressed adults with HIV (STRATEGY-NNRTI): 48 week results of a randomised, open-label,

- phase 3b non-inferiority trial. *The Lancet Infectious Diseases*. 2014;14(7):590–9. doi:10.1016/S1473-3099(14)70796-0. [PubMed: 24908550]
93. Pozniak A, Flamm J, Antinori A, Bloch M, Ward D, Berenguer J et al. Switching to the single-tablet regimen of elvitegravir, cobicistat, emtricitabine, and tenofovir DF from non-nucleoside reverse transcriptase inhibitor plus coformulated emtricitabine and tenofovir DF regimens: Week 96 results of STRATEGY-NNRTI. *HIV Clin Trials*. 2017;18(4):141–8. doi:10.1080/15284336.2017.1338844. [PubMed: 28689453]
94. Arribas JR, DeJesus E, van Lunzen J, Zurawski C, Doroana M, Towner W et al. Simplification to single-tablet regimen of elvitegravir, cobicistat, emtricitabine, tenofovir DF from multi-tablet ritonavir-boosted protease inhibitor plus coformulated emtricitabine and tenofovir DF regimens: week 96 results of STRATEGY-PI. *HIV Clin Trials*. 2017;18(3):118–25. doi:10.1080/15284336.2017.1330440. [PubMed: 28555519]
95. Arribas JR, Pialoux G, Gathe J, Di Perri G, Reynes J, Tebas P et al. Simplification to coformulated elvitegravir, cobicistat, emtricitabine, and tenofovir versus continuation of ritonavir-boosted protease inhibitor with emtricitabine and tenofovir in adults with virologically suppressed HIV (STRATEGY-PI): 48 week results of a randomised, open-label, phase 3b, non-inferiority trial. *Lancet Infect Dis*. 2014;14(7):581–9. doi:10.1016/S1473-3099(14)70782-0. [PubMed: 24908551]
96. Mills A, Crofoot G, Ortiz R, Rashbaum B, Towner W, Ward D et al. Switching from twice-daily raltegravir plus tenofovir disoproxil fumarate/Emtricitabine to once-daily Elvitegravir/Cobicistat/Emtricitabine/ Tenofovir disoproxil fumarate in virologically suppressed, HIV-1-infected subjects: 48 weeks data. *HIV Clinical Trials*. 2014;15(2):51–6. doi:10.1310/hct1502-51. [PubMed: 24710918]
97. Sax PE, Wohl D, Yin MT, Post F, DeJesus E, Saag M et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate, coformulated with elvitegravir, cobicistat, and emtricitabine, for initial treatment of HIV-1 infection: Two randomised, double-blind, phase 3, non-inferiority trials. *Lancet*. 2015;385(9987):2606–15. doi:10.1016/S0140-6736(15)60616-X. [PubMed: 25890673]
98. Wohl D, Oka S, Clumeck N, Clarke A, Brinson C, Stephens J et al. A randomized, double-blind comparison of tenofovir alafenamide versus tenofovir disoproxil fumarate, each coformulated with elvitegravir, cobicistat, and emtricitabine for initial HIV-1 treatment: Week 96 results. *Journal of acquired immune deficiency syndromes*. 2016;72(1):58–64. doi:10.1097/QAI.0000000000000940. [PubMed: 26829661]
99. Arribas JR, Thompson M, Sax PE, Haas B, McDonald C, Wohl DA et al. Brief Report: Randomized, Double-Blind Comparison of Tenofovir Alafenamide (TAF) vs Tenofovir Disoproxil Fumarate (TDF), Each Coformulated With Elvitegravir, Cobicistat, and Emtricitabine (E/C/F) for Initial HIV-1 Treatment: Week 144 Results. *J Acquir Immune Defic Syndr*. 2017;75(2):211–8. doi:10.1097/QAI.0000000000001350. [PubMed: 28282300]
100. Mills A, Arribas JR, Andrade-Villanueva J, DiPerri G, Van Lunzen J, Koenig E et al. Switching from tenofovir disoproxil fumarate to tenofovir alafenamide in antiretroviral regimens for virologically suppressed adults with HIV-1 infection: a randomised, active-controlled, multicentre, open-label, phase 3, non-inferiority study. *Lancet Infect Dis*. 2016;16(1):43–52. doi:10.1016/S1473-3099(15)00348-5. [PubMed: 26538525]
101. DeJesus E, Haas B, Segal-Maurer S, Ramgopal MN, Mills A, Margot N et al. Superior Efficacy and Improved Renal and Bone Safety After Switching from a Tenofovir Disoproxil Fumarate- to a Tenofovir Alafenamide-Based Regimen Through 96 Weeks of Treatment. *AIDS Res Hum Retroviruses*. 2018;34(4):337–42. doi:10.1089/AID.2017.0203. [PubMed: 29368537]
102. Walmsley SL, Antela A, Clumeck N, Duiculescu D, Eberhard A, Gutie rez F et al. Dolutegravir plus Abacavir-Lamivudine for the treatment of HIV-1 infection; Bristol Myers Squibb; Gilead; ViiV. *New England Journal of Medicine*. 2013;369(19):1807–18. doi:10.1056/NEJMoa1215541. [PubMed: 24195548]
103. Walmsley S, Baumgarten A, Berenguer J, Felizarta F, Florence E, Khuong-Josses MA et al. Dolutegravir plus abacavir/lamivudine for the treatment of HIV-1 infection in antiretroviral therapy-naïve patients: Week 96 and week 144 results from the SINGLE randomized clinical trial. *Journal of acquired immune deficiency syndromes*. 2015;70(5):515–9. doi:10.1097/QAI.0000000000000790. [PubMed: 26262777]

104. Clotet B, Feinberg J, Van LJ, Khuong-Josses MA, Antinori A, Dumitru I et al. Once-daily dolutegravir versus darunavir plus ritonavir in antiretroviral-naive adults with HIV-1 infection (FLAMINGO): 48 week results from the randomised open-label phase 3b study. *Lancet*. 2014;383(9936):2222–31. doi:10.1016/S0140-6736(14)60084-2. [PubMed: 24698485]
105. Molina JM, Clotet B, van Lunzen J, Lazzarin A, Cavassini M, Henry K et al. Once-daily dolutegravir versus darunavir plus ritonavir for treatment-naive adults with HIV-1 infection (FLAMINGO): 96 week results from a randomised, open-label, phase 3b study. *Lancet HIV*. 2015;2(4):e127–36. doi:10.1016/S2352-3018(15)00027-2. [PubMed: 26424673]
106. Raffi F, Rachlis A, Stellbrink HJ, Hardy WD, Torti C, Orkin C et al. Once-daily dolutegravir versus raltegravir in antiretroviral-naive adults with HIV-1 infection: 48 week results from the randomised, double-blind, non-inferiority SPRING-2 study. *Lancet*. 2013;381(9868):735–43. doi:10.1016/S0140-6736(12)61853-4. [PubMed: 23306000]
107. Raffi F, Jaeger H, Quiros-Roldan E, Albrecht H, Belonosova E, Gatell JM et al. Once-daily dolutegravir versus twice-daily raltegravir in antiretroviral-naive adults with HIV-1 infection (SPRING-2 study): 96 week results from a randomised, double-blind, non-inferiority trial. *The Lancet Infectious Diseases*. 2013;13(11):927–35. doi:10.1016/S1473-3099(13)70257-3. [PubMed: 24074642]
108. Cahn P, Pozniak AL, Mingrone H, Shuldyakov A, Brites C, Andrade-Villanueva JF et al. Dolutegravir versus raltegravir in antiretroviral-experienced, integrase-inhibitor-naive adults with HIV: Week 48 results from the randomised, double-blind, non-inferiority SAILING study. *Lancet*. 2013;382(9893):700–8. doi:10.1016/S0140-6736(13)61221-0. [PubMed: 23830355]
109. Eron JJ, Clotet B, Durant J, Katlama C, Kumar P, Lazzarin A et al. Safety and efficacy of dolutegravir in treatment-experienced subjects with raltegravir-resistant HIV type 1 infection: 24-week results of the VIKING Study. *J Infect Dis*. 2013;207(5):740–8. doi:10.1093/infdis/jis750. [PubMed: 23225901]
110. Castagna A, Maggiolo F, Penco G, Wright D, Mills A, Grossberg R et al. Dolutegravir in antiretroviral-experienced patients with raltegravir- and/or elvitegravir-resistant HIV-1: 24-week results of the phase III VIKING-3 study. *J Infect Dis*. 2014;210(3):354–62. doi:10.1093/infdis/jiu051. [PubMed: 24446523]
111. Llibre JM, Hung CC, Brinson C, Castelli F, Girard PM, Kahl LP et al. Efficacy, safety, and tolerability of dolutegravir-rilpivirine for the maintenance of virological suppression in adults with HIV-1: phase 3, randomised, non-inferiority SWORD-1 and SWORD-2 studies. *Lancet*. 2018;391(10123):839–49. doi:10.1016/S0140-6736(17)33095-7. [PubMed: 29310899]
112. Aboud M, Orkin C, Podzamczar D, Bogner JR, Baker D, Khuong-Josses MA et al. Efficacy and safety of dolutegravir-rilpivirine for maintenance of virological suppression in adults with HIV-1: 100-week data from the randomised, open-label, phase 3 SWORD-1 and SWORD-2 studies. *Lancet HIV*. 2019;6(9):e576–e87. doi:10.1016/S2352-3018(19)30149-3. [PubMed: 31307948]
113. Cahn P, Madero JS, Arribas JR, Antinori A, Ortiz R, Clarke AE et al. Dolutegravir plus lamivudine versus dolutegravir plus tenofovir disoproxil fumarate and emtricitabine in antiretroviral-naive adults with HIV-1 infection (GEMINI-1 and GEMINI-2): week 48 results from two multicentre, double-blind, randomised, non-inferiority, phase 3 trials. *Lancet*. 2019;393(10167):143–55. doi:10.1016/S0140-6736(18)32462-0. [PubMed: 30420123]
114. Dovato® [package insert]. Research Triangle Park, NC Viiv Healthcare 3 2020.
115. Cahn P, Madero JS, Arribas JR, Antinori A, Ortiz R, Clarke AE et al. Durable Efficacy of Dolutegravir Plus Lamivudine in Antiretroviral Treatment-Naive Adults With HIV-1 Infection: 96-Week Results From the GEMINI-1 and GEMINI-2 Randomized Clinical Trials. *J Acquir Immune Defic Syndr*. 2020;83(3):310–8. doi:10.1097/QAI.0000000000002275. [PubMed: 31834000]
116. Gallant J, Lazzarin A, Mills A, Orkin C, Podzamczar D, Tebas P et al. Bictegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir, abacavir, and lamivudine for initial treatment of HIV-1 infection (GS-US-380–1489): a double-blind, multicentre, phase 3, randomised controlled non-inferiority trial. *Lancet*. 2017;390(10107):2063–72. doi:10.1016/S0140-6736(17)32299-7. [PubMed: 28867497]
117. Wohl DA, Yazdanpanah Y, Baumgarten A, Clarke A, Thompson MA, Brinson C et al. Bictegravir combined with emtricitabine and tenofovir alafenamide versus dolutegravir, abacavir, and

- lamivudine for initial treatment of HIV-1 infection: week 96 results from a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet HIV*. 2019;6(6):e355–e63. doi:10.1016/S2352-3018(19)30077-3. [PubMed: 31068270]
118. Sax PE, Pozniak A, Montes ML, Koenig E, DeJesus E, Stellbrink HJ et al. Coformulated bicitegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir with emtricitabine and tenofovir alafenamide, for initial treatment of HIV-1 infection (GS-US-380–1490): a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet*. 2017;390(10107):2073–82. doi:10.1016/S0140-6736(17)32340-1. [PubMed: 28867499]
119. Stellbrink HJ, Arribas JR, Stephens JL, Albrecht H, Sax PE, Maggiolo F et al. Co-formulated bicitegravir, emtricitabine, and tenofovir alafenamide versus dolutegravir with emtricitabine and tenofovir alafenamide for initial treatment of HIV-1 infection: week 96 results from a randomised, double-blind, multicentre, phase 3, non-inferiority trial. *Lancet HIV*. 2019;6(6):e364–e72. doi:10.1016/S2352-3018(19)30080-3. [PubMed: 31068272]
120. Molina JM, Ward D, Brar I, Mills A, Stellbrink HJ, López-Cortés L et al. Switching to fixed-dose bicitegravir, emtricitabine, and tenofovir alafenamide from dolutegravir plus abacavir and lamivudine in virologically suppressed adults with HIV-1: 48 week results of a randomised, double-blind, multicentre, active-controlled, phase 3, non-inferiority trial. *Lancet HIV*. 2018;5(7):e357–e65. doi:10.1016/S2352-3018(18)30092-4. [PubMed: 29925489]
121. Daar ES, DeJesus E, Ruane P, Crofoot G, Oguchi G, Creticos C et al. Efficacy and safety of switching to fixed-dose bicitegravir, emtricitabine, and tenofovir alafenamide from boosted protease inhibitor-based regimens in virologically suppressed adults with HIV-1: 48 week results of a randomised, open-label, multicentre, phase 3, non-inferiority trial. *Lancet HIV*. 2018;5(7):e347–e56. doi:10.1016/S2352-3018(18)30091-2. [PubMed: 29925490]
122. Sax PE, Rockstroh JK, Luetkemeyer AF, Yazdanpanah Y, Ward D, Trottier B et al. Switching to bicitegravir, emtricitabine, and tenofovir alafenamide in virologically suppressed adults with HIV. *Clin Infect Dis*. 2020. doi:10.1093/cid/ciaa988.
123. Overton ET, Richmond GJ, Rizzardini G, et al. Cabotegravir + rilpivirine every 2 months is noninferior to monthly: ATLAS-2M study Conference on Retroviruses and Opportunistic Infections (CROI). 3 8–11, 2020 Boston Abstract 34. .
124. Landovitz RJ, Li S, Grinsztejn B, Dawood H, Liu AY, Magnus M et al. Safety, tolerability, and pharmacokinetics of long-acting injectable cabotegravir in low-risk HIV-uninfected individuals: HPTN 077, a phase 2a randomized controlled trial. *PLoS Medicine*. 2018;15(11). doi:10.1371/journal.pmed.1002690.
125. Markowitz M, Frank I, Grant RM, Mayer KH, Elion R, Goldstein D et al. Safety and tolerability of long-acting cabotegravir injections in HIV-uninfected men (ECLAIR): a multicentre, double-blind, randomised, placebo-controlled, phase 2a trial. *Lancet HIV*. 2017;4(8):e331–e40. doi:10.1016/S2352-3018(17)30068-1. [PubMed: 28546090]
126. Libre JM, Montoliu A, Miró JM, Domingo P, Riera M, Tiraboschi J et al. Discontinuation of dolutegravir, elvitegravir/cobicistat and raltegravir because of toxicity in a prospective cohort. *HIV Medicine*. 2019;20(3):237–47. doi:10.1111/hiv.12710. [PubMed: 30688007]
127. Lennox JL, DeJesus E, Berger DS, Lazzarin A, Pollard RB, Ramalho Madruga JV et al. Raltegravir versus Efavirenz regimens in treatment-naïve HIV-1-infected patients: 96-week efficacy, durability, subgroup, safety, and metabolic analyses. *J Acquir Immune Defic Syndr*. 2010;55(1):39–48. doi:10.1097/QAI.0b013e3181da1287. [PubMed: 20404738]
128. Tepler H, Brown DD, Leavitt RY, Sklar P, Wan H, Xu X et al. Long-term safety from the raltegravir clinical development program. *Curr HIV Res*. 2011;9(1):40–53. doi:10.2174/157016211794582650. [PubMed: 21198432]
129. Nguyen A, Calmy A, Delhumeau C, Mercier I, Cavassini M, Mello AF et al. A randomized cross-over study to compare raltegravir and efavirenz (SWITCH-ER study). *AIDS*. 2011;25(12):1481–7. doi:10.1097/QAD.0b013e328348dab0. [PubMed: 21593661]
130. Lennox JL, Landovitz RJ, Ribaldo HJ, Ofotokun I, Na LH, Godfrey C et al. Efficacy and tolerability of 3 nonnucleoside reverse transcriptase inhibitor-sparing antiretroviral regimens for treatment-naïve volunteers infected with HIV-1: A Randomized, controlled equivalence trial; Abbott; Bristol Myers Squibb; Gilead; janssen therapeutics; Merck. *Annals of Internal Medicine*. 2014;161(7):461–71. doi:10.7326/M14-1084. [PubMed: 25285539]

131. Martinez E, Larrousse M, Llibre JM, Gutierrez F, Saumoy M, Antela A et al. Substitution of raltegravir for ritonavir-boosted protease inhibitors in HIV-infected patients: The SPIRAL study. *AIDS*. 2010;24(11):1697–707. doi:10.1097/QAD.0b013e32833a608a. [PubMed: 20467288]
132. Curtis L, Nichols G, Stainsby C, Lim J, Aylott A, Wynne B et al. Dolutegravir: clinical and laboratory safety in integrase inhibitor-naïve patients. *HIV Clin Trials*. 2014;15(5):199–208. doi:10.1310/hct1505-199. [PubMed: 25350958]
133. Lee FJ, Amin J, Bloch M, Pett SL, Marriott D, Carr A. Skeletal muscle toxicity associated with raltegravir-based combination antiretroviral therapy in HIV-infected adults. *J Acquir Immune Defic Syndr*. 2013;62(5):525–33. doi:10.1097/QAI.0b013e3182832578. [PubMed: 23274936]
134. Tsai WJ, Lee SS, Tsai HC, Sy CL, Chen JK, Wu KS et al. Rapid onset of rhabdomyolysis after switching to a raltegravir-based antiretroviral regimen. *J Microbiol Immunol Infect*. 2016;49(2):286–8. doi:10.1016/j.jmii.2013.02.008. [PubMed: 23612027]
135. Masia M, Enriquez R, Sirvent A, Gutierrez F. Severe acute renal failure associated with rhabdomyolysis during treatment with raltegravir. A call for caution. *J Infect*. 2010;61(2):189–90. doi:10.1016/j.jinf.2010.04.011. [PubMed: 20447415]
136. Zembower TR, Gerzenshtein L, Coleman K, Palella FJ Jr. Severe rhabdomyolysis associated with raltegravir use. *AIDS*. 2008;22(11):1382–4. doi:10.1097/QAD.0b013e328303be40. [PubMed: 18580624]
137. Croce F, Vitello P, Dalla Pria A, Riva A, Galli M, Antinori S. Severe raltegravir-associated rhabdomyolysis: a case report and review of the literature. *Int J STD AIDS*. 2010;21(11):783–5. doi:10.1258/ijsa.2010.010246. [PubMed: 21187364]
138. Dori L, Buonomini AR, Viscione M, Sarmati L, Andreoni M. A case of rhabdomyolysis associated with raltegravir use. *AIDS*. 2010;24(3):473–5. doi:10.1097/QAD.0b013e328334cc4a. [PubMed: 20087078]
139. Lepik KJ, Yip B, Ulloa AC, Wang L, Toy J, Akagi L et al. Adverse drug reactions to integrase strand transfer inhibitors. *AIDS*. 2018;32(7):903–12. doi:10.1097/QAD.0000000000001781. [PubMed: 29424784]
140. Lepist EI, Zhang X, Hao J, Huang J, Kosaka A, Birkus G et al. Contribution of the organic anion transporter OAT2 to the renal active tubular secretion of creatinine and mechanism for serum creatinine elevations caused by cobicistat. *Kidney Int*. 2014;86(2):350–7. doi:10.1038/ki.2014.66. [PubMed: 24646860]
141. Eron J, Kalayjian R, Wurapa A, Stephens J, McDonald C, Wilkin A et al. Safety and efficacy of elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (E/C/F/TAF) in HIV-infected adults on chronic haemodialysis. *HIV Medicine*. 2018;19:S16.
142. Sax PE, DeJesus E, Crofoot G, Ward D, Benson P, Dretler R et al. Bictegravir versus dolutegravir, each with emtricitabine and tenofovir alafenamide, for initial treatment of HIV-1 infection: a randomised, double-blind, phase 2 trial. *Lancet HIV*. 2017;4(4):e154–e60. doi:10.1016/S2352-3018(17)30016-4. [PubMed: 28219610]
143. Hill AM, Mitchell N, Hughes S, Pozniak AL. Risks of cardiovascular or central nervous system adverse events and immune reconstitution inflammatory syndrome, for dolutegravir versus other antiretrovirals: Meta-analysis of randomized trials. *Current Opinion in HIV and AIDS*. 2018;13(2):102–11. doi:10.1097/COH.0000000000000445. [PubMed: 29278532]
144. Hayes E, Derrick C, Smalls D, Smith H, Kremer N, Weissman S. Adverse events with Biktarvy: post-marketing study ID Week. 10 2–6, 2019 Washington, DC Abstract #2489.
145. Brehm TT, Franz M, Hufner A, Hertling S, Schmiedel S, Degen O et al. Safety and efficacy of elvitegravir, dolutegravir, and raltegravir in a real-world cohort of treatment-naïve and -experienced patients. *Medicine (Baltimore)*. 2019;98(32):e16721. doi:10.1097/MD.00000000000016721. [PubMed: 31393378]
146. Cuzin L, Pugliese P, Katlama C, Bani-Sadr F, Ferry T, Rey D et al. Integrase strand transfer inhibitors and neuropsychiatric adverse events in a large prospective cohort. *Journal of Antimicrobial Chemotherapy*. 2019;74(3):754–60. doi:10.1093/jac/dky497. [PubMed: 30534993]
147. Hoffmann C, Llibre JM. Neuropsychiatric adverse events with dolutegravir and other integrase strand transfer inhibitors. *AIDS Reviews*. 2019;21(1):4–10. doi:10.24875/AIDSRev.19000023. [PubMed: 30899113]

148. Hoffmann C, Welz T, Sabranski M, Kolb M, Wolf E, Stellbrink HJ et al. Higher rates of neuropsychiatric adverse events leading to dolutegravir discontinuation in women and older patients; Gilead(United States); Merck Sharp and Dohme(United States); ViiV. *HIV Medicine*. 2017;18(1):56–63. doi:10.1111/hiv.12468. [PubMed: 27860104]
149. Cid-Silva P, Llibre JM, Fernández-Bargiela N, Margusino-Framiñán L, Balboa-Barreiro V, Pernas-Souto B et al. Clinical Experience with the Integrase Inhibitors Dolutegravir and Elvitegravir in HIV-infected Patients: Efficacy, Safety and Tolerance. *Basic and Clinical Pharmacology and Toxicology*. 2017;121(5):442–6. doi:10.1111/bcpt.12828. [PubMed: 28627771]
150. Antiretroviral Pregnancy Registry Steering Committee. Antiretroviral Pregnancy Registry international interim report for 1 January 1989–31 January 2019. Wilmington, NC: Registry Coordinating Center 2019 Available at: apregistry.com/forms/exec-summary.pdf.
151. Centers for Disease C, Prevention. Update on overall prevalence of major birth defects--Atlanta, Georgia, 1978–2005. *MMWR Morb Mortal Wkly Rep* 2008;57(1):1–5.
152. Sibiude J, Warszawski J, Blanchard S, et al. Evaluation of the risk of birth defects among children exposed to raltegravir in utero in the ANRS-French perinatal cohort EPF. Presented at: International AIDS Society; 2017; Paris, France.
153. Zash R, Makhema J, Shapiro RL. Neural-tube defects with dolutegravir treatment from the time of conception. *New England Journal of Medicine*. 2018;379(10):979–81. doi:10.1056/NEJMc1807653. [PubMed: 30037297]
154. Zash R, Holmes L, Diseko M, Jacobson DL, Brummel S, Mayondi G et al. Neural-Tube Defects and Antiretroviral Treatment Regimens in Botswana. *N Engl J Med*. 2019;381(9):827–40. doi:10.1056/NEJMoa1905230. [PubMed: 31329379]
155. Bakal DR, Coelho LE, Luz PM, Clark JL, De BR, Cardoso SW et al. Obesity following ART initiation is common and influenced by both traditional and HIV-/ART-specific risk factors. *Journal of Antimicrobial Chemotherapy*. 2018;73(8):2177–85. doi:10.1093/jac/dky145. [PubMed: 29722811]
156. Menard A, Meddeb L, Tissot-Dupont H, Ravaux I, Dhiver C, Mokhtari S et al. Dolutegravir and weight gain: An unexpected bothering side effect? *AIDS*. 2017;31(10):1499–500. doi:10.1097/QAD.0000000000001495. [PubMed: 28574967]
157. Sax PE, Erlandson KM, Lake JE, McComsey GA, Orkin C, Esser S et al. Weight Gain Following Initiation of Antiretroviral Therapy: Risk Factors in Randomized Comparative Clinical Trials. *Clin Infect Dis*. 2019. doi:10.1093/cid/ciz999.
158. Kouanfack C, Mpoudi-Etame M, Bassega PO, Eymard-Duvernay S, Leroy S, Boyer S et al. Dolutegravir-based or low-dose efavirenz-based regimen for the treatment of HIV-1. *New England Journal of Medicine*. 2019;381(9):816–26. doi:10.1056/NEJMoa1904340. [PubMed: 31339676]
159. Venter WDF, Moorhouse M, Sokhela S, Fairlie L, Mashabane N, Masenya M et al. Dolutegravir plus two different prodrugs of tenofovir to treat HIV; Gilead; Macleods; Mylan; ViiV. *New England Journal of Medicine*. 2019;381(9):803–15. doi:10.1056/NEJMoa1902824. [PubMed: 31339677]
160. Bourgi K, Rebeiro PF, Turner M, Castilho JL, Hulgán T, Raffanti SP et al. Greater Weight Gain in Treatment Naive Persons Starting Dolutegravir-Based Antiretroviral Therapy. *Clin Infect Dis*. 2019. doi:10.1093/cid/ciz407.
161. Norwood J, Turner M, Bofill C, Rebeiro P, Shepherd B, Bebawy S et al. Brief Report: Weight Gain in Persons With HIV Switched From Efavirenz-Based to Integrase Strand Transfer Inhibitor-Based Regimens. *J Acquir Immune Defic Syndr*. 2017;76(5):527–31. doi:10.1097/QAL.0000000000001525. [PubMed: 28825943]
162. Landovitz RJ, Zangeneh SZ, Chau G, Grinsztejn B, Eron JJ, Dawood H et al. Cabotegravir Is Not Associated With Weight Gain in Human Immunodeficiency Virus-uninfected Individuals in HPTN 077. *Clin Infect Dis*. 2020;70(2):319–22. doi:10.1093/cid/ciz439. [PubMed: 31125395]
163. Landovitz RJ, Donnell D, Clement M, Hanscom B, Cottle L, Coelho L et al. HPTN083 interim results: Pre-exposure prophylaxis (PrEP) containing long-acting injectable cabotegravir (CAB-LA) is safe and highly effective for cisgender men and transgender women who have sex with

- men (MSM,TGW). 23rd International HIV Conference (AIDS 2020: Virtual), abstract OAXLB0101, 2020.
164. Yuh B, Tate J, Butt AA, Crothers K, Freiberg M, Leaf D et al. Weight change after antiretroviral therapy and mortality. *Clin Infect Dis*. 2015;60(12):1852–9. doi:10.1093/cid/civ192. [PubMed: 25761868]
165. Achhra AC, Mocroft A, Reiss P, Sabin C, Ryom L, de Wit S et al. Short-term weight gain after antiretroviral therapy initiation and subsequent risk of cardiovascular disease and diabetes: the D:A:D study. *HIV Med*. 2016;17(4):255–68. doi:10.1111/hiv.12294. [PubMed: 26216031]
166. Herrin M, Tate JP, Akgun KM, Butt AA, Crothers K, Freiberg MS et al. Weight Gain and Incident Diabetes Among HIV-Infected Veterans Initiating Antiretroviral Therapy Compared With Uninfected Individuals. *J Acquir Immune Defic Syndr*. 2016;73(2):228–36. doi:10.1097/QAI.0000000000001071. [PubMed: 27171741]
167. Hill A, McCann KM, Pilkington V, et al. Risks of metabolic syndrome, diabetes, and cardiovascular disease in ADVANCE Trial Conference on Retroviruses and Opportunistic Infections (CROI). 3 8–11, 2020 Boston Abstract 81.
168. Fong PS, Flynn DM, Evans CD, Korthuis PT. Integrase strand transfer inhibitor-associated diabetes mellitus: A case report. *Int J STD AIDS*. 2017;28(6):626–8. doi:10.1177/0956462416675107. [PubMed: 27733708]
169. McLaughlin M, Walsh S, Galvin S. Dolutegravir-induced hyperglycaemia in a patient living with HIV. *J Antimicrob Chemother*. 2018;73(1):258–60. doi:10.1093/jac/dkx365. [PubMed: 29077869]
170. Lamorde M, Atwiine M, Owarwo NC, Ddungu A, Laker EO, Mubiru F et al. Dolutegravir-associated hyperglycaemia in patients with HIV. *Lancet HIV*. 2020. doi:10.1016/S2352-3018(20)30042-4.
171. Bhagwat P, Ofotokun I, McComsey GA, Brown TT, Moser C, Sugar CA et al. Changes in Waist Circumference in HIV-Infected Individuals Initiating a Raltegravir or Protease Inhibitor Regimen: Effects of Sex and Race. *Open Forum Infect Dis*. 2018;5(11):ofy201. doi:10.1093/ofid/ofy201. [PubMed: 30465010]
172. Bernardino JI, Mocroft A, Wallet C, de Wit S, Katlama C, Reiss P et al. Body composition and adipokines changes after initial treatment with darunavir-ritonavir plus either raltegravir or tenofovir disoproxil fumarate-emtricitabine: A substudy of the NEAT001/ANRS143 randomised trial. *PLoS One*. 2019;14(1):e0209911. doi:10.1371/journal.pone.0209911. [PubMed: 30689664]
173. Triumeq® [package insert]. Research Triangle Park, NC Viiv Healthcare 3 2020.
174. New York Department of Health AIDS Insite. Clinical Guidelines Program. When to initiate ART, with protocol for rapid initiation. Available at: <https://www.hivguidelines.org/antiretroviral-therapy/when-to-start-plus-rapid-start/> Accessed April 5, 2020.
175. Juluca® [package insert]. Research Triangle Park, NC Viiv Healthcare 10 2019.

Key points

- Bictegravir and dolutegravir are both co-formulated as a single tablet regimen, and the choice of agent may be driven based on the preferred nucleoside reverse transcriptase inhibitor.
- Elvitegravir/cobicistat is available as a single tablet regimen, though it has a higher propensity for drug-drug interactions and must be administered with food, increasing complexity compared with other integrase strand transfer inhibitors (INSTI).
- Raltegravir has the most safety data associated with its use; however, it is the only INSTI not available as single tablet antiretroviral regimen, increasing the pill burden associated with a raltegravir-containing regimen.
- Cabotegravir is part of the first long-acting, injectable antiretroviral regimen, recently approved in Canada for treatment of individuals currently virologically suppressed on another antiretroviral regimen.

Table 1. Available formulations, dosing, administration, and indications for products containing integrase strand transfer inhibitors

INSTI component	Formulation components	Brand Name	Dosing in patients without resistance ^a	Dosing in patients with resistance ^b	Administration instructions	Pediatric Use	Use in pregnancy [43]	Use in mild or moderate hepatic impairment ^c	Use in renal impairment
Bictegravir [16]	FDC tablet containing: Bictegravir 50 mg / Tenofovir AF 25 mg / Emtricitabine 200 mg	Biktarvy®	1 tablet daily	-	May be taken with or without food	25 kg: Standard dose	Not recommended: Insufficient data	Standard dosing	Standard dosing for CrCl ≥ 30 mL/min
	Cabotegravir 30 mg tablet	Vocabria®	1 tablet daily ^d	-	May be taken with or without food	No Data in individuals < 18 years	Not recommended: Insufficient data	Standard dosing	Standard dosing 15 mL/min
Dolutegravir [173]	Co-packaged loading dose injection: Cabotegravir 600 mg / Rilpivirine 900 mg	Cabenuva®	Loading dose: 600mg/3mL IM x 1 dose ^e (Plus RPV: 900mg/3mL IM X 1 dose)	-	IM gluteus medius administration ^f	No Data in individuals < 18 years	Not recommended: Insufficient data and two drug regimens are not recommended	Standard dosing	Standard dosing 30 mL/min
	Co-packaged maintenance dose injection: Cabotegravir 400 mg / Rilpivirine 600 mg		Maintenance dose: 400mg/2mL IM every 4 weeks ^e (Plus RPV: 600mg/2mL every 4 weeks)	-	May be taken with or without food	Weight based dosing for pediatrics 3 kg - 20 kg: Standard dose	Standard dosing; take with food	Standard dosing	Standard dosing ^h
	Dolutegravir 50 mg, 25 mg, 10 mg tablets, 5 mg dispersible tablet for oral suspension ^g		Tivicay®	50 mg tablet daily	1 tablet twice daily	May be taken with or without food	40 kg: Standard dose	Standard dosing; take with food	Standard dosing
Dolutegravir 50 mg / Abacavir 300 mg / Lamivudine 300 mg	FDC tablet containing: Dolutegravir 50 mg / Abacavir 300 mg / Lamivudine 300 mg	Triumeq®	1 tablet daily	-	May be taken with or without food	-	Standard dosing; take with food	Standard dosing	Standard dosing > 50 mL/min
	FDC tablet containing: Dolutegravir 50 mg / Lamivudine 300 mg	Dovato®	1 tablet daily	-	May be taken with or without food	-	Two drug regimens are not recommended	Standard dosing	Standard dosing > 50 mL/min
	FDC tablet containing: Dolutegravir 50 mg / Rilpivirine 25 mg	Juluca®	1 tablet daily ⁱ	-	Take with food	-	Two drug regimens are not recommended	Standard dosing	Standard dosing ^{h,j}

INSTI component	Formulation components	Brand Name	Dosing in patients without resistance ^a	Dosing in patients with resistance ^b	Administration instructions	Pediatric Use	Use in pregnancy [43]	Use in mild or moderate hepatic impairment ^c	Use in renal impairment
Elvitegravir [13, 18]	FDC tablet containing: Elvitegravir 150 mg / Cobicistat 150 mg / Emtricitabine 200 mg / Tenofovir DF 300 mg	Stribild®	1 tablet daily	-	Take with food	35 kg: Standard dose	Not recommended: suboptimal exposure observed in PK studies	Standard dosing	Initiation of therapy. Standard dosing 70 mL/min. Therapy should be switched in patients with a CrCl <50 mL/min
Raltegravir [12]	FDC tablet containing: Elvitegravir 150 mg / Cobicistat 150 mg / Emtricitabine 200 mg / Tenofovir AF 10 mg	Genvoya®	1 tablet daily	-	Take with food	25 kg: Standard dose	Not recommended: suboptimal exposure observed in PK studies	Standard dosing	Standard dosing 30 mL/min or in patients with ESRD receiving hemodialysis
	Raltegravir 400 mg, 100 mg scored tablet, 100 mg packet for oral solution	Isentress®	400 mg tablet twice daily	1 tablet twice daily	May be taken with or without food	Weight based dosing for pediatrics 3 kg: 25 kg: Standard dose	Standard dosing	Standard dosing	Standard dosing
	Raltegravir HD 600 mg	Isentress HD®	2 tablets daily	-	May be taken with or without food	40 kg: Standard dose	Not recommended: potential for suboptimal exposure	Standard dosing	Standard dosing

Abbreviations: AF, alafenamide; CAB, cabotegravir; CrCl, creatinine clearance; DF, disoproxil fumarate; ESRD, end stage renal disease; FDC, fixed dose combination; HD, high dose; IM, intramuscular; INSTI, Integrase strand transfer inhibitors; PK, pharmacokinetic; RPV, rilpivirine.

^a Package inserts indicate this dose for adults and adolescents with HIV who are antiretroviral-naïve or to simplify a regimen in individuals currently virologically suppressed on another antiretroviral regimen who do not have resistance to the new regimen components.

^b Indicated as part of a new regimen in patients switching therapy due to virologic failure. Refer to the package insert regarding specific drug resistance mutations relevant to each INSTI formulation.

^c All product labels state that there are insufficient evidence to support the use of INSTIs in patient with severe hepatic impairment, except raltegravir which has data to support its use at standard dosing in patients with severe hepatic dysfunction.

^d Oral cabotegravir may be used as part of an oral lead-in therapy for 28 days prior to initiation of the long-acting, injectable therapy, or as a bridging therapy to accommodate up to two missed doses of the injection. The final oral dose is taken on the same day as the cabotegravir injection is initiated.

^e Administered on the same day as the final oral dose.

^f Administered separately from the rilpivirine IM companion injection.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

The dispersible tablet for suspension is not interchangeable with other oral products on a mg per mg basis. Refer to product labeling for dosing of each product. The dispersible tablet is approved for use in children at least 4 weeks of age and weighing at least 3 kg.

Caution in patients with severe renal insufficiency plus INSTI resistance due to lower concentrations observed in a healthy volunteer study of severe renal insufficiency.

Only indicated for treatment simplification in individuals virologically suppressed on another antiretroviral regimen.

Product labeling recommends increased monitoring for adverse effects in patients with severe renal impairment.

Table 2. Comparative rates of virologic suppression in selected clinical trials evaluating INSTI efficacy in adults living with HIV.

INSTI	Study	Participant characteristics	Regimens (n for primary endpoint)	Primary endpoint / Difference (95% CI)	Final published data / Difference (95% CI)	Summary
Raltegravir	STARTMRK [77, 78]	ART-naïve	Raltegravir 400mg bid + TDF/FTC (n=281) vs. Efavirenz/TDF/FTC FDC daily (n=282)	Week 48: 4.2% (-1.9%, 10.3%)	Week 240: 9.5% (1.7%, 17.3)	Non-inferior through week 192; raltegravir was superior to efavirenz-based therapy at week 240
	QDMRK [83]	ART-naïve	Raltegravir 800mg daily + TDF/FTC daily (n=382) vs. Raltegravir 400mg bid + TDF/FTC daily (n=389)	Week 48: -5.7% (-10.7%, -0.83%)	-	Raltegravir once daily failed to achieve non-inferiority with raltegravir twice daily
	ONCEMRK [84, 85]	ART-naïve	Raltegravir HD 1200mg daily + TDF/FTC daily (n=531) vs. Raltegravir 400mg bid + TDF/FTC daily (n=266)	Week 48: 0.5% (-4.2%, 5.2%)	Week 96: 1.4% (-4.4%, 7.3%)	Non-inferior through week 96
	SWITCHMRK [79]	Switch study; ART-experienced, virologically suppressed	Raltegravir 400mg bid + 2 NRTIs (n=350) vs. Lopinavir/ritonavir-containing ART (n=352)	Week 24: -6.2% (-11.2%, -1.3%)	-	Terminated at week 24 due to inferior outcomes in raltegravir 400mg bid
	BENCHMARK [8082]	ART-experienced, INSTI-naïve, with triple drug-class resistance	Raltegravir 400mg bid + OBR (n=462) vs. Placebo + OBR (n=237)	Week 16; proportion with HIV-RNA <400 copies/mL: Raltegravir: 77.5% Placebo: 41.9% P<0.001	Week 156; proportion with HIV-RNA <50 copies/mL: Raltegravir: 51% Placebo: 22% P<0.0001	Raltegravir was superior to placebo for highly treatment-experience participants through week 156
	GS102 [86, 87]	ART-naïve	Elvitegravir/cobicistat/TDF/FTC FDC daily (n=348) vs. Efavirenz/TDF/FTC FDC daily (n=352)	Week 48: 3.6% (-1.6%, 8.8%)	Week 144: 4.9% (-1.3%, 11.1%)	Non-inferior through week 144
Elvitegravir	GS103 [89-91]	ART-naïve	Elvitegravir/cobicistat/TDF/FTC FDC daily (n=353) vs. Atazanavir/ritonavir daily + TDF/FTC daily (n=355)	Week 48: 3% (-1.9%, 7.8%)	Week 144: 3.1% (-3.2%, 9.4%)	Non-inferior through week 144
	GS104 and GS111 [97-99]	ART-naïve	Elvitegravir/cobicistat/TAF/FTC FDC daily (n=866) vs. Elvitegravir/cobicistat/TDF/FTC FDC daily (n=867)	Week 48: 2.0% (-0.7%, 4.7%)	Week 144: 4.2% (0.6%, 7.8%)	Non-inferior through week 96; TAF superior to TDF at week 144
	STRATEGY- NNRTI [92, 93]	Switch study; ART-experienced, virologically suppressed	Elvitegravir/cobicistat/TDF/FTC FDC daily (n=291) Vs. TDF/FTC + NNRTI containing ART (n=143)	Week 48: 5.3% (-0.5%, 12.0%)	Week 96: 6.1% (-1.3%, 14.2%)	Non-inferior through week 96
	STRATEGY-PI [94, 95]	Switch study; ART-experienced, virologically suppressed	Elvitegravir/cobicistat/TDF/FTC FDC daily (n=293) vs. TDF/FTC + PI containing ART (n=140)	Week 48: 6.7% (0.4%, 13.7%)	Week 96: 17% (8.7%, 26.0%)	Non-inferior at week 48, but the elvitegravir FDC was superior to continuing PI-containing ART at week 96
	GS123 [96]	Switch study; ART-experienced, virologically suppressed	Elvitegravir/cobicistat/TDF/FTC FDC daily (n=48) Single arm study: switched from raltegravir 400mg bid containing ART	Week 48: 48 (100%) remained virologically suppressed	-	All participants remained virologically suppressed at week 48 after switching to the

INSTI	Study	Participant characteristics	Regimens (n for primary endpoint)	Primary endpoint / Difference (95% CI)	Final published data / Difference (95% CI)	Summary
	GS109 [100, 101]	Switch study; ART-experienced, virologically suppressed	Elvitegravir/cobicistat/TAF/FTC FDC daily (n=959) vs. TDF containing regimen + efavirenz, boosted atazanavir, or Elvitegravir/cobicistat FDC (n=477)	Week 48: 4.1% (1.6, 6.7%)	Week 96: 3.7% (0.4%, 7.0%)	once daily, single tablet regimen Non-inferior through week 48; statistically favors elvitegravir FDC at week 96
Dolutegravir	SINGLE [102, 103]	ART-naive	Dolutegravir/ABC/3TC FDC daily (n=414) vs. Efavirenz/TDF/FTC FDC daily (n=419)	Week 48: 7.4% (2.5%, 12.3%)	Week 144: 8.3% (2.0%, 14.6%)	Dolutegravir FDC superior to efavirenz FDC through week 144
	SPRING-2 [106, 107]	ART-naive	Dolutegravir 50 mg daily + NRTIs (n=403) vs. Raltegravir 400mg bid + NRTIs (n=405)	Week 48: 2.5% (-2.2%, 7.1%),	Week 96: 4.5% (-1.1%, 10.0%)	Non-inferior through week 96
	FLAMINGO [104, 105]	ART-naive	Dolutegravir 50 mg daily + NRTIs (n=243) vs. Darunavir/ritonavir daily + NRTIs (n=242)	Week 48: 7.1% (0.9%, 13.2%)	Week 96: 12.4% (4.7%, 20.2%)	Dolutegravir was superior to darunavir/ritonavir containing regimens through 96 weeks.
	SAILING [108]	ART-experienced, INSTI-naive, 2-drug class resistance	Dolutegravir 50 mg daily + OBR (n=354) vs. Raltegravir 400 mg bid + OBR (n=361)	Week 48: 7.4% (0.7%, 14.2%)	-	Dolutegravir-containing ART was superior at week 48
	VIKING-3 [110]	ART-experienced, failing a raltegravir or elvitegravir-based regimen with resistance	Dolutegravir 50mg bid + OBR (n=183) Single arm study; switched from a failing regimen containing raltegravir or elvitegravir to dolutegravir containing ART	Week 24: 69% (62%, 76%)	-	Dolutegravir twice daily was effective in OBR for participants with first generation INSTI resistance.
Dolutegravir/ Rilpivirine FDC	SWORD-1 and SWORD-2 [111]	Switch study; ART-experienced, virologically suppressed	Dolutegravir/rilpivirine daily (n=513) vs. NRTIs plus either a PI, NNRTI, or INSTI containing ART (n=511)	Week 48: -0.2% (-3.0, 2.5%)	-	Non-inferior through week 48
Dolutegravir/ Lamivudine FDC	GEMINI-1 and GEMINI-2 [113, 115]	ART-naive	Dolutegravir 50mg daily + 3TC daily (n=716) vs. Dolutegravir 50mg daily + TDF/FTC daily (n=717)	Week 48: -1.7% (-4.4, 1.1%)	Week 96: -3.4% (-6.7, 0.0%)	Non-inferior through week 96
Bictegravir	GS1489 [116, 117]	ART-naive	Bictegravir/TAF/FTC FDC daily (n=314) vs. Dolutegravir/ABC/3TC FDC daily (n=315)	Week 48: -0.6% (-4.8%, 3.6%)	Week 96: -1.9% (-6.9%, 3.1%)	Non-inferior through week 96
	GS1490 [118, 119]	ART-naive	Bictegravir/TAF/FTC FDC daily (n=320) vs. Dolutegravir 50mg daily + TAF/FTC daily (n=325)	Week 48: -3.5% (-7.9%, 1.0%)	Week 96: -2.3%, (-7.9%, 3.2%)	Non-inferior through week 96
	GS1844 [120]	Switch study; ART-experienced, virologically suppressed	Bictegravir/TAF/FTC FDC daily (n=282) vs. Dolutegravir/ABC/3TC FDC daily (n=281)	Week 48: 0.7% (-1.0%, 2.8%)	-	Non-inferior through week 48
	GS1878 [121]	Switch study; ART-experienced, virologically suppressed	Bictegravir/TAF/FTC FDC daily (n=290) vs. Atazanavir/ritonavir, or darunavir/ritonavir containing ART (n=287)	Week 48: 0.0% (-2.5%, 2.5%)	-	Non-inferior through week 48

INSTI	Study	Participant characteristics	Regimens (n for primary endpoint)	Primary endpoint ¹ Difference (95% CI)	Final published data ² Difference (95% CI)	Summary
	GS4030 [122]	Switch study; ART-experienced, resistance other than INSTI allowed, virologically suppressed	Bictegravir/TAF/FTC vs. Dolutegravir 50mg daily + TAF/FTC daily (n=281)	Week 48: -0.7% (-2.8%, 1.0%)	-	Non-inferior through week 48
Cabotegravir	FLAIR [24]	ART-naive	Cabotegravir/riplivirine long-acting injectable every 4 weeks (n=283) vs. Dolutegravir/ABC/3TC FDC daily (n=283)	Week 48: -0.4% (-2.8 to 2.1%)	-	Non-inferior through week 48
	ATLAS [25]	Switch study; ART-experienced, virologically suppressed	Cabotegravir/riplivirine long-acting injectable every 4 weeks (n=308) vs. NNRTI, PI or INSTI containing ART (n=308)	Week 48: 0.6% (-1.2, 2.5%)	-	Non-inferior through week 48

Abbreviations: 3TC, lamivudine; ABC, abacavir; ART, antiretroviral therapy; bid, twice daily; CI, confidence interval; FDC, fixed dose combination; FTC, emtricitabine; INSTI, integrase strand transfer inhibitor; NNRTI, non-nucleoside reverse transcriptase inhibitor; NRTI, nucleoside reverse transcriptase inhibitor; OBR, optimized background regimen; PI, protease inhibitor; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate

¹Endpoint is HIV-RNA < 50 copies/mL unless indicated.

²HIV-RNA < 40 copies/mL.

Table 3. Drug-related adverse events of integrase strand transfer inhibitors reported in clinical trials^a

Adverse Event	Raltegravir [77, 81, 84]	Elvitegravir/cobicistat [86, 89, 96, 98]	Dolutegravir [102, 106, 111, 113, 116]	Bictegravir [116, 118]	Cabotegravir [24, 25, 123-125]
Fatigue	0-2%	4-5%	<1-3%	2-3%	2-4%
Gastrointestinal					
Nausea	0-3%	11-16%	<1-17%	3-5%	5-6%
Diarrhea	-	10-16%	<1-4%	3-6%	7-11%
Neuropsychiatric					
Headache	<1-4%	5-7%	<1-5%	4-5%	11-14%
Abnormal dreams	-	9%	<1-3%	<1-3%	<1-2%
Insomnia	<1-4%	3%	<1-3%	2%	2-5%
Somnolence	0-2%	3%	-	-	<1%
Dizziness	-	1%	<1-3%	2%	3-5%
Depression	-	<1%	<1-1%	<1%	<1-2%
Suicidality	-	<1%	<1%	<1%	<1%
Rash	-	4%	0-<1%	-	2-4%
Pyrexia	-	-	-	-	7-8%
Elevated Serum Creatinine					
Grade 3 (3x ULN)	0%	-	-	-	<1%
Mean change from baseline (mg/dL)	-	+<0.1 to 0.1	+ 0.09 to 0.15	+ 0.1 ^b	-
Hyperglycemia					
(Grade 3; serum glucose ≥ 251 mg/dL)	0-3%	-	<1-2%	-	<1%
Transaminases (Grade 3; 5.1x ULN)					
ALT	<1-2%	2-3%	<1-4%	1-2%	<1%
AST	<1-5%	3-4%	<1-5%	1-2%	<1-1%
Creatinine Kinase (Grade 3; 10x ULN)	3-4%	8-11%	1-7%	4%	8-9%
Lipid Markers (mean change from baseline, mg/dL)					
LDL-Cholesterol	+10	+8 to 20	+4.0 to 16.0	+7 to 9	-
HDL-Cholesterol	+5	+3 to 7	+2 to 5.4	+4 to 5	-
Total Cholesterol	+16	+14 to 31	+8.1 to 24.0	+12 to 13	-

Adverse Event	Raltegravir [77, 81, 84]	Elvitegravir/cobicistat [86, 89, 96, 98]	Dolutegravir [102, 106, 111, 113, 116]	Bictegravir [116, 118]	Cabotegravir [24, 25, 123-125]
Triglycerides	+2	+12 to 29	+3.0 to 13.6	+3 to 9	-
Total Cholesterol:HDL Ratio	-	+0.1 to 0.2	-	-	-
Injection Site Reaction	-	-	-	-	11-71%

Abbreviations: ULN, upper limits of normal; ALT, alanine aminotransferase; AST, aspartate aminotransferase; LDL, low-density lipoprotein; HDL, high-density lipoprotein.

^aReported as range of values observed in clinical trials unless otherwise specified

^bReported as median (interquartile range)