



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



A clinical trial of IL-15 and IL-21 combination therapy for COVID-19 is warranted

Stephen W. Wilz

Boston VA Healthcare System, Pathology and Laboratory Medicine, 1400 VFW Parkway, West Roxbury, MA, 02132, United States

ARTICLE INFO

Keywords:

COVID-19
SARS-CoV-2
Immune cells
Interleukin-15
Interleukin-21

ABSTRACT

Previous studies of SARS-CoV-2 viral infection suggest that both the humoral and cytotoxic arms of the immune system are weak in patients with severe COVID-19 disease when compared to mild disease. A cytokine storm is also induced in severe disease. IL-15 has been shown to support the cytotoxic arm of the immune response. IL-21 has been shown to support both the cytotoxic and humoral arms of the immune response. In addition, in some settings, IL-21 has been shown to actually decrease IL-6 and TNF-alpha production, reducing the inflammatory proteins involved in the cytokine storm. Furthermore, in other settings, the combination of IL-15 and IL-21 has been shown to be more effective than either interleukin alone in promoting an effective immune response. Therefore, a clinical trial that examines the use of the combination of IL-15 and IL-21 for COVID-19 patients is warranted.

In “*IL-15 immunotherapy is a viable strategy for COVID-19*”, Kandikattu *et al.* urged initiation of a clinical trial that examines the use of IL-15 for SARS-CoV-2 patients that present with diminished NK cell or CD8 + T cell counts [1]. While this is certainly a worthwhile endeavor, the combination of IL-15 and IL-21 should be given in one arm of the study. There have been published reports that examined the differences between the immune response that occurred in patients with severe symptoms versus mild symptoms of the viral infection. The research has shown that hallmarks of severe SARS-CoV-2 disease include a cytokine storm, involving elevation of IL-6 and TNF-alpha, elevation of IL-10, a decreased absolute B cell count, and decreased NK cell and/or CD8 + T cell counts [2]. In addition to the decreased B cell count, there is evidence that patients with severe SARS-CoV-2 disease are unable to mount a strong IgG response against the spike proteins on the virus [3]. These findings suggest that both the cytotoxic (NK-cell and CD8 + T cell) and humoral (B cell) arms of the immune system are weak in patients with severe disease. IL-15 has been shown to stimulate the cytotoxic CD8 + T-cell response and the NK cell response, as well as promote memory-phenotype CD8 + T cells [4–7]. Also, in contrast to IL-2, which is already in clinical trials, IL-15 does not cause capillary leak syndrome [7]. IL-21 supports both the cytotoxic and humoral arms of the immune response. It has been shown that IL-21 promotes B cell and plasma cell development, upregulating IgG1 antibody formation. IL-21 also promotes CD8 effector cell proliferation and costimulation of T and NK cell proliferation and function [9–11]. In addition, IL-21 has been shown to

actually decrease IL-6 and TNF-alpha production [12]. Furthermore, in other settings, the combination of IL-15 and IL-21 has been shown to be more effective than IL-15 alone in promoting an effective immune response [13,14]. Also, while concerning side effects have been identified in some of the studies of these interleukins [8], they occurred in the relatively long course of treatment required for chronic conditions. These side effects may not be as much of a concern in the short time frame required to treat an acute viral infection.

So, given all of these findings, I urge the combination of IL-21 and IL-15 be given to SARS-CoV-2 patients, particularly to patients that present with diminished B cell and/or CD8 + T cell counts. I would also urge that the trial be conducted on patients early in the course of the disease. If severe disease can be avoided by directing the immune response to arms of the immune system that are known to be most effective in treating the disease, the devastating effects of the infection may be avoided.

Declaration of Competing Interest

The authors report no declarations of interest.

References

- [1] H.K. Kandikattu, S.U. Venkateshaiah, S. Kumar, A. Mishra, IL-15 immunotherapy is a viable strategy for COVID-19, *Cytokine Growth Factor Rev.* (Jun) (2020), <https://doi.org/10.1016/j.cytogfr.2020.06.008>.

E-mail address: stephen.wilz@va.gov.

<https://doi.org/10.1016/j.cytogfr.2020.10.005>

Received 7 October 2020; Accepted 20 October 2020

Available online 23 October 2020

1359-6101/© 2020 The Author.

Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

- [2] Y. Zhang, X. Wang, X. Li, et al., Potential contribution of increased soluble IL-2R to lymphopenia in COVID-19 patients, *Cell. Mol. Immunol.* 17 (2020) 878–880, <https://doi.org/10.1038/s41423-020-0484-x>.
- [3] Baoqing Sun, Ying Feng, Xiaoneng Mo, Peiyan Zheng, Qian Wang, Pingchao Li, Ping Peng, Xiaoqing Liu, Zhilong Chen, Huimin Huang, Fan Zhang, Wenting Luo, Xuefeng Niu, Peiyu Hu, Longyu Wang, Hui Peng, Zhifeng Huang, Liqiang Feng, Feng Li, Fuchun Zhang, Fang Li, Nanshan Zhong, Ling Chen, Kinetics of SARS-CoV-2 specific IgM and IgG responses in COVID-19 patients, *Emerg. Microbes Infect.* 9 (1) (2020) 940–1948, <https://doi.org/10.1080/22221751.2020.1762515>.
- [4] T.O. Robinson, K.S. Schluns, The potential and promise of IL-15 in immunoncogenic therapies, *Immunol. Lett.* 190 (2017) 159–168, <https://doi.org/10.1016/j.imlet.2017.08.010>.
- [5] Karin M. Knudson, James W. Hodge, Jeffrey Schlom, R. Sofia, D. Gameiro, Rationale for IL-15 superagonists in cancer immunotherapy, *Expert Opin. Biol. Ther.* (2020), <https://doi.org/10.1080/14712598.2020.1738379>.
- [6] R. Romee, S. Cooley, M.M. Berrien-Elliott, et al., First-in-human phase 1 clinical study of the IL-15 superagonist complex ALT-803 to treat relapse after transplantation, *Blood* 131 (23) (2018) 2515–2527, <https://doi.org/10.1182/blood-2017-12-823757>.
- [7] X. Zhang, S. Sun, I. Hwang, D.F. Tough, J. Sprent, Potent and selective stimulation of memory-phenotype CD8+ T cells in vivo by IL-15, *Immunity* 8 (5) (1998) 591–599, [https://doi.org/10.1016/s1074-7613\[00\]80564-6](https://doi.org/10.1016/s1074-7613[00]80564-6).
- [8] Kevin Conlon, Milos Miljkovic, Thomas Waldmann, Cytokines in the treatment of cancer, *J. Interferon Cytokine Res.* 39 (2018), <https://doi.org/10.1089/jir.2018.0019>.
- [9] K. Ozaki, R. Spolski, C.G. Feng, et al., A critical role for IL-21 in regulating immunoglobulin production, *Science* 298 (5598) (2002) 1630–1634, <https://doi.org/10.1126/science.1077002>.
- [10] M. Croce, V. Rigo, S. Ferrini, IL-21: a pleiotropic cytokine with potential applications in oncology, *J. Immunol. Res.* 2015 (2015), 696578, <https://doi.org/10.1155/2015/696578>.
- [11] S. Kim-Schulze, H.S. Kim, Q. Fan, D.W. Kim, H.L. Kaufman, Local IL-21 promotes the therapeutic activity of effector T cells by decreasing regulatory T cells within the tumor microenvironment, *Mol. Ther.* 17 (2) (2009) 380–388, <https://doi.org/10.1038/mt.2008.249>.
- [12] S.N. Li, W. Wang, S.P. Fu, et al., IL-21 modulates release of proinflammatory cytokines in LPS-stimulated macrophages through distinct signaling pathways, *Mediators Inflamm.* 2013 (2013) 548073, <https://doi.org/10.1155/2013/548073>.
- [13] R. Zeng, R. Spolski, S.E. Finkelstein, et al., Synergy of IL-21 and IL-15 in regulating CD8+ T cell expansion and function, *J. Exp. Med.* 201 (1) (2005) 139–148, <https://doi.org/10.1084/jem.20041057>.
- [14] Tsunao Kishida, Hidetsugu Asada, Yoshiki Itokawa, Feng-De Cui, Masaharu Shin-Ya, Satoshi Gojo, Kakei Yasutomi, Yuji Ueda, Hisakazu Yamagishi, Jiro Imanishi, Osam Mazda, Interleukin [IL]-21 and IL-15 genetic transfer synergistically augments therapeutic antitumor immunity and promotes regression of metastatic lymphoma, *Mol. Ther.* 8 (2003) 552–558, [https://doi.org/10.1016/S1525-0016\[03\]00222-3](https://doi.org/10.1016/S1525-0016[03]00222-3).



Stephen W. Wilz, MD is Associate Chief of Pathology and Laboratory Medicine for the Boston VA Healthcare System, and Assistant Professor of Medicine at Boston University School of Medicine. Dr. Wilz's research includes developing simple screening tests for common epithelial cancers, such as lung, colon, breast and bladder cancer, using the modalities of mass spectroscopy and flow cytometry. He recently published a paper that identifies the presence of bladder cancer in the urine, with high sensitivity and specificity, by categorizing proteins present in cells in the urine sediment by their sub-cellular localization. In addition to his Doctor of Medicine degree from Tufts University, he obtained a Master's degree from Harvard University, where he contributed to the development of a Molecular Dynamics computer program for analysis of large biomolecules. While undertaking his residency at Massachusetts General Hospital, he also studied the T cell immune response to pyelonephritis in mice. He is a fellow of the College of American Pathologists.