



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



JAMDA

journal homepage: [www.jamda.com](http://www.jamda.com)

## Original Study

# Clinical Suspicion of COVID-19 in Nursing Home Residents: Symptoms and Mortality Risk Factors



Jeanine J.S. Rutten MD, MSc\*, Anouk M. van Loon PhD, Janine van Kooten MD, PhD, Laura W. van Buul PhD, Karlijn J. Joling PhD, Martin Smalbrugge MD, PhD, Cees M.P.M. Hertogh MD

Department of Medicine for Older People, Amsterdam Public Health Research Institute, Amsterdam University Medical Center, Amsterdam, the Netherlands

## A B S T R A C T

**Keywords:**

Long-term care facility  
SARS-CoV-2  
cohort  
survival analyses

**Objectives:** To describe the symptomatology, mortality, and risk factors for mortality in a large group of Dutch nursing home (NH) residents with clinically suspected COVID-19 who were tested with a reverse transcription–polymerase chain reaction (RT-PCR) test.

**Design:** Prospective cohort study.

**Setting and Participants:** Residents of Dutch NHs with clinically suspected COVID-19 and who received RT-PCR test.

**Methods:** We collected data of NH residents with clinically suspected COVID-19 via electronic health records between March 18 and May 13, 2020. Registration was performed on diagnostic status [confirmed (COVID-19+)/ruled out (COVID-19–)] and symptomatology (typical and atypical symptoms). Information on mortality and risk factors for mortality were extracted from usual care data.

**Results:** In our sample of residents with clinically suspected COVID-19 (N = 4007), COVID-19 was confirmed in 1538 residents (38%). Although symptomatology overlapped between residents with COVID-19+ and COVID-19–, those with COVID-19+ were 3 times more likely to die within 30 days [hazard ratio (HR), 3.1, 95% confidence interval (CI) 2.7–3.6]. Within this group, mortality was higher for men than for women (HR 1.8, 95% CI 1.5–2.2), and we observed a higher mortality for residents with dementia, reduced kidney function, and Parkinson's disease, even when corrected for age, gender, and comorbidities.

**Conclusions and Implications:** About 40% of the residents with clinically suspected COVID-19 actually had COVID-19, based on the RT-PCR test. Despite an overlap in symptomatology, mortality rate was 3 times higher for residents with COVID-19+. This emphasizes the importance of using low-threshold testing in NH residents, which is an essential prerequisite to using limited personal protective equipment and isolation measures efficiently.

© 2020 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

A few weeks after the first reported coronavirus case in the Netherlands on February 27,<sup>1</sup> reports emerged of COVID-19 cases in nursing homes (NHs). Because of a lack of reverse transcription–polymerase chain reaction (RT-PCR) test facilities and the limited available personal protective equipment, isolation

measures were used in NH residents with clinically suspected but not necessarily confirmed COVID-19. By April, one-third of the Dutch NHs reported at least 1 resident with COVID-19,<sup>2</sup> and the number of deaths in NHs doubled during the COVID-19 pandemic.<sup>3</sup>

It is evident that severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) acts differently in older adults compared with younger adults.<sup>4</sup> First, older adults have been reported to develop signs and symptoms more often, after being in contact with SARS-CoV-2.<sup>4,5</sup> Next, there are also indications for atypical disease presentation in older adults with symptoms such as delirium and falls,<sup>6–8</sup> and even for asymptomatic presentation.<sup>9</sup> In addition, shortness of breath and a high respiratory rate are reported to be more common in older adults.<sup>5</sup> Indisputably, older age is associated with a higher mortality.<sup>4,5,10–12</sup>

This study was funded by the Dutch Ministry of Health, Welfare and Sport. The authors declare no conflicts of interest.

J.J.S. Rutten and A. M. van Loon contributed equally to the study.

\* Address correspondence to Jeanine J. S. Rutten, MD, MSc, Department of Medicine for Older People, Amsterdam UMC, van der Boerhorststraat 7, 1081 BT, Amsterdam, the Netherlands.

E-mail address: [j.rutten1@amsterdamumc.nl](mailto:j.rutten1@amsterdamumc.nl) (J.J.S. Rutten).

<https://doi.org/10.1016/j.jamda.2020.10.034>

1525-8610/© 2020 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

However, accurate knowledge is lacking about the spectrum of symptomatology of COVID-19, mortality, and risk factors for mortality in NH residents. Furthermore, it is unknown to what extent COVID-19 can be distinguished based on symptomatology from other infections (such as influenza and other viral and bacterial infections) in NH residents, and whether an RT-PCR test is always necessary to make the distinction.

From March 18 onwards, we set-up a COVID-19 registration, via an electronic health record (EHR) used by more than half of the NHs in the Netherlands. Combining data from this registration with usual care data extracted from the same EHR, we conducted a cohort study to

- describe symptomatology and analyze mortality in residents with a clinical suspicion of COVID-19 and compare within this group the residents with confirmed COVID-19 to those where COVID-19 was ruled out as assessed by an RT-PCR test; and
- analyze risk factors for mortality in residents with confirmed COVID-19.

## Methods

### Study Design and Setting

We conducted a prospective cohort study in Dutch NHs using a COVID-19 registration form linked to the EHR Ysis. Ysis, managed by software developer GeriMedica, is the most widely used EHR in NHs in the Netherlands and provides pseudonymized data on more than 61,000 of a total of 115,000 residents.<sup>13</sup>

### Selection of Study Population

We included residents with a clinical suspicion of COVID-19 based on the physician's assessment and for whom we had the results of the RT-PCR test. We excluded residents if results of follow-up diagnostics were not (yet) available. The Dutch Association of Elderly Care Physicians (Verenso) created a guideline that describes when to consider COVID-19 and when to perform an RT-PCR test.<sup>14</sup> Because of scarcity, until 26 March it was advised to use a PCR test only in NH residents with at least 2 of the following symptoms: fever or feverish feeling, coughing, and shortness of breath. In addition, it was advised only to test at wards without proven cases of COVID-19. Other NH residents with clinically suspected COVID-19 were treated as patients with COVID-19 upfront. From 26 March onwards, it was advised to perform an RT-PCR test in NH residents with any possibly COVID-19 related symptoms. From the first week of April, it was advised to test all NH residents with clinically suspected COVID-19, also at wards with proven cases of COVID-19.

### Data Collection

The study started on March 18, 2020, and is ongoing. For this article, we analyzed data up to May 13. If "Covid" or "corona" were registered in the EHR Ysis, a standardized assessment form appeared automatically on the EHR for the physician to complete. This standardized assessment form contains questions about the presence of "typical symptoms" (fever, coughing, and shortness of breath), "atypical symptoms" (sore throat and delirium, confusion, or drowsiness), other symptoms, oxygen saturation (normal or decreased), and body temperature. All the symptoms were determined and registered by the physician(s) caring for the resident.

For all subsequent entries in the EHR for participants included in our cohort, a follow-up form was presented. The follow-up form contained a categorical question on follow-up diagnostics (COVID-19 confirmed, COVID-19 ruled out, no test outcome yet available).

Age, gender, type of ward, mortality, date of death, and comorbidity were derived from usual care data in the EHR. Data on comorbidity was registered by the physician(s) in open text fields in the medical history. We selected comorbidities that were considered risk factors for severe illness or mortality with COVID-19 in the general population by the Dutch National Institute for Public Health and the Environment at that moment. These comorbidities—dementia (including other cognitive disorders), chronic respiratory diseases, chronic cardiovascular disease, cerebrovascular diseases, diabetes mellitus, reduced kidney function, and Parkinson's disease (PD)—were extracted from the usual care data by an extensive search using MATLAB (The MathWorks, Natick, MA). Search entries for each comorbidity were formed by 3 medical doctors (M.S., J.v.K., and J.R.), taking into account multiple variations in the way of registration. For example, the search entry for "reduced kidney function" contained the following keywords: *impaired renal/kidney function, renal/kidney insufficiency, renal/kidney failure, dialysis, nephropathy*, etc. The search entries for each comorbidity were programmed in MATLAB. MATLAB then automatically searched the open text fields in which the physicians registered the medical history. For the first 200 residents, the automated search was compared to a manual coding, and the programmed coding scheme was adjusted accordingly.

The usual care data and data from the standardized assessment form and follow-up form were linked and pseudonymized by GeriMedica (the software development company that developed and manages the EHR).

### Ethics

The Medical Ethics Committee of the Vrije Universiteit Medical Centre in Amsterdam reviewed and approved the study protocol.

### Statistical Analysis

Symptomatology of residents with a clinical suspicion of COVID-19 was analyzed descriptively and separate for residents with COVID-19 confirmed (COVID-19+) and residents in whom COVID-19 was ruled out (COVID-19−) as assessed with a RT-PCR test. We used *t* tests for continuous variables and chi-square tests for categorical variables to compare demographic characteristics and symptomatology. Survival curves on 30-day mortality were estimated based on the days between the date of clinically suspected COVID-19 and the date of death using Kaplan-Meier curves in both residents with COVID-19+ and COVID-19−. The mortality rates in residents with COVID-19+ and COVID-19− were compared using a Cox proportional hazards model. The association between demographic characteristics, clinical characteristics, and mortality rate was analyzed in residents with COVID-19+ using 3 models: model 1 unadjusted, model 2 included gender and age, model 3 included gender, age, and comorbidities. All model covariates were selected a priori on the basis of clinical relevance or results of the unadjusted model 1. Results are presented with 95% confidence intervals (CIs) and all reported *P* values are 2-sided. To indicate the number of missing values, the *N* on which percentages were calculated are reported. All analyses were performed with the use of the SPSS statistical package, version 20.0 (IBM Corp, Armonk, NY).

## Results

Between March 18 and May 13, 2020, a clinical suspicion of COVID-19 was registered for 5425 NH residents (see [Supplementary Figure 1](#)). Because of the lack of (results of) follow-up diagnostics, 1418 residents were excluded. Follow-up diagnostics confirmed COVID-19 in 1538 of the 4007 residents (38%); COVID-19 was ruled out in 2469 (62%) residents.

**Table 1**  
Resident Characteristics

	COVID-19 Confirmed (n = 1538)			COVID-19 Ruled Out (n = 2469)	Total Study Population (N = 4007)		
	Men (n = 554)	Woman (n = 984)	P Value (Men vs Woman)	Total (n = 1538)	Total	P Value (Confirmed vs Ruled Out)	Total
Age, y mean (SD)	82 (8.8)	85 (8.4)	<.001	84 (8.7)	83 (10.4)	.02	84 (9.8)
Women	—	—	—	64	61	.07	62
Ward type							
Psychogeriatric	42	50	.003	47	39	<.001	42
Somatic	18	19	.81	19	21	.12	20
Rehabilitation or short-term care	18	16	.35	16	17	.48	17
Other	1	1	.46	1	2	.25	2
Unknown	21	14	.001	17	21	.001	19
Comorbidity	(n = 547*)	(n = 978*)		(n = 1525*)	(n = 2436*)		(n = 3961*)
Dementia	58	64	.02	62	51	<.001	59
Cardiovascular disease	56	47	<.001	50	49	.28	53
Cerebrovascular disease	47	38	.001	41	41	.98	44
Diabetes Mellitus	25	27	.53	26	25	.28	27
Chronic respiratory disease	20	16	.09	18	21	.02	21
Reduced kidney function	19	18	.61	18	19	.66	20
Parkinson disease	9	5	.001	6	7	.53	7

Unless otherwise noted, values are percentages.

\*Total number of residents with data on comorbidity.

*Resident Characteristics*

Residents with a clinical suspicion of COVID-19 had a mean age of 84 years (standard deviation 9.8), were mostly women (62%) and resided mostly on psychogeriatric wards (42%) (Table 1). Residents with COVID-19+ were more likely to reside in a psychogeriatric ward (47% vs 39%) and to have dementia (62% vs 51%) and were less likely to have chronic respiratory disease (21% vs 18%).

Except for these differences, no important clinical differences were found between the characteristics of residents with COVID-19+ and COVID-19-.

Of the 1538 COVID-19+ residents, 554 were men (36%). Compared with women, men were younger (82 vs 85 years) and more often had cardiovascular disease (56% vs 47%), cerebrovascular disease (47% vs 38%), and PD (9% vs 5%). Women with COVID-19+ had dementia (64% vs 58%) more often than men.

**Table 2**  
Symptomatology of NH Residents With a Clinical Suspicion of COVID-19

	COVID-19 Confirmed	COVID-19 Ruled Out	
	Total (n = 1538)	P value (Confirmed vs Ruled Out)	Total (n = 2469)
Typical symptoms			
Cough	63 (900/1431)	.75	62 (1440/2309)
Shortness of breath	30 (417/1373)	<.001	39 (882/2249)
Fever	63 (917/1455)	<.001	42 (969/2316)
Atypical symptoms			
Sore throat	10 (94/976)	.007	13 (233/1699)
Delirium, confusion, or drowsiness	29 (372/1288)	.16	27 (570/2138)
Other symptoms*			
Fatigue	22 (93/417)	<.001	13 (112/869)
Diarrhea	18 (74/417)	.43	16 (139/869)
Malaise	18 (73/417)	.004	12 (101/869)
Rhinorrhea	12 (52/417)	.23	15 (130/869)
Nausea or vomiting	12 (48/417)	.001	19 (163/869)
Common cold	12 (52/417)	.58	14 (118/869)
Signs			
Decreased oxygen saturation	44 (453/1023)	.19	47 (820/1751)
Temperature			
<36.5 °C	4 (54/1213)	<.001	12 (220/1883)
36.5–37.5 °C	17 (204/1213)	<.001	33 (613/1883)
37.5–38.0 °C	17 (201/1213)	.64	16 (310/1883)
>38.0 °C	62 (754/1213)	<.001	39 (740/1883)

Values are % (n/group n).

\*Not assessed via categorical questions, but reported in an open text field.

## Symptomatology

### Typical symptoms

Most residents with COVID-19+ had 1 (42%) or 2 (35%) typical symptoms (ie, cough, shortness of breath, or fever); 9% did not have any typical symptoms.

Of the residents with COVID-19+, 63% presented with cough and 30% suffered from shortness of breath (Table 2). For residents in whom COVID-19 was ruled out, this was 62% and 39%, respectively. Fever was more common in residents with COVID-19+ compared to residents where COVID-19 was ruled out (63% vs 42%), whereas shortness of breath and sore throat were reported more frequently in residents with COVID-19– (39% vs 30% and 13% vs 10%).

### Atypical and other symptoms

The risk of delirium, confusion, or drowsiness and sore throat in residents with COVID-19+ was comparable to the incidence of these symptoms in residents with COVID-19– (29% and 10% vs 27% and 13%) (Table 2). Other symptoms, which could be outlined in an open text field, were reported for 417 (27%) residents with COVID-19+ and for 869 (35%) of residents with COVID-19–. Fatigue was most commonly reported in residents with COVID-19+ (22%), followed by diarrhea (18%), malaise (18%), rhinorrhea (12%), nausea or vomiting (12%), and common cold (12%) (Table 2). Other symptoms such as headache and anosmia were reported in less than 10% of cases. Fatigue and malaise were reported

more often in residents with COVID-19+ than in residents with COVID-19– (22% vs 13% and 18% vs 12%), whereas nausea or vomiting was reported less often in residents with COVID-19+ (12% vs 19%).

### Signs


Of the residents with COVID-19+, 44% had decreased oxygen saturation, vs 47% of the residents with COVID-19–. Of the residents with COVID-19+, most had a temperature of  $>38$  °C (62%) compared to the residents with COVID-19– (39%).

### Mortality in NH Residents With a Clinical Suspicion of COVID-19

The median number of days on which NH residents with COVID-19 died was 22 days after the day of the COVID-19 suspicion [interquartile range (IQR) = 21] vs 28 (IQR = 15) for COVID-19. Of the residents with COVID-19+, 42% had died in 30 days (95% CI 39%–44%), vs 15% of the residents with COVID-19– (95% CI 14%–17%) (see Figure 1A). Residents with COVID-19+ were 3 times more likely to die within 30 days than residents with COVID-19– [adjusted hazard ratio (HR) 3.14, 95% CI 2.7–3.6;  $P < .001$ ].

### Risk Factors for Mortality in Residents With Confirmed COVID-19

Gender, age, and comorbidity were associated with a higher mortality in residents with COVID-19+ (Table 3).



**Fig. 1.** Kaplan-Meier estimates of survival: (A) NH residents with confirmed and ruled out COVID-19. The blue line indicates the 30-day survival probability of NH residents in whom COVID-19 was ruled out; the red line indicates the 30-day survival probability of NH residents with confirmed COVID-19. The triangles indicate censored data points. (B) Men and women with confirmed COVID-19. The dashed line indicates the 30-day survival probability of women with confirmed COVID-19; the solid line indicates the 30-day survival probability of men with confirmed COVID-19. The triangles indicate censored data points.

**Table 3**  
Cox Proportional Hazard Model for Risk Factors for Mortality in Residents With Confirmed COVID-19

	Model 1: unadjusted		Model 2: adjusted for gender and age		Model 3: Adjusted for Gender, Age, and Relevant Comorbidity*	
	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value
Gender						
Women	1.00 (ref)		1.00 (ref)		1.00 (ref)	
Men	1.70 (1.45-2.00) <sup>†</sup>	<.001 <sup>†</sup>	1.84 (1.56-2.17) <sup>†</sup>	<.001 <sup>†</sup>	1.82 (1.54-2.15) <sup>†</sup>	<.001 <sup>†</sup>
Age, y						
≤80	1.00 (ref)		1.00 (ref)		1.00 (ref)	
81-85	1.08 (0.85-1.37)	.55	1.12 (0.88-1.42)	.37	1.07 (0.84-1.37)	.57
86-90	1.37 (1.11-1.69) <sup>†</sup>	.004 <sup>†</sup>	1.56 (1.26-1.94) <sup>†</sup>	<.001 <sup>†</sup>	1.49 (1.19-1.85) <sup>†</sup>	<.001 <sup>†</sup>
>90	1.20 (0.96-1.51)	.11	1.41 (1.12-1.78) <sup>†</sup>	.004 <sup>†</sup>	1.38 (1.09-1.75) <sup>†</sup>	.008 <sup>†</sup>
Comorbidity						
Dementia	1.23 (1.04-1.46) <sup>†</sup>	.02 <sup>†</sup>	1.25 (1.05-1.48) <sup>†</sup>	.01 <sup>†</sup>	1.26 (1.06-1.50) <sup>†</sup>	.008 <sup>†</sup>
Chronic cardiovascular disease	1.22 (1.04-1.43) <sup>†</sup>	.02 <sup>†</sup>	1.15 (0.98-1.36)	.08	1.15 (0.97-1.35)	.10
Cerebrovascular disease	1.10 (0.94-1.29)	.24	1.08 (0.92-1.27)	.33	1.06 (0.89-1.25)	.53
Diabetes mellitus	0.99 (0.83-1.19)	.91	1.03 (0.86-1.23)	.79	0.99 (0.82-1.19)	.90
Chronic respiratory disease	0.97 (0.79-1.20)	.79	0.94 (0.76-1.16)	.55	0.92 (0.75-1.14)	.46
Reduced kidney function	1.39 (1.15-1.69) <sup>†</sup>	.001 <sup>†</sup>	1.36 (1.13-1.65) <sup>†</sup>	.002 <sup>†</sup>	1.35 (1.11-1.64) <sup>†</sup>	.002 <sup>†</sup>
Parkinson's disease	1.44 (1.08-1.91) <sup>†</sup>	.01 <sup>†</sup>	1.40 (1.05-1.88) <sup>†</sup>	.02 <sup>†</sup>	1.49 (1.11-2.00) <sup>†</sup>	.007 <sup>†</sup>

\*Relevant comorbidity includes dementia, chronic cardiovascular disease, reduced kidney function and Parkinson's disease.

<sup>†</sup>Statistically significant ( $P < .05$ ).

**Gender differences**

In comparison with women, men were more likely to have experienced severe symptoms such as shortness of breath (35% vs 28%); fever (68% vs 60%); delirium, confusion, or drowsiness (35% vs 25%); and decreased oxygen saturation (49% vs 42%) in the COVID-19+ group (Table 4).

Of men with COVID-19+, 52% had died within 30 days (95% CI 48%-56%) vs 36% of women with COVID-19+ (95% CI 33%-39%). The median number of days, after the day of the COVID-19 suspicion, on which male NH residents with COVID-19 died was 16 days (IQR = 23), vs 25 days (IQR = 9) for women with COVID-19. We found an HR of 1.8 (95% CI 1.6-2.2;  $P < .001$ ) when adjusted for age and comorbidities (see Figure 1B).

**Comorbidities**

Dementia (HR 1.26, 95% CI 1.06-1.50;  $P = .008$ ), reduced kidney function (HR 1.35, 95% CI 1.11-1.64;  $P = .002$ ), and PD (HR 1.49, 95% CI

1.11-2.00;  $P = .007$ ) were associated with a higher mortality rate in residents with COVID-19+ when adjusted for age, gender, and comorbidities. For other comorbidities, no statistically significant associations were found.

**Discussion**

COVID-19 was only confirmed in 1538 (38%) of the 4007 residents with a clinically suspected COVID-19. Many NH residents with confirmed COVID-19 had “typical symptom(s)” such as fever, cough, or shortness of breath. Atypical symptoms were also reported regularly (ie delirium, confusion, or drowsiness; fatigue; diarrhea; malaise; rhinorrhea; nausea or vomiting; and common cold). We observed an overlap in symptoms between residents with and without confirmed COVID-19 following clinically suspected COVID-19. However, fever occurred more frequently in the COVID-19 confirmed group (63% vs 42%). Furthermore, the mortality rate in residents with confirmed

**Table 4**  
Symptomatology of Men and Women With a Clinical Suspicion of COVID-19

	COVID-19 Confirmed		
	Men (n = 554)	Woman (n = 984)	P value (Men vs Woman)
Typical symptoms			
Cough	62 (321/515)	63 (579/916)	.74
Shortness of breath	35 (174/498)	28 (243/875)	.005
Fever	68 (361/533)	60 (556/922)	.005
Atypical symptoms			
Sore throat	8 (26/339)	11 (68/637)	.13
Delirium, confusion, or drowsiness	35 (170/482)	25 (202/806)	<.001
Other symptoms*			
Fatigue	19 (26/136)	24 (67/281)	.28
Diarrhea	18 (24/136)	18 (50/281)	.97
Malaise	17 (23/136)	18 (50/281)	.82
Rhinorrhea	11 (15/136)	13 (37/281)	.54
Nausea or vomiting	8 (11/136)	13 (37/281)	.13
Common cold	14 (19/136)	12 (33/281)	.52
Signs			
Decreased oxygen saturation	49 (187/383)	42 (266/640)	.02
Temperature			
<36.5 °C	4 (17/450)	5 (37/763)	.48
36.5-37.5 °C	14 (64/450)	18 (140/763)	.14
37.5-38.0 °C	14 (64/450)	18 (137/763)	.19
>38.0 °C	68 (305/450)	59 (449/763)	<.001

Values are % (n/group n).

\*Not assessed via categorical questions, but reported in an open text field.



COVID-19 was 3 times higher than in residents where COVID-19 was ruled out. For residents with confirmed COVID-19, the 30-day mortality in men was higher than in women. In addition, dementia, reduced kidney function, and PD were all associated with a higher mortality rate in residents with confirmed COVID-19, even when corrected for age, gender, and comorbidities.

In our sample, 63% of the NH residents with confirmed COVID-19 presented with fever, 63% with cough, and 30% with shortness of breath. This is in line with a previous study that showed that fever (75%) was most common in older adults, followed by cough (44%) and shortness of breath (25%).<sup>11</sup> In contrast, another study in older people showed that cough (80%) was more common than fever (60%).<sup>15</sup> However, the sample sizes of the age groups (>80 years) was much smaller in both studies (n = 16 and n = 20). Our study showed that COVID-19 in NH residents presents itself with both typical and atypical symptoms.

In our study, almost half of the residents with confirmed COVID-19 (42%) died within 30 days. This percentage is higher than the 34% reported by McMichael et al,<sup>16</sup> and the 26% reported by Arons et al.<sup>9</sup> A possible explanation could be that in our sample, in part because of the RT-PCR test shortage in the early weeks of patient enrollment, only residents with symptoms were tested, whereas in the above-mentioned studies residents were also tested in the absence of symptoms. In comparison with asymptomatic presentation of COVID-19, symptomatic presentation will result in a higher mortality. Because of the high mortality in this vulnerable group, it is necessary to protect NH residents from COVID-19, especially now that the flu season has arrived and the virus may spread even faster.<sup>17</sup>

We observed that in residents with confirmed COVID-19, men were more likely to die than women. This difference in mortality has also been found in other studies on COVID-19,<sup>10,18</sup> and preliminary data indicate an association between comorbidities, such as cardiovascular disease, and severity of COVID-19.<sup>5,19</sup> However, we adjusted our survival analyses for comorbidities, yet the difference in mortality between men and women remained. It has been suggested that this gender difference might be related to gender differences in SARS-CoV-2 entry receptors.<sup>20</sup> Nonetheless, it remains unclear whether increased mortality for men is specific to COVID-19. In general, men are sick more often and more severely sick because of gender differences in hormones, immune responses and immune aging, and gender-specific differences in lifestyle and health behavior.<sup>21</sup> Therefore, the observed difference in COVID-19 mortality between men and woman may be explained by the men-women disability survival paradox.<sup>22</sup> This is the phenomenon that higher life expectancy is accompanied by higher rates of disability in women compared with men, but disability increases risk of mortality more in men than women. Therefore, men might be less resilient to acute infectious diseases as COVID-19.

Furthermore, within our NH residents with confirmed COVID-19, we observed a higher mortality in those with dementia, reduced kidney function, and PD than those without these comorbidities, even when corrected for age, gender, and comorbidities. Similarly, Williamson et al showed that dementia/stroke and kidney diseases are risk factors.<sup>23</sup> In contrast to our study, they also found an association between diabetes mellitus, cardiovascular, neurological, and respiratory diseases, and mortality. Prior to our study, PD had been suggested as a comorbid risk factor for COVID-19 mortality, but empirical data were lacking.<sup>24</sup> We show, for the first time in a large study population, that PD increased mortality risk of COVID-19 in NH residents. One possibility for this is an increased chance of death arising from pneumonia in patients with PD.<sup>24,25</sup> although further data are required to test this possibility.

This study had several limitations. Based on the Verenso guideline, we assumed that follow-up diagnostics consists of an RT-PCR test. A RT-PCR test has relatively low sensitivity (63%–78%).<sup>26</sup> Consequently, it

could be that the group of residents in whom COVID-19 was ruled out partly consisted of residents who actually had COVID-19. Moreover, the group of residents in whom COVID-19 was ruled out did have symptoms, which also may have caused by other underlying infections such as influenza or other viruses. It would be useful to conduct an additional case-control study in which a cohort of residents with confirmed covid-19 would be compared with a pre-COVID-19 era cohort to compare the mortality outcomes.

Second, our study was set up in the very early stages of the COVID-19 outbreak in the Netherlands (March 18, 2020) and our standardized assessment form was created with available knowledge at the time. As a result, not all relevant symptoms were listed in the standard form. However, we included a question about delirium, confusion, or drowsiness in the standardized form, because we hypothesized that NH residents could have atypical presentations. Moreover, physicians had the opportunity to report other symptoms in an open text field and did so for 32% of the participants in this study. Of note, because of the shortage of RT-PCR tests in the beginning of the epidemic in the Netherlands, initially only residents who met the criteria as described in the Verenso guidelines (ie residents with fever, cough, or shortness of breath) were tested. As a result, we were not able to describe the occurrence of an atypical or asymptomatic presentation of COVID-19 properly. A study in US NH residents found that almost half of residents with COVID-19 were asymptomatic at the time of testing, but developed symptoms later.<sup>9</sup>

## Conclusions and Implications

In a large cohort of clinically suspected COVID-19 NH residents, 38% of the residents with clinically suspected COVID-19 actually had confirmed COVID-19. There was an overlap in symptoms between residents with confirmed COVID-19 and those for whom COVID-19 was ruled out, but we found that mortality rate in residents with confirmed COVID-19 was 3 times higher. This emphasizes the importance of using a low threshold to test NH residents for SARS-CoV-2. This is of vital importance because detecting positive COVID-19 diagnoses in NHs is extremely important, and an essential prerequisite to using limited personal protective equipment and quarantine and isolation measures efficiently. Furthermore, our data demonstrated that COVID-19 affects male residents disproportionately, and mortality was almost twice as high for men than for women. It has been suggested that gender differences in comorbidity could explain this effect. However, our data demonstrate that the increased mortality risk for men remained even when corrected for age and comorbidities.

## Acknowledgments


We thank the facility residents, the physicians who work in the long-term care facilities for their time to fill in the COVID-19 registration and their ongoing efforts to provide care in the face of these challenges, Prof Dr Jos Twisk and Dr Birgit Lissenberg-Witte for their help with the survival analyses, Dr James Kirkbride for proofreading our manuscript, and the staff of Gerimedita for their support.

## References

1. RIVM. Patient with new coronavirus in the Netherlands; 2020. Available at: <https://www.rivm.nl/nieuws/patient-met-nieuw-coronavirus-in-nederland>. Accessed June 15, 2020.
2. Kruse F, Abma I, Jeurissen P. The Impact of COVID-19 on Long-Term Care in the Netherlands. LTCcovid, International Long-Term Care Policy Network. CPEC-LSE; 2020.
3. RIVM. Excess mortality caused by novel coronavirus (Covid-19); 2020. Available at: <https://www.rivm.nl/node/155011>. Accessed November 23, 2020.
4. Liu K, Chen Y, Lin R, et al. Clinical features of COVID-19 in elderly patients: A comparison with young and middle-aged patients. *J Infect* 2020;80:e14–e18.

5. Chen T, Dai Z, Mo P, et al. Clinical characteristics and outcomes of older patients with coronavirus disease 2019 (COVID-19) in Wuhan, China (2019): A single-centered, retrospective study. *J Gerontol A Biol Sci Med Sci* 2020;75: 1788–1795.
6. Cesari M, Montero-Odasso M. COVID-19 and older adults. Lessons learned from the Italian epicenter. *Can Geriatr J* 2020;23:155–159.
7. Olde Rikkert MGM, Vingerhoets RW, van Geldorp N, et al. Atypical clinical picture of COVID-19 in older patients [in Dutch]. *Ned Tijdschr Geneesk* 2020; 164:D5004.
8. Tay HS, Harwood R. Atypical presentation of COVID-19 in a frail older person. *Age Ageing* 2020;49:523–524.
9. Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *N Engl J Med* 2020;382: 2081–2090.
10. Li LQ, Huang T, Wang YQ, et al. COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *J Med Virol* 2020;92: 577–583.
11. Niu S, Tian S, Lou J, et al. Clinical characteristics of older patients infected with COVID-19: A descriptive study. *Arch Gerontol Geriatr* 2020;89:104058.
12. Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: A model-based analysis. *Lancet Infect Dis* 2020;20: 669–677.
13. CBS. Number of residents of care and nursing homes. Available at: <https://www.cbs.nl/nl-nl/maatwerk/2020/13/aantal-bewoners-van-verzorgings-en-verpleeghuizen-2019>; 2019. Accessed June 15, 2020.
14. Verenso. Treatment advice acute phase and postcare COVID-19. Available at: <https://www.verenso.nl/themas-en-projecten/infectieziekten/covid-19-coronavirus/behandeladvies-acute-fase-en-nazorg>; 2020. Accessed June 15, 2020.
15. Guo T, Shen Q, Guo W, et al. Clinical characteristics of elderly patients with COVID-19 in Hunan Province, China: A multicenter, retrospective study. *Gerontology* 2020;66:465–475.
16. McMichael TM, Currie DW, Clark S, et al. Epidemiology of COVID-19 in a long-term care facility in King County, Washington. *N Engl J Med* 2020;382:2005–2011.
17. de Celles MD, Casalegno JS, Lina B, et al. Influenza may facilitate the spread of SARS-CoV-2. medRxiv; 2020.
18. Zhao S, Cao P, Chong MKC, et al. COVID-19 and gender-specific difference: Analysis of public surveillance data in Hong Kong and Shenzhen, China, from January 10 to February 15, 2020. *Infect Control Hosp Epidemiol* 2020;41:750–751.
19. Li M, Dong Y, Wang H, et al. Cardiovascular disease potentially contributes to the progression and poor prognosis of COVID-19. *Nutr Metab Cardiovasc Dis* 2020;30:1061–1067.
20. Conti P, Younes A. Coronavirus COV-19/SARS-CoV-2 affects women less than men: Clinical response to viral infection. *J Biol Regul Homeost Agents* 2020;34:339–343.
21. Gebhard C, Regitz-Zagrosek V, Neuhauser HK, et al. Impact of sex and gender on COVID-19 outcomes in Europe. *Biol Sex Differ* 2020;11:29.
22. Kingston A, Davies K, Collerton J, et al. The contribution of diseases to the male-female disability-survival paradox in the very old: Results from the Newcastle 85+ study. *PLoS One* 2014;9:e88016.
23. Williamson E, Walker AJ, Bhaskaran KJ, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;584:430–436.
24. Helmich RC, Bloem BR. The impact of the COVID-19 pandemic on Parkinson's disease: Hidden sorrows and emerging opportunities. *J Parkinsons Dis* 2020; 10:351–354.
25. Beyer MK, Herlofson K, Arslan D, et al. Causes of death in a community-based study of Parkinson's disease. *Acta Neurol Scand* 2001;103:7–11.
26. Zitek T. The appropriate use of testing for COVID-19. *West J Emerg Med* 2020; 21:470–472.



**Appendix****Supplementary Figure 1.** Flow chart.